

Q101  
o2564  
1903-0  
V.36

THE  
JOURNAL  
OF  
THE LINNEAN SOCIETY.

---

BOTANY.

---

VOL. XXXVI.

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.,

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1903-1905.

11. Bot. Garden  
1905

Dates of Publication of the several Numbers included in this Volume.

No. 249, pp. 1- 72, published January 1, 1903.

„ 250, „ 73-136, „ April 1, „

„ 251, „ 137-216, „ June 30, „

„ 252, „ 217-296, „ October 31, „

„ 253, „ 297-376, „ March 31, 1904.

„ 254, „ 377-456, „ July 25, „

Nos. 255-256, 457-686,

Index to Names and Synonyms, Vols. **23**, **26**, and **36**,  
Titlepage, Contents, Historical Note, etc.

published December 20, 1905.

PRINTED BY TAYLOR AND FRANCIS,

RED LION COURT, FLEET STREET.

## CONTENTS.

---

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.— (Hydrocharidaceæ—Gramineæ) . . . . .	1-449
List of the Genera and Species discovered in China since the publication of the various parts of the "Enumeration," from 1886 to March 1904, alphabetically arranged. . . . .	451-530
Index to Names and Synonyms, Vols. 23, 26, and 36 . . . . .	531-686

Titlepage, Contents, Historical Note, etc.

## HISTORICAL NOTE.

---

THE completion of an undertaking which has been on hand for some twenty years, has far exceeded the limits originally assigned to it, and must, I fear, have long ago exhausted the patience of the Linnean Society, invites, if it does not almost demand, a few words of explanation as to its history.

As long ago as 1878 I was invited to deliver before the Royal Geographical Society a lecture which was in substance an attempt to review the knowledge existing at the time of the Earth's flora. When I came to the vast territory occupied in the Old World by the Chinese Empire, I could only quote the statement made four years earlier by the well-known botanist, the late Dr. Hance:—

“While M. Maximowicz's excellent and very complete ‘*Index Floræ Pekinensis*’ provides a good catalogue of the flora of the Chinese metropolis and its vicinity, and Mr. Bentham's classical ‘*Flora Hongkongensis*’ has acquainted us with the principal constituents of that of the extreme South-east of the Empire, nothing whatever of a scientific character has yet, to my knowledge, been written on the vegetation of the districts intermediate to those two points, which are separated by 17° of latitude, or of the various ports of trade along the coast or on the Yangtse.”

It seemed to me that a beginning might at any rate be made to remedy this conspicuous defect in our knowledge of the vegetation of the Old World, and that a list of Chinese plants which had actually been collected would throw some light on the character of the Chinese Flora and would afford a starting-point for fresh research.

I accordingly in December 1883 made the following appeal to the Government Grant Committee of the Royal Society:—

“To ask for appointment of a Committee to report on our present knowledge of the Flora of China. It is believed that

the national herbaria contain a considerable accumulation of material, which it is desirable should be catalogued after the manner of the Botany of Godman and Salvin's 'Biologia Centrali-Americana.' Such a catalogue would embody descriptions of all undescribed species of which material is available, and references to the widely scattered published notices of Chinese plants would be intercalated in their proper place. The report would therefore give a complete view from all readily accessible sources of our present knowledge of the Chinese Flora. . . . Our present ignorance of the vegetation of China is an insuperable bar to any rational attempt at generalisation with regard to the distribution of the plants of the Palæarctic region."

The application was acceded to: a Committee was appointed consisting of Mr. J. Ball, Mr. Carruthers, Mr. Thiselton-Dyer, and Prof. Oliver, and a grant of £200 was placed at its disposal.

The first meeting of the Committee was held at the Royal Society on February 7, 1884. The following passage is extracted from the Minutes:—

"It was stated that Mr. F. B. Forbes, F.L.S., had as early as 1875, with the paid assistance of Mr. W. B. Hemsley and others, made considerable progress in cataloguing the Chinese plants preserved in the Herbaria of the Royal Botanic Gardens, Kew, and of the British Museum, and that he was actively engaged in arranging his material."

Mr. Forbes attended, by invitation, the next meeting of the Committee on February 14 following. He offered very liberally to cooperate with it, and it was agreed to employ Mr. Hemsley to carry on the work.

At subsequent meetings the form in which the Catalogue should be drawn up was settled, and in 1885 the Chairman, Mr. John Ball, addressed on its behalf the following letter to the President of the Linnean Society:—

10 Southwell Gardens, London, S.W.

13 May, 1885.

SIR,

A Committee consisting of Mr. Ball, Professor Oliver, Mr. Carruthers, and Mr. Thiselton-Dyer, was appointed last year by the Government Grant Committee of the Royal Society to draw up a Report on our present knowledge of the Flora of China.

The Committee has since been joined, at its invitation, by Mr. Forbes, F.L.S., who has most liberally offered his personal

assistance, and placed at its disposal his valuable and extensive manuscript collections and notes\*.

The Committee has further engaged the services of Mr. Hemsley, A.L.S., to cooperate in drawing up the Report, and he has prepared a specimen dealing with the Ranunculaceæ to its entire satisfaction.

The Committee now finds itself in a position to press on the work with considerable despatch, and feels confident that, by affording for the first time a comprehensive view of the vegetation of one of the most interesting of existing botanical regions, it will supply invaluable aid for further research, as well to the student of physiography, as to travellers, diplomatic agents and missionaries abroad.

The Committee is anxious that the printing should proceed *pari passu* with the preparation of the Report, and would deem it an advantage that it should be issued to the public under the auspices of the Linnean Society.

I, therefore, on behalf of the Committee, beg to offer the Report to the Council of the Linnean Society on the following conditions:—

1. The Committee to have placed at its disposal an entire volume of the botanical series of the Journal of the Society.

2. The Committee to pay the entire cost of setting the Report in type, and of correcting the press.

3. The Committee to be at liberty to print off at its own cost 150 copies to remain at its disposal.

4. The Linnean Society to bear the cost of press-work for copies issued to Fellows, and for stock for sale by the Society.

5. Plates illustrating species of exceptional interest may be included in the Report, at the discretion of the Committee, on the same terms as those above stated with regard to the letterpress †.

The Committee suggests that the Report may be issued in parts to the Fellows.

I have the honour to be, Sir,

Your obedient Servant,

(Signed) JOHN BALL.

*The President of the Linnean Society.*

---

\* [Owing to unforeseen private engagements, he was unable to take any further active part in the preparation of the "Enumeration."]

† [The expense of the map and drawing of the first three plates in Volume 23 was defrayed by Mr. Forbes, afterwards the entire cost of the plates was borne by the Linnean Society.]

To these proposals the Council agreed, and Parts 1 and 2 of the Enumeration were issued in 1886. Copies of these and subsequent ones were freely distributed amongst English residents in China, with the result of inducing many to assist in the work of collecting specimens. Amongst the earlier was Dr. Henry, at the time an officer in the Chinese Imperial Maritime Customs. Down to the time of his leaving China in 1900, the collections made by this indefatigable botanist reached 15,700 numbers, each represented by numerous duplicates and amounting in all to some 150,000 sheets. Henry's collections revealed the existence of a flora of surprising and unexpected richness, and raised problems of geographical distribution of the highest interest.

The Committee had hoped that the Catalogue they contemplated might be contained in a single volume of the Society's Journal. But it speedily became clear that it would far exceed those limits. It was further evident that the whole undertaking would be more costly and laborious than was originally contemplated. In addition to three grants amounting in all to £700 obtained from the Government Grant Committee of the Royal Society, further aid amounting to £175 was therefore obtained from the British Association, and progress reports were presented to that body in 1887, 1888, and 1889. The total sum received and expended by the Committee on the undertaking up to 1891 amounted to £875.

In 1890 Mr. Hemsley was appointed to the post of Principal Assistant in the Herbarium of the Royal Botanic Gardens, Kew, and his official duties precluded his devoting himself any longer to the work. The completion therefore only became possible by the cooperation, under Mr. Hemsley's general supervision, of various botanists whose names stand at the heads of their several contributions. From this point onwards the cost of printing and publication has been exclusively borne by the Linnean Society.

During their lifetime (for both unhappily passed away during the progress of the work) the Committee met with the kindest sympathy and assistance from M. C. J. Maximowicz of the Académie Impériale of St. Petersburg, who had long been engaged on the elaboration of the collections made by Russian travellers in China, and from M. Franchet of the Muséum d'Histoire Naturelle at Paris, who was occupied in describing and publishing the extremely rich collections made by the French missionaries in Yunnan.

The following letter affords an interesting testimony to the importance Maximowicz attached to the Enumeration :—

Petersburg Botanic Garden,  
15 September, 1885.

DEAR SIR,

You have caused me a most agreeable surprise by the gift of the first fasciculus of Messrs. Forbes and Hemsley's most important enumeration of the flora of China. I need not tell you that it will be a constant and most useful source of reference to me and that I am extremely grateful for this laborious and well-executed undertaking which will save me an immense deal of work, when occupied with my own work on the neighbour-floras. The interest I take in it is still more lively through the important set of Dr. Henry's Hupeh plants, recently received from the Kew Herbarium, in which I find quite a number of the novelties. It appears, however, that the first orders were already printed when Dr. Henry's collection reached Kew.

What a pity it is that our esteemed Dr. Hance did not live to see the commencement of a work which would have elicited his most lively interest and approval.

Believe me, etc.,

(Signed) C. J. MAXIMOWICZ.

The late Baron Richthofen, the greatest authority on the physical geography of China, was no less appreciative of the value of the work :—

Berlin W., Kurfürstenstrasse, 117,  
January 20, 1889.

My dear Mr. THISELTON-DYER,

It has been a very pleasant surprise to me, this morning, to receive from you the "Index Floræ Sinensis." I thank you sincerely for this kind token of remembrance.

It is of great value to have now a Flora of China embodying all the species known from that country. You have evidently succeeded at Kew to get up a very complete collection. At the same time, in looking over the localities mentioned in the book, it strikes me that large portions of China are still unexplored botanically. There remains a splendid field for a good collector in the Tsinking mountains, the province of Sz'-chwan, and chiefly its elevated regions west of Chêng-tu-fu. Work in those parts will be greatly facilitated by the solid foundation laid through the work of Forbes and Hemsley.

Yours very truly,

(Signed) F. RICHTHOFEN.



From the circumstances of the case the enumeration of the species, of which we now possess material or know of the existence in other herbaria, is admittedly unequal. The number of those for instance contained in Part I would probably from present knowledge have to be increased by a third. This has been as far as possible remedied by the list drawn up by Miss M. Smith of the new species published during the progress of the work and of those already described whose area has since been found to extend to China.

The usefulness of the whole work has been enormously enhanced by the complete index of all names cited including synonyms. It contains some 17,000 entries, and is due to the indefatigable industry of Mr. Daydon Jackson, the Society's General Secretary.

No useful purpose would be served by an attempt to specify all those who have supplied the material which has been used in preparing the Enumeration. Their names are given throughout under the species they contributed. Full particulars for the most part will be found about them in Bretschneider's exhaustive "History of European Botanical Discoveries in China," published in 1898.

A few exceptions must, however, be made. Dr. Hance who, to use the words of Bretschneider, "has connected his name for all time with the Flora of China," died June 22, 1886, as already noticed, after the appearance of the first part. His Herbarium of Asiatic plants containing upwards of 22,000 species was acquired by the British Museum.

Dr. Henry's important collections were, as also already noticed, not available for the first part and only to a small extent for the second. From that point they were continuously drawn upon as they successively reached this country.

Dr. Henry also obtained for the Kew Herbarium Dr. Ernst Faber's important collection from Mt. Omei as well as Morse's from Kwangsi and Ducloux's from Yunnan. Faber's own herbarium was afterwards destroyed by fire. His plants are taken up in the Enumeration from the fifth part onwards.

The extraordinary richness of the flora of Western and Central China as revealed by Henry's collections, induced Messrs. Veitch to send out Mr. E. H. Wilson to make further botanical explorations. He made two journeys—the first in 1899, when he arrived at Szemao to see Henry, and the second in 1903. Messrs. James

Veitch and Sons, with a liberality worthy of their distinguished firm, have presented to the Kew Herbarium a complete set of Wilson's plants. Some are taken up in the later parts of the Enumeration.

In 1875 Mr. Hemsley had drawn up a rough list of the Chinese plants in the Kew Herbarium for Mr. Forbes. It contained 3064 species, including ferns. The present Enumeration contains 8271, of which 4230 are endemic or not known to occur outside the Chinese Empire. The most moderate estimate cannot put the whole flora as containing less than 12,000 species.

It only remains in bringing this note to a conclusion to add a few particulars as to the bibliography, which have been supplied by Mr. Hemsley.

With regard to the literature cited, it may be useful to explain two or three points which might not otherwise be quite clear. Throughout, the "Mélanges Biologiques" are cited for Maximowicz's species, though in most instances they were originally published in the "Bulletin de l'Académie Impériale des Sciences de St. Pétersbourg." The references to Franchet's "Plantæ Davidianæ" are to the repaged issue of the "Première Partie," and not to the original pages in the "Nouvelles Archives du Muséum." On the other hand, the pages are given of the "Annales Musei Botanici Lugduno-Batavi" for Miquel's "Pro-lusio Floræ Japonicæ," of which there is a repaged edition. Maximowicz's "Ad Floræ Asiæ Orientalis cognitionem meliorem Fragmenta" is sometimes cited under this title, and sometimes the references are direct to the "Bulletin de la Société Impériale des Naturalistes de Moscou," but the pagination is the same in both. Siebold and Zuccarini's "Floræ Japonicæ Familiæ Naturales . . ." originally appeared in the Münchener "Abhandlungen," vol. iv. 1844-6, and the references are sometimes to the pages of the re-issue and sometimes to the consecutive numbers of the plants enumerated therein. Bunge's "Enumeratio Plantarum quas in China Boreali collegit" was published in the "Mémoires présentées à l'Académie de St. Pétersbourg par divers Savans," ii. 1835, but the references in the following "Index" are to a repaged copy. Lastly, the original edition of Loureiro's "Flora Cochinchinensis" is the one cited where not otherwise stated.

W. T. THISELTON-DYER.

Kew,  
December, 1905.

75 3/8

2241-2

JANUARY 1.

Price 5s.

# THE JOURNAL

OF

# THE LINNEAN SOCIETY.

VOL. XXXVI.

BOTANY.

No. 249. ✓

## CONTENTS.

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.—	
PART XIV. ....	1

**See Notice on last page of Wrapper.**

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1903.

814.00

112

# LINNEAN SOCIETY OF LONDON.

## LIST OF THE OFFICERS AND COUNCIL.

Elected 24th May, 1902.

### PRESIDENT.

Prof. Sydney Howard Vines, M.A., D.Sc., F.R.S.

### VICE-PRESIDENTS.

William Carruthers, F.R.S.  
Frank Crisp, LL.B., B.A.

Herbert Druce, F.Z.S.  
Rev. Thomas R. R. Stebbing, F.R.S.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

Prof. G. B. Howes, D.Sc., LL.D., F.R.S. | D. H. Scott, M.A., Ph.D., F.R.S.

### COUNCIL.

William Carruthers, F.R.S.  
Frank Crisp, LL.B., B.A.  
Herbert Druce, F.Z.S.  
Prof. J. Reynolds Green, M.A., D.Sc.,  
F.R.S.  
W. Botting Hemsley, F.R.S.  
Prof. G. B. Howes, D.Sc., LL.D., F.R.S.  
B. Daydon Jackson, Esq.  
George Masee, Esq.

G. S. Saunders, Esq.  
Dukinfield H. Scott, M.A., Ph.D.,  
F.R.S.  
Rev. Thomas R. R. Stebbing, M.A.,  
F.R.S.  
Col. C. Swinhoe, F.Z.S., F.E.S.  
A. G. Tansley, B.A.  
Prof. Sydney H. Vines, D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*. The former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meet at 4 p.m., as required during the Session. The Members for 1902-1903, in addition to the officers, are:—

Herbert Goss, F.G.S.  
Albert C. L. G. Günther, M.D., F.R.S.  
W. Botting Hemsley, F.R.S.  
G. R. M. Murray, F.R.S.  
P. Chalmers Mitchell, M.A., F.Z.S.

Prof. F. W. Oliver, D.Sc.  
A. B. Rendle, M.A., D.Sc.  
Howard Saunders, F.Z.S.  
F. N. Williams, L.R.C.P.

ENUMERATION  
OF  
ALL THE PLANTS KNOWN  
FROM  
CHINA PROPER, FORMOSA, HAINAN, THE COREA,  
THE LUCHU ARCHIPELAGO,  
AND  
THE ISLAND OF HONGKONG;  
TOGETHER WITH THEIR  
DISTRIBUTION AND SYNONYMY.

(CONTINUED FROM VOL. XXVI.)

133. HYDROCHARIDACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Hydrilla verticillata*, *Royle, Ill. Bot. Himal.* t. 376 (1839); *Presl, Bemerk.* p. 112 (1844); *Caspary, in Pringsh. Jahrb.* i. (1858) p. 494; *Maxim. Fl. As. Or. Fragm.* p. 59; *Hook. f. Fl. Brit. Ind.* v. p. 659.

*Hydrilla verticillata*, var. *Roxburghii*, *Casp. in Pringsh. Jahrb.* i. (1858) p. 494.

*Hydrilla verticillata*, var. *crispa*, *Casp. l. c.*

*Hydrilla dentata*, *Casp. in Bot. Zeit.* xii. (1854) p. 56.

*Hydrilla japonica*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 271.

*Serpicula verticillata*, *Roxb. Pl. Corom.* ii. t. 164.

CHIHLI: Peking Plain (*Bretschneider ex Maximowicz; Moellendorf* in hb. *Hance*, 20353!); KIANGSU: Kiangnan (*Staunton*!); HUPEH: Ichang (*A. Henry*, 4368!); KWANGTUNG (*Ford*!), Lienchau River (*Graves*!). *Mus. Brit.*; *Herb. Kew.*

Central Europe, Tropical Asia and Australia, Mascarene Islands.

1. **Lagarosiphon Roxburghii**, *Benth. in Benth. & Hook. f. Gen. Pl.* iii. p. 451; *Hook. f. Fl. Brit. Ind.* v. p. 659.

*Vallisneria alternifolia*, *Roxb. Hort. Beng.* p. 71; *Fl. Ind.* iii. p. 750; *Wight, in Hook. Bot. Misc.* ii. (1831) p. 344, t. 11.

*Nechamandra Roxburghii*, *Planch. in Ann. Sc. Nat.* 3<sup>me</sup> sér. xi. (1849) p. 78, excl. syn. *Roxb.*; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 125.

*Nechamandra alternifolia*, *Thw. Enum. Pl. Zeyl.* p. 332.

KWANGTUNG: Whampoa (*Hance, 1517, 5411!*). Mus. Brit.; Herb. Kew.

Tropical Asia.

1. **Vallisneria spiralis**, *Linn. Sp. Pl.* ed. 1, p. 1015; *Hook. Bot. Misc.* iii. (1833) p. 87; *Suppl.* tt. 23-24; *Maxim. Fl. As. Or. Fragm.* p. 59; *Hook. f. Fl. Brit. Ind.* v. p. 660.

CHIHLI: Peking, Wanshoushan (*Bretschneider, 718!*); KIANGSU: Kiangnan (*Staunton!*). Mus. Brit.; Herb. Kew.

The warmer regions of both hemispheres.

1. **Blyxa Roxburghii**, *Rich. in Mém. Inst. Fr.* (1811) p. 77, t. 5; *Benth. Fl. Hongk.* p. 347; *Fl. Austral.* vi. 258.

*Blyxa octandra*, *Planch. ex Thw. Enum. Pl. Zeyl.* p. 332.

*Diplosiphon oryzetorum*, *Decne. in Jacquem. Voy.* iv. p. 166, t. 167.

*Vallisneria octandra*, *Roxb. Pl. Corom.* t. 165.

KWANGTUNG: Swatow (*Sampson!*); HONGKONG (*Wilford, 285! Wright, 541! Hance, 1517!*). Mus. Brit.; Herb. Kew.

Tropical Asia and Australia.

1. **Hydrocharis Morsus-ranæ**, *Linn. Sp. Pl.* ed. 1, p. 1036; *Lam. Encycl.* t. 820; *Schnizl. Icon.* i. 59; *Hook. f. Fl. Brit. Ind.* v. p. 662.

*Hydrocharis asiaticus*, *Miq. Fl. Ind. Bat.* iii. p. 239; *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 194; *Franch. et Savat. Enum. Pl. Jap.* ii. 19; *Maxim. Fl. As. Or. Fragm.* p. 59; *Debeaux, Fl. Shangh.* p. 60.

CHIHLI: Wanshoushan, Peking (*Bretschneider in Herb. Hance, 18137!*); KIANGSU: Shanghai (*Carles, 384!*); CHEKIANG: Ningpo Creek (*Faber, 59!*); HUPEH: Ichang (*A. Henry, 127! 2251!*). Mus. Brit.; Herb. Kew.

Europe, North Asia, North India, Japan, Australia, "cultivated in Java."

1. *Ottelia alismoides*, *Pers. Syn. Pl.* i. p. 400; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 272 & iii. p. 201; *Benth. Fl. Austral.* vi. p. 257; *Hook. f. Fl. Brit. Ind.* v. p. 662.

*Ottelia japonica*, *Miq. in Verh. Akad. Wetensch.* ii. ii. (1868) p. 89, & *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 271.

*Ottelia indica*, *Planch. in Dalz. & Gibs. Bomb. Fl.* p. 278.

*Stratiotes alismoides*, *Linn. Sp. Pl.* ed. 1, p. 535.

*Damasonium indicum*, *Willd. Sp. Pl.* ii. p. 276; *Roxb. Fl. Ind.* ii. p. 216; *Pl. Corom.* ii. p. 45, t. 185.

KIANGSU: Kianang (*Staunton*!); Kiangsi: Kiukiang (*Shearer*!); Szechuen: Hokiang (*Fuber*, 966!); Hupeh: Ichang (*A. Henry*, 165!); Kwangtung (*Ford*, 241!), Lantao Is. (*Ford*, 61!); Hainan (*A. Henry*, 8291! 8662!); Whampoa (*Hance*, 6056!); Fukwing (*Krone*, 1539!). *Mus. Brit.*; *Herb. Kew.*

Tropical Asia, North Australia.

1. *Enhalus Kœnigii*, *Rich. in Mém. Inst. Par.* ii. (1797) p. 78; *Miq. Fl. Ind. Bat.* iii. p. 237; *Hook. f. Fl. Brit. Ind.* v. p. 663.

*Enhalus acoroides*, *Rich. ex Cnatin, Anat. Pl. Aquat.* p. 15, t. 6; *Tokutaro Ito, in Ann. Bot.* xiii. (1899) p. 465.

*Enhalus*, *Griff. Notul.* iii. p. 175; *Ic.* tt. 249-250.

*Stratiotes acoroides*, *Linn. f. Suppl.* p. 268.

LUCHU ARCHIPELAGO: between the islands of Irumuti and Uchibanari (ex *Tokutaro Ito*).

Malaya, Ceylon, Arabia, and North Australia.

1. *Thalassia Hemprichii*, *Aschers. in Neumayer, Anleit. Wiss.* p. 361; *Durand & Schinz, Consp. Fl. Afr.* v. p. 5.

*Schizotheca Hemprichii*, *Ehrenb. in Abh. Akad. Berl.* i. (1832) p. 429; *Solms-Laubach, in Schweinf. Beitr. Fl. Aethiop.* p. 194.

LUCHU ARCHIPELAGO (*Wright*!). *Herb. Kew.*

Red Sea, Indian and Pacific Oceans.

1. *Halophila ovata*, *Gaud. in Freyc. Voyage, Bot.* p. 430, t. 40. f. 1; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 129; *Tokutaro Ito, in Ann. Bot.* xiii. (1899) p. 465.

*Halophila ovalis*, *Hook. f. Fl. Tism.* ii. p. 45.

*Caulinia ovalis*, *R. Br. Prodr.* p. 339.

HONGKONG: Kowlung Bay (*Harland*, 282!); PRATAS REEF (*Wilford*, 453!); LUCHU ARCHIPELAGO (*Wright*, 327!). *Mus. Brit.*; *Herb. Kew.*

Tropical Asia, Australia.

## 134. BURMANNIACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Burmannia cœlestis*, *D. Don, Prodr. Fl. Nepal.* p. 44; *Royle, Ill. Bot. Himal.* p. 373, t. 91. f. 1; *Hance, in Journ. Bot.* xvi. (1878) p. 111; *Hook. f. Fl. Brit. Ind.* v. p. 665.

*Burmannia azurea*, *Griff. Notul.* iii. p. 236, &  *Ic. t. 272.* fig. 1; *Becc. Malesia*, i. p. 242, t. 15. figs. 1-3.

*Cryptonema malaccensis*, *Turcz. in Bull. Soc. Nat. Mosc.* xxi. (1848) i. p. 591; *Kränzl. in Engl. Bot. Jahrb.* vi. (1885) p. 55.

*Nephrocœlium malaccense*, *Turcz. in Bull. Soc. Nat. Mosc.* (1853) i. p. 287.

*Nephrocodium malaccense*, *C. Muell. in Walp. Ann.* vi. p. 40; *Becc. Malesia*, i. p. 242.

KIANGSI (*Staunton!*); KWANGTUNG: Foot of the White Cloud Hill, near Canton (*Sampson, 677! Hance, 7933! Lamont, 1086!*), Macao (*Nelson!*); LUCHU ARCHIPELAGO: Amami Osima (*Döderlein ex Kränzlin*). *Mus. Brit.*; *Herb. Kew.*

India and Malaya.

2. *Burmannia Dalzieli*, *Rendle, in Journ. Bot.* xl. (1902) p. 311, pl. 441 B.

KWANGTUNG: near Canton (*Dalziel!*). *Mus. Brit.*

3. *Burmannia disticha*, *Linn. Sp. Pl.* ed. 1, p. 287; *Roxb. Pl. Corom.* iii. t. 242; *Benth. Fl. Austral.* vi. p. 397; *Hook. f. Fl. Brit. Ind.* v. p. 664.

*Burmannia distachya*, *R. Br. Prodr.* p. 265.

*Burmannia spica gemina Linnæi*, *Burm. Thes. Zeyl.* p. 50, t. 20. fig. 1.

KWANGTUNG: Shing Hing Pass, West River (*Ford!*); HONGKONG (*Champion!*). *Herb. Kew.*

India, Australia.

4. *Burmannia nepalensis*, *Hook. f. Fl. Brit. Ind.* v. p. 666; *Hemsl. in Ann. Bot.* v. (1891) p. 407; *Tokutaro Ito, Burmanniaceæ of Japan*, p. 14, t. 2. figs. 3-4.

*Cyanotis nepalensis*, *Miers, in Wall. Cat.* n. 9006.

*Gonyanthes nepalensis*, *Miers, in Trans. Linn. Soc.* xviii. (1841) p. 537, t. 38. fig. 1.

KWANGTUNG: Lofaushan (*Ford, 68!*). *Herb. Kew.*

North India, Cochin China.



5. **Burmannia Wallichii**, *Hook. f. Fl. Brit. Ind.* v. p. 666.  
*Gonyanthes Wallichii*, *Miers, in Trans. Linn. Soc.* xviii. (1841) p. 537,  
 t. 38. fig. 2; *Benth. Fl. Hongk.* p. 364.  
*Burmannia* sp., *Griff. Ic.* t. 272. fig. 2.  
 HONGKONG: Mount Gough (*Wilford*, 281!). Herb. Kew.  
 India.

## 135. ORCHIDACEÆ: EPIDENDREÆ.

(By R. A. FOLFE, A.L.S.)

1. **Microstylis congesta**, *Reichb. f. in Walp. Ann.* vi. p. 206;  
*Hance, in Journ. Linn. Soc., Bot.* xiii. (1871) p. 127; *Ridl. in Journ.*  
*Linn. Soc., Bot.* xxiv. (1888) p. 334; *Hook. f. Fl. Brit. Ind.* v.  
 p. 689.

*Dienia congesta*, *Lindl. Gen. & Sp. Orch.* p. 22; *Benth. Fl. Hongk.*  
 p. 352.

FORMOSA: Bankinsing Mountains and South Cape (*A. Henry*,  
 835! 1342!); HONGKONG (*Hance!* *Harland*, 1057! *Lamont!*)  
 Mus. Brit.; Herb. Kew.

North India, from Nepal eastward, Malaya, and Queensland.

2. **Microstylis cylindrostachya**, *Reichb. f. in Walp. Ann.* vi.  
 p. 207; *Ridl. in Journ. Linn. Soc., Bot.* xxiv. (1888) p. 333;  
*Hook. f. Fl. Brit. Ind.* v. p. 689.

*Dienia cylindrostachya*, *Lindl. Gen. & Sp. Orch.* p. 22.

SZECHUEN: Tatsienlou (*Soulié*, 633!).

Northern and Central India

3. **Microstylis monophyllos** *Lindl. Gen. & Sp. Orch.* p. 19;  
*Ridl. in Journ. Linn. Soc., Bot.* xxiv. (1888) p. 318.

*Malaxis monophyllos*, *Sw. in Vet.-Akad. Nya Handl. Stockh.* xxi. (1800)  
 p. 234; *Franch. Pl. David.* i. p. 204.

CHIHLI: Pohnashan (*David*); HUPEH: Fang (*A. Henry*,  
 6869, 6875!); SZECHUEN: Mount Omei, at 9000 ft. (*Faber*, 958!).  
 Herb. Kew.

Widely diffused through the temperate regions of the northern  
 hemisphere.

1. **Oberonia iridifolia**, *Lindl. (?) Gen. & Sp. Orch.* p. 15;  
*Hook. f. Fl. Brit. Ind.* v. p. 675.

*Malaxis iridifolia*, *Reichb. f. in Walp. Ann.* vi. p. 208.

HAINAN (*Ford*, 575!). Herb. Kew.

India.

The specimens are passing into the fruiting stage, hence there  
 remains a slight uncertainty as to their identity with the Indian

plant. The habit, however, is very similar, and the lip distinctly toothed.

2. *Oberonia yunnanensis*, Rolfe, n. sp.

*Caules* brevi, 5-6-phylli. *Folia* lanceolato-linearia, paululo falcata, acuta,  $\frac{3}{4}$ - $1\frac{1}{4}$  poll. longa,  $1-1\frac{3}{4}$  lin. lata. *Scapi* graciles, 4-6 poll. longi. Flores numerosi, minuti. *Bracteæ* subulato-filiformes,  $\frac{3}{4}$  lin. longæ. *Pedicelli*  $\frac{3}{4}$ -1 lin. longi. *Sepala* ovata, subobtusata, concava,  $\frac{1}{3}$  lin. longa. *Petala* oblonga, obtusa,  $\frac{1}{3}$  lin. longa. *Labellum*  $\frac{3}{4}$  lin. longum, ad medium in lobos oblongos subobtusos paulo divergentes fissum, basi latum; lobi integri. *Columna* brevissima.

YUNNAN: Manme mountains, south of Red River, at 7000 ft. (*A. Henry*, 9152!). Flowers yellow. Herb. Kew.

Allied to *O. pyrulifera*, Lindl., a native of Khasia and Munnipore, differing in its longer stems, narrower leaves, and the divergent lobes of the lip.

1. *Liparis chloroxantha*, Hance, in *Journ. Bot.* xxi. (1883) p. 231; *Ridl. in Journ. Linn. Soc., Bot.* xxii. (1886) p. 284.

HONGKONG (*Ford!*). Mus. Brit.; Herb. Kew.

2. *Liparis Henryi*, Rolfe, in *Kew Bulletin* (1896), p. 193.

FORMOSA: South Cape (*A. Henry*, 2074!). Herb. Kew.

Allied to *Liparis acuminata*, Hook. f., from the Khasia Hills. The flowers are considerably smaller than in *L. macrantha*, Rolfe, and the colour purple, with the front and margin of the lip whitish.

3. *Liparis latilabris*, Rolfe, n. sp.

*Caules* diphylli. *Folia* subcoriacea, lanceolata, acuta, basi attenuata, 10-15 poll. longa, 7-12 lin. lata. *Scapi* graciles, 9-15 poll. longi; racemi 5-6 poll. longi, multiflori. *Bracteæ* anguste lineari-lanceolatae, acuminatae,  $2\frac{1}{2}$ -3 lin. longæ. *Pedicelli* 4-5 lin. longi. *Sepala* lineari-lanceolata, acuta,  $3-3\frac{1}{2}$  lin. longa. *Petala* anguste linearia,  $3-3\frac{1}{2}$  lin. longa. *Labellum* reniformi-orbiculare, crenulatum, 4-5 lin. latum, basi bituberculatum. *Columna*  $1-1\frac{1}{4}$  lin. longa.

YUNNAN: Mengtze, rocks in deep shade at 5500 ft. (*Hancock*, 255!). Herb. Kew.

A very distinct species of the section *Coriifoliae*, allied to *L. plantaginea*, Lindl., but with more numerous flowers of about half the size.

4. *Liparis longipes*, *Lindl. in Wall. Pl. Asiat. Rar.* i. p. 31, t. 35; *Lindl. Gen. & Sp. Orch.* p. 30; *Benth. Fl. Hongk.* p. 352; *Ridl. in Journ. Linn. Soc., Bot.* xxii. (1886) p. 293 (excl. syn. *L. elegans*); *Hook. f. Fl. Brit. Ind.* v. p. 703.

*Sturmia longipes*, *Reichb. f. in Bonplandia*, iii. (1855) p. 250.

FORMOSA: Bankinsing Mountains (*A. Henry*, 1592!); South Cape (*A. Henry*, 1979!); KWANGTUNG: Lofaushan (*Faber*!); HONGKONG (*Champion*, 335! 523! *Harland*, 812! *Bowring*! *Lamont*, 749!). Mus. Brit.; Herb. Kew.

India, Malaya, Fiji, and Samoa.

5. *Liparis macrantha*, *Rolfe in Ann. Bot.* ix. (1895) p. 156.

FORMOSA: Tamsui (*A. Henry*, 1695!). Herb. Kew.

6. *Liparis nervosa*, *Lindl. Gen. & Sp. Orch.* p. 26; *Benth. Fl. Hongk.* p. 352; *Franch. et Savat. Enum. Pl. Japon.* ii. p. 21; *Ridl. in Journ. Linn. Soc.* xxii. p. 262.

*Ophrys nervosa*, *Thunb. Fl. Japon.* p. 27.

*Epidendrum nervosum*, *Thunb. Ic. Pl. Japon.* i. t. 10.

*Malaxis nervosa*, *Sw. in Vet.-Akad. Nya Handl. Stockh.* xxi. (1800) p. 235.

*Sturmia nervosa*, *Reichb. f. in Bonplandia*, iii. (1855) p. 250.

*Liparis formosana*, *Reichb. f. in Gard. Chron.* (1880) i. p. 394.

*Liparis bituberculata*, var. ? *formosana*, *Ridl. in Journ. Linn. Soc., Bot.* xxii. (1886) p. 263.

FORMOSA: Tamsui (*Oldham*, 555!); South Cape (*A. Henry*, 646!); HONGKONG (*Hance*, 1193!); LUCHU ARCHIPELAGO (*Wright*, 342!). Mus. Brit.; Herb. Kew.

Japan.

There remains a slight doubt about *Liparis formosana*, *Reichb. f.*, which was based on a cultivated plant introduced from Formosa by *Maries*. The author compares it with *L. bituberculata*, *Lindl.*, but describes the lip as acute, which certainly does not apply to *Oldham's* specimen, which otherwise seems identical. The only Japanese specimen I have seen is marked "cultivated."

7. *Liparis odorata*, *Lindl. Gen. & Sp. Orch.* p. 26; *Reichb. f. in Walp. Ann.* vi. p. 217; *Benth. Fl. Hongk.* p. 352.

*Malaxis odorata*, *Willd. Sp. Pl.* iv. p. 91.

*Malaxis lancifolia*, *Sm. in Rees, Cyclop.* xxii. n. 7.

*Empusa paradoxa*, *Lindl. Bot. Reg.* sub t. 825; *Gen. & Sp. Orch.* p. 17.

*Liparis paradoxa*, *Reichb. f. in Walp. Ann.* vi. p. 218; *Ridl. in Journ. Linn. Soc., Bot.* xxii. (1886) p. 26; *Hook. f. Fl. Brit. Ind.* v. p. 697.

CHEKIANG (*Fortune*, 39!); Ningpo (*Faber*, 300!); FORMOSA:

Bankinsing Mountains (*A. Henry*, 896!); SZECHUEN: Mount Omei (*Faber*, 952! 954!); YUNNAN: Mengtze, mountains at 6000–7000 ft. (*Hancock*, 217!); 5000–5500 ft. (*A. Henry*, 11105!); HONGKONG (*Champion*, 266!). Mus. Brit.; Herb. Kew.

India, Malaya, and Japan.

The specific name *odorata* is much older than *paradoxa*, under which the species is generally known. Ridley says “*L. odorata*, Benth., Flora Hongkong. p. 352, sine dubio *L. nervosa* est,” but in this I cannot agree.

8. *Liparis pauciflora*, *Rolfe*, in *Kew Bulletin* (1896), p. 193.

SZECHUEN: S. Wushan (*A. Henry*, 5675, 5675 a!). Herb. Kew.

Allied to the Indian *L. rostrata*, Reichb. f., which also has an ecallose lip and short bracts, though the Indian species has larger and more numerous flowers.

9. *Liparis plicata*, *Franch. et Savat. Enum. Pl. Japon.* ii. pp. 22, 509; *Somoku Zusetsu*, xviii. p. 21; *Ridl. in Journ. Linn. Soc., Bot.* xxii. (1886) p. 285.

FORMOSA: Bankinsing Mountains (*A. Henry*, 138! 1627!); HONGKONG (*Champion*, 521!). Herb. Kew.

Japan.

Ridley identifies the Hongkong plant (which he attributes to “Fortune?”) with the Japanese one.

10. *Liparis yunnanensis*, *Rolfe*, n. sp.

*Caules* diphylli. *Folia* subcoriacea, elongato-lanceolata, acuta, basi attenuata, 10–15 poll. longa, 10–16 lin. lata. *Scapi* graciles, 1¼ poll. longi; racemi 6 poll. longi, multiflori. *Bracteæ* lineares, acuminatæ, 3–8 lin. longæ. *Pedicelli* 5–8 lin. longi. *Sepala* lineari-lanceolata, acuta, 5–6 lin. longa. *Petala* anguste linearia, 5–6 lin. longa. *Labellum* obovato-orbiculare, minutissime crenulatum, 5 lin. latum, basi bituberculatum. *Columna* incurva, 2½–3 lin. longa.

YUNNAN: Mengtze, mountains at 5000 ft., among rocks (*A. Henry*, 10485!). Flowers yellow. Herb. Kew.

This is allied to *L. plantaginea*, Lindl., of the section *Coriifoliae*, but has longer leaves, much narrower bracts, and narrower sepals; the unopened buds are also only half as stout. From *L. latilabris*, Rolfe, it differs in having fewer flowers, of twice the size.

1. *Corallorhiza innata*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 209; *Maxim. Ind. Fl. Pekin. in Prim. Fl. Amur.* p. 477.

*Ophrys corallorhiza*, *Linn. Sp. Pl.* ed. 1, p. 945.

*Cymbidium Corallorhiza*, *Sw. in Vet.-Akad. Handl. Nya Handl. Stockh.* xxi. (1800) p. 238.

CHILHI: Peking (ex *Maximowicz*).

Europe, N. Asia, and N. America.

*Maximowicz* enumerates this with a query in the above work, but without giving any clue to his authority. He, however, confirms the locality in a letter to Mr. Forbes dated Dec. 16, 1894.

1. *Oreorchis foliosa*, *Lindl. in Journ. Linn. Soc.* iii. (1859) p. 27; *Hook. f. Fl. Brit. Ind.* v. p. 709.

"*Corallorhiza* with leaves," *Lindl. ex Royle, Ill. Bot. Himal.* p. 362.

SZECHUEN: without locality (*Pratt*, 19, 55!). *Herb. Kew.*

Temperate Himalaya.

1. *Dendrobium aduncum*, *Wall. in Bot. Reg.* xxviii., *Misc.* p. 58; xxxii. t. 15; *Bot. Mag.* t. 6784; *Hook. f. Fl. Brit. Ind.* v. p. 730.

KWANGTUNG: Lofaushan (*Ford*, 91!). *Herb. Kew.*

North India, from Bhotan to Assam.

2. *Dendrobium aggregatum*, *Rowb. Fl. Ind.* iii. p. 477; *Lindl. Bot. Reg.* t. 1695; *Bot. Mag.* t. 3643; *Hook. f. Fl. Brit. Ind.* v. p. 749.

KWANGTUNG: Lofaushan, at 800 ft. (*Ford*!). *Herb. Kew.*

North-eastern India and Malayan Peninsula.

3. *Dendrobium atropurpureum*, *Miq. Fl. Ind. Batav.* iii. p. 644; *Hook. f. Fl. Brit. Ind.* v. p. 724.

*Oxystophyllum atropurpureum*, *Blume, Rumphia*, iv. p. 41, t. 193, fig. 4, t. 198, fig. C.

*Oxystophyllum carnosum*, *Blume, Bijdr.* p. 336, fig. 38; *Lindl. Gen. & Sp. Orch.* p. 72.

*Aporum concinnum*, *Lindl. in Wall. Cat.* n. 2019.

HAINAN: Taihui (*Ford*, 570!). Flowers dark claret. *Herb. Kew.*

Burma, Malaya Peninsula and Archipelago.

4. *Dendrobium aurantiacum* *Reichb. f. in Gard. Chron.* (1887) ii. p. 98; *Hook. f. Fl. Brit. Ind.* v. p. 751.

*Dendrobium chryseum*, *Rolfe, in Gard. Chron.* (1888) i. p. 233; *Hook. f. Fl. Brit. Ind.* v. p. 748.

SZECHUEN: without locality (*Pratt*, 26!). Herb. Kew.  
Eastern Himalaya.

When describing *Dendrobium chryseum* I inadvertently overlooked Reichenbach's *D. aurantiacum*.

[*Dendrobium auriferum*, Lindl., belongs to *Sarcochilus*.]

5. *Dendrobium* (§ *Formosæ*) *bellatulum*, *Rolfe*, n. sp.

*Planta* circa 3 poll. alta, insignis, nigro-hirsuta. *Pseudobulbi* fusiformi-oblongi, 3-4-phylli. *Folia* oblonga v. elliptico-oblonga, inæqualiter et brevissime biloba, 1-1 $\frac{3}{4}$  poll. longa, 5-9 lin. lata. *Flores* 1-3, subterminales, pulchelli, albidi. *Bracteæ* elliptico-oblongæ, acutæ, 4-5 lin. longæ. *Pedicelli* 9-12 lin. longi. *Sepalum* posticum elliptico-oblongum, subobtusum, 9-10 lin. longum; lateralia triangularia, subacuta, 10-12 lin. longa. *Petala* elliptico-obovata, obtusa v. minute apiculata, 10-12 lin. longa. *Labellum* 13-15 lin. longum, pandurato-trilobum; lobi laterales rotundati; lobum intermedium latissime obcordato-bilobum, 5-6 lin. latum; discus medio obtuse 5-carinatus, carinis tuberculoso-verrucosis. *Columna* clavato-oblonga, 3 lin. longa; alæ oblongæ, obtusæ. *Mentum* late saccatum, subobtusum, 4-5 lin. longum.

YUNNAN: Mengtze, S.E. mountain-forests, at 5000 ft., on trees (*A. Henry*, 11109!). Flowers whitish. Herb. Kew.

A beautiful little plant, much like a miniature edition of *D. formosum*, Roxb.

6. *Dendrobium Ceraia*, *Lindl. Gen. & Sp. Orch.* p. 89.

*Ceraia simplicissima*, *Lour. Fl. Cochinch.* p. 518.

CHINA: without locality (ex *Loureiro*).

Cochin China.

7. *Dendrobium chrysanthum*, *Wall. in Lindl. Gen. & Sp. Orch.* p. 80; *Bot. Reg.* t. 1299; *Hook. f. Fl. Brit. Ind.* v. p. 747.

YUNNAN: on hot sandy rocks at Mengtze; very rare (*Hancock*, 119!); grassy banks, rocks, &c., at 5000 ft. (*A. Henry*, 11094!). Herb. Kew.

Tropical Himalaya and Khasia.

8. *Dendrobium clavatum*, *Wall. Cat.* n. 2004; *Lindl. in Paxt. Fl. Gard.* ii. p. 104; *Bot. Mag.* t. 6993; *Hook. f. Fl. Brit. Ind.* v. p. 746.

YUNNAN: on rocks in sandy gulches facing south, at Mengtze

(*Hancock*, 118!); mountain gulches, 6000–7000 ft. (*Hancock*, 584!). Herb. Kew.

Tropical Himalaya and Khas a.

9. *Dendrobium Gibsoni*, *Part. Mag. Bot.* v. p. 169, with plate?; *Hook. f. Fl. Brit. Ind.* v. p. 746.

YUNNAN: Mengtze, S.E. mountains (*A. Henry*, 10613!). Herb. Kew.

Northern India.

The specimen is in fruit, but so far as it goes is not distinguishable from the Indian plant.

10. *Dendrobium hainanense*, *Rolfe*, in *Kew Bulletin* (1896), p. 193.

HAINAN: Lingmen (*A. Henry!* *Ford*, 272!); Hongting (*Ford*, 473!). Herb. Kew.

A species belonging to Lindley's section *Onychium*, and closely allied to the Philippine *D. aciculare*, Lindl., which, however, has longer internodes, more slender and straighter leaves, a shorter mentum, and a shorter and narrower lip.

11. *Dendrobium* (§ *Eudendrobium*) *Hancockii*, *Rolfe*, n. sp.

*Caules* graciles, parce ramosi,  $\frac{1}{2}$ –1 ped. alti,  $1\frac{1}{2}$ –3 lin. lati. *Folia* linearia, obtusa, inæqualiter bidentata, 2–3 poll. longa, 2–3 lin. lata. *Racemi* axillares, breves, biflori v. uniflori. *Bracteæ* ovatæ, acutæ, 1 lin. longæ. *Pedicelli* 6 lin. longi. *Sepala* et *petala* subæqualia, oblonga, subobtusa, v. apiculata, 10–11 lin. longa. *Labellum* trilobum, breviter et late unguiculatum, 9 lin. longum, 8 lin. latum, facie velutino-pubescent; lobi laterales rotundati, obtusi; lobum intermedium reniformi-orbiculare, subobtusum, 5–6 lin. latum. *Columna* lata, 2 lin. longa. *Mentum* breviter saccatum, obtusum,  $1\frac{1}{2}$ –2 lin. longum.

YUNNAN: Mengtze, on rocks at 6000–7000 ft. (*Hancock*, 581!). Flowers yellow, strongly fragrant of honey. Herb. Kew.

A very distinct species, most nearly allied to *D. velutinum*, Rolfe (from Upper Burma), in floral details, but very different in its slender, branching habit.

[*Dendrobium hercoglossum*, Reichb. f. (*Gard. Chron.* (1886) ii. p. 487; *Hamb. Gartenz.* xlii. p. 487), is stated by Sir Joseph Hooker, in the 'Flora of British India,' vol. v. p. 730, to be Chinese. The statement is based on a specimen in the Kew Herbarium labelled "Canton, Mr. Ford," but on referring to

Mr. Ford's letter relating to it we find that it is "a *Dendrobium* which is cultivated about Canton." As Reichenbach originally stated that it was discovered in Malacca by Försterman, we do not enumerate it here.]

12. *Dendrobium Linawianum*, *Reichb. in Walp. Ann.* vi. p. 284; *Warn. & Will. Orch. Album*, iii. t. 141; *Veitch, Man. Orch.* pt. iii. p. 53.

*Dendrobium moniliforme*, *Lindl. Bot. Reg.* xvi. t. 1314, excl. syn. (non Sw.); *Pact. Mag. Bot.* iii. p. 77, cum ic.; *Bot. Mag.* t. 4153; *Maund, Botanist*, iv. t. 194; *Ann. Bot. de Gand*, iii. (1847) t. 137.

CHINA: without locality. Herb. Kew.

The precise habitat of this species remains unknown. It was introduced from China by the Horticultural Society of London, in 1824, and for a long time was confused with *Dendrobium moniliforme*, Sw.

13. *Dendrobium Loddigesii*, *Rolfe, in Gard. Chron.* (1887) ii. p. 155, also p. 786; *Veitch, Man. Orch.* iii. p. 55.

*Dendrobium pulchellum*, *Lodd. Bot. Cab.* xx. t. 1935 (non Roxb.); *Maund, Botanist*, i. t. 5; *Bot. Mag.* t. 5037.

KWANGTUNG: 150 miles north-west of Canton, on rocks (*Ford!*); HAINAN (*A. Henry*, 8136!). Herb. Kew.

For many years this species was confounded with the Indian *Dendrobium pulchellum*, Roxb. (*D. Dalhousieanum*, Wall.).

14. *Dendrobium moniliforme*, *Sw. in Kongl. Vet.-Akad. Nya Handl. Stockh.* xxi. (1800) p. 246; *Lindl. Gen. & Sp. Orch.* p. 84.

*Epidendrum moniliforme*, *Linn. Sp. Pl.* ed. 1, p. 954.

*Epidendrum monile*, *Thunb. Fl. Jap.* p. 30.

*Dendrobium catenatum*, *Lindl. Gen. & Sp. Orch.* p. 84; *id. Bot. Reg.* xxx., *Misc.* p. 60; *Reichb. f. in Walp. Ann.* vi. p. 300.

*Dendrobium japonicum*, *Lindl. Gen. & Sp. Orch.* p. 89.

*Onychium japonicum*, *Blume, Bijdr.* p. 328.

CHINA: without locality (*Reeves ex Lindley*); COREA: Peel Island (*Oldham*, 832!). Herb. Kew.

Japan.

15. *Dendrobium nobile*, *Lindl. Gen. & Sp. Orch.* p. 79; *Sert. Orch.* t. 3; *Bot. Reg.* xxx., *Misc.* p. 48; *Hook. f. Fl. Brit. Ind.* v. p. 740.

HUPEH: Ichang (*A. Henry*, 2365!); SZECHUEN: Chingking (*A. Henry*, 3656!). Herb. Kew.

N.E. India, from Sikkim eastward.



“Sent from Chingking to Hankow as a ‘medicine’ export. The dried stems are also an article of export. It is *Shih-hu*: *Hut’ou*: *Ya-t’ou* of the Chinese.”—*Dr. A. Henry*.

Var. *formosanum*, *Reichb. f. in Gard. Chron.* (1883) i. p. 432.  
FORMOSA (*B. S. Williams*, cult.).

The species was originally described from Chinese material sent home by Mr. Reeves. A living plant, which was bought in the market at Macao, was brought home by him, and presented to Messrs. Loddiges, of Hackney. The variety I have not seen.

16. *Dendrobium yunnanense*, *Finet, in Bull. Soc. Bot. Fr.* xliv. (1897) p. 419, t. 13, fig. A–H.

YUNNAN: Lami, near the River Namna or Laniouho (*Prince Henri d’Orleans, ex Finet*).

17. *Dendrobium zonatum*, *Rolfe, n. sp.*

*Pseudobulbi* graciles, 7–12 poll. longi, foliati. *Folia* oblongo-lanceolata, subacuta,  $2\frac{1}{4}$ –3 poll. longa, 3–6 lin. lata. *Racemi* laterales, breves, 2–4-flori, vaginis membranaceis obtecti. *Bracteæ* late oblongæ, obtusæ, concavæ, membranaceæ, medio brunneo-zonatæ,  $2\frac{1}{2}$ – $3\frac{1}{2}$  lin. longæ. *Pedicelli* graciles, 7–8 lin. longi. *Sepala* 9–10 lin. longa; posticum lanceolato-oblongum, subobtusum; lateralia triangulari-lanceolata, subacuta. *Petala* elliptico-oblonga, subacuta, 9–10 lin. longa. *Labellum* 7–8 lin. longum, pandurato-oblongum, basi latum, apice breviter acuminatum; discus medio pubescens. *Columna* brevissima. *Mentum* late saccatum, 2 lin. longum.

YUNNAN: Fengchenlin, mountain-forest at 7000 ft., on trees (*A. Henry, 10668!*). Flowers white. *Herb. Kew.*

Allied to *D. candidum*, Wall., but readily distinguished by the different bracts, which are remarkable on account of a broad brown zone about the middle.

1. *Bulbophyllum odoratissimum*, *Lindl. Gen. & Sp. Orch.* p. 55; *Wall. Cat.* n. 1987; *Hook. f. Fl. Brit. Ind.* v. p. 758.

YUNNAN: Mengtze, on mountains at 6000–7000 ft. (*Hancock, 586!*). *Herb. Kew.*

Northern India, from Nepal eastward to Tenasserim.

2. *Bulbophyllum radiatum*, *Lindl. Gen. & Sp. Orch.* p. 55; *Wall. Cat.* n. 1986; *Benth. Fl. Hongk.* p. 353; *Hook. f. Fl. Brit. Ind.* v. p. 758.

HUPEH: Ichang (*A. Henry*, 4548!); HONGKONG (*Champion*, 281! *Wright*, 515! *Ford*!). Mus. Brit.; Herb. Kew.  
Burma.

3. *Bulbophyllum Watsonianum*, *Reichb. f. in Flora*, lxxi. (1888) p. 155.

*Eria ambrosia*, *Hance*, in *Journ. Bot.* xxi. (1883) p. 232.

HONGKONG (*Ford*!). Mus. Brit.; Herb. Kew.

This species flowered at Kew a few years ago from the above source. Its identity with *Eria ambrosia*, *Hance*, has only now been discovered; it is certainly not an *Eria*.

4. *Bulbophyllum* (§ *Sestochilos*) *yunnanense*, *Rolfe*, n. sp.

*Rhizoma* repens, validum. *Pseudobulbi* oblongi, monophylli,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. longi. *Folia* petiolata, elliptico-oblonga, subobtusa,  $1\frac{3}{4}$ –5 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{4}$  poll. lata; petiolus  $\frac{1}{4}$ –1 poll. longus. *Scapi* breves,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. longi, 1–2-flori. *Bracteæ* ovato-oblongæ, apiculatæ, 4 lin. longæ. *Pedicelli* 1– $1\frac{1}{2}$  poll. longi. *Sepala* ovata, subobtusa, 5 lin. longa, lateralia basi cum columnæ pedimentum saccatum obtusum extensum. *Petala* ovata, subobtusa, 4 lin. longa, 7-nervia. *Labellum* unguiculatum, ovato-oblongum, subobtusum, recurvum, 4 lin. longum; unguis  $1\frac{1}{2}$  lin. longus. *Columna* lata,  $1\frac{1}{2}$  lin. longa. *Mentum* 4 lin. longum.

YUNNAN: Mengtze, rocks on Great Black mountains, at 6000 ft. (*Hancock*, 587!); North mountains at 7000 ft. (*A. Henry*, 11370!). Herb. Kew.

Near the Indian *B. affine*, *Lindl.*, but the flowers have broader, more obtuse segments.

1. *Cirrhopetalum bicolor*, *Rolfe*.

*Bulbophyllum bicolor*, *Lindl. Gen. & Sp. Orch.* p. 49.

CHINA: without locality (*Reeves ex Lindley*); HONGKONG (*Ford*, 83!). Herb. Kew.

This species was originally described from a drawing.

2. *Cirrhopetalum chinense*, *Lindl. Bot. Reg.* xxviii. *Misc.* p. 37; xxix. t. 49.

*Bulbophyllum chinense*, *Reichb. f. in Walp. Ann.* vi. p. 260.

CHINA: without locality. Herb. Kew.

The precise locality of this species is not recorded. It was introduced to cultivation by Messrs. Loddiges, of Hackney, who flowered it in 1842.

3. *Cirrhopetalum delitescens*, Rolfe, in *Gard. Chron.* (1882) ii. p. 461.

*Bulbophyllum delitescens*, Hance, in *Journ. Bot.* xiv. (1876) p. 44.

HONGKONG (*Ford! Lamont!*). Mus. Brit.; Herb. Kew.

4. *Cirrhopetalum Fordii*, Rolfe, in *Kew Bulletin* (1896), p. 193.

KWANGTUNG (*Ford, 359!*). Herb. Kew.

This species has the general appearance of *C. gamosepalum*, Griff., which, however, has strongly ciliate petals and dorsal sepal.

5. *Cirrhopetalum Henryi*, Rolfe, n. sp.

*Rhizoma* repens, validum. *Pseudobulbi*  $2\frac{1}{4}$ –3 poll. distantes, ovoideo-oblongi,  $\frac{3}{4}$ –1 poll. longi, monophylli. *Folia* oblonga, obtusa, basi in petiolum brevem attenuata, 4–6 poll. longa, 10–16 lin. lata. *Scapi* graciles, 5–8 poll. longi, vaginis 2 lineari-lanceolatis obtecti. *Flores* umbellati. *Bracteæ* lineari-lanceolatæ, acutæ,  $2\frac{1}{2}$ –3 lin. longæ. *Pedicelli* 4–6 lin. longi. *Sepalum* posticum ellipticum, apiculatum, subintegrum, 5-nervium,  $2\frac{1}{2}$ –3 lin. longum; lateralia basi libera, dein fere omnino connata, apice breviter bidentata, 9–11 lin. longa, 4 lin. lata. *Petala* ovata, apiculata, denticulata,  $1\frac{1}{2}$  lin. longa. *Labellum* 2 lin. longum, recurvum, deltoideo-oblongum, obtusum, integrum, canaliculatum. *Columna* 1 lin. longa; dentes subulati, incurvi,  $\frac{1}{4}$  lin. longi.

YUNNAN: Mengtze, mountain rocks, at 5000 ft. (*A. Henry, 11264!*). Flowers salmon-coloured. Herb. Kew.

6. *Cirrhopetalum retusiusculum*, Hemsl. in *Gard. Chron.* (1882) ii. p. 365; *Hook. f. Fl. Brit. Ind.* v. p. 776.

*Bulbophyllum retusiusculum*, Reichb. f. in *Gard. Chron.* (1869) p. 1182; *Trans. Linn. Soc.* xxx. (1874) p. 138.

YUNNAN: Mengtze, south-east mountain forests, at 7000 ft., on trees (*A. Henry, 11106 a!*). Herb. Kew.

Tenasserim.

7. *Cirrhopetalum tigridum*, Rolfe.

*Bulbophyllum tigridum*, Hance in *Journ. Bot.* xxi. (1883) p. 232.

KWANGTUNG: Lofaushan, at 3000 ft. alt. (*Faber, 151!*). Herb. Kew.

8. *Cirrhopetalum Wallichii*, Lindl. in *Wall. Pl. As. Rar.* i. p. 53, t. 67; *Gen. & Sp. Orch.* 59; *Hook. f. Fl. Brit. Ind.* v. p. 776.

YUNNAN: south-east mountain forests, 40 miles from Mengtze, at 7000 ft., on trees (*A. Henry*, 11106!). Herb. Kew.

Temperate Himalaya.

1. *Eria cæspitosa*, *Rolfe*, in *Kew Bulletin* (1896), p. 194.

HAINAN. Flowered at Kew in August 1894, the plant having been received from the Hongkong Botanic Garden, with other Hainan orchids.

An anomalous little species, differing from every other in its tufted habit and absence of pseudobulbs. The flowers are white, with a pair of maroon stripes at the base of the petals and lateral sepals, the dorsal sepal having four. The front lobe of the lip is yellow, and the angles of the side-lobes maroon-purple. It approaches nearest to the section *Bulbodium*.

2. *Eria Corneri*, *Reichb. f.* in *Gard. Chron.* (1878) ii. p. 106.

FORMOSA (*Corner*); KWANGTUNG: Lofaushan (*Ford*, 149!). Herb. Kew.

I have not seen Formosan specimens, but according to Reichenbach it was introduced to cultivation from this island by Mr. Arthur Corner, and flowered with Mr. Charles Leach, of King's Road, Clapham Park.

3. *Eria* (§ *Dendrolirium*) *formosana*, *Rolfe*, in *Kew Bulletin* (1896), p. 194.

FORMOSA: South Cape (*A. Henry*, 1978!). Herb. Kew.

4. *Eria* (§ *Dendrolirium*) *hainanensis*, *Rolfe*, n. sp.

*Rhizoma* crassum. *Pseudobulbi* elliptico-oblongi, subcompressi,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. longi,  $1$ – $1\frac{1}{4}$  poll. lati. *Folia* oblongo-lanceolata, acuta, 5–9 poll. longa,  $\frac{3}{4}$ – $1\frac{1}{2}$  poll. lata. *Scapi* validi,  $1$ – $1\frac{1}{4}$  ped. alti, cum pedicellis et floribus crasse tomentosis; racemi breves, multiflori. *Bracteæ* ovatæ v. ovato-lanceolatæ, acutæ, 6–9 lin. longæ. *Pedicelli* 12–15 lin. longi. Sepalum posticum oblongo-lanceolatum, obtusum, 6–7 lin. longum; lateralia falcato-lanceolata, obtusa, 6–7 lin. longa, basi obliqua. *Petala* lineari-lanceolata, subacuta, 5–6 lin. longa. *Labellum* recurvum, trilobum, 6–7 lin. longum; lobi laterales erecti, semioblongi, angusti, subobtusius; lobus intermedius obovatus, subobtusius v. apiculatus, undulatus, basi angustus; discus tricarinatus, carinis antice verrucosis. *Columna* lata,  $1\frac{1}{2}$  lin. longa. *Mentum* saccato-oblongum, 2 lin. longum.

HAINAN: Hongta, on trees (*Ford*, 474! 574!). Herb. Kew.

Nearly allied to *E. ornata*, Lindl., from Java, *E. armeniaca*, Lindl., from the Philippines, and *E. tomentosa*, Hook. f., from India.

5. *Eria luchuensis*, *Yatabe*, in *Tokyo Bot. Mag.* vii. (1892) p. 131, t. 6.

LUCHU ARCHIPELAGO: at Nishi-omotesima (*Tashiro!*). Herb. Kew.

Allied to the Indian *E. myosurensis*, Lindl.

6. *Eria rosea*, *Lindl. Bot. Reg.* xii. t. 978; *et in Journ. Linn. Soc., Bot.* iii. (1859) p. 50; *Benth. Fl. Hongk.* p. 353.

HONGKONG (*Wilford*, 381! *Champion*, 275!). Mus. Brit.; Herb. Kew.

This species flowered in the garden of the Horticultural Society, at Chiswick, in 1824. It is said to have been introduced from China, by Mr. J. D. Parks. Its Hongkong habitat seems to be the only one certainly known.

7. *Eria pannea*, *Lindl. Bot. Reg.* 1842, *Misc.* p. 74; *et in Journ. Linn. Soc., Bot.* iii. (1859) p. 50; *Hook. f. Fl. Brit. Ind.* v. p. 804.

HAINAN (*Ford*, 571!). Herb. Kew.

India.

8. *Eria sinica*, *Lindl. in Journ. Linn. Soc., Bot.* iii. (1859) p. 48. *Conchidium sinicum*, *Lindl. in Hook. Kew Journ.* vii. (1855) p. 34.

*Eria pusilla*, *Benth. Fl. Hongk.* p. 353 (non *Lindl.*).

HONGKONG (*Champion*, 278! *Hance*, 435!). Mus. Brit.; Herb. Kew.

Quite distinct from the Khasian species, with which Bentham confounds it.

1. *Phreatia elegans*, *Lindl. Gen. & Sp. Orch.* p. 63; *Journ. Linn. Soc., Bot.* iii. (1859) p. 61; *Hook. f. Fl. Brit. Ind.* v. p. 810.

*Thelasis elegans*, *Blume, Mus. Lugd.-Bat.* ii. p. 187.

YUNNAN: Mengtze, S.E. mountain forests, at 5000 ft., on trees (*A. Henry*, 11093!). Flowers white. Herb. Kew.

N. India, Ceylon, and Java.

2. *Phreatia formosana*, *Rolfe, in Ann. of Bot.* ix. (1895) p. 156. FORMOSA: South Cape (*A. Henry*, 1349!). Herb. Kew.

1. *Pachystoma chinense*, *Reichb. f. in Bonplandia*, iii. (1855) p. 251; *Benth. Fl. Hongk.* p. 356.

*Apaturia chinensis*, *Lindl. Gen. & Sp. Orch.* p. 131.

FORMOSA: Tangkang (*A. Henry*, 384!); KWANGTUNG:

near Macao (*Vachel!*); HONGKONG (*Champion, 280! Wright, 521! Tate! Urquhart, 148! Hance, 260! Lamont, 759!*). Mus. Brit.; Herb. Kew.

1. *Spathoglottis pubescens*, *Lindl. Gen. & Sp. Orch.* p. 120; *Journ. Linn. Soc., Bot.* iii. (1859) p. 22; *Kurz, in Journ. of Bot.* xi. (1873) p. 193; *Hook. f. Fl. Brit. Ind.* v. p. 814.

*Spathoglottis Fortunei*, *Lindl. Bot. Reg.* xxxi. t. 19; *Benth. Fl. Hongk.* p. 355.

*Pachystoma Fortunei*, *Reichb. f. in Walp. Ann.* vi. p. 464.

YUNNAN (*Anderson*): on mountain grassy moor at 6500 ft. alt.; at Mengtze, very rare (*Hancock, 84!*), 5000 ft. (*A Henry, 11096!*); HONGKONG (*Fortune, 79! Champion, 279! Hance, 383! 862! Wilford, 285! Wright, 518! Lamont, 758!*). Mus. Brit.; Herb. Kew.

Khasia and Naga Hills to the mountains of Arracau and Burma.

There seems no character to separate the Hongkong plant. The villose lip is found in some Indian specimens.

1. *Phaius grandifolius*, *Lour. Fl. Cochinch.* p. 729; *Lindl. Gen. & Sp. Orch.* p. 126; *Benth. Fl. Hongk.* p. 355; *Benth. Fl. Austral.* vi. p. 304; *Griseb. Fl. Brit. W. Ind.* p. 622.

*Limodorum Tancarvilleæ*, *Banks, ex L'Hérit. Sert. Angl.* p. 28.

*Limodorum Tankervilleæ*, *Dryand., Ait. Hort. Kew.* ed. 1, p. 302, t. 12.

*Limodorum Incarvillei*, *Sw. ex Pers. Syn.* ii. p. 520.

*Bletia Tankervilleæ*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 205; *Bot. Mag.* t. 1924.

*Pachyne spectabilis*, *Salisb. in Trans. Hort. Soc.* i. (1812) p. 299.

*Phaius Tankervillei*, *Blume, Mus. Bot. Lugd.-Bat.* ii. p. 177.

*Calanthe speciosa*, *Vieill. in Ann. Sc. Nat. sér. 4, xvi.* (1861) p. 44.

FORMOSA: Tamsui (*Oldham, 556!*); HONGKONG (*Champion, 283! Urquhart, 151! Wilford, 419! Wright, 519! Hance, 723! Lamont!*). Mus. Brit.; Herb. Kew.

New Caledonia, Rarotonga, and North Australia, both in Queensland and New South Wales; also naturalized in Jamaica and Cuba in the West Indies.

1. *Thunia Marshalliana*, *Reichb. f. in Linnæa*, xli. (1877) p. 65; *Gartenfl.* xxxi. p. 321, t. 1098; *Warn. & Will. Orch. Alb.* iii. t. 130.

YUNNAN: Mengtze, wet mountains at 5000 ft. (*A. Henry, 11091!*). Herb. Kew.

Burma.

1. *Bletia hyacinthina*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 206; *Lindl. Gen. & Sp. Orch.* p. 122; *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 17, t. 6, fig. 1.

*Limodorum striatum*, *Thunb. Fl. Jap.* p. 28.

*Epidendrum tuberosum*, *Lour. Fl. Cochinch.* p. 523, excl. syn.

*Epidendrum striatum*, *Thunb. in Trans. Linn. Soc.* ii. (1794) p. 327; *Thunb. Ic. Jap.* i. t. 9.

*Cymbidium striatum*, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) p. 77.

*Cymbidium hyacinthinum*, *Sm. Exot. Bot.* i. p. 117, t. 60; *Bot. Mag.* t. 1492.

*Gyas humilis*, *Salisb. in Trans. Hort. Soc.* i. (1812) p. 300.

*Bletilla striata*, *Reichb. f. in Bot. Zeit.* xxxvi. (1878) p. 75.

CHEKIANG: Ningpo (*Cooper*!); KIANGSI: Kiukiang (*Shearer*! *Carles*, 242!); HUPEH: Ichang, Nanto, and mountains to northward (*A. Henry*, 492! 1909!); SZECHUEN: Min River and Mt. Omei, at 4500 ft. alt. (*Faber*, 940!); YUNNAN: Mengtze, 6000 ft. (*A. Henry*, 11104 & 11104 C!); KWANGTUNG (*Ford*, 377!). *Mus. Brit.*; *Herb. Kew.*

Japan and Eastern Tibet.

Var. *Gebina*, *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 17, t. 6. fig. 2.

*Bletia Gebina*, *Lindl. in Journ. Hort. Soc.* ii. (1847) p. 307; *Lindl. Bot. Reg.* xxxiii. t. 60.

*Bletilla Gebinæ*, *Reichb. f. in Fl. des Serres*, sér. 1, viii. (1851-53) p. 246.

*Bletilla striata*, var. *Gebinæ*, *Reichb. f. in Bot. Zeit.* xxxvi. (1878) p. 75.

YUNNAN: grassy slopes at 5000-6500 ft., Mengtze (*Hancock*, 98!), Flowers yellowish white; 4700-6000 ft. (*A. Henry*, 11104 A & B!); KWANGSI (*Bourne*!). *Herb. Kew.*

Japan.

1. *Nephelaphyllum chinense*, *Rolfe, in Kew Bulletin* (1896), p. 194.

KWANGTUNG: in rupibus ad Tingushan, West River, Prov. Canton (*Hance*, 17733!). *Mus. Brit.*

2. *Nephelaphyllum cristatum*, *Rolfe, in Kew Bulletin* (1896), p. 194.

HONGKONG (*Ford*, 48 & 254!). *Herb. Kew.*

An interesting outlying species of this small Indo-Malayan genus, known only from the Victoria Peak, where it grows among grasses. It is allied to *N. cordifolium*, *Lindl.*, but has a

much more developed crest. According to Mr. Ford, the sepals and petals are green with purplish streaks near the base, the lip white, passing into purple and then green near the base, the column pinkish, and the leaves and flower-stems mottled with dull purple on an olive-green ground. A plant in the Kew collection flowered in May, 1893.

HANCOCKIA, *Rolfe*.

(Genus novum affine *Nephelaphyllo*.)

*Sepala* subæqualia, libera, subconniventia, omnia angusta. *Petala* sepalis subsimilia. *Labellum* ad basin columnæ breviter adnatum, erectum, longe calcaratum, elliptico-oblongum, subtrilobum, lobis lateralibus subinvolutis; discus tricarinatus. *Columna* erecta, longiuscula, 2-alata, apoda, vertice in clinandrium angustum producta. *Anthera* operculata, apice subobtusa, loculis 2, bene distinctis; pollinia 8, cerea, in quoque loculo 4, compressa, parallela, 2-seriata, appendiculæ lineari sursum adnata.—*Herba* terrestris, repens, epseudobulbosa. *Folia* solitaria, ovata, submembranacea, breviter petiolata. *Scapus* brevis, in vagina scariosa inclusus, uniflorus. *Flores* mediocres.

1. *Hancockia uniflora*, *Rolfe*, sp. unica.

*Rhizoma* repens, gracilis. *Folia* ovata, acuta,  $1\frac{1}{4}$ – $1\frac{3}{4}$  poll. longa, 7–9 lin. lata; petioli 4–8 lin. lati. *Scapi* 2 poll. longi. *Sepala* et *petala* lineari-lanceolata, 9–10 lin. longa. *Labellum* 8 lin. longum, 5 lin. latum; calcar tubulosum, 8 lin. longum. *Columna* 6 lin. longa.

YUNNAN: Mengtze, S.E. mountain forests, at 5000 ft., on ground (*A. Henry*, 11112!). Flowers pink. Herb. Kew.

Allied to *Nephelaphyllum*, but differing in its short, 1-flowered scapes, connivent sepals and petals, and long slender spur. It is dedicated to W. Hancock, Esq., F.L.S., who has sent to Kew several collections of dried plants from China.

1. *Tainia cordifolia*, *Hook. f. in Hook. Ic. Pl.* xix. t. 1861.

FORMOSA: Kelung (*Ford*, 56!). Mus. Brit.; Herb. Kew.

2. *Tainia hongkongensis*, *Rolfe, in Kew Bulletin* (1896), p. 195.

*Ania angustifolia*, *Benth. Fl. Hongk.* p. 356 (non *Lindl.*!).

HONGKONG (*Wilford*, 384! *Wright*, 522! *Hance!* *Ford!*). Mus. Brit.; Herb. Kew.



Very easily distinguished from the Indian *Ania angustifolia*, Lindl., with which Bentham confounded it, by its entire, not trilobed lip.

3. *Tainia macrantha*, *Hook. f. in Hook, Ic. Pl.* xix. t. 1860.

KWANGTUNG: Lofaushan, at 3100 ft. (*Ford*, 153!). Herb. Kew.

1. *Agrostophyllum formosanum*, *Rolfe, in Ann. Bot.* ix. (1895) p. 157.

FORMOSA: South Cape (*A. Henry*, 1350!). Herb. Kew.

1. *Anthogonium gracile*, *Lindl. in Wall. Cat.* n. 7398; *Gen. & Sp. Orch.* p. 426; *Journ. Linn. Soc., Bot.* iii. (1859) p. 22; *Hook. f. Fl. Brit. Ind.* v. p. 822; *Allg. Gartenz.* (1856) p. 218.

*Anthogonium Griffithii*, *Reichb. f. in Bonplandia*, ii. (1854) p. 90; *Walp. Ann.* vi. p. 471.

*Anthogonium* sp., *Griff. Notul.* iii. p. 383; *Ic. Pl. Asiat.* t. 345, fig. 2.

YUNNAN: Mengtze, among grass at 5000 ft. (*Hancock*, 366!), Flowers rose-coloured; grassy mountains at 5500 ft. (*A. Henry*, 11099!), Flowers pink. Herb. Kew.

North India and Burma.

1. *Cœlogyne* (§ *Pleione*) *bulbocodioides*, *Franch. Pl. David.* ii. p. 122 (excl. pl. Delavay).

KWEICHAU (*Faurie ex Franchet*).

Eastern Tibet.

Delavay's Yunnan plant, formerly included here by M. Franchet, belongs to the next species.

I have only seen specimens from Eastern Tibet, but M. Franchet, in a letter, says that no doubt the Kweichau plant is identical.

2. *Cœlogyne* (§ *Pleione*) *Delavayi*, *Rolfe, in Kew Bulletin* (1896), p. 195.

YUNNAN (*Delavay*!). Herb. Kew.

Distinguished from the preceding, with which M. Franchet formerly identified it, by its larger flowers, broader, less acute sepals and petals, and distinctly toothed lamellæ. The specimens were communicated by M. Franchet, with the remark that he now questioned his former determination, because of the characters just pointed out. Another difference is that in *C. bulbocodioides* the petals are only half as broad as the sepals.

3. *Cœlogyne corymbosa*, *Lindl. Fol. Orch., Cœlog.* p. 7; *Reichb. f. in Gard. Chron.* (1876) ii. p. 8; *Rolfe, op. cit.* 1887, ii. p. 73, fig. 15; *Bot. Mag.* t. 6955; *Hook. f. in Fl. Brit. Ind.* v. p. 831.

YUNNAN: Mengting, S. of Red River, at 8000 ft., on trees  
 (A. Henry, 11114!). Herb. Kew.

Temperate Himalaya.

4. *Cœlogyne elata*, *Lindl. Gen. & Sp. Orch.* p. 40; *Kurz, in Journ. Bot.* xi. (1873) p. 193; *Bot. Mag.* t. 5001; *Hook. f. Fl. Brit. Ind.* v. p. 838.

YUNNAN: Momien (*Anderson!*). Herb. Kew.

Tropical Himalaya and Upper Burma.

5. *Cœlogyne fimbriata*, *Lindl. in Bot. Reg.* xi. t. 868; *id. Fol. Orch., Cœl.* p. 12; *Benth. Fl. Hongk.* p. 354; *Hook. f. Fl. Brit. Ind.* p. 836.

HONGKONG (*Champion, 527! Wilford, 101! Hance, 401! Lamont, 755!*); KWANGTUNG: near Canton (*Faber!*). Mus. Brit.; Herb. Kew.

Khasia Hills.

6. *Cœlogyne* (§ *Pleione*) *grandiflora*, *Rolfe*, n. sp.

*Rhizoma* validum. *Pseudobulbi* ovoidei v. ovoideo-oblongi,  $1\frac{1}{4}$ – $1\frac{1}{2}$  poll. longi. *Folia* immatura tantum visa. *Scapi* erecti, 4–5 poll. longi, basi vaginis imbricatis obtecti, uniflori. *Bracteæ* lanceolato-oblongæ, obtusæ,  $1\frac{1}{4}$ – $1\frac{1}{2}$  lin. longæ. *Sepala et petala* subsimilia, lanceolato-oblonga, obtusa,  $2\frac{1}{4}$  poll. longa, 5–6 lin. lata. *Labellum* obscure trilobum, late elliptico-oblongum,  $2\frac{1}{4}$  poll. longum,  $1\frac{3}{4}$  poll. latum, valde fimbriatum, disco medio fere ad apicem 5-carinatum, carinis fimbriatis. *Columna* clavata,  $1\frac{1}{2}$  poll. longa, alata, apice 4 lin. lata.

YUNNAN: Mengtze, northern mountain forests, at 8500 ft.  
 (A. Henry, 11116). Flowers white. Herb. Kew.

A handsome species, apparently most nearly allied to the Himalayan *C. præcox*, Lindl.

7. *Cœlogyne* (§ *Pleione*) *Henryi*, *Rolfe, in Kew Bulletin* (1896), p. 195.

HUPEH: South Patung (A. Henry, 6068, 6068 a!); SZECHUEN (A. Henry, 8826!). Mus. Brit.; Herb. Kew.

Larger in all its parts than the other Chinese species of the

section *Pleione*, and the leaves well developed when the flowers appear.

8. *Cœlogyne* (§ *Pleione*) *pogonoides*, *Rolfe*, in *Kew Bulletin* (1896), p. 196.

*Pogonia* (§ *Eupogonia*), sp., *Hance*, in *Journ. Bot.* xxiii. (1885) p. 247.

ANWHEI: wet rocks near Wuhu, at 3000 ft. alt. (*Bullock*, hb. *Hance*, 22038!); HUPEH: Patung, in high mountains (*A. Henry*, 1473, 3785!). Mus. Brit.; Herb. Kew.

This species differs from *C. bulbocodioides*, which it otherwise much resembles, in having the petals as broad as the sepals, the keels of the lip distinctly crenate, and the leaves partially developed at the time of flowering. *Hance*, in the place above cited, calls it a *Pogonia*, which it is not, though the resemblance is perhaps sufficient to justify the specific name. Mr. *Bullock* records the colour of the flowers as bright scarlet, and Dr. *A. Henry* as pink. The latter states that it is known as "*Pen mu*" drug, the bulbs being used in medicine.

9. *Cœlogyne* (§ *Pleione*) *yunnanensis*, *Rolfe*, n. sp.

*Rhizoma* validum. *Pseudobulbi* parvi, approximati. *Folia* non vidi. *Scapi* graciles, erecti, 4-5 poll. longi, basi vaginis oblongis obtecti, uniflori. *Bracteæ* tubuloso-oblongæ, obtusæ, 10-12 lin. longæ. *Sepala et petala* subsimilia, oblongo-lanceolata, subacuta, 1½ poll. longa, 4 lin. lata. *Labellum* obovato-orbiculare, subintegrum v. obscure trilobum, obtusum, 14 lin. longum et latum, apice fimbriatum; disco medio fere ad apicem acute 5-lamellatum, lamellis subintegris. *Columna* clavata, alata, 9-10 lin. longa, apice 3 lin. lata.

YUNNAN: Mengtze, grassy mountains to north, at 5500 ft. (*A. Henry*, 11113!), Flowers pink; Mountain pastures, 6000-7000 ft. (*Hancock*, 173!), Flowers red to purple. Herb. Kew.

Allied to *C. Delavayi*, *Rolfe*, but different both in the shape of the pseudobulbs and details of the flowers. *A. Henry*, 11115, from south of Red River, at 9000 ft., on rocks, is very similar in the scape and bract, but the pseudobulbs are narrowly oblong. The flowers appear to have been over when specimens were collected, and the species remains doubtful.

1. *Pholidota cantonensis*, *Rolfe*, in *Kew Bulletin* (1896), p. 196.

KWANGTUNG: North River, Canton (*Ford*, 139!). Herb. Kew.

2. **Pholidota chinensis**, *Lindl. in Journ. Hort. Soc.* ii. (1847) p. 308; *Benth. Fl. Hongk.* p. 354.

*Cœlogyne chinensis*, *Reichb. f. in Walp. Ann.* vi. p. 237.

*Cœlogyne Pholas*, *Reichb. f. in Walp. Ann.* vi. p. 237.

*Pholidota Pholas*, *Reichb. f. in Walp. Ann.* vi. p. 237 (in note).

HONGKONG (*Champion*, 276! *Wright*, 516! *Fortune*! *Seemann*, *Hance*, 971!); HAINAN (*B. C. Henry*, 136!). Mus. Brit.; Herb. Kew.

3. **Pholidota Lugardi**, *Rolfe, in Kew Bulletin* (1893), p. 6.

YUNNAN: Mengtze, on a bank at 5000 ft. (*A. Henry*, 11126!). Herb. Kew.

Upper Burma.

The Mengtze specimen is in fruit, but is so similar in general character to the original specimen from Burma as to leave scarcely a doubt as to their identity. The withered flowers persist on the capsules, and proved quite perfect when soaked in hot water and spread out.

4. **Pholidota yunnanensis**, *Rolfe*, n. sp.

*Rhizoma* validum, lignosum. *Pseudobulbi* oblongi v. ovoideo-oblongi,  $\frac{3}{4}$ – $1\frac{1}{2}$  poll. longi, diphylli,  $\frac{1}{2}$ –1 poll. distantes. *Folia* lanceolata, subobtusa, coriacea, 3–5 poll. longa, 4–8 lin. lata. *Scapi* arcuati, graciles, 3–4 poll. longi, basi vaginis oblongis imbricatis obtecti; racemi flexuosi, multiflori. *Bracteæ* elliptico-oblongæ, subobtusæ, convoluto-concavæ, 3–4 lin. longæ. *Pedicelli*  $2\frac{1}{2}$ –4 lin. longi. *Sepala* late ovato-oblonga, obtusa, concava, 2 lin. longa. *Petala* subsimilia. *Labellum* obovatum,  $2\frac{1}{2}$  lin. longum; basi globoso-saccatum, 1 lin. longum; limbus obscure 4-lobus, 2 lin. latus. *Columna* clavata, 1 lin. longa.

YUNNAN: Mengtze, on rocks at 6000–7000 ft. (*Hancock*, 585!), Flowers cream-coloured; Rocky mountains at 6500 ft., on exposed rocks (*A. Henry*, 11144!). Herb. Kew.

Allied to the Indian *Pholidota articulata*, Lindl., but the leaves much narrower, pseudobulbs less distant, and the lip very different in structure.

1. **Calanthe angusta**, *Lindl. Fol. Orch., Calanthe*, p. 7; *Hook. f. Fl. Brit. Ind.* v. p. 849.

YUNNAN: marshy hill-slopes at Mengtze (*Hancock*, 38!). Rare or very local. Flowers pure white. Herb. Kew.

Khasia Hills and Munnipore.

2. *Calanthe arcuata*, Rolfe, in *Kew Bulletin* (1896), p. 196.

HUPEH: Hsingshan (*A. Henry*, 6514!). Herb. Kew.

Allied to the Himalayan *C. brevicornu*, Lindl., but readily distinguished by the structure of the lip, the more slender spur, and other differences. The name is given in allusion to the character of the spur. According to Dr. Henry, the flowers are yellow and purple.

3. *Calanthe buccinifera*, Rolfe, in *Journ. Linn. Soc., Bot.* xxix. (1892) p. 318.

HUPEH: N. Patung (*A. Henry*, 7161!), S. Patung, at 6000 ft. elevation (*A. Henry*, 6064!); SZECHUEN: Mt. Omei, at 5000 ft. elevation, in dense jungle (*Faber*, 953!); without locality (*Pratt*, 72, 102, 683!). Mus. Brit.; Herb. Kew.

Flowers pink, according to Dr. Henry.

4. *Calanthe ecarinata*, Rolfe, in *Journ. Linn. Soc., Bot.* xxix. (1892) p. 318.

SZECHUEN: without locality (*Pratt*, 765!). Herb. Kew.

5. *Calanthe ensifolia*, Rolfe, in *Kew Bulletin* (1896), p. 197.

HUPEH: Chienshih (*A. Henry*, 6005!); SZECHUEN: Mt. Omei, at 4500 ft. (*Faber*, 945!). Herb. Kew.

Remarkable for its long narrow leaves, which much resemble those of *C. Davidi*, Franch., but the lip is very different in shape. Dr. Henry describes the flowers as yellow.

6. *Calanthe formosana*, Rolfe, in *Ann. Bot.* ix. (1895) p. 157.

FORMOSA: South Cape (*A. Henry*, 1347!). Herb. Kew.

7. *Calanthe gracilis*, Lindl. *Gen. & Sp. Orch.* p. 251; *Bot. Mag.* t. 4714; *Hook. f. Fl. Brit. Ind.* v. p. 855.

*Limatodes gracilis*, Lindl. *Fol. Orch., Limatod.* p. 1; *Hook. Kew Journ. Bot.* vii. (1855) p. 36; *Benth. Fl. Hongk.* p. 359.

HONGKONG (*Champion*, 520!). Mus. Brit.; Herb. Kew.

Khasia Hills.

8. *Calanthe Hancockii*, Rolfe, in *Kew Bulletin* (1896), p. 197.

YUNNAN: under shady blocks of rock at 6600 ft., Mengtze (*Hancock*, 78!). Flowers yellow. Very rare. Herb. Kew.

Allied to *C. striata*, R. Br., but with much smaller lip.

9. *Calanthe Henryi*, Rolfe, in *Kew Bulletin* (1896), p. 197.

HUPEH: Changyang (*A. Henry*, 5253 A, 5253 D, 5958 A!). Herb. Kew.

Allied to the Himalayan *Calanthe plantaginea*, Lindl., but with a more lax raceme and a shorter spur. Dr. Henry records the flowers as whitish with a little yellow.

10. *Calanthe herbacea*, Lindl. *Fol. Orch., Cal.* p. 10; *Hook. f. Fl. Brit. Ind.* v. p. 852; *Hook. f. in Ann. Roy. Bot. Gard. Calc.* v. p. 30, t. 44; *King & Pantl., op. cit.* viii. p. 172, t. 233.

*Calanthe elytroglossa*, Reichb. f. ex *Hook. f. Fl. Brit. Ind.* v. p. 853.

YUNNAN: Mengtze, S.E. mountain forests, at 5000 ft. (*A. Henry*, 11130!). Flowers yellowish. *Herb. Kew.*

Sikkim Himalaya.

There remains a slight doubt about this determination, but I fail to distinguish the Yunnan plant from the two cited above, which themselves are considered distinct in the 'Flora of British India,' where, however, they are separated by the interpolation of four other quite distinct species, and are not compared together.

11. *Calanthe lamellosa*, Rolfe, in *Kew Bulletin* (1896), p. 197.

HUPEH: Chiensih (*A. Henry*, 5958!). *Mus. Brit.*; *Herb. Kew.*

Allied to the Himalayan *C. brevicornu*, Lindl., but readily distinguished by its narrower, more elevated membranaceous plates to the lip. Dr. Henry records the flowers as white with a little red and yellow.

12. *Calanthe puberula*, Lindl. *Gen. & Sp. Orch.* p. 252; *Fol. Orch., Cal.* p. 2; *Hook. f. Fl. Brit. Ind.* v. p. 848.

SZETCHUEN: Mount Omei, in dense jungle at 5000 ft. (*Faber*, 953!). *Herb. Kew.*

Subtropical Himalaya from Simla eastward.

13. *Calanthe tricarinata*, Lindl. *Gen. & Sp. Orch.* p. 252; *Fol. Orch., Cal.* p. 2; *Hook. f. Fl. Brit. Ind.* v. p. 847.

YUNNAN: Mengtze, mountain forests, at 7000 ft. (*A. Henry*, 11297!). *Herb. Kew.*

Temperate Himalaya, from Kashmir to Nepal.

I fail to distinguish this from the Himalayan plant. The oldest bud was just on the point of expanding when gathered.

14. *Calanthe veratrifolia*, R. Br. in *Bot. Reg.* vii. sub t. 573; *Bot. Mag.* t. 2615; *Hook. f. Fl. Brit. Ind.* v. p. 851; *Kränzl. in Engl. Bot. Jahrb.* vi. (1885) p. 54.

FORMOSA: South Cape (*A. Henry*, 1327!); HONGKONG

(*Ford*, 89! an *culta*?); LUCHU ARCHIPELAGO: Amami Osima (*Döderlein ex Kränzlin*). Herb. Kew.

Mr. Ford's specimen is unlocalized, and therefore it is not quite clear whether it is indigenous or cultivated.

South India, Ceylon, Malay Islands, and Tropical Australia.

15. *Calanthe yunnanensis*, *Rolfe*, n. sp.

*Folia* 2 v. 3, elliptico- v. oblanceolato-oblonga, acuta, basi in petiolum vaginatum attenuata, 8-14 poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{4}$  poll. lata, basi vaginis latis imbricatis obtecta. *Scapi* 1-2 poll. alti, erecti; racemi 5-8 poll. longi, 7-12-flori. *Bracteæ* ovato-lanceolatæ, acuminatæ, 3-6 lin. longæ. *Pedicelli* 8-12 lin. longi. *Sepala* ovato-lanceolata, acuminata, 9-11 lin. longa. *Petala* oblongo-lanceolata, apiculato-acuminata, 8-10 lin. longa. *Labellum* basi columnæ adnatum, breviter unguiculatum, trilobum, 8-9 lin. longum, 6 lin. latum; lobi laterales reniformi-orbiculares, truncati; lobus intermedius transverse oblongus, emarginatus, subundulatus, basi valide trilamellatus, lamellis crenulatis; calcar oblongum, subobtusum,  $1-1\frac{1}{2}$  lin. longum. *Columna* lata, 3-4 lin. longa.

YUNNAN: Mengtze, mountain forests at 8000-9000 ft. (*Hancock*, 589!), Flowers creamy green, with lilac lip; fragrant of grated cocoa-nut; mountain forests at 8000 ft. (*A. Henry*, -11107!). Herb. Kew.

Allied to the Indian *Calanthe brevicornu*, Lindl., but having larger flowers, with broader segments.

16. *Calanthe* sp.

YUNNAN: Mengtze, E. mountain forests, at 7000 ft. (*A. Henry*, n. 11359!). Herb. Kew.

In fruit only, and therefore indeterminable.

1. *Arundina chinensis*, *Blume*, *Bijdr.* p. 402; *Lindl. Gen. & Sp. Orch.* p. 125; *Hook. & Arn. Bot. Beech. Voy.* p. 217; *Benth. Fl. Hongk.* p. 355; *Hook. f. Fl. Brit. Ind.* v. p. 857.

*Cymbidium Meyenii*, *Schauer*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 433.

*Arundina Meyenii*, *Reichb. f. in Linnæa*, xxv. (1852) p. 227.

*Arundina Philippii*, *Reichb. f. in Linnæa*, xxv. (1852) p. 227; *Seem. Bot. Voy. Herald*, p. 419.

*Arundina pulchella*, *Teijsm. et Binn. in Nederl. Kruidk. Arch.* iii. (1855) p. 400.

*Arundina pulchra*, *Miq. Journ. Bot. Néerl.* i. (1861) p. 90.

CHINA: without locality (*Fortune*, 21, 73!); FOKIEN (*Carles*, 687!), Amoy (*Swinhoe*!), Foochow (*Hurst* ex *A. Henry*, 2078!); FORMOSA: Siko Hill, Tamsui (*A. Henry*, 1386!, 1386 a!); SZECHUEN (*A. Henry*, 8915!); YUNNAN: on mountain grassy slopes at 5700–6000 ft. (*Hancock*, 83!), 5500 ft. (*A. Henry*, 9988!); HONGKONG (*Champion*, 22! *Seemann*, 2402! *Wright*, 517! *Wilford*, 74! *Lamont*, 756! *Urquhart*!); KWANGTUNG: Macao and the islands adjacent (*Vachell*, 82! *Nelson*!). Mus. Brit.; Herb. Kew.

North India, from Sikkim to Khasia.

[*Epidendrum tuberosum*, *Lour. Fl. Cochinch.* p. 523, mentioned as being cultivated in China, is a species of *Phaius*, probably *P. grandifolius*, *Lour.*]

[*Broughtonia chinensis*, *Lindl. ex Benth. in Hook. Loud. Journ. Bot.* i. (1842) p. 192; *Læliopsis chinensis*, *Lindl. in Paxt. Fl. Gard.* iii. p. 156; *Benth. in Hook. Kew Journ. Bot.* vii. (1855) p. 38; is *Epidendrum non-chinense*, *Reichb. f. in Walp. Ann.* vi. p. 324, a Central American species which through error was assigned a Chinese habitat.]

#### ORCHIDACEÆ: VANDEÆ.

1. *Eulophia Faberi*, *Rolfe, in Kew Bulletin* (1896), p. 198.

HUPEH: Lukan gorge, Yangtse-kiang (*Faber*, 55, 946!), Ichang (*A. Henry*, 494, 3589!); YUNNAN: Red River Valley, Manpan, on grassy hills at 2000 ft. (*A. Henry*, 10612!). Flowers red. Herb. Kew.

This species resembles *E. campestris*, *Wall.*, from the plains of India and Afghanistan. The flowers, however, are rather larger and less numerous, the details of the lip somewhat different, and the spur longer.

2. *Eulophia flava*, *Hook. f. Fl. Brit. Ind.* vi. p. 7.

*Cyrtopera flava*, *Lindl. Gen. & Sp. Orch.* p. 189; *Royle, Illustr.* p. 370, t. 88. fig. 1; *Benth. Fl. Hongk.* p. 357; *Reichb. f. in Gard. Chron.* (1870) p. 1407.

HONGKONG (*Wright*, 520!). Herb. Kew.

India, from near Garhwal to Nepal, also Travancore.

3. *Eulophia formosana*, *Rolfe.*

*Cyrtopera formosana*, *Rolfe, in Kew Bulletin* (1896), p. 198.

FORMOSA: South Cape (*A. Henry*, 1974!). Herb. Kew.

Allied to the Indian *C. bicarinata*, *Lindl.*



4. *Eulophia* ? *sinensis*, *Miq. Journ. Bot. Néerl.* i. (1861) p. 91.  
KWANGTUNG: Fukwing (*Krone ex Miquel*).  
Only known to me from the original description.

5. *Eulophia nuda*, *Lindl. Gen. & Sp. Orch.* p. 180; *Hook. f. Fl. Brit. Ind.* vi. p. 5.

YUNNAN: Mengtze, grassy mountains at 5000 ft. (*A. Henry*, 11141!). Flowers pink. *Herb. Kew.*

North and South India and Ceylon.

6. *Eulophia yunnanensis*, *Rolfe*, n. sp.

*Scapi* erecti, 1-1½ ped. alti, basi vaginis oblongo-lanceolatis acutis subimbricatis tecti; racemi 6-9-flori. *Bracteæ* lanceolatae, acutae, ½-1¼ poll. longae. *Pedicelli* 8-10 lin. longi. *Sepala* oblongo-lanceolata, subacuta, 10-11 lin. longa. *Petala* subsimilia. *Labellum* oblongum, subobtusum, obscure trilobum, 9 lin. longum, 4-5 lin. latum, ecalcaratum; lobi laterales minuti, obtusi; discus bicarinatus. *Columna* clavata, 6 lin. longa, subapoda.

YUNNAN: Mengtze, rocky mountains at 6000 ft. (*A. Henry*, 11125!). Flowers yellow. *Herb. Kew.*

This apparently belongs to the section *Cyrtopera*, but the foot of the column is nearly obsolete, by which character, together with the spurless lip, it is readily distinguished from other Asiatic species.

[*Cymbidium eburneum*, *Lindl.*, is erroneously recorded from China, in *Walp. Ann.* i. p. 784.]

1. *Cymbidium ensifolium*, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) p. 77; *Lindl. Gen. & Sp. Orch.* p. 162; *Bot. Mag.* t. 1751; *Bot. Reg.* t. 1976 (var. *estriatum*, *Lindl.*); *Benth. Fl. Hongk.* p. 357; *Debeaux, Fl. de Shangh.* p. 62; *Kränzl. in Engl. Bot. Jahrb.* vi. (1885) p. 54 (*excl. syn. C. sinense*).

*Epidendrum ensifolium*, *Linn. Sp. Pl.* ed. 1, p. 954; *Smith, Spicileg.* p. 22, t. 24; *Lour. Fl. Cochinch.* p. 524; *Andr. Bot. Rep.* v. t. 344.

*Limodorum ensatum*, *Thunb. Fl. Jap.* p. 29.

*Epidendrum sinense*, *Red. Lil.* ii. t. 113 (non *Andr.*).

*Cymbidium micans*, *Schauer, in Nov. Act. Nat. Cur.* xix. *Suppl.* i: (1843) p. 433.

*Cymbidium xiphiifolium*, *Lindl. Bot. Reg.* vii. t. 529; *Gen. & Sp. Orch.* p. 162; *Reichb. f. in Bonplandia*, iii. (1855) p. 250; *Benth. Fl. Hongk.* p. 357.

CHINA: without locality (*Osbeck*); KWANGTUNG: Macao (*Meyen!*), Lienchau, Canton (*Lamont!*); HONGKONG (*Champion*),

525! *Lamont*!); LUCHU ARCHIPELAGO: Amami Osima (*Döderlein ex Kränzlin*). Mus. Brit.; Herb. Kew.

Cambodia, CochinChina, and Japan.

*Cymbidium xiphiifolium* is only a form of *C. ensifolium* with unstriped sepals and petals, with which it is connected by intermediate forms. It is absolutely identical in other respects. But *C. sinense*, Willd., which has also been united with it by several authors, has far broader leaves, various floral differences, quite different geographical distribution, and is certainly distinct. There is no evidence of the occurrence of *C. ensifolium* proper in India.

2. **Cymbidium Faberi**, *Rolfe, in Kew Bulletin* (1896), p. 198.

CHEKIANG: Tientai Mt., 2000 ft. (*Faber, 94*!); SZECHUEN: S. Wushan (*A. Henry, 5515*!). Herb. Kew.

This has the habit of the Himalayan *C. cyperifolium*, Wall., but the flowers are quite distinct from it and from every other species known to me. Dr. Henry records the flowers as yellow.

3. **Cymbidium floribundum**, *Lindl. Gen. & Sp. Orch.* p. 162 (in note). Described from a drawing formerly in the Horticultural Society's Library.

CHINA: without precise locality; YUNNAN: Mengtze, mountain gullies, at 6000 ft. (*Hancock, 580*!). Flowers chocolate and lilac. Herb. Kew.

This species, hitherto only known from an unlocalized Chinese drawing, has now been rediscovered in Yunnan. It is allied to *C. sinense*, Willd., but has a shorter, dense raceme, of more numerous flowers.

4. **Cymbidium grandiflorum**, *Griff. Notul.* iii. p. 342;  *Ic. Pl. Asiat.* t. 321;  *Hook. f. Fl. Brit. Ind.* vi. p. 12.

*Cymbidium Hookerianum*, *Reichb. f. in Gard. Chron.* (1866) p. 7; *Batem. in Bot. Mag.* t. 5574.

*Cymbidium giganteum*, *Lindl. in Journ. Linn. Soc., Bot.* iii. (1859) p. 29, in part.

*Cymbidium* sp., *Griff. Itin. Notes*, p. 145.

YUNNAN: Mengtze (*Hancock, 583*!); wooded cliff at 5000 ft., on a tree (*A. Henry, 11097*!). Herb. Kew.

Eastern Himalaya.

5. **Cymbidium Leachianum**, *Reichb. f. in Gard. Chron.* (1878) ii. p. 106.

FORMOSA (*Corner ex Reichenbach*).

Only known to me from the original description.

6. *Cymbidium longifolium*, *D. Don, Prodr. Fl. Nepal.* p. 36; *Lindl. Gen. & Sp. Orch.* p. 163; *Hook. f. Fl. Brit. Ind.* vi. p. 13.

*Cymbidium erythræum*, *Lindl. in Journ. Linn. Soc., Bot.* iii. (1859) p. 30.

SZECHUEN: without locality (*Pratt!*); YUNNAN: Mengtze, mountains to the south-west, at 5000 ft. (*A. Henry, 11000, 11371!*), on cliffs in forest, at 6000 ft. (*A. Henry, 11371!*). *Herb. Kew.*

Subtropical Himalaya, from Kumaon eastwards.

7. *Cymbidium pendulum*, *Sw. in Nov. Act. Soc. Sc. Upsal.* vi. (1799) p. 73; *Lindl. Gen. & Sp. Orch.* p. 165.

*Epidendrum pendulum*, *Roxb. Pl. Coromand.* i. p. 35, t. 44.

*Cymbidium aloifolium*, *Lodd. Bot. Cab.* x. t. 967, excl. syn.; *Hook. f. Fl. Brit. Ind.* vi. p. 10, excl. syn.; *Epidendrum aloifolium*, *Linn., non Sw.*

CHINA: without locality (*Potts!*). Cult. sp. in *Herb. Lindl.*, at *Kew.*

India from Nepal eastward to Assam, and southward to Tenasserim and the Andaman Islands.

This species has been much confused with *C. aloifolium*, *Sw.*, which is limited to South-western India and Ceylon, and readily distinguished by the longer, acute front lobe of the lip. The Chinese habitat remains doubtful. Messrs. Loddiges received it from China in 1790, and Potts also sent it to the Horticultural Society, where it flowered in 1821, a specimen being preserved in Lindley's Herbarium. Besides which there are Chinese drawings in the collection at Kew. It may, however, be only a cultivated plant in China, as there are no dried specimens at Kew, and the collection of Chinese drawings contains some undoubted garden plants.

8. *Cymbidium sinense*, *Willd. Sp. Pl.* iv. p. 111; *Lodd. Bot. Cab.* i. t. 37; *Lindl. Gen. & Sp. Orch.* p. 162; *Lindl. in Journ. Linn. Soc., Bot.* iii. (1859) p. 30.

*Epidendrum sinense*, *Andr. Bot. Rep.* iii. t. 216; *Bot. Mag.* t. 888.

*Cymbidium fragrans*, *Salisb. in Trans. Hort. Soc.* i. (1812) p. 298.

*Cymbidium ensifolium*, *Hook. f. Fl. Brit. Ind.* vi. p. 13, in part.

CHINA: without locality (*Slater, ex Andrews*).

Khasia Hills and Ceylon.

This species has been confused with *C. ensifolium*, *Sw.*, by various authors, but is easily distinguished by its much broader leaves, and larger, more numerous and differently coloured

flowers. I have not seen wild Chinese specimens, but there is a fine coloured drawing from that country in the collection of drawings at Kew. It appears to be common in the Khasia Hills. The Ceylon plant has slightly narrower leaves.

9. *Cymbidium* sp.

FORMOSA: South Cape (*A. Henry*, 1352!). Herb. Kew.

Allied to *C. pendulum*, Sw., but with leaves less than half as broad. The specimen is in fruit, and the two remaining flowers have lost their lips.

1. *Geodorum dilatatum*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 207; *Lodd. Bot. Cab.* t. 1797; *Bot. Reg.* t. 675; *Lindl. Gen. & Sp. Orch.* p. 175; *Fol. Orch., Geod.* p. 1; *Hook. f. Fl. Brit. Ind.* vi. p. 17.

*Geodorum fucatum*, *Lindl. Bot. Reg.* t. 1687; *Fol. Orch., Geod.* p. 2.

YUNNAN: Hsinkai, Red River, in shady woods (*A. Henry*, 11103!), Flowers white; HONGKONG (*Ford*!). Herb. Kew.

India and Ceylon.

There has been some confusion between this and the allied species, and Lindley considered the above-cited names to represent two distinct species, though I fail to separate them. It is quite possible that plants belonging to the allied *G. purpureum*, *R. Br.*, may be included in some of the above references.

2. *Geodorum formosanum*, *Rolfe, in Ann. Bot.* ix. (1895) p. 157.

FORMOSA: Bankinsing (*A. Henry*, 1137!); South Cape (*A. Henry*, 1375!). Herb. Kew.

3. *Geodorum semicristatum*, *Lindl. Fol. Orch., Geod.* p. 2.

*Tropidia grandis*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 128; *Journ. Bot.* xi. (1873) p. 153.

FORMOSA: Ape's Hill, Takow (*Playfair*, n. 96!); HONGKONG (*Hance*, 13343!). Mus. Brit.; Herb. Kew.

Philippines.

1. *Plocoglottis*? sp.

SZECHUEN: Mount Omei, at 4000 ft. (*Faber*, 947!). Herb. Kew.

Apparently belongs to this genus, but all the flowers are badly eaten.

1. **Luisia Hancockii**, *Rolfe*, in *Kew Bulletin* (1896) p. 199.

*Luisia teres*, *Maxim.* in *Bull. Soc. Nat. Mosc.* (1879) p. 60 (non *Blume*).

CHEKIANG: western hills, Ningpo, growing on a wax-tree (*Stillingia sebifera*) beside the river—very rare (*Hancock*, 22!); on bare rocks (*Hickin*!), *Rock Lan* of the natives. Mus. Brit.; Herb. Kew.

In habit similar to *L. teres*, *Blume*, but more slender, the flowers smaller, and the lip differently shaped.

2. **Luisia Morsei**, *Rolfe*, n. sp.

Herba circa  $\frac{1}{2}$  ped. alta. *Caulis* validus. *Folia* rigida, teretia, 2–4 poll. longa; *vaginæ* corrugatæ. *Racemi* breves, pauciflori. *Bracteæ* late ovatæ, subacutæ, divaricatæ, rigidæ, 1 poll. longæ. *Pedicelli* circa 4 lin. longi. *Sepala* et *petala* subæqualia, oblonga, subobtusata, 3 lin. longa. *Labellum*  $3\frac{1}{2}$  lin. longum; *hypochilium* late triangulare,  $2\frac{1}{2}$  lin. latum; *epichilium* late rhomboideo-ovatum, retusum,  $2\frac{1}{2}$  lin. latum. *Columna* brevissima. *Capsula* clavata,  $\frac{1}{2}$  poll. longa.

KWANGSI: Lungchao (*Morse*, 101, herb. *A. Henry*!). Herb. Kew.

Nearest allied to *L. teretifolia*, *Gaud.* (at least as to the Indian plant so named), but the leaves are shorter and stouter, and the flowers much larger.

3. **Luisia teres**, *Blume*, *Rumphia*, iv. p. 50, in nota; *Blume*, *Mus. Bot. Lugd.-Bat.* i. p. 64; *Lindl. Fol. Orch., Luisia*, p. 2; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 28.

*Epidendrum teres*, *Thunb. Fl. Jap.* p. 30; *Thunb. Ic. Pl. Jap.* i. t. 7.

*Cymbidium triste*, *Lindl. Gen. & Sp. Orch.* p. 167, ex parte, non *Willd.*

SZECHUEN: without locality (*Pratt*!). Herb. Kew.

Japan.

Readily distinguished by the bifurcate lip with acute lobes.

4. **Luisia teretifolia**, *Gaud. in Freyc. Voy., Bot.* p. 427, *Atlas*, t. 37; *Blume, Rumphia*, iv. p. 50, t. 194, fig. 3, t. 197, fig. D; *Lindl. Fol. Orch., Luisia*, p. 2; *Hook. f. Fl. Brit. Ind.* vi. p. 22.

HAINAN (*A. Henry*, 8686!). Herb. Kew.

India and Malaya, Marianne Islands, and perhaps New Caledonia.

The specimen is in young fruit, but the lip seems identical with that of the Indian plant. The New Caledonian locality remains doubtful, as the specimen has lost all its flowers.

5. **Luisia** sp.

FORMOSA: South Cape (*A. Henry*, 695!). Herb. Kew.

Possibly *L. teretifolia*, Gaud., but impossible to determine certainly without flowers.

[**Luisia** sp., *Benth. Fl. Hongk.* p. 358, from rocks on Mount Victoria, Hongkong (*Champion*, 526!), is without flowers, but is evidently *Sarcanthus teretifolius*, Lindl.]

1. **Diploprora Championi**, *Hook. f. Fl. Brit. Ind.* vi. p. 26; *Hook. Ic. Pl.* t. 2120.

*Cotonia Championi*, *Lindl. in Hook. Kew Journ. Bot.* vii. (1855) p. 35; *Journ. Linn. Soc., Bot.* iii. (1859) p. 39; *Benth. Fl. Hongk.* p. 357.

*Luisia bicaudata*, *Thw. Enum. Pl. Zeyl.* p. 302.

*Vanda bicaudata*, *Thw., l. c.* p. 429.

*Sarcanthus rostratus*, *Reichb. f. in Seem. Bot. Herald*, p. 419, ? non *Lindl.*

FORMOSA: Balkinsing mountains (*A. Henry*, 898, 1606!); HONGKONG (*Champion*, 227! *Harland*, 438!). Mus. Brit.; Herb. Kew.

Bhotan, Khasia, Tenasserim, and Ceylon.

The *Sarcanthus* above cited is doubtfully included by Bentham, but there is no specimen so named in Hance's herbarium.

1. **Phalænopsis Aphrodite**, *Reichb. f. in Hamb. Gartenz.* 1862, p. 35; *Xen. Orch.* ii. p. 6.

*Phalænopsis amabilis*, *Lindl. Bot. Reg.* 1838, t. 34; *Bot. Mag.* t. 4297, non *Blume*.

FORMOSA: South Cape (*A. Henry*, 1705!). Herb. Kew.

Philippines.

1. **Doritis Wightii**, *Benth. in Benth. et Hook. f. Gen. Pl.* iii. p. 574; *Hook. f. Fl. Brit. Ind.* vi. p. 32.

*Phalænopsis Wightii*, *Reichb. f. in Bot. Zeit.* xx. (1862) p. 214.

*Aërides latifolium*, *Thw. Enum. Pl. Zeyl.* p. 429.

HAINAN: Hongtai, on trees (*Ford*, 475!). Herb. Kew.

India and Ceylon.

1. **Sarcochilus auriferus**, *Rolfe*.

*Dendrobium auriferum*, *Lindl. Gen. & Sp. Orch.* p. 83; *Bot. Reg.* xx., *Misc.* p. 59.

CHINA: without locality (*Reeves*!). Herb. Kew.

Based on a drawing formerly in the possession of the Horticultural Society. According to a copy in Lindley's Herbarium it is evidently a *Sarcochilus* of the section *Dendrocolla*, though

I am unable to identify it with any dried specimen, or indeed to find any other evidence about it.

2. *Sarcochilus hainanensis*, Rolfe, in *Kew Hand-list, Orch.* p. 201; *Bot. Mag.* t. 7489; Rolfe, in *Kew Bulletin* (1896), p. 199.

HAINAN (*Ford*, 572!). Flowered also at Kew in July 1891, the plant having been received from the Hongkong Botanic Garden, with other Hainan Orchids. *Herb. Kew.*

Allied to *S. Arachnites*, Reichb. f., but the side-lobes of the lip are more truncate and the front lobe distinctly triangular. The flowers are light yellow, except the front lobe of the lip, which is white, and a few buff markings which occur in the throat and on the sides of the lip.

3. *Sarcochilus luniferus*, Benth. ex Hook. f. *Bot. Mag.* t. 7044; *Hook. f. Fl. Brit. Ind.* vi. p. 37.

*Thrixspermum luniferum*, Reichb. f. in *Gard. Chron.* (1868) p. 786.

YUNNAN: Mengtze, south-east mountain forests, at 5000 ft., on trees (*A. Henry*, 11129!). *Herb. Kew.*

1. *Aërides odoratum*, Lour. *Fl. Cochinch.* p. 525; *Bot. Mag.* t. 4139; *Hook. f. Fl. Brit. Ind.* vi. p. 47.

CHINA: without locality (ex *Loureiro*).

*Loureiro* gives China as a habitat of this species, but I have seen no specimen from there.

India, Malaya, and Cochin China.

1. *Renanthera coccinea*, Lour. *Fl. Cochinch.* p. 521; *Lindl. Gen. & Sp. Orch.* p. 217; *Bot. Reg.* t. 1131; *Bot. Mag.* t. 2997-8; *Pact. Mag. Bot.* iv. p. 49, with plate; *Hook. f. Fl. Brit. Ind.* vi. p. 48.

CHINA: without locality (*Millett*!); HAINAN: 20 miles north of Hoihow, on rocks (*A. Henry*!). *Herb. Kew.*

Tenasserim and Cochin China.

1. *Vanda concolor*, Blume, *Rumphia*, iv. p. 49, in nota; *Lindl. in Pact. Fl. Gard.* ii. p. 20; *Lindl. Fol. Orch., Vanda*, p. 3; *Sallier, in Orchidophile*, 1887, p. 144, cum ic.

*Vanda Roxburghii*, var. *unicolor*, *Hook. Bot. Mag.* t. 3416.

*Vanda furva*, *Lindl. Bot. Reg.* xxx., *Misc.* p. 39, ex parte.

CHINA: without locality (*Loddiges*!); YUNNAN: on lofty trees in a hot rocky glen at Mengtze (*Hancock*, 117!); woods at 4000-4700 ft. (*A. Henry*, 11092!). *Mus. Brit.*; *Herb. Kew.*

The precise locality of this plant is not known, but there are

several Chinese drawings of it in the Kew collection. It has been confused with the Indian *V. Roxburghii*, R. Br., and the Moluccan *V. furva*, Lindl.

2. **Vanda hainanensis**, Rolfe, in *Kew Bulletin* (1896), p. 199.  
HAINAN (*B. C. Henry*, 37!). Herb. Kew.

A very interesting species, belonging to the section *Anota*, which only contains two others, namely, the Indian *V. densiflora*, Lindl., and the Philippine *V. violacea*, Lindl. According to the Rev. B. C. Henry, the flowers are white and purple, and fragrant.

1. **Saccolabium hainanense**, Rolfe, in *Kew Bulletin* (1895), p. 284.

HAINAN (*B. C. Henry*!). Herb. Kew.

Flowered at Kew in March, 1895.

1. **Acampe multiflora**, Lindl. *Fol. Orch.*, *Acampe*, p. 1; *Reichb. in Walp. Ann.* vi. p. 872; *Benth. Fl. Hongk.* p. 358.

*Vanda multiflora*, Lindl. *Collect. Bot.* t. 38.

CHINA: without locality (*Reeves*!); HONGKONG (*Champion*, 528! *Lamont*, 764! *Hance*, 1334!). Mus. Brit.; Herb. Kew.

Allied to the Indian *A. longifolia*, Lindl.

1. **Sarcanthus cerinus**, Rolfe.

*Cleisostoma cerinum*, *Hance*, in *Journ. Bot.* xx. (1882) p. 359.

KWANGTUNG: Lofaushan (*Faber*!). Mus. Brit.

The spur has the longitudinal plate of *Sarcanthus*, as is also the case with the two following species.

2. **Sarcanthus elongatus**, Rolfe, n. sp.

*Caules* 1 ped. longi v. ultra. *Folia* teretia, gracilia,  $3\frac{1}{2}$ - $5\frac{1}{2}$  poll. longa. *Racemi* pauce ramosi, 8-9 poll. longi, multiflori. *Bracteæ* subulato-lanceolatæ, acutæ, divaricatæ,  $\frac{3}{4}$ -1 lin. longæ. *Pedicelli* 3 lin. longi. *Sepalum* posticum ovatum, subobtusum, 1 lin. longum; lateralibus oblique ovato-oblonga, subacuta,  $1\frac{1}{2}$  lin. longa. *Petala* lineari-oblonga, subacuta,  $1\frac{1}{4}$  lin. longa. *Labellum* trilobum; lobi laterales triangulares, subobtusi; lobus intermedius triangulari-ovatus, acutus,  $1\frac{1}{4}$  lin. longus; discus subcarnosus; calcar late saccatum, obovoideum,  $1\frac{1}{2}$  lin. longum. *Columna*  $\frac{3}{4}$  lin. longa.

HAINAN (*Ford*, 573!). Flowers lilac. Herb. Kew.

Allied to *S. filiformis*, Lindl., but markedly different in the details of the lip.



3. *Sarcanthus Fordii*, Rolfe.

*Cleisostoma Fordii*, Hance, in *Journ. of Bot.* xiv. (1876) p. 45.

HONGKONG (*Ford!*). Mus. Brit.

4. *Sarcanthus formosanus*, Rolfe, in *Kew Hand-list, Orch.* p. 201.

*Cleisostoma formosana*, Hance, in *Journ. Bot.* xxii. (1884) p. 364.

FORMOSA (*Oldham*, 557!); near Tamsui (*Ford!*); Kelung (*Ford*, 93!). Mus. Brit.; Herb. Kew.

5. *Sarcanthus hongkongensis*, Rolfe, in *Kew Hand-list, Orch.* p. 201 (name only); *Kew Bulletin* (1898), p. 198.

HONGKONG (*Ford!*). Herb. Kew.

Allied to the Indian *S. filiformis*, Lindl. The flowers are pale lilac, with the column, the side-lobes of the lip, and base of the front lobe bright purple.

6. *Sarcanthus paniculatus*, Lindl. in *Bauer Ill. Orch. Pl., Genera*, t. 9; *Lindl. Gen. & Sp. Orch.* p. 233.

*Aërides paniculatum*, Ker, in *Bot. Reg.* iii. t. 220.

*Vanda paniculata*, R. Br. in *Bot. Reg.* vi. sub t. 506 et *Notis ad calcem vol.* [p. 3].

CHINA: without locality (*Banks ex Lindley*). Cultivated specimens in Herb. Kew.

Introduced by Sir Joseph Banks, and flowered in his collection about the year 1817. Living plants have since been received at Kew from China, but we are still ignorant of its precise habitat.

7. *Sarcanthus rostratus*, Lindl. *Collect. Bot.* p. 39, fig. B; *Bot. Reg.* t. 981.

*Vanda recurva*, Hook. *Exot. Fl.* iii. t. 187.

*Vanda rostrata*, Lodd. *Bot. Cab.* t. 1008.

CHINA: without locality, ex *Lindley*. Cultivated specimens in Herb. Kew.

Received from China by the Horticultural Society of London in 1821. It is quite distinct from the Indian *Sarcanthus oxyphyllus*, Wall., with which Lindley (*Bot. Reg.* xxx. *Misc.* p. 58) confounded it.

8. *Sarcanthus succisus*, Lindl. *Bot. Reg.* xii. t. 1014.

CHINA: without locality (*Parks ex Lindley*).

Brought from China for the Horticultural Society of London by Mr. John Damper Parks, in 1824. I have not seen a specimen.

9. **Sarcanthus teretifolius**, *Lindl. Gen. & Sp. Orch.* p. 234; *Bot. Mag.* t. 3571; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 127; *Belg. Hort.* x. p. 129, t. 9, fig. 3.

*Vanda teretifolia*, *Lindl. Collect. Bot.* t. 6; *Bot. Reg.* viii. t. 676.

*Luisia* sp., *Lindl. Fol. Orch., Luisia*, p. 1; *Benth. Fl. Hongk.* p. 358.

CHINA: without locality (*Brooks*); HONGKONG (*Hance*, 833! *Lamont*, 764 A! *Ford*, 568!). *Mus. Brit.*; *Herb. Kew.*

Introduced from China by W. Brooks, and flowered in the collection of Mr. Cattley, at Barnet, about the year 1820.

There can be no doubt that the *Luisia* above cited belongs here, as Hance suspected. Champion's specimen is flowerless, but the roots are produced all the way up the stem, as in this, not as in *Luisia teres*, Blume; while the habit of the plant and the texture and striæ of the leaf-sheaths are equally sufficient to settle the matter, in spite of the general resemblance of the two plants. Champion's and Hance's specimens were gathered at the same spot, while the *Luisia* is not known from Hongkong.

1. **Cleisostoma virginale**, *Hance, in Journ. Bot.* xv. (1877) p. 38. *Appendicula* sp., *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 128 (species secunda).

HONGKONG (*Bowring*! *Ford*!). *Mus. Brit.*; *Herb. Kew.*

1. **Ornithochilus fuscus**, *Wall. ex Lindl. Gen. & Sp. Orch.* p. 242; *Hook. f. Fl. Brit. Ind.* vi. p. 76.

*Ornithochilus eublepharon*, *Hance, in Journ. Bot.* xxii. (1884) p. 364.

*Aërides difforme*, *Wall. ex Lindl. Gen. & Sp. Orch.* 1884, p. 242.

*Aërides Hystrix*, *Lindl. in Journ. Linn. Soc., Bot.* iii. (1859) p. 42 (non *Gen. & Sp. Orch.*).

KWANGTUNG: Lofaushan (*Ford*, 92!). *Mus. Brit.* *Herb. Kew.*

Tropical Himalaya and Tennasserim.

I can detect no difference between the Chinese and Indian plants.

1. **Angræcum falcatum**, *Lindl. Gen. & Sp. Orch.* p. 237 (in syn.); *Benth. et Hook. f. Gen. Plant.* iii. p. 583.

*Orchis falcata*, *Thunb. Fl. Jap.* p. 26.

*Limodorum falcatum*, *Thunb. Ic. Pl. Japon.* dec. 1, t. 6; *Bot. Reg.* t. 283; *Bot. Mag.* t. 2097.

*Œceoclades falcata*, *Lindl. Gen. & Sp. Orch.* p. 237; *S. Moore, in Journ. Bot.* xv. (1878) p. 138; *Franch. et Savat. Enum. Pl. Japon.* ii. p. 28 *excl. syn.* *Aërides japonicum*).

*Aërides Thunbergii*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 205.

CHEKIANG: Hangchau (*Moule*!); Ningpo (*Everard*!); Ning-kongjao, near Ningpo (*Medhurst*!). *Herb. Kew.*

Also found in Japan.

There has been some confusion with this species in the books, but the above synonymy I believe is correct.

1. **Appendicula bifaria**, *Lindl. ex Benth. in Kew Journ. Bot.* vii. (1855) p. 35; *Benth. Fl. Hongk.* p. 358; *Reichb. f. in Walp. Ann.* vi. p. 893; *Hook. f. Fl. Brit. Ind.* vi. p. 82.

HONGKONG (*Champion*, 282! *Ford*! *Bowring*!). *Mus. Brit.*; *Herb. Kew.*

India, Tenasserim, and Singapore.

1. **Thelasis hongkongensis**, *Rolfe, in Kew Bulletin* (1896), p. 199.

*Thelasis pygmæa*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 127, non *Lindl.*

HONGKONG (*Hance*, 1287! *Ford*, 13!). *Mus. Brit.*; *Herb. Kew.*

This is nearly allied to the Philippine *T. triptera*, *Reichb. f.*, which, however, has larger flowers, and the sheathing bracts of the scape loose and twice as broad. The Indian *T. pygmæa*, *Lindl.*, to which it has been doubtfully referred, is still more distinct.

#### ORCHIDACEÆ: NEOTTIEÆ.

1. **Galeola Faberi**, *Rolfe, in Kew Bulletin* (1896), p. 200.

SZECHUEN: Mount Omei, 7000 ft. alt. (*Faber*!). *Herb. Kew.*

A very distinct species allied to the Himalayan *G. Lindleyana*, *Reichb. f.*, but with far narrower segments, a much shorter ovary, and quite different aspect generally.

1. **Vanilla** sp.

FORMOSA: Bankinsing (*A. Henry*, 479!). *Herb. Kew.*

1. **Tropidia angulosa**, *Blume, Coll. Orch. Archip. Ind.* p. 122; *Hook. f. Fl. Brit. Ind.* vi. p. 92.

*Cnemidia angulosa*, *Lindl. Gen. & Sp. Orch.* p. 463.

*Decaisnea angulosa*, *Lindl. in Wall. Cat.* n. 7388.

*Tropidia Govindooi*, *Blume, Coll. Orch. Archip. Ind.* p. 122.

*Govindooia nervosa*, *Wight, Ic.* vi. t. 2090.

*Tropidia semilibera*, *Blume, Coll. Orch. Archip. Ind.* p. 122.

*Cnemidia semilibera*, *Lindl. Gen. & Sp. Orch.* p. 463.

FORMOSA: Ape's Hill, Takow, in dark situations (*A. Henry*, 1905). Flowers white. *Herb. Kew.*

India and Philippines.

2. *Tropidia formosana*, *Rolfe, in Ann. of Bot.* ix. (1895) p. 158.

FORMOSA: Bankinsing (*A. Henry*, 1573!). *Herb. Kew.*

3. *Tropidia hongkongensis*, *Rolfe.*

*Tropidia curculigoides*, *Benth. Fl. Hongk.* p. 359, excl. syn., non *Lindl.*

HONGKONG (*Champion*, 522!). *Herb. Kew.*

Bentham unites both this and the Philippine *T. septemnervis*, *Reichb. f. (Ptychochilus septemnervis, Meyen)*, with the Himalayan *T. curculigoides*, *Lindl.*, and in a note further confounds it with species from Malaya and Ceylon. The inflorescence is not in good condition, but the leaves are distinctly narrower than in both the Himalayan and Philippine plants.

[*Tropidia grandis*, *Hance*, is *Geodorum semicristatum*, *Lindl.*]

1. *Neottia camtschatica*, *Spreng. Syst. Veg.* iii. p. 707; *Ledeb. Fl. Ross.* iv. p. 82.

*Ophrys camtschatea*, *Linn. Sp. Pl.* ed. 1, p. 948.

*Epipactis camtschatea*, *Sw. in Vet.-Acad. Nya Handl. Stockh.* xxi. (1800) p. 232.

*Serapias camtschatea*, *Steud. Nomencl. Bot.* ed. 1, p. 766.

*Neottia kamtschatica*, *Lindl. Gen. & Sp. Orch.* p. 458.

*Neottia camtschatea*, *Reichb. Ic. Fl. Germ.* xiii-xiv. p. 146, t. 478. fig. 1, 2.

KANSUH: Tangut (*Przewalski*!). *Herb. Kew.*

Siberia.

2. *Neottia micrantha*, *Lindl. Gen. & Sp. Orch.* p. 458; *Ledeb. Fl. Ross.* iv. p. 82.

HUPEH (*A. Henry*, 6877! 6877 a!). *Herb. Kew.*

Siberia.

A rather anomalous species.

1. *Listera grandiflora*, *Rolfe, in Kew Bulletin* (1896), p. 200.

HUPEH: Fang (*A. Henry*, 6876!); SZECHUEN: Mount Omei, in dark damp place at 8000-9000 ft. (*Faber*, n. 948!). *Herb. Kew.*

A most distinct species, the largest-flowered one in the genus.

2. *Listera puberula*, Maxim. in *Bull. Acad. Pétersb.* xxix. (1884) p. 204; *Mél. Biol.* xi. p. 843.

KANSUH: Tangut (*Przewalski*!). Herb. Kew.

1. *Spiranthes australis*, Lindl. *Bot. Reg.* x. sub t. 823; *Gen. & Sp. Orch.* p. 464; *Swinhoe, List Pl. Formosa*, p. 4; *Hook. f. Fl. Brit. Ind.* vi. p. 102; *Franch. Pl. David.* i. p. 295; *Palibin, in Act. Hort. Petrop.* xiv. (1895) p. 140.

*Aristotelea spiralis*, Lour. *Fl. Cochinch.* p. 522.

*Neottia sinensis*, Pers. *Syn. Pl.* ii. p. 511.

*Neottia australis*, R. Br. *Prodr.* p. 319.

*Spiranthes pudica*, Lindl. *Collect. Bot.* t. 30.

*Neottia australis*  $\beta$  *chinensis*, Edwards, in *Bot. Reg.* t. 602.

*Spiranthes amœna*, Bunge, *Enum. Pl. Chin. Bor.* p. 63.

*Spiranthes australis*, var. *pudica*, Lindl. *Gen. & Sp. Orch.* p. 465; *Benth. in Hook. Kew Journ. Bot.* vii. (1855) p. 38; *Benth. Fl. Hongk.* p. 360; *Miq. in Journ. Bot. Néerl.* i. (1861) p. 91.

*Gyrostachys australis*, Blume, *Coll. Orch. Archip. Ind.* p. 128, t. 37, fig. 3, t. 38, fig. 3.

CHILHI (*Putjata & Borodowsky, ex Palibin*); COREA (*Wilford, 923*!); KANSUH (*Przewalski*!); SZECHUEN (*Pratt, 626*!); YUNNAN (*Hancock, 516*!); HONGKONG (*Champion, 268*! *Wilford, 397*! *Urquhart*! *Wright, 526*!); FORMOSA (*Oldham, 562*! *Swinhoe*!); LUCHU ARCHIPELAGO (*Beechey*! *Wright, 335-338*!). Also various other collectors. Mus. Brit.; Herb. Kew.

This is probably the most widely distributed Orchid known, as it extends from Jekaterinenburg, on the southern part of the Ural Mountains, and Afghanistan on the west, to Ceylon and South India, Japan, Java, Australia, and New Zealand.

2. *Spiranthes exigua*, Rolfe, in *Kew Bulletin* (1896), p. 200.

HUPEH (*A. Henry, 6585*!). Herb. Kew.

A curious little leafless plant, not nearly allied to any other Old World species. The minute hastate lobes of the lip are easily overlooked, unless very carefully searched for; and the rostellum is curiously recurved at the apex. A single specimen is at present known.

1. *Physurus chinensis*, Rolfe, in *Kew Bulletin* (1896), p. 200.

KWANGTUNG: Lienchow River (*Ford, 134, 240*!). Herb. Kew.

It is also cultivated at the Hongkong Botanic Garden and at Kew.

1. *Anæctochilus Roxburghii*, *Lindl. Gen. & Sp. Orch.* p. 499; *Blume, Coll. Orch. Archip. Ind.* p. 50, t. 12 B, fig. 2; *Hook. f. Fl. Brit. Ind.* vi. p. 95; *Hance, in Journ. Bot.* xxv. (1887) p. 13.

*Anæctochilus* sp., *Hance, in Journ. Bot.* xx. (1882) p. 39.

CHEKIANG: Ningpo (*Hancock!*); FORMOSA: Bankinsing Mountains (*A. Henry, 1626!*); KWANGTUNG: North-west River (*Ford, 253!*), Lofaushan Mts. (*Ford!*). Mus. Brit.; Herb. Kew.

Tropical Himalaya.

2. *Anæctochilus Tashiroi*, *Maxim. in Bull. Acad. St. Pétersb.* xxxi. (1887) p. 103; *Mél. Biol.* xii. p. 546.

LUCHU ARCHIPELAGO (*Tashiro misit Tanaka*).

Only known to me from the original description.

1. *Vrydagzynea nuda*, *Blume, Coll. Orch. Archip. Ind.* p. 74, t. 20, fig. 3; *Hance, in Journ. Bot.* xxiii. (1885) p. 247.

HONGKONG (*Ford! ex hb. Hance, 22306!*). Mus. Brit.

Java.

This agrees remarkably well with Blume's figure, though I do not feel quite certain of the determination.

1. *Zeuxine formosana*, *Rolfe, in Ann. Bot.* ix. (1895) p. 158.

FORMOSA: South Cape (*A. Henry, 644!*). Herb. Kew.

2. *Zeuxine sulcata*, *Lindl. Gen. & Sp. Orch.* p. 485; *Journ. Linn. Soc., Bot.* i. (1857) p. 186; *Benth. Fl. Hongk.* p. 360; *Hook. f. Fl. Brit. Ind.* vi. p. 106.

*Zeuxine emarginata*, *Lindl. Gen. & Sp. Orch.* p. 485; *Benth. in Kew Journ. Bot.* vii. (1855) p. 38.

*Zeuxine Tripleura*, *Lindl. in Journ. Linn. Soc., Bot.* i. (1857) p. 186; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 128.

*Tripleura pallida*, *Lindl. Gen. & Sp. Orch.* p. 452.

FORMOSA: near Tamsui (*Walters!*), Ape's Hill, Taiwan (*Playfair, 364!*); HUPEH: Fang (*A. Henry, 6589!*); HONGKONG (*Champion, 524! Wilford, 284! Hance!*). Mus. Brit.; Herb. Kew.

Afghanistan, widely dispersed in India, to CochinChina, South Japan, Java, and the Philippines.

1. *Cheirostylis chinensis*, Rolfe, in *Ann. Bot.* ix. (1895) p. 158.  
FORMOSA: South of island (*A. Hance*, 390!), summit of Ape's Hill, Takow (*A. Henry*, 320!). Herb. Kew.

2. *Cheirostylis flabellata*, Wight,  *Ic. Pl. Ind. Or.* v. p. 16, t. 1727; *Lindl. in Journ. Linn. Soc., Bot.* i. (1857) p. 187, *excl. syn.* Griff.; *Hook. f. Fl. Brit. Ind.* vi. p. 105.

*Goodyera flabellata*, *A. Rich. in Ann. Sc. Nat. sér. 2*, xv. (1841) p. 79, t. 12.

HONGKONG (*Ford*, 130!). Herb. Kew.

North and South India.

3. *Cheirostylis yunnanensis*, Rolfe, in *Kew Bulletin* (1896), p. 201.

YUNNAN: shady rocky places at Mengtze (*Hancock*, 25!), Flowers white; very rare; Woods at 4600 ft., in shade, on stones (*A. Henry*, 10423!). Herb. Kew.

Habit of *C. flabellata*, Wight, but with much larger flowers and petals much longer than the sepals.

1. *Odontochilus Elwesii*, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 100; *Hook. Ic. Pl.* t. 2167.

SZECHUEN: Mount Omei, 5000 ft. (*Faber*, 939!). Herb. Kew.  
Sikkim, Assam, Upper Burma.

The Chinese plant has slightly larger flowers, but the whole plant is similar in other respects to the Indian one.

2. *Odontochilus yunnanensis*, Rolfe, n. sp.

*Rhizoma* repens. *Caulis* erectus, 4-8 poll. longus. *Folia* petiolata, elliptico-oblonga, acuta, 2-3 poll. longa, 9-18 lin. lata; petiolus 9-10 lin. longus, basi in vaginam spathaceam amplexicaulem expansus. *Racemus* 2-4 poll. longus, multiflorus. *Bracteæ* lanceolatæ, acutæ, 6-8 lin. longæ. *Pedicelli* 4-5 lin. longi. *Sepalum* posticum ovato-oblongum, subobtusum, 3 lin. longum; lateralia patentia, elliptico-oblonga, obtusa, 4 lin. longa. *Petala* oblique semiovata, subobtusa, cum sepalo postico in galeam conniventia. *Labellum* 7 lin. longum, basi angustum, apice in laminam bilobam dilatatum; lobi oblongi, subdivaricati; unguis utrinque appendiculis linearibus 3-4 instructus, basi in sacculum globosum 1½ lin. latum dilatatus. *Columna* 2 lin. longa.

YUNNAN: Mengtze, mountain forests, at 5000 ft. (*A. Henry*, 11110!). Flowers yellow. Herb. Kew.

Allied to *O. Clarkei*, Hook. f., a native of Sikkim, but differing in numerous details.

1. *Myrmechis chinensis*, Rolfe, n. sp.

*Caules* graciles, repentes, ascendentes. *Folia* orbicularia, obtusa v. apiculata,  $1\frac{1}{2}$ –2 lin. lata; petiolus 1 lin. longus, basi in vaginam tubulosam brevem dilatatus. *Scapus* terminalis, 1– $1\frac{1}{2}$  poll. longus, sparse pilosus, apice 2–3-florus. *Bracteæ* oblonge-lanceolatæ, acutæ v. apiculatæ,  $1\frac{1}{2}$  lin. longæ. *Pedicelli*  $1\frac{1}{2}$ –2 lin. longi. *Sepala* et *petala* ovato-oblonga, apiculata, 2 lin. longa. *Labellum* basi saccatum, limbus oblongus, obtusus, 2 lin. longus, apice vix dilatatus; calli oblongi, obtusi. *Columna* generis.

HUPEH (*A. Henry*, 6841!). Herb. Kew.

Allied to *M. japonica*, Rolfe (*M. gracilis*, Franch. et Savat. Enum. Pl. Jap. ii. p. 39 (non Blume); *Rhamphidia japonica*, Reichb. f. in Bot. Zeit. xxvi. (1878) p. 39), but the leaves and flowers considerably smaller, and the lip not dilated at the apex.

1. *Hæmaria discolor*, Lindl. *Gen. & Sp. Orch.* p. 490; *Benth. Fl. Hongk.* p. 361; *Hook. f. Fl. Brit. Ind.* vi. p. 101, in part.

*Goodyera discolor*, Ker, in *Bot. Reg.* iv. t. 271; *Bot. Mag.* t. 2055; *Lodd. Bot. Cab.* ii. t. 143.

*Ludisia discolor*, A. Rich. in *Dict. Clas. Nat.* vii. p. 437; *Blume, Orch. Archip. Ind.* p. 113, t. 42, fig. D.

*Ludisia Furetii*, *Blume, Orch. Archip. Ind.* p. 114, t. 34, fig. 3.

*Ludisia odorata*, *Blume, Orch. Archip. Ind.* p. 114, t. 34, fig. 2.

KWANGTUNG (*Furet*); HONGKONG (*Champion*, 274! *Wright*, 527! *Hance*, 1168! *Lamont*, 771!). Mus. Brit.; Herb. Kew.

Cochinchina.

This is *Goodyera discolor*, which has been cultivated in gardens ever since 1818, and was at first reported to have come from Brazil. The leaves are uniformly deep dull purplish green above and reddish purple beneath. Recently one or more Malayan species with silvery veined leaves have been cited as synonymous, though they appear to represent distinct but closely allied species. *Hæmaria argyroneura*, Miq. *Choix Pl.* t. 16, cited in the *Flora of British India*, vi. p. 101, is, however, *Macodes argyroneura*, Rolfe, in *Kew Bulletin* (1896), p. 47.

1. *Goodyera foliosa*, *Benth. ex Hook. f. in Fl. Brit. Ind.* vi. p. 113, *excl. syn.* Griff.

*Georchis foliosa*, *Lindl. Gen. & Sp. Orch.* p. 496.



*Goodyera* spp. nn. 10, 11, *Griff. Notul.* iii. pp. 394, 396; *Ic. Pl. Asiat.* t. 346, fig. 1.

KWANGTUNG: Lofaushan (*Ford*, 328!). Herb. Kew.  
Sikkim to Burma and Perak.

2. *Goodyera formosana*, *Rolfe*, in *Ann. Bot.* ix. (1895) p. 159.  
FORMOSA: Bankinsing Mountains (*A. Henry*, 409!). Herb. Kew.

3. *Goodyera Henryi*, *Rolfe*, in *Kew Bulletin* (1896), p. 201.  
HUPEH: Ichang (*A. Henry*, 6878!). Herb. Kew.  
Allied to the Indian *G. foliosa*, Lindl.

4. *Goodyera nebulorum*, *Rolfe*.  
*Cystorchis*? *nebulorum*, *Hance*, in *Journ. Bot.* xxi. (1883) p. 232.  
KWANGTUNG: in summis montibus Lofaushan, Canton (*Faber*; hb. *Hance*, 22147!). Mus. Brit.  
This proves on examination to be a *Goodyera*.

5. *Goodyera procera*, *Hook. Exot. Flora*, i. t. 39; *Lindl. Gen. & Sp. Orch.* p. 493; *Benth. Fl. Hongk.* p. 360; *Hook. f. Fl. Brit. Ind.* vi. p. 111.  
*Neottia procera*, *Ker*, in *Bot. Reg.* viii. t. 639.

FORMOSA: Tamsui (*Oldham*, 558!), Bankinsing (*A. Henry*, 165, 1826!); KWANGTUNG: Macao (*Vachell*, 79!); HONGKONG (*Champion*, 267! *Wilford*, 375! *Hance*, 533! *Lamont*!); LUCHU ARCHIPELAGO (*Wright*, 339!). Mus. Brit.; Herb. Kew.  
India and Java.

6. *Goodyera repens*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 198; *Lindl. Gen. & Sp. Orch.* p. 492; *Nyman, Conspect. Fl. Europ.* p. 689; *Boiss. Fl. Orient.* v. p. 90; *Ledeb. Fl. Ross.* iv. p. 86; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 38 (*excl. syn.*) et p. 520; *Hook. f. Fl. Brit. Ind.* vi. p. 111; *Macoun, Cat. Pl. Canad.* ii. p. 9; *A. Gray, Bot. N. U. States*, ed. 5, p. 503; *Chapm. Fl. S. U. States*, p. 463.

KIANGSU: Tangut (*Przewalski*!); HUPEH: Hsingshan (*A. Henry*, 6513!). Herb. Kew.

A very widely diffused species, ranging from Britain and Portugal to the Himalayas, Japan, and North-eastern Asia, and from Canada along the Eastern United States to the mountains of North Carolina.

7. **Goodyera Schlechtendaliana**, *Reichb. f. in Linnæa*, xxii. (1849) p. 861; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 38; *Somoku Zusetsu*, xviii. t. 79.

*Goodyera secundiflora*, *Griff. Notul.* iii. p. 394; *Ic. Pl. Asiat.* iii. t. 346, fig. 1; *Lindl. in Journ. Linn. Soc., Bot.* i. (1857) p. 182; *Hook. f. Fl. Brit. Ind.* vi. p. 113.

*Goodyera japonica*, *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 38, t. 9, fig. 1, t. 11, fig. C.

*Goodyera similis*, *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 39, t. 9, fig. 2, t. 11, fig. D.

CHEKIANG (*Fortune!*): woods on top of hills at 3000 ft. (*Hickin!*), Ningpo Mts., under bamboos (*Faber, 199!*); HUPEH: Nanto and mountains to northward (*A. Henry, 2605!*), without precise locality (*A. Henry, 6133, 7580!*). *Herb. Kew.*

Northern India, Japan.

The above all appear to me to belong to one variable species, which sometimes has the leaves variegated, at others uniformly green.

8. **Goodyera velutina**, *Maxim. in Regel's Gartenflora*, xvi. (1867) p. 38, t. 533, fig. 1; *Fl. des Serres*, t. 1779; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 38.

HUPEH: Fang (*A. Henry, 6638, 7682!*). *Herb. Kew.*

Japan.

1. **Microtis parviflora**, *R. Br. Prodr.* p. 321; *Lindl. Gen. & Sp. Orch.* p. 395; *Benth. Fl. Austral.* vi. p. 347; *Hance, in Journ. Bot.* xvii. (1879) p. 15.

FORMOSA: Tamsui (*Oldham, 561!*); LUCHU ARCHIPELAGO (*Wright, 340!*). *Mus. Brit.*; *Herb. Kew.*

Australia, New Caledonia, Java, and Japan.

1. **Arethusa sinensis**, *Rolfe*, n. sp.

*Tuber* subglobosum, parvum. *Folia* radicalia, 2-3, subsessilia, lanceolata v. elliptico-lanceolata, acuta v. acuminata, 2-4 poll. longa,  $\frac{1}{3}$ -1 poll. lata. *Scapus* erectus, 4-6 poll. longus, 2-3-florus, glaber. *Bracteæ* oblongo-lanceolatæ, acutæ, 2-3 $\frac{1}{2}$  lin. longæ. *Pedicelli* 3-4 $\frac{1}{2}$  lin. longi. *Sepala* lanceolato-lineararia, subacuta, 5-6 lin. longa. *Petala* lanceolata, acuta, 5-6 lin. longa. *Labellum* 5-6 lin. longum, basi columnæ breviter adnatum, et subsaccatum, limbus obscure trilobus, 3 lin. latus, apice

apiculatus, margine fimbriato-denticulatus, discus trilamellatus, lamellæ fimbriatæ. *Columna* clavata, 4 lin. longa.

YUNNAN: Mengtze (*A. Henry*, sine numero!). Herb. Kew.

An interesting addition to the genus, most resembling the section *Crybe* in habit.

1. *Pogonia Fordii*, *Hance*, in *Journ. Bot.* xxiii. (1885) p. 247.

KWANGTUNG: Lofaushan (*Ford!*). Mus. Brit.

2. *Pogonia ophioglossoides*, *Ker*, in *Bot. Reg.* ii. t. 148; *Nutt. Gen. N. Amer. Pl.* ii. p. 192; *Hook. Exot. Fl.* i. t. 70; *Lindl. Gen. & Sp. Orch.* p. 413; *Hance*, in *Journ. Bot.* xxiii. (1885) p. 247; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 34.

*Pogonia similis*, *Blume*, *Coll. Orch. Archip. Ind. et Jap.* p. 148, t. 32, fig. 4, et t. 54, fig. A.

*Pogonia japonica*, *Reichb. f.* in *Linnaea*, xxv. (1852) p. 228.

HUPEH: Ichang (*A. Henry*, 1777, 4044!), Chiensih (*A. Henry*, 5811!); KWANGTUNG: Lofaushan at 2500 ft. (*B. C. Henry!*); COREA: Chemulpo (*Carles*, 52!), Soul Mountains (*Carles*, 134!). Mus. Brit.; Herb. Kew.

Japan and Eastern North America.

3. *Pogonia pulchella*, *Hook. f.* *Bot. Mag.* t. 6851.

HONGKONG (*Ford!*). Herb. Kew.

4. *Pogonia yunnanensis*, *Finet*, in *Bull. Soc. Bot. Fr.* xliv. (1897) p. 419, t. 13, figs. K-Z.

YUNNAN: west of Talifou, between the Salween and Mekong Rivers (*Prince Henri of Orleans*, ex *Finet*).

1. *Didymoplexis pallens*, *Griff.* in *Calc. Journ. Nat. Hist.* iv. (1844) p. 383, t. 17; *Hemsl.* in *Journ. Linn. Soc., Bot.* xx. (1883) p. 308, t. 28; *Hook. f. Fl. Brit. Ind.* vi. p. 122.

*Leucorchis sylvatica*, *Blume*, *Mus. Bot.* i. p. 31; *Coll. Orch. Arch. Ind. et Jap.* p. 147, with fig.

*Apetalon minutum*, *Wight*, *Ic.* t. 1758.

*Arethusa ecristata*, *Griff. Notul.* iii. p. 378; *Ic. Pl. Asiat.* tt. 343, 344.

*Epiphanes pallens*, *Reichb. f.* in *Seem. Fl. Vit.* p. 296.

FORMOSA: Takow (*A. Henry*, 1878!). Parasite on roots, flowers white. Herb. Kew.

India.

An interesting discovery. I fail to distinguish it from the Indian plant, which is rather widely diffused.

1. *Gastrodia elata*, *Blume, Mus. Bot.* ii. p. 174; *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 143, t. 53, fig. 1.

HUPEH: Ichang (*A. Henry*, 868, 4659!), Changyang, S. Patung, at 3000–5000 ft. (*A. Henry*, 6078!); SZECHUEN: without locality (*Pratt*, 100!). *Mus. Brit.*; *Herb. Kew.*

Japan and Tibet.

Native name “*Tien-ma*.” The root is a drug and is often eaten roasted like a potato (*A. Henry*).

1. *Cephalanthera erecta*, *Lindl. Gen. & Sp. Orch.* p. 412; *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 188, t. 65, fig. 2; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 35.

*Serapias erecta*, *Thunb. Fl. Jap.* p. 27;  *Ic. Pl. Jap.* i. t. 4.

HUPEH: Patung (*A. Henry*, 1429, 3761!); SZECHUEN: S. Wushan (*A. Henry*, 5464 A!). *Mus. Brit.*; *Herb. Kew.*

Japan.

2. *Cephalanthera falcata*, *Lindl. Gen. & Sp. Orch.* p. 412; *Blume, Coll. Orch. Archip. Ind. et Jap.* p. 187, t. 65, fig. 1, et t. 66, fig. E; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 35.

*Serapias falcata*, *Thunb. Fl. Jap.* p. 28;  *Ic. Pl. Jap.* i. t. 5.

*Cephalanthera platycheila*, *Reichb. f. in Bot. Zeit.* iii. (1845) p. 335.

*Cephalanthera japonica*, *A. Gray, Pl. Jap.* p. 319.

CHEKIANG: Ningpo (*Everard!* *Fortune!*); SZECHUEN: S. Wushan (*A. Henry*, 5621!). *Mus. Brit.*; *Herb. Kew.*

Japan.

1. *Epipactis consimilis*, *D. Don, Prodr. Fl. Nepal.* p. 28; *Wall. Cat.* n. 7403; *Hook. f. Fl. Brit. Ind.* vi. p. 126; *King & Pantl., in Ann. Roy. Bot. Gard. Calc.* viii. p. 272, t. 364.

*Epipactis veratrifolia*, *Boiss. et Hoh. ex Boiss. Diagn.* ser. 1, xiii. p. 11; *Lindl. in Journ. Linn. Soc., Bot.* i. (1857) p. 174; *Boiss. Fl. Orient.* v. p. 87.

YUNNAN: Lunan, at riverside (*A. Henry*, 11108!). Flowers brownish. *Herb. Kew.*

Sikkim, Western Temperate Himalaya, Afghanistan, and westward to Syria.

Lindley, relying rather on the locality than the characters given, has referred Don's plant to *E. latifolia*, Linn.; but Wallich, on whose specimens it was based, credits the species to Don, and the characters, “*floribus bracteis triplo brevioribus, labello cucullato, lamina acuminata, perianthii foliolis longiore,*” agree

with the type-specimen, and not at all with Wallich's specimen of *E. latifolia*.

2. *Epipactis gigantea*, Dougl. in Hook. *Fl. Bor.-Amer.* ii. p. 202, t. 202; *S. Watson, Bot. Calif.* ii. p. 136.

*Epipactis americana*, Lindl. in *Ann. Nat. Hist.* ser. 1, iv. (1840) p. 385; *Lindl. Gen. & Sp. Orch.* p. 462.

*Epipactis Royleana*, Lindl. *Gen. & Sp. Orch.* p. 461; *Journ. Linn. Soc., Bot.* i. (1857) p. 174; *Hook. f. Fl. Brit. Ind.* vi. p. 126.

*Cephalanthera Royleana*, Regel, in *Act. Hort. Petrop.* vi. (1879) p. 490; *Boiss. Fl. Orient.* v. p. 85.

HUPEH: in Wushau, Chienshub, Patung, Changyang, Fang, and Nanto, common at about 5000 ft. (*A. Henry*, 705, 1880, 3953, 4518, 5633!); SZECHUEN: Tatsienlou (*Soulié*, 113, 788!), without locality (*Pratt*, 23, 76, 212, 711!); COREA: Chemulpo (*Carles*, 150!). *Herb. Kew.*

Mandshuria, Himalayas, and Western North America.

The Japanese *E. Thunbergii*, A. Gray, which has been referred here by Franchet & Savatier, has rather small flowers and some differences in the lip, from which I should question its specific identity with this widely diffused plant.

#### ORCHIDACEÆ: OPHRYDEÆ.

1. *Orchis aristata*, Fisch. ex Lindl. *Gen. & Sp. Orch.* p. 262; *Reichb. Fl. Germ.* xiii.-xiv. t. 410; *Hook. Fl. Bor.-Amer.* ii. p. 195; *Macoun, Cat. Canad. Pl.* ii. p. 12; *Somoku Zusetsu*, xviii. t. 51.

*Orchis latifolia*, var. *Beeringiana*, Cham. in *Linnæa*, iii. (1828) p. 26; *Hook. et Arn. Bot. Beechey, Voy.* p. 117; *Reichb. Fl. Germ.* xiii.-xiv. p. 60; *Ledeb. Fl. Ross.* iv. p. 54; *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 206; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 29.

*Orchis camtschatica*, Cham. in *Linnæa*, iii. (1828) p. 27?

SZECHUEN: Mount Omei, at 9000 ft. (*Faber*, 949!). *Herb. Kew.*

Kamtschatka, Unalaska, Sachalin, and Japan.

The aristate segments and different habit readily separate this species from *O. latifolia*, Linn. *O. camtschatica*, Cham., may represent a luxuriant state of the same, but in the absence of specimens the point remains doubtful.

2. *Orchis Chusua*, D. Don, *Prodr. Fl. Nepal.* p. 23; *Hook. f. Fl. Brit. Ind.* vi. p. 127.

*Gymnadenia Chusua*, Lindl. *Gen. & Sp. Orch.* p. 280.

SZECHUEN: Tatsienlou (*Soulié*, 210!); without locality (*Pratt*, 690, 697!). Mus. Brit.; Herb. Kew.

Alpine Himalaya.

3. *Orchis monophylla*, *Rolfe*, in *Orch. Rev.* (1898) p. 144; *Bot. Mag.* t. 7601.

*Habenaria monophylla*, *Coll. et Hemsl. in Journ. Linn. Soc., Bot.* xxviii. (1890) p. 134; *Hook. f. Fl. Brit. Ind.* vi. p. 143.

YUNNAN: Mengtze, grassy uplands, 6000–7000 ft. (*Hancock*, 590!), Flowers pink, leaves mottled; mountains, among rocks, 6000–7000 ft. (*A. Henry*, 11118, 11118 A!), Flowers pinkish-lilac and purple-pink. Herb. Kew.

Shan States.

A beautiful little species.

4. *Orchis Roborovskii*, *Maxim. in Bull. Acad. St. Pétersb.* xxxi. (1887) p. 104; *Mél. Biol.* xii. p. 547.

KANSUH: Tangut (*Przewalski*, ex *Maximowicz*).

Only known to me from the original description.

5. *Orchis salina*, *Turez. ex Lindl. Gen. & Sp. Orch.* p. 259; *Ledeb. Fl. Ross.* iv. p. 53.

*Orchis incarnata*, var. *lanceolata*, *Reichb. f. in Reichb. Fl. Germ.* xiii.–xiv. p. 176, t. 522, fig. 2.

KANSUH: Tangut (*Przewalski!*). Herb. Kew.

Siberia and Mongolia.

6. *Orchis spathulata*, *Reichb. f. ex Hook. f. Fl. Brit. Ind.* vi. p. 127; *Hook. Ic. Pl.* t. 2197, fig. A.

*Gymnadenia spathulata*, *Lindl. Gen. & Sp. Orch.* p. 280.

SZECHUEN: Tongolo (*Soulié*, 349!); without locality (*Pratt*, 11!). Herb. Kew.

Alpine Himalaya.

7. *Orchis Szechenyiana*, *Reichb. f. in Kanitz, Pl. Exped. Szechenyi Asia Centr.* (1891) p. 58.

KANSUH: Sziningfu, 9000 ft. (*Szechenyi*, ex *Reichenbach*).

Only known to me from the original description.

1. *Herminium angustifolium*, *Benth. ex Hook. f. Fl. Brit. Ind.* vi. p. 129.

*Aceras angustifolia*, *Lindl. Gen. & Sp. Orch.* p. 232; *Bot. Reg.* sub t. 1525; *Journ. Linn. Soc., Bot.* iii. (1859) p. 43; *Royle, Ill. Bot. Himal.* p. 370, t. 87, fig. 1; *Wight, Ic.* v. p. 11, t. 1691.

*Aceras longicruris*, C. Wright, in *Mem. Amer. Acad.* ser. 2, vi. (1858-9) p. 411.

FORMOSA (*Oldham*, 248, 560!); HUPEH: Ichang and Nanto (*A. Henry*, 856, 2042, 3912, 6202, 6421, 7383!). Mus. Brit.; Herb. Kew.

Japan, Mandshuria, and Northern India; also collected in Java and Timor by H. O. Forbes.

2. *Herminium biporosum*, Maxim. in *Bull. Acad. St. Pétersb.* xxxi. (1887) p. 106; *Mél. Biol.* xii. p. 549.

KANSUH: Tangut (*Przewalski*).

Only known to me from the original description.

3. *Herminium Monorchis*, R. Br. in *Ait. Hort. Kew.* ed. 2, v. p. 191; *Lindl. Gen. & Sp. Orch.* p. 305; *Reichb. Fl. Germ.* xiii.-xiv. p. 105, t. 415; *Boiss. Fl. Orient.* v. p. 82; *Hook. f. Fl. Brit. Ind.* vi. p. 128.

*Ophrys Monorchis*, *Linn. Sp. Pl.* ed. 1, p. 947.

CHIHLI: Peking (*David!*), Jehol (*David!*); KIANGSU: Peikou Pass (*Pierson!*); KANSUH: Tangut (*Przewalski!*). Mus. Brit.; Herb. Kew.

Widely diffused in Europe and N. Asia.

4. *Herminium Souliei*, Rolfe, n. sp.

*Planta* circa  $\frac{1}{2}$  ped. alta. *Folia* 2-3, suberecta, linearia v. oblongo-linearia, subacuta,  $1\frac{3}{4}$ -3 poll. longa,  $2\frac{1}{2}$ -4 lin. lata. *Racemus*  $2\frac{1}{4}$ - $2\frac{1}{2}$  poll. longus, multiflorus. *Bracteæ* ovato-lanceolatae, acuminatae, 2- $2\frac{1}{2}$  lin. longae. *Pedicelli*  $1\frac{1}{2}$ -2 lin. longi. *Sepalum* posticum orbiculare, obtusum, 1 lin. longum; lateralia orbiculari-ovata, obtusa v. apiculata, subobliqua,  $1\frac{1}{4}$  lin. longa. *Petala* lineari-oblonga, obtusa, 1 lin. longa. *Labellum*  $1\frac{3}{4}$  lin. longum, ecalcaratum, basi latum, planum, medio constrictum, apice tricuspido-trilobum; lobi laterales falcato-lineares, 1 lin. longi; lobus intermedius oblongus, obtusus,  $\frac{1}{2}$  lin. longus. *Columna* lata,  $\frac{1}{3}$  lin. longa.

SZECHUEN: Tongolo (*Soulié*, 407!). Herb. Kew.

Allied to *H. angustifolium*, Benth., but dwarfer, and quite different in the details of the flower.

5. *Herminium tanguticum*, Rolfe, n. sp.

*Tuber* ovoideo-globosum, 4-5 lin. longum. *Folia* 2-3, subradicalia, ovato-oblonga, subacuta,  $\frac{3}{4}$ -2 poll. longa, 4-7 lin. lata. *Scapus* 2-6 poll. altus; racemus densiflorus,  $\frac{1}{2}$ -1 poll. longus. *Bracteæ* triangulari-lanceolatae, acutae,  $1\frac{1}{2}$ -2 lin. longae. *Sepala*

ovato-oblonga, subobtusata,  $1\frac{1}{2}$ – $1\frac{3}{4}$  lin. longa. *Petala* subtriloba, 2– $2\frac{1}{2}$  lin. longa, basi lata, apice caudato-acuminata. *Labellum* 2– $2\frac{1}{2}$  lin. longum, tricuspido-trilobum, basi latum, concavum, ecalcaratum, lobis lateralibus divaricatis oblongis subobtusis brevibus, lobo intermedio caudato-acuminato. *Columna* brevissima.—*Herminium alaschanicum* var. *tangutica*, Maxim. in Bull. Acad. St. Pétersb. xxxi. (1887) p. 105; Mém. Biol. xii. p. 548.

KANSUH: Tangut, in rupibus, 10,100 ft. (*Przewalski!*). Herb. Kew.

Very distinct from the Mongolian *Herminium alaschanicum*, Maxim., to which it has been referred as a variety, both in habit and structure.

1. *Gymnadenia conopsea*, R. Br. in Ait. Hort. Kew. ed. 2, v. p. 191; Lindl. Gen. & Sp. Orch. p. 275; Ledeb. Fl. Ross. iv. p. 64; Boiss. Fl. Orient. v. p. 81; Franch. Pl. David. i. p. 294.

*Orchis conopsea*, Linn. Sp. Pl. ed. 1, p. 942.

*Habenaria conopsea*, Benth. in Journ. Linn. Soc., Bot. xviii. (1880) p. 354 (non Reichb. f.).

CHIHLI: Pohuashan (*David*); SZECHUEN: Tatsienlou (*Soulié*, 100, 506!); without locality (*Pratt*, 672!). Herb. Kew.

Widely diffused in Europe and Siberia.

2. *Gymnadenia cucullata*, Rich. in Mém. Mus. Par. iv. (1818) p. 57; Lindl. Gen. & Sp. Orch. p. 279; Maxim. Ind. Fl. Pekin., in Prim. Fl. Amur. p. 477.

*Orchis cucullata*, Linn. Sp. Pl. ed. 1, p. 939.

*Habenaria cucullata*, Höfft, Cat. Kursk. p. 56, ex Ledeb. Fl. Ross. iv. p. 66.

*Himantoglossum cucullatum*, Reichb. Fl. Germ. Excurs. p. 120.

CHIHLI: Pekin (*Maximowicz*); HUPEH: Fang, at 8000–9000 ft. (*A. Henry*, 6873, 6890, 6899 B!); KANSUH: Tangut (*Przewalski!*). Mus. Brit.; Herb. Kew.

Central Russia from Moscow eastward, through Siberia to China; also in Sikkim.

3. *Gymnadenia Faberi*, Rolfe.

*Habenaria Faberi*, Rolfe, in Kew Bulletin (1896), p. 201.

SZECHUEN: Mt. Omei, on rocks at 9000 ft. alt. (*Faber*, 319!). Herb. Kew.

Allied to *G. Pinguicula*, S. Moore, but the flowers are only about a fourth as large. It also approaches *G. gracilis*, Miq.,



which, however, has numerous flowers, a differently-shaped lip, and a spur of less than half the size. *G. Faberi* is described from a single specimen, with one flower, but there is a second bract with the remains of another flower.

4. **Gymnadenia gracile**, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 207; *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) p. 60.

*Mitostigma gracile*, *Blume, Mus. Bot.* ii. p. 190.

*Gymnadenia tryphiæformis*, *Reichb. f. ex Hemsl. in Journ. Bot.* xxiv. (1876) p. 209.

KIANGSI: Kiukiang (*Shearer! F. B. Forbes*, 151! *Faber*, 1090!); HUPEH: Patung district (*A. Henry*, 4660!), Tunghu, at 4000 ft. (*A. Henry*, 6434!); COREA: Chemulpo (*Carles*, 88!), Paul, Herschel, and Tracey Islands (*Oldham*, 1027!). *Mus. Brit.*; *Herb. Kew*

Japan.

5. **Gymnadenia lepida**, *Reichb. f. Otia Bot. Hamb.* (1878) p. 51; *Xen. Orch.* iii. p. 28.

LUCHU ARCHIPELAGO (*Nilson, ex Reichenbach!*).

6. **Gymnadenia Pinguicula**, *S. Moore, in Journ. Bot.* xxvi. (1878) p. 135; *Maxim. in Bull. Soc. Hist. Nat. Mosc.* liv. (1879) p. 60.

*Habenaria Pinguicula*, *Benth. in Journ. Linn. Soc., Bot.* xviii. (1880) p. 355.

CHEKIANG: Ningpo (*Everard! Hancock! hb. Hance*, 21232!). *Mus. Brit.*; *Herb. Kew.*

1. **Peristylus chloranthus**, *Lindl. in Hook. Kew Journ. Bot.* vii. (1855) p. 37.

*Coeloglossum lacertiferum et acuminatum*, *Lindl. Gen. & Sp. Orch.* p. 163.

*Habenaria lacertifera*, *Benth. Fl. Hongk.* p. 362; *Hook. f. Fl. Brit. Ind.* v. p. 163.

*Habenaria tentaculata*, *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 160, in part.

*Chæradoplectron Spiranthes*, *Schauer, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 436, t. 6, fig. C.

CHINA: Cape Syngmoon (*Meyen*); FORMOSA: Bankinsing Mountains (*A. Henry*, 895, 1572 A!); SZECHUEN: Mt. Omei, at 5000 ft. (*Faber*, 938, 941, 944!); HONGKONG (*Champion*, 269! *Ford*, 46, 1289!; *Hance! Lamont!*). *Mus. Brit.*; *Herb. Kew.*

India.

2. **Peristylus goodyeroides**, *Lindl. Gen. & Sp. Orch.* p. 299; *Dalz. & Gibs. Bomb. Fl.* p. 270; *Royle, Ill. Himal. Pl.* t. 87, fig. 2.

*Habenaria goodyeroides*, *D. Don, Prodr. Fl. Nepal.* p. 25; *Hook. f. Fl. Brit. Ind.* vi. p. 161.

*Peristylis grandis*, *Blume, Bijdr.* p. 405.

FORMOSA: Ape's Hill, Takow (*Playfair*, 154!), Flowers yellowish white (*A. Henry*, 1126!), South Cape (*A. Henry*, 1211!), Bankinsing (*A. Henry*, 1572!); YUNNAN: Mengtze, mountains and damp places (*Hancock*, 218!), 4800–5500 ft. (*A. Henry*, 11127!). *Herb. Kew.*

Widely diffused in India, also found in Java and the Philippines.

There seems a slight doubt as to whether there may not be more than one species included under this name.

3. **Peristylus Sampsoni**, *Hance, in Journ. Bot.* vi. (1868) p. 371.

*Habenaria Sampsoni*, *Hance, in Journ. Bot.* vii. (1869) p. 163 (in note).

KWANGTUNG: Pakwan, near Canton (*Sampson*, herb. *Hance*, 14493!), White Cloud Hill (*Sampson*, 1158!); without locality (*Ford*, 361!). *Mus. Brit.*; *Herb. Kew.*

4. **Peristylus viridis**, *Lindl. Syn. Brit. Fl.* ed. 2, p. 261; *Gen. & Sp. Orch.* p. 299.

*Satyrium viride*, *Linn. Sp. Pl.* ed. 1, p. 944.

*Habenaria viridis*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 192; *Syme, Engl. Bot.* ed. 3, ix. p. 105, t. 1462.

*Orchis viridis*, *Crantz, Stirp. Austr.* ed. 2, p. 491.

*Gymnadenia viridis*, *A. Rich. in Mém. Mus. Par.* iv. (1818) p. 57.

*Platanthera viridis*, *Lindl. Syn. Brit. Fl.* ed. 1, p. 261; *Reichb. Fl. Germ.* xiii.–xiv. p. 129, t. 434.

*Himantoglossum viride*, *Reichb. Fl. Germ. Excurs.* p. 120.

*Orchis bracteata*, *Willd. Sp. Pl.* iv. p. 34.

*Satyrium bracteatum*, *Pers. Syn.* ii. p. 507.

*Habenaria bracteata*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 192.

*Peristylus bracteatus*, *Lindl. Gen. & Sp. Orch.* p. 298; *Palibin, in Act. Hort. Petrop.* xiv. (1895) p. 140.

*Platanthera viridis*, var. *bracteata*, *Reichb. f. in Reichb. Fl. Germ.* xiii.–xiv. p. 130, t. 434, fig. 1.

CHIHLI: Weichang (*Putjata & Borodowsky, ex Palibin*); HUPEH: Hsingshan, at 8000–9000 ft. alt. (*A. Henry*, 6874, 6996!). *Mus. Brit.*; *Herb. Kew.*

Widely diffused in the Northern Hemisphere.

1. *Platanthera chlorantha*, *Cust. ex Reichb. in Moessl. Handb.* ed. 2, ii. p. 1565; *Reichb. Ic. Crit.* ix. t. 853; *Lindl. Gen. & Sp. Orch.* p. 285.

*Habenaria chlorantha*, *Bab. in Trans. Linn. Soc.* xvii. (1836) p. 462; *Syme, Engl. Bot.* ed. 3, ix. p. 107, t. 1664 ("H. eu-bifolia, t. 1663" on plate, by error).

SHINGKING: between Mukden and Tungcheshien (*James!*); SZECHUEN: without locality (*Pratt*, 689!); COREA: Soul Mountains (*Carles*, 136!). *Herb. Kew.*

Western Europe and Northern Asia, from Britain to Mandshuria.

2. *Platanthera flava*, *Lindl. Gen. & Sp. Orch.* p. 293.

*Orchis fuscescens*, *Linn. Sp. Pl.* ed. 1, p. 943.

*Habenaria herbiola*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 193; *Spreng. Syst. Veg.* iii. p. 688.

*Habenaria virescens*, *Spreng. Syst. Veg.* iii. p. 688; *A. Gray, Bot. N. U. St.* ed. 5, p. 499.

*Perularia fuscescens*, *Lindl. Gen. & Sp. Orch.* p. 281; *Spreng. Syst. Veg.* iii. p. 688; *Ledeb. Fl. Ross.* iv. p. 66; *Franch. Pl. David.* i. p. 294; *Reichb. Fl. Germ.* xiii.-xiv. p. 116, t. 426.

CHIHLI: Puhuashan (*David*, 2253); SHINGKING: between Tungcheshien and Maorshan, Yaloo River (*James!*). *Herb. Kew.* Siberia and North America.

3. *Platanthera Henryi*, *Rolfe*.

*Habenaria Henryi*, *Rolfe, in Kew Bulletin* (1896), p. 202.

CHEKIANG (*Hickin!*); KIANGSI: Kiukiang (*Shearer!*); FOKIEN: about 2000 ft. (*Carles*, 685!); HUPEH: Patung (*A. Henry*, 4716, 6148!), Kuei (*A. Henry*, 7663!); SZECHUEN: S. Wushan (*A. Henry*, 7453!), Mt. Omei (*Faber*, 941!). *Mus. Brit.*; *Herb. Kew.*

Allied to *Platanthera mandarinorum*, *Reichb. f.*, but taller, with more lax racemes and smaller flowers.

4. *Platanthera hologlottis*, *Maxim. Prim. Fl. Amur.* p. 268.

SHINGKING: Mukden (*Webster!*); COREA: Port Chusan (*Wilford*, 962!). *Herb. Kew.*

Dahuria and Japan.

5. *Platanthera iantha*, *Wight, Ic.* v. p. 11, t. 1692.

*Habenaria iantha*, *Hook. f. Fl. Brit. Ind.* vi. p. 164.

YUNNAN: hill-slopes at Mengtze, at 5700 ft. (*Hancock*, 41! *A. Henry*, 11111!). Apparently rare. Flowers purple-red. *Herb. Kew.*

South India.

6. *Platanthera interrupta*, Maxim. in *Bull. Acad. Sc. St. Pétersb.* xxxi. (1887) p. 106.

KWANGTUNG: Lantao Island (*Ford*, 46!); COREA: without locality (*Oldham*, 843!), Port Chusan (*Wilford*, 944!), Chemulpo (*Carles*, 96!). Mus. Brit.; Herb. Kew.

Mandshuria and Japan.

7. *Platanthera japonica*, Lindl. *Gen. & Sp. Orch.* p. 290; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 31.

*Habenaria japonica*, A. Gray, in *Mem. Am. Acad.* n. s. vi. (1858-9) p. 410; *Somoku Zusetsu*, xviii. t. 52.

*Orchis japonica*, Thunb. *Fl. Jap.* p. 26.

HUPEH: Patung (*A. Henry*, 1835, 4082, 5041!), Fang (*A. Henry*, 5889, 5889 A!); SZECHUEN: S. Wushan (*A. Henry*, 5713!), without locality (*Pratt*, 104!). Mus. Brit.; Herb. Kew.

Japan.

8. *Platanthera mandarinorum*, Reichb. f. in *Linnæa*, xxv. (1852) p. 226.

*Habenaria Keiskei*, Miq. *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 207.

*Platanthera Keiskei*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 31.

*Platanthera oreades*, *Franch. et Savat. l. c.* p. 514.

CHEKIANG: Chusan Hills (*Fortune*, 48!); Ningpo (*Hancock*, herb. *Hance*, 21214!); FOKIEN: Amoy (*Fortune*, 79!). Mus. Brit.; Herb. Kew.

Japan.

9. *Platanthera obcordata*, Lindl. *Gen. & Sp. Orch.* p. 290.

*Orchis obcordata*, Buch.-Ham. in D. Don, *Prodr. Fl. Nepal.* p. 23.

*Gymnadenia obcordata*, Reichb. f. *Otia Bot. Hamb.* p. 32.

*Platanthera Galeandra*, Reichb. f. in *Linnæa*, xxv. (1852) p. 226.

*Habenaria Galeandra*, Benth. *Fl. Hongk.* p. 363; *Hook. f. Fl. Brit. Ind.* vi. p. 163.

*Gymnadenia Galeandra*, Reichb. f. *Otia Bot. Hamb.* p. 33.

*Platanthera Championi*, Lindl. in *Hook. Kew Journ. Bot.* vii. (1855) p. 37.

FORMOSA: Bankinsing Mountains (*A. Henry*, 850!); HONG-KONG (*Champion*, 270! *Hance*, 105, 738! *Wright*, 525! *Fortune*, 78!); KWANGTUNG: Lantao Island (*Ford*, 47!). Mus. Brit.; Herb. Kew.

Western Himalaya and Central India.

10. *Platanthera stenostachya*, Lindl. in *Hook. Kew Journ. Bot.* vii. (1855) p. 37.

*Habenaria stenostachya*, Benth. *Fl. Hongk.* p. 362; *Hook. f. Fl. Brit. Ind.* vi. p. 156.

*Habenaria peristyloides*, *Wight, Ic. Pl. Ind. Or.*, v. p. 13, t. 1702 (non *A. Rich.*).

*Cœloglossum densum*, *Lindl. Gen. & Sp. Orch.* p. 302; *Reichb. f. in Bonplandia*, iii. (1855) p. 250, iv. p. 321.

*Cœloglossum peristyloides*, *Reichb. f. in Bonplandia*, iv. (1856) p. 321.

HONGKONG (*Champion! Hance, 740! Lamont!*). Mus. Brit.; Herb. Kew.

North and South India.

11. *Platanthera Susannæ*, *Lindl. Gen. & Sp. Orch.* p. 295.

*Orchis Susannæ*, *Linn. Sp. Pl.* ed. 1, p. 939.

*Habenaria Susannæ*, *R. Br. Prodr.* p. 312; *Blume, Bijdr.* p. 402; *Benth. Fl. Hongk.* p. 363; *Hook. f. Fl. Brit. Ind.* vi. p. 137.

*Orchis gigantea*, *Sm. Exot. Bot.* ii. p. 79, t. 100.

*Habenaria gigantea*, *D. Don, Prodr. Fl. Nepal.* p. 24; *Bot. Mag.* t. 3374.

*Platanthera robusta*, *Lindl. Gen. & Sp. Orch.* p. 295.

YUNNAN: Mengtze, rare, on grassy mountain slopes (*Hancock, 120!*), 5500 ft. (*A. Henry, 11,095!*); KWANGTUNG: près de Fukwing (*Krone*); HONGKONG (*Champion, 272! Fortune, 126! Hance, 741!*). Mus. Brit.; Herb. Kew.

India and Malaya.

12. *Platanthera ussuriensis*, *Maxim. in Bull. Acad. St. Pétersb.* xxxi. (1887) p. 107; *Mél. Biol.* xii. p. 551.

*Platanthera tipuloides*, var. *ussuriensis*, *Regel, Fl. Ussur.* p. 157, t. 10, figg. 7-9.

CHEKIANG: Ningpo Mts. (*Faber, 57, 1707!*); KIANGSI: Lushan Mts., Kiukiang (*Bullock, 93!*); HUPEH: Tunghu (*A. Henry, 6415!*), Hsingshan (*A. Henry, 6415 a!*), Chienshih (*A. Henry, 7470!*). Mus. Brit.; Herb. Kew.

Mandshuria and Japan.

1. *Habenaria acuifera*, *Lindl. Gen. & Sp. Orch.* p. 325; *Hook. f. Fl. Brit. Ind.* vi. p. 146; *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 155.

*Habenaria linguella*, *Lindl. Gen. & Sp. Orch.* p. 325; *Benth. Fl. Hongk.* p. 262; *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 196.

CHINA: without locality (*Fortune, 86! Millett!*); KWANGTUNG: without locality (*Wenyon!*); Lofaushan (*Ford!*); HONGKONG (*Champion, 217! Urquhart! Hance, 545! Lamont, 766! Forbes!*); HAINAN (*B. Henry!*). Mus. Brit.; Herb. Kew.

Northern India and Tavoy.

Lindley specifically separated the Chinese and Indian plants, though placing them next to each other, but Sir Joseph Hooker afterwards united them, and I believe correctly. Dr. Kränzlin, however, places them far apart, in different sections of the genus, which is certainly erroneous.

2. *Habenaria camptoceras*, Rolfe, in *Journ. Linn. Soc., Bot.* xxix. (1892) p. 319.

SZECHUEN: Tatsienlou (*Soulié*, 610!); without locality (*Pratt*, 305!). Mus. Brit.; Herb. Kew.

3. *Habenaria ciliolaris*, Kränzl. in *Engl. Bot. Jahrb.* xvi. (1892) p. 169.

CHEKIANG: Ningpo Mts. (*Faber*, 201!); HUPEH: S. Tunghu (*A. Henry*, 7487!); HONGKONG (*Ford*, 98!), Little Hongkong (*Ford*, 95!). Mus. Brit.; Herb. Kew.

4. *Habenaria endothis*, Miq. *Journ. Bot. Néerl.* i. (1861) p. 92.

KWANGTUNG: Hoan (*Krone*, ex *Miquel*).

The author describes this species as an ally of *H. linguella*, Lindl. (i. e. *H. acutifera*, Lindl.).

5. *Habenaria ensifolia*, Lindl. *Gen. & Sp. Orch.* p. 321.

*Habenaria pectinata*, Hook. f. *Fl. Brit. Ind.* vi. p. 137, partim (non *Don*); *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 193, partim; *Orch. Gen. et Sp.* p. 405, partim, & excl. var.; *King & Pantl. in Ann. Roy. Bot. Gard. Calc.* viii. p. 310, partim, t. 406.

HUPEH: Tunghu, at 4000 ft. (*A. Henry*, 6446!), S. Patung (*A. Henry*, 7384!); YUNNAN: Mengtze, rocky and grassy mountains, 5000-6500 ft. (*A. Henry*, 11119!). Herb. Kew.

Temperate Himalaya.

6. *Habenaria Fordii*, Rolfe, in *Kew Bulletin* (1896), p. 202.

YUNNAN: Mengtze, woods at 4700 ft. (*A. Henry*, 11101!); KWANGTUNG (*Ford*, 360!). Flowers white. Herb. Kew.

Allied to the Indian *H. commelynifolia*, Wall., but having much larger flowers, and the leaves not cauline.

7. *Habenaria geniculata*, D. Don, *Prodr. Fl. Nepal.* p. 25; *Lindl. Gen. & Sp. Orch.* p. 324; *Hook. f. Fl. Brit. Ind.* vi. p. 138; *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 198.

*Orchis dentata*, Sw. in *Vet. Akad. Nya Handl. Stock.* xxi. (1800) p. 207.  
*Platanthera dentata*, Lindl. *Gen. & Sp. Orch.* p. 296.

CHINA (*Swartz*).

Northern India to Burma.

Inserted on the authority of *Swartz*, whose habitat, however, has not been confirmed.

8. *Habenaria glaucifolia*, *Bur. et Franch. in Journ. de Bot.* v. (1891) p. 152.

SZECHUEN: Tatsienlou (*Bonvalot & Prince Henri d'Orleans, ex Bur. et Franch.*; *Soulié*, 159!); Tongolo (*Soulié*, 324!), without locality (*Pratt*, 587!); YUNNAN: near Tali (*Delavay*, 124, ex *Bur. et Franch.*). Mus. Brit.; Herb. Kew.

9. *Habenaria Hancockii*, *Rolfe, in Kew Bulletin* (1896), p. 202.

YUNNAN: damp grassy slopes at Mengtze, at 5500–6000 ft. (*Hancock*, 85!), Flowers flesh-colour; grassy mountains, 4600–6000 ft. (*A. Henry*, 11122!). Herb. Kew.

Allied to *H. acuifera*, Wall., but with much longer side-lobes to the lip, more clavate spur, &c.

10. *Habenaria humidicola*, *Rolfe, in Kew Bulletin* (1896), p. 202.

CHEKIANG: Ningpo Mts., in damp places in the shade of rocks (*Faber*, 200!). Herb. Kew.

Nearly allied to *H. reniformis*, Hook. f., but the leaves are longer, the flowers smaller, and the different parts much more slender.

11. *Habenaria intermedia*, *D. Don, Prodr. Fl. Nepal.* p. 24; *Lindl. Gen. & Sp. Orch.* p. 322; *Reichb. f. in Gard. Chron.* (1879) ii. p. 136; *Hook. f. Fl. Brit. Ind.* vi. p. 138.

SZECHUEN: Mt. Omei, at 3500–4500 ft. (*Faber*, 950!); Yangtzekiang (*Faber*, 976!); YUNNAN: Mengtze, grassy mountains and uplands (*Hancock*, 588!). Herb. Kew.

Temperate Himalaya.

12. *Habenaria leptoloba*, *Benth. Fl. Hongk.* p. 362.

HONGKONG (*Hance*, 1288!). Mus. Brit.; Herb. Kew.

Kränzlin (in *Engl. Bot. Jahrb.* xvi. (1892) p. 160) doubtfully reduces this to *H. tentaculata*, Reichb. f. (i. e. *Glossula tentaculata*,

Lindl.), but it is a very different plant, with a long spur, and more nearly allied to *H. khasiana*, Hook. f.

13. **Habenaria Miersiana**, *Champ. in Hook. Kew Journ. Bot.* vii. (1855) p. 37; *Benth. Fl. Hongk.* p. 363; *Hance, in Journ. Bot.* vii. (1869) p. 161, partim.

*Habenaria Sieboldiana*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 208.

*Habenaria radiata*, *Kränzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 200, partim (non *Miq.*).

FOKIEN: Amoy (*Fortune*, 40!), Foochow (*Carles*, 688!); FORMOSA: South Cape (*A. Henry*, 1212!), Bankinsiug Mountains *A. Henry*, 1607!); HUPEH: Nanto (*A. Henry*, 4576, 4611!); YUNNAN: Mengtze, grassy mountains at 5000–6000 ft. (*A. Henry*, 11098, 11123, 11128!), Flowers white; KWANGTUNG: Macao (*Gaudichaud*, 59!), Lienchau River (*Ford*, 329!), Pakwan (*Sampson*, 546!); HONGKONG (*Champion*, 336! *Harland!* *Urquhart*, 231! *Lamont!*). *Mus. Brit.*; *Herb. Kew.*

Japan.

I cannot follow Hance and Kränzlin in referring this plant to *Habenaria radiata*, *Miq.*

Var. **yunnanensis**, *A. Finet, in Journ. de Bot.* xii. (1898) p. 341.

YUNNAN: Tapintze (*Delavay*), Salouen Valley (*Prince Henri d'Orleans, ex Finet*).

A dwarf alpine form of the species.

14. **Habenaria omeiensis**, *Rolfe, in Kew Bulletin* (1896), p. 203.

SZECHUEN: Mount Omei, at 8000 ft. (*Faber*, 951!). *Herb. Kew.*

Allied to the Indian *H. latilabris*, Hook. f., and *H. stenantha*, Hook. f., but with a more lax raceme of larger flowers.

15. **Habenaria polytricha**, *Rolfe, in Hook. Ic. Pl.* t. 2496.

FORMOSA: South Cape (*A. Henry*, 1246!). *Herb. Kew*

A remarkable species, belonging to the small group *Medusæ-formes*, and allied to *H. ternatea*, *Reichb. f.*

16. **Habenaria reniformis**, *Hook. f. Fl. Brit. Ind.* vi. p. 152.

*Listeria reniformis*, *D. Don, Prodr. Fl. Nepal.* p. 208.

*Neottia reniformis*, *Spreng. Syst. Veg.* iii. p. 707.

*Herminium reniforme*, *Lindl. in Wall. Cat.* n. 7067.



*Aopla reniformis*, *Lindl. in Bot. Reg.* vii. sub t. 1701; *Gen. & Sp. Orch.* pp. 304, 457; *Benth. Fl. Hongk.* p. 363.

KWANGTUNG: Pakwan (*Sampson*!); HONGKONG (*Harland*, 1063! *Lamont*, 780! *Hance*, 1279!). Mus. Brit.; Herb. Kew.

Nepal and Khasia Hills.

This species often occurs in a peloriate state, with entire lip and no spur; but in the normal form the lip is three-lobed, and the spur linear.

17. *Habenaria rhodocheila*, *Hance, in Ann. Sc. Nat. sér. 5*, v. (1866) p. 243.

*Habenaria militaris*, *Kränzln. in Engl. Bot. Jahrb.* xvi. (1892) p. 203, partim (non *Reichb. f.*).

KWANGTUNG: Canton, North River, on rocks (*Sampson*, hb. *Hance*, 11332!), Lofaushan, at 600 ft. (*Ford*, hb. *Hance*, 17376!). Mus. Brit.; Herb. Kew.

Quite distinct from *H. militaris*, *Reichb. f.*, with which *Kränzlin* has united it.

18. *Habenaria sagittifera*, *Reichb. f. in Bot. Zeit.* iii. (1845) p. 334; *Hance, in Journ. Bot.* vi. (1878) p. 233; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 33; *Somoku Zusetsu*, xviii. t. 60; *Franch. in Mém. Soc. Nat. Cherb.* xxiv. (1884) p. 257; *Debeaux, Fl. Tien-tsin*, p. 69; *Kränzln. in Engl. Bot. Jahrb.* xvi. (1892) p. 205.

*Habenaria linearifolia*, *Maxim. Prim. Fl. Amur.* p. 269; *Regel, Fl. Ussur.* p. 157, t. 10, figg. 14-18.

CHIHLI: Peking (*Bretschneider*!); SHANTUNG: Chefoo (*Stuhlmann, Mollendorff*, hb. *Hance*, 11332! *Fauvel*, ex *Franchet*); CHEKIANG: Ningpo mts. (*Faber*, 55, 198, 298!); COREA: Chemulpo (*Carles*, 113!), Gensan (*Hickin*!). Mus. Brit.; Herb. Kew.

Japan and Eastern Asia.

19. *Habenaria yunnanensis*, *Rolfe*, n. sp.

*Planta*  $\frac{3}{4}$ - $1\frac{1}{4}$  ped. alta. *Folia* 3 v. 4, radicalia, subsessilia, ovata, apiculata,  $1-1\frac{1}{4}$  poll. longa, 8-11 lin. lata. *Scapus* vaginis lanceolatis acuminatis obtectus. *Racemus* 2-4 poll. longus, laxiflorus. *Bracteæ* lanceolatæ, acuminatæ, 2-4 lin. longæ. *Pedicelli* 5-7 lin. longi. *Sepalum* posticum ovatum, cucullatum, obtusum,  $1\frac{1}{2}$  lin. longum; lateralia patentia, oblongo-lanceolata, acuta,  $2\frac{1}{2}$  lin. longa. *Petala* linearia,  $1\frac{1}{2}$  lin. longa, cum sepalo

postico in galeam conniventia. *Labellum* tripartitum, 4 lin. longum; lobi lineares, æquales; calcar elongatum, incurvum, 6-7 lin. longum, apice paullo fusiformi-incrassatum. *Columna* 1 lin. longa; processus stigmaticus  $\frac{1}{4}$  lin. longus.

YUNNAN: Mengtze, rocky mountains at 6000 ft. (*A. Henry*, 11121!). Flowers white. Herb. Kew.

A very distinct species belonging to the section *Trimeroglossa*.

1. *Diplomeris chinensis*, *Rolfe*, in *Kew Bulletin* (1896), p. 203.

CHEKIANG: Tientai mt., at 1000 ft., on damp rocks (*Faber*, 95!). Herb. Kew.

Remarkable for the inflated base of the spur.

1. *Hemipilia brevicarata*, *Finet*, in *Bull. Soc. Bot. Fr.* xliv. (1897) p. 420, t. 14, figg. A-G.

YUNNAN: west of Talifou, between the rivers Yangpi and Tieueultsin (*Prince Henri d'Orleans*, ex *Finet*).

2. *Hemipilia cruciata*, *Finet*, in *Bull. Soc. Bot. Fr.* xliv. (1897) p. 421, t. 14, figg. H-P.

YUNNAN: Menghoutin, south of Talifou (*Prince Henri d'Orleans*, ex *Finet*).

3. *Hemipilia flabellata*, *Bur. et Franch.* in *Journ. de Bot.* v. (1891) p. 152.

SZETCHUEN (probably): without locality (*Bonvalot & Prince Henri d'Orleans*).

The locality is not stated, but the majority of the plants of this collection were from this province.

4. *Hemipilia Henryi*, *Reichb. f. ex Bur. et Franch.* in *Journ. de Bot.* vi. (1891), p. 152 (nomen tantum); *Rolfe*, in *Kew Bulletin* (1896) p. 203.

HUPEH: Ichang (*A. Henry*, 1534!), Nanto (*A. Henry*, 6347!), Hsingshan (*A. Henry*, 6347 A!), Fang (*A. Henry*, 6347 B!). Mus. Brit.; Herb. Kew.

Allied to the Himalayan *H. cordifolia*, Lindl.

1. *Glossula tentaculata*, *Lindl. Bot. Reg.* x. (1824) t. 862.

*Glossaspis tentaculata*, *Spreng. Syst. Veg.* iii. p. 694; *Benth. Fl. Hongk.* p. 361.

*Glossaspis antennifera*, *Reichb. f. in Linnæa*, xxv. (1852) p. 225.

*Habenaria tentaculata*, *Reichb. f. Otia Bot. Hamb.* p. 34; *Trans. Linn. Soc.* xxx. (1874) p. 139; *Krünzl. in Engl. Bot. Jahrb.* xvi. (1892) p. 160, partim.

CHINA: without locality (*Fortune*, 182!); KWANGTUNG: Maisaou, at 1800 ft. (*Tate!*), Lofaushan, at 600 ft. (*Ford!*), Lienchau, Canton (*Lamont!*); HONGKONG (*Champion*, 30! *Wright*, 523! *Wilford*, 282! *Lamont*, 772! *Bowring!* *Hance*, 524!). *Herb. Kew.*

Siam and Cambodia.

This plant has been confused with *Habenaria lacertifera*, *Benth.*, and *H. leptoloba*, *Benth.* Whether it can be retained as a separate genus is perhaps doubtful, but its specific distinctness can hardly be called in question.

✓ 1. *Satyrium nepalense*, *D. Don, Prodr. Fl. Nepal.* p. 26; *Lindl. Gen. & Sp. Orch.* p. 340; in *Journ. Linn. Soc., Bot.* iii. (1859) p. 44; *Bot. Mag.* t. 6625; *Hook. f. Fl. Brit. Ind.* vi. p. 168.

SZECHUEN: Tatsienlou (*Soulié*, 62, 433!); YUNNAN: Mengtze, S.E. mts., among grass at 6000 ft. (*A. Henry*, 9192, 9192 B!); south of Red River, from Mamnei, at 6000 ft. (*A. Henry*, 9192 A!). *Herb. Kew.*

North and South India and Ceylon.

A widely diffused and variable species.

#### ORCHIDACEÆ: CYPRIPEDEÆ.

1. *Cypripedium arietinum*, *R. Br. in Ait. Hort. Kew.* ed. 2, v. p. 222; *Bot. Mag.* t. 1567; *Lindl. Gen. & Sp. Orch.* p. 529; *Franch. in Bull. Soc. Bot. Fr.* xxxiii. (1886) p. 206; *Bull. Soc. Philom. Paris*, sér. 7, xii. (1891) p. 140; in *Journ. Bot.* viii. (1894) p. 252.

*Cypripedium plectrochilon*, *Franch. in Bull. Soc. Bot. Fr.* xxxii. (1885) p. 27.

HUPEH: S. Patung (*A. Henry*, 5474!); SZECHUEN: without locality (*Pratt*, 87, 763!); YUNNAN: Mt. Chetchotze (*Delavay*, 1020!). *Mus. Brit.*; *Herb. Kew.*

Canada and North-east United States.

2. *Cypripedium Calceolus*, *Linn. Sp. Pl.* p. 951; *Lindl. Gen. & Sp. Orch.* p. 527; *Reichb. Fl. Germ.* xiii.–xiv. p. 167, t. 496; *Palibin, in Act. Hort. Petrop.* xiv. (1895) p. 140.

CHIHLI: Huangtuliangtze (*Putjata et Borodowsky, ex Palibin*).  
Europe and Central Asia.

3. *Cypripedium corrugatum*, *Franch. in Journ. de Bot.* viii. (1894) p. 251.

YUNNAN: calcareous hills at the sources of the Peechaha, at 3500 m. (*Delavay, 3478*); calcareous rocks at Maculchan (*Delavay*).

Var. *obesa*, *Franchet, l. c.*

YUNNAN: woods at Lienyn, at 2800 m.; also Koutoui and Fangyangtchang (*Delavay*).

Only known to me from description.

4. *Cypripedium ebracteatum*, *Rolfe, in Kew Bulletin* (1896), p. 204.

HUPEH (*A. Henry, 1404 (2)!*). Mus. Brit.

A remarkable ebracteate species, differing from *C. micranthum*, *Franch.*, in its much larger flowers, and from *C. margaritaceum*, *Franch.*, and *C. Fargesii*, *Franch.*, in its saccate lip. The only specimen seen is at the British Museum, and it has been mixed with *C. japonicum*, *Thunb.*

5. *Cypripedium Fargesii*, *Franch. in Journ. de Bot.* viii. (1894) p. 267.

SZECHUEN: Tchenkéoutin, at 2000 m. (*Farges, 585!*). Herb. Kew.

Much larger than *C. margaritaceum*, of which it may possibly be a luxuriant state.

6. *Cypripedium fasciolatum*, *Franch. in Journ. de Bot.* viii. (1894) p. 232.

SZECHUEN: woods of Heoupin, near Tchenkeou, at 2200 m. (*Farges, 922!*). Herb. Kew.

7. *Cypripedium guttatum*, *Sw. in Vet. Akad. Nya Handl. Stockh.* xxi. (1800) p. 251; *Lindl. Gen. & Sp. Orch.* p. 529; *Ledeb. Fl. Ross.* iv. p. 88; *Maxim. Prim. Fl. Amur.* p. 270; *F. Schmidt, Reis. in Amurl. und Ins. Sachalin*, p. 184; *Reichb. Fl. Germ.* xiii.–xiv. p. 166, t. 495; *Fl. des Serres*, t. 573; *Franch. in Journ. de Bot.* viii. (1894) p. 253.

CHIHLI: Peking mountains, at 6000 ft. alt. (*Hancock*, 14! *Bretschneider*!); Siaowutaishan (*Möllendorff*!); Ipehoashan (*David*); Sylingchan (*Provot*, 136); SHINGKING: Toonghwahein (*Ross*, 33!); SZECHUEN: without locality (*Pratt*, 264!); Tongolo (*Soulié*, 311, ex *Franchet*, 327, 328!). Mus. Brit.; Herb. Kew.

Widely diffused from Central Europe, through temperate Asia, to Alaska, in the neighbourhood of the Mackenzie River.

8. *Cypripedium Henryi*, *Rolfe*, in *Kew Bulletin* (1892), p. 211; *Franch. in Journ. de Bot.* viii. (1894) p. 271.

*Cypripedium chinense*, *Franch. in Journ. de Bot.* viii. (1894) p. 230:

HUPEH: S. Patung (*A. Henry*, 5391 A!), Chienshih (*A. Henry*, 5391 D!); SZECHUEN: S. Wushan (*A. Henry*, 5391 B!), Kimise near Tchenkéoutin, at 1200 m. (*Farges*, 1036, ex *Franchet*). Herb. Kew.

9. *Cypripedium himalaicum*, *Rolfe*, in *Journ. Linn. Soc., Bot.* xxix. (1892) p. 319; *Franch. in Journ. de Bot.* viii. (1894) p. 249.

*Cypripedium macranthum*, *Hook. f. Fl. Brit. Ind.* vi. p. 170, partim (non *Sw.*).

SZECHUEN: without locality (*Pratt*, 748!). Herb. Kew.  
Sikkim and Nepal.

10. *Cypripedium japonicum*, *Thunb. Fl. Jap.* p. 30; *Ic. Pl. Jap.* i. t. 1; *Lindl. Gen. & Sp. Orch.* p. 529; *Blume, Coll. Orch.* p. 169, t. 59; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 40; *Franchet, in Journ. de Bot.* viii. (1894) p. 254.

HUPEH: Patung (*A. Henry*, 1404, 3777!); SZECHUEN: Tchenkéoutin (*Farges*, 649, ex *Franchet*). Mus. Brit.; Herb. Kew.

Japan.

11. *Cypripedium luteum*, *Franch. Pl. David.* ii. p. 126; *id. in Journ. de Bot.* viii. (1894) p. 228.

HUPEH: Fang, at 7000 ft. (*A. Henry*, 5391 E, 6740!); SZECHUEN: Tatsienlou (*Soulié*, 413! *Prince Henri d'Orleans & R. P. Soulié*, 576), without locality (*Pratt*, 263!); Tchenkéoutin (*Farges*, 134 bis). Herb. Kew.

Tibet.

12. *Cypripedium macranthum*, Sw. in *Vet. Akad. Nya Handl. Stockh.* xxi. (1800) p. 251; *Bot. Mag.* t. 2938; *Bot. Reg.* t. 1534; *Lindl. Gen. & Sp. Orch.* p. 528; *Ledeb. Fl. Ross.* iv. p. 87; *Reichb. Fl. Germ.* xiii.–xiv. p. 169, t. 498; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 40; *Hook. f. Fl. Brit. Ind.* vi. p. 170, partim; *Franch. in Journ. de Bot.* viii. (1894) p. 232.

CHIHLI: Puhuashan (*David*, 2216! *Möllendorff*, 431!), Siaowutaishan, at 5000 ft. alt. (*Möllendorff*!), Peking mountains, at 5000 ft. (*Hancock*, 13! *Bretschneider*! *David*, 3!), Sylingchan (*Bodinier et Provot*); SHINGKING: Toonghwabien (*Ross*, 54!), Mukden to Yaloo River (*Webster*, 34!), between Mukden and Tungcheshien (*James*!); SZECHUEN: Tchenkéoutin (*Farges*, 509, ex *Franchet*). Mus. Brit.; Herb. Kew.

Central Russia to Eastern Siberia and Japan.

13. *Cypripedium margaritaceum*, *Franch. in Bull. Soc. Phil. Par. sér. 7*, xii. (1891) p. 141; — *Reprint*, p. 8; *Orchidophile*, 1888, p. 368, with plate; *Rolfe, in Gard. Chron.* (1889) i. p. 43; *Franch. in Journ. de Bot.* viii. (1894) p. 265.

YUNNAN: on mount Tsangchan, above Tali (*Delavay*, 282); also above Mosoyn, and at Hec-chan-men (*Delavay*).

14. *Cypripedium micranthum*, *Franch. in Journ. de Bot.* viii. (1894) p. 265.

SZECHUEN: in woods at Heoupin, near Tchenkéoutin, at 2000 m. (*Farges*, 1286!).

15. *Cypripedium tibeticum*, *King, ex Rolfe, in Journ. Linn. Soc., Bot.* xxix. (1892) p. 320; *Franch. in Journ. de Bot.* viii. (1894) p. 250 (*thibeticum*).

*Cypripedium macranthum*, *Hook. f. Fl. Brit. Ind.* vi. p. 170, partim (non Sw.).

SZECHUEN: Tongolo (*Soulié*, 953!), without locality (*Pratt*, 14, 42, 301, 736!), between Batang and Litang (*Prince Henri d'Orleans*), Tatsienlou (*Soulié*, 312). Herb. Kew.

Tibet.

16. *Cypripedium yunnanense*, *Franch. in Journ. de Bot.* viii. (1894) p. 231.

YUNNAN: forests of Hec-chan-men (*Delavay*, 2480); Mosoyn, in the woods of Koutoui, at 2800 m. (*Delavay*, ex *Franchet*).

Only known to me from the description.

1. **Paphiopedilum Parishii**, *Pfitz. in Engl. Bot. Jahrb.* xix. (1894) p. 41.

*Cypripedium Parishii*, *Reichb. f. in Flora*, lii. (1869), p. 322, et in *Gard. Chron.* (1869) pp. 814, 1158, cum fig.; *Bot. Mag.* t. 5791; *Williams, Orch. Alb.* ii. t. 86; *Ill. Hort.* xxii. t. 214.

*Selenipedium Parishii*, *Jolibois, in Rev. Hort.* (1885) p. 132.

YUNNAN: Mengtze, on mountains at 6000–7000 ft. (*Hancock*, 582!). *Herb. Kew.*

Burma.

2. **Paphiopedilum purpuratum**, *Pfitz. in Engl. Bot. Jahrb.* xix. (1894) p. 40.

*Cypripedium purpuratum*, *Lindl. in Bot. Reg.* xxiii. t. 1991; *Gen. & Sp. Orch.* p. 530; *Bot. Mag.* t. 4901; *Benth. Fl. Hongk.* p. 364; *Godefr.-Lebeuf & Brown, Les Cyrip.* t. 5.

*Cypripedium sinicum*, *Hance, in Walp. Ann.* iii. p. 602.

HONGKONG (*Champion*, 29!, *Wilford*, 283! *Ford*, 443!). *Mus. Brit.*; *Herb. Kew.*

An endemic species. In the original description it is said to have been introduced to cultivation from the Malayan Archipelago, which is clearly an error; while in the 'Flora Hongkongensis,' Khasia is given as one of its habitats; but the specimen on which this remark is based is a piece of *C. venustum*, Wall., collected by Griffith, and mounted on the same sheet as *purpuratum*.

### 136. SCITAMINEÆ.

(By C. H. WRIGHT, A.L.S.)

1. **Globba bulbosa**, *Gagnep. in Bull. Soc. Bot. France*, xlvi. (1901) p. 202, pl. iv. figg. 1–6.

KWEICHAU: environs of Kweiyang (*Bodinier, ex Gagnepain*).

1. **Roscoea alpina**, *Royle, Ill. Him. Bot.* p. 361, t. 89. fig. 1; *Wight, Ic. Pl. Ind. Or.* t. 2013, fig. 1; *Horan. Monogr. Scit.* p. 20; *Hook. f. Fl. Brit. Ind.* vi. p. 207.

*Roscoea purpurea*, *Royle, Ill. Him. Bot.* p. 361, t. 89. fig. 3.

*Roscoea purpurea*, var. *minor*, *Wall. Cat.* n. 6528, D & E.

SZECHUEN: Tachienlou (*Pratt*, 518!).

Northern India.

2. **Roscoea capitata**, var. **purpurata**, *Gagnep. in Bull. Soc. Bot. France*, xlvi. (1901) p. lxxiv.

YUNNAN: woods in the gorges of San-tchang-kiou (*Delavay*, 4491, *ex Gagnepain*).

Var. *scillifolia*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxiv.

YUNNAN: Kanhaytze on the Hec-chan-men at 9000 ft. (*Delavay*, 2685, 2685 bis, *ex Gagnepain*), moist places below the pass of Hec-chan-men at 9000 ft. (*Delavay*, 3283, *ex Gagnepain*).

3. *Roscoea cautleoides*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxv.

YUNNAN: Hec-chan-men (*Delavay*, 92, 231 ter, *ex Gagnepain*), environs of Lake Er'Hai (*Prince Henry of Orleans*, *ex Gagnepain*).

4. *Roscoea Chamæleon*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxvi.

YUNNAN: sides of the Hec-chan-men (*Delavay*, 2659, *ex Gagnepain*).

5. *Roscoea debilis*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxvi.

YUNNAN: in the wood of Lan-ngy-tien (*Ducloux*, 688, *ex Gagnepain*).

6. *Roscoea intermedia*, var. *minuta*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxiii.

YUNNAN: Naeulchan (*Delavay*, *ex Gagnepain*).

Var. *macrorhiza*, *Gagnep. l. c.*

YUNNAN: environs of Yunnan-sen (*Ducloux*, 601, *ex Gagnepain*).

Var. *anomala*, *Gagnep. l. c.*

YUNNAN: environs of Yunnan-sen (*Ducloux*, 597, *ex Gagnepain*).

1. *Kæmpferia cœnobialis*, *C. H. Wright*.

*Monolophus cœnobialis*, *Hance, in Journ. Bot.* viii. (1870) p. 75.

KWANGTUNG: North River, Canton (*Sampson*, *ex Hance*).

2. *Kæmpferia elegans*, *Wall. Pl. As. Rar.* i. p. 24, t. 27; *Hook. f. Fl. Brit. Ind.* vi. p. 222.

*Monolophus?* *elegans*, *Wall. Cat.* n. 6593.

SZECHUEN (*Mesny*, 8).

India and Malay Peninsula.



3. *Kæmpferia fongyuensis*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxviii.

YUNNAN: environs of Lake Er'Hai, at Fungyu (*Prince Henry of Orleans*, ex *Gagnepain*).

4. *Kæmpferia Galanga*, *Linn. Sp. Pl.* ed. 1, p. 3; *Lour. Fl. Cochinch.* p. 12; *Rosc. Scitam.* t. 92; *Bot. Mag.* t. 850; *Hook. f. Fl. Brit. Ind.* vi. p. 219.

CHINA (ex *Loureiro*); KWANGTUNG: Whampoa (*Hance*, 22423! cult.). Mus. Brit.

India, Malaya.

5. *Kæmpferia pandurata*, *Roxb. in As. Res.* xi. (1810) p. 328, t. 4; *Rosc. Scitam.* t. 96; *Bot. Reg.* t. 173; *Bretschn. Early Res.* p. 176; *Hook. f. Fl. Brit. Ind.* vi. p. 220; *Horan. Monogr. Scit.* p. 21.

*Curcuma rotunda*, *Linn. Sp. Pl.* ed. 1, p. 2; *Lour. Fl. Cochinch.* p. 9.

CHINA (ex *Loureiro*).

India, Malaya.

6. *Kæmpferia rotunda*, *Linn. Sp. Pl.* ed. 1, p. 3; *Rosc. Scitam.* t. 97; *Bot. Mag.* tt. 920 & 6054.

FORMOSA: Bankinsing (*A. Henry*, 851!). Herb. Kew.

India, Malaya.

7. *Kæmpferia* (§ *Pyrgophyllum*) *yunnanensis*, *Gagnep. in Bull. Soc. Bot. France*, xlviii. (1901) p. lxxvii.

YUNNAN: gorges of Tsongso, north of Tali (*Delavay*, 2721, ex *Gagnepain*).

[*Hedychium coronarium*, *Koen.* (*Retz. Obs.* iii. p. 73), is said in *Bot. Mag.* sub t. 708 to be "probably of Chinese origin." It has, however, been collected only in India and the Malayan Islands.]

1. *Curcuma longa*, *Linn. Sp. Pl.* ed. 1, p. 2; *Lour. Fl. Cochinch.* p. 8; *Horan. Monogr. Scit.* p. 23; *Bot. Reg.* t. 886; *Flück. & Hanbury, Pharmac.* p. 577; *Bentl. & Trim. Med. Pl.* t. 269; *Hook. f. Fl. Brit. Ind.* vi. p. 214; *Bretschn. Early Res.* p. 176.

*Amomum Curcuma*, *Jacq. Hort. Vind.* iii. t. 4.

CHINA (ex *Loureiro*): FORMOSA: Bankinsing, cultivated (*A. Henry*, 89, 1606!). Herb. Kew.

Cultivated throughout the Tropics.

“The radical tubers when dried constitute the medicine known as Yu-chin by the Chinese. The rhizomes are very yellow and undoubtedly turmeric.”—*Dr. A. Henry.*

2. *Curcuma pallida*, *Lour. Fl. Cochinch.* p. 9; *Horan. Monogr. Scit.* p. 24; *Bretschn. Early Res.* p. 176.

KWANGTUNG: Canton (ex *Loureiro*).

[*Amomum medium*, *Lour. Fl. Cochinch.* p. 4, said to be from Yunnan, has been doubtfully referred to *Alpinia alba*, *Rosc.*, which in its turn is probably only a form of *Alpinia Galanga*, *Willd.*]

1. *Zingiber confine*, *Miq. Journ. Bot. Néerl.* i. (1861) p. 94.

KWANGTUNG (*Krone*, ex *Miquel*).

2. *Zingiber corallinum*, *Hance, in Journ. Bot.* xviii. (1880) p. 301, & 1883, p. 357.

HAINAN (*Hancock!*). Mus. Brit.

3. *Zingiber integrilabrum*, *Hance, in Journ. Bot.* xx. (1882) p. 80.

HONGKONG: Mount Gough (*Ford*, 5! *Hance*, 21812!). Mus. Brit.; Herb. Kew.

4. *Zingiber officinale*, *Rosc. in Trans. Linn. Soc.* viii. (1807) p. 348; *Scitam.* t. 83; *Bentl. & Trim. Med. Pl.* t. 270; *Hook. f. Fl. Brit. Ind.* vi. p. 246.

*Amomum Zingiber*, *Linn. Sp. Pl.* ed. 1, p. 1; *Lour. Fl. Cochinch.* p. 2; *Bretschn. Early Res.* p. 176; *Jacq. Hort. Vind.* i. t. 75.

CHINA (ex *Loureiro* and *Bretschneider*); FORMOSA: Bankinsing (*A. Henry*, 491, 1575!). “Without any doubt a native of China.”—*Bretschneider.*

Widely cultivated in the Tropics.

5. *Zingiber Zerumbet*, *Rosc. in Sm. Exot. Bot.* ii. p. 105, t. 112, and *Trans. Linn. Soc.* viii. (1807) p. 348; *Scitam.* t. 84; *Bot. Mag.* t. 2000; *Horan. Monogr. Scit.* p. 27; *Hook. f. Fl. Brit. Ind.* vi. p. 247; *Hance, in Journ. Bot.* xvii. (1879) p. 15.

*Amomum Zerumbet*, *Linn. Sp. Pl.* ed. 1, p. 1; *Jacq. Hort. Vind.* iii. t. 54.

*Zerumbet Zingiber*, *Lestib. in Ann. Sc. Nat.* 2<sup>me</sup> sér. xv. (1841) p. 329.

KWANGTUNG: Tingushan, Canton (*Hance*, 17695!). Mus. Brit.; Herb. Kew.

Widely cultivated in the Tropics of the Old World.

6. *Zingiber* sp. indescript. ?

HUPEH: Tunghu (*A. Henry*, 7594, 7671!); SZECHUEN:  
N. Wushan (*A. Henry*, 7126!). Herb. Kew.

7. *Zingiber* sp. indescript. ?

HUPEH: Ichang (*A. Henry*, 7104 A!). Herb. Kew.

8. *Zingiber* sp. indescript. ?

FORMOSA: Bankinsing (*A. Henry*, 147, 1607!). Herb. Kew.

1. *Costus speciosus*, *Sm. Trans. Linn. Soc.* i. (1791) p. 249;  
*Roxb. in As. Res.* xi. p. 349; *Wight, Ic. Pl. Ind. Or.* t. 2014;  
*Horan. Monogr. Scit.* p. 37; *Hook. f. Fl. Brit. Ind.* vi. p. 249.

*Costus arabicus*, *Jacq. Ic.* t. 1, non *Linn.*

*Banksia speciosa*, *Koenig, in Retz. Obs.* iii. p. 75; *Rheede, Hort. Malab.*  
xi. t. 8.

FORMOSA: Takow (*A. Henry*, 1204!). Herb. Kew.

India and Malaya.

1. *Alpinia calcarata*, *Rosc. in Trans. Linn. Soc.* viii. (1807)  
p. 347; *Scitam.* t. 68; *Bot. Reg.* t. 141; *Hook. f. Fl. Brit.*  
*Ind.* vi. p. 254; *Horan. Monogr. Scitam.* p. 34.

*Alpinia cernua*, *Sims, in Bot. Mag.* t. 1900.

*Renealmia calcarata*, *Haw. in Andr. Rep.* t. 421.

*Globba erecta*, *Red. Lil.* t. 174.

HONGKONG (*Urquhart*, 227, 261! *Hance*, 1231!); HAINAN  
(*Swinhoe*!). Herb. Kew.

India and Malay Peninsula.

2. *Alpinia chinensis*, *Rosc. in Trans. Linn. Soc.* viii. (1807)  
p. 346; *Benth. Fl. Hongk.* p. 349.

*Languas chinensis*, *Koenig, in Retz. Obs.* iii. p. 65.

*Heritiera chinensis*, *Retz. Obs.* vi. p. 18.

*Hellenia chinensis*, *Willd. Sp. Pl.* i. p. 5.

*Hellenia abnormis*, *Lindl. in Trans. Hort. Soc.* vii. (1830) p. 60?

*Monocystis abnormis*, *Lindl. Introd. Bot.* ed. 2, p. 445; *Horan. Monogr.*  
*Scitam.* p. 36?

FOKIEN: Amoy (*Swinhoe*!); HONGKONG (*Wilford*, 403!  
*Wright*, 512, 513!); CHINA (*Fortune*, 65!). Herb. Kew.

Bonin Islands and Cambodia.

3. *Alpinia Galanga*, *Willd. Sp. Pl.* i. p. 12; *Horan. Monogr.*  
*Scitam.* p. 33; *Roxb. Fl. Ind.* i. p. 58; *Benth. Fl. Hongk.*  
p. 348; *Hook. f. Fl. Brit. Ind.* vi. p. 253.

*Maranta Galanga*, Swartz, *Obs. Bot.* p. 8.

*Amomum Galanga*, Lowr. *Fl. Cochinch.* p. 5.

*Alpinia viridiflora*, Griff. *Notul.* iii. p. 423;  *Ic.* t. 353.

FORMOSA: Bankinsing (*A. Henry*, 804!); HONGKONG (*Champion*, *Hance ex Benth*); HAINAN (*A. Henry*, 8522!). Herb Kew.

India and Malaya.

4. *Alpinia globosa*, Horan. *Monogr. Scit.* p. 34.

*Amomum globosum*, Lowr. *Fl. Cochinch.* p. 4; *Bretschn. Early Res.* p. 176.

CHINA (ex *Loureiro*).

"The plant is unknown to modern botanists."—*Bretschneider*.

5. *Alpinia japonica*, Miq. in *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 140; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 20; *Maxim. in Bull. Soc. Nat. Mosc.* liv. (1879) p. 59.

*Globba japonica*, Thunb. *Fl. Jap.* p. 23.

CHEKIANG: Ningpo (*Hancock ex Maximowicz*); HUPEH: South Tunghu (*A. Henry*, 7710!), Ichang (*Aldridge*!); SZECHUEN: Mount Omei (*Faber*, 956!); FORMOSA (*A. Henry*, 2079!). Herb. Kew.

Japan.

6. *Alpinia nutans*, Rosc. in *Trans. Linn. Soc.* viii. (1807) p. 346; *Scitam.* t. 73; *Smith, Exot. Bot.* ii. t. 106; *Bot. Mag.* t. 1903; *Wight, Ic. Pl. Ind. Or.* t. 2027; *Benth. Fl. Hongk.* p. 348; *Hook. f. Fl. Brit. Ind.* vi. p. 256.

*Globba nutans*, Linn. *Mant.* ii. p. 170; *Red. Lil.* t. 60.

*Renealmia nutans*, Andr. *Bot. Rep.* t. 360.

*Zerumbet speciosum*, Schrad. & *Wendland, Sert. Hannov.* iii. t. 19 (err. typ. 18); *Jacq. Fragm.* t. 68.

CHINA (*Fortune*, 79!); FORMOSA: Takow (*A. Henry*, 782!); HONGKONG (*Urquhart*, 158! *Wright*, 514!); SZECHUEN: above Chung Ching (*Faber*, 957!); HAINAN (*A. Henry*, 8594, 8731!), Ma Augza Volcano, Hoihow (*Hancock*, 34!). Herb. Kew.

Eastern Himalaya, Malayan Peninsula and Archipelago.

7. *Alpinia officinarum*, Hance, in *Journ. Linn. Soc., Bot.* xiii. (1872) p. 6, et *Journ. Bot.* xi. (1873) p. 175; *Bentl. & Trim. Med. Pl.* t. 271; *Flück. & Hanbury, Pharm.* p. 580; *Bot. Mag.* t. 6995; *Bretschneider, Hist. Europ. Bot. Disc.* p. 660.

## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they would appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

Part I. for 1901, consisting of 378 pages, price 21 shillings, was issued on July 10, 1902; Part II., completing the volume, is now in course of compilation.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.

---

VOL. XXXV. is still in course of issue, the Parts already published being Nos. 242-245. Those still unpublished are Parts 246-248 inclusive.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."

APRIL 1.

Price 5s.

# THE JOURNAL

OF

# THE LINNEAN SOCIETY.

---

VOL. XXXVI.

BOTANY.

No. 250.

---

## CONTENTS.

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.— PART XV. ....	73

---

**See Notice on last page of Wrapper.**

---

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1903.

12

# LINNEAN SOCIETY OF LONDON.

## LIST OF THE OFFICERS AND COUNCIL.

Elected 24th May, 1902.

### PRESIDENT.

Prof. Sydney Howard Vines, M.A., D.Sc., F.R.S.

### VICE-PRESIDENTS.

William Carruthers, F.R.S.  
Frank Crisp, LL.B., B.A.

Herbert Druce, F.Z.S.  
Rev. Thomas R. R. Stebbing, F.R.S.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

Prof. G. B. Howes, D.Sc., LL.D., F.R.S. | D. H. Scott, M.A., Ph.D., F.R.S.

### COUNCIL.

William Carruthers, F.R.S.  
Frank Crisp, LL.B., B.A.  
Herbert Druce, F.Z.S.  
Prof. J. Reynolds Green, M.A., D.Sc.,  
F.R.S.  
W. Botting Hemsley, F.R.S.  
Prof. G. B. Howes, D.Sc., LL.D., F.R.S.  
B. Daydon Jackson, Esq.  
George Masee, Esq.

G. S. Saunders, Esq.  
Dukinfield H. Scott, M.A., Ph.D.,  
F.R.S.  
Rev. Thomas R. R. Stebbing, M.A.,  
F.R.S.  
Col. C. Swinhoe, F.Z.S., F.E.S.  
A. G. Tansley, M.A.  
Prof. Sydney H. Vines, D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*. The former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meet at 4 p.m., as required during the Session. The Members for 1902-1903, in addition to the officers, are :—

Herbert Goss, F.G.S.  
Albert O. L. G. Günther, M.D., F.R.S.  
W. Botting Hemsley, F.R.S.  
G. R. M. Murray, F.R.S.  
P. Chalmers Mitchell, M.A., F.Z.S.

Prof. F. W. Oliver, D.Sc.  
A. B. Rendle, M.A., D.Sc.  
Howard Saunders, F.Z.S.  
F. N. Williams L.R.C.P.



HAINAN (*Taintor* in herb. *Hance*, 16866!), opposite Hainan (*Sampson* ex *Bretschneider*). Mus. Brit.; Herb. Kew.

8. *Alpinia oxyphylla*, *Miq. in Journ. Bot. Néerl.* i. (1861) p. 93.  
KWANGTUNG: HOAN (*Krone* ex *Miquel*).

9. *Alpinia?* *pumila*, *Hook. f. Bot. Mag.* t. 6832.  
*Amomum vittatum*, *Hance, in Journ. Bot.* xxiv. (1886) p. 53.  
KWANGTUNG: LOFAUSHAN (*Ford!* *Faber* ex *Hance*). Mus. Brit.; Herb. Kew.

10. *Alpinia stachyodes*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1876) p. 126.  
HONGKONG (*Hance*, 1231!). Mus. Brit.; Herb. Kew.

1. *Phrynium capitatum*, *Willd. Sp. Pl.* i. 17; *Roxb. in Asiat. Res.* xi. p. 325, t. 3; *Wight, Ic. Pl. Ind. Or.* t. 2016; *Horan. Monogr. Scitam.* p. 11; *Hook. f. Fl. Brit. Ind.* vi. p. 258.

*Pontederia ovata*, *Linn. Sp. Pl.* ed. 1, p. 288.

*Phyllodia placentaria*, *Lour. Fl. Cochinch.* p. 13.

KWANGTUNG: FUKWING (*Krone* ex *Hance*), West River (ex *Hance*); HAINAN (*A. Henry*, 8677!). Mus. Brit.; Herb. Kew.

India, Malay Peninsula and Archipelago.

2. *Phrynium parviflorum*, *Roxb. Hort. Beng.* p. 1; *Fl. Ind.* i. p. 7; *Rosc. Scitam.* t. 34; *Horan. Monogr.* p. 11; *Hook. f. Fl. Brit. Ind.* vi. p. 259.

*Maranta parviflora*, *A. Dietr. Sp. Pl.* i. p. 30.

KWANGTUNG: LOFAUSHAN (*Ford!*); HAINAN (*A. Henry*, 8732!). Herb. Kew.

India and Malayan Peninsula.

3. *Phrynium sinicum*, *Miq. in Journ. Bot. Néerl.* i. (1861) p. 94.  
KWANGTUNG: PHULUWEI (*Krone* ex *Miquel*).

1. *Canna indica*, var. *orientalis*, *Hook. f. Fl. Brit. Ind.* vi. p. 260.

*Canna indica*, *Linn. Sp. Pl.* p. 1, ex parte; *Roxb. Fl. Ind.* i. p. 1; *Asiat. Res.* xi. p. 322; *Lour. Fl. Cochinch.* p. 10; *Benth. Fl. Hongk.* p. 349.

*Canna orientalis*, *Rosc. Scitam.* t. 12; *Horan. Monogr. Scit.* p. 16.

*Canna chinensis*, Willd. in *Ges. Naturf. Fr. Berl. Mag.* ii. (1808) p. 170.

SZECHUEN (*A. Henry*, 8828 !), near Tachienlu (*Pratt*, 378 !), Chung Ching (*Bourne*, 78 !), Hokiang (*Faber*, 308 !); HONGKONG (*Urquhart*, 69 !); HAINAN (*A. Henry*, 8030 !). *Herb. Kew.*  
India and Malaya.

2. *Canna juncea*, *Retz. Obs. fasc. i.* p. 9.  
CHINA (*Bladh ex Retzio*).

1. *Musa Basjoo*, *Sieb. in Verh. Batav. Gen.* xii. (1830) p. 18; *Miq. Ann. Mus. Bat.* iii. p. 140; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 21; *Baker, in Bot. Mag.* t. 7182; *Ann. Bot.* vii. p. 210; *Kew Bulletin*, 1894, p. 248.

LUCHU ARCHIPELAGO (*ex Siebold*).  
Cultivated in Southern Japan.

2. *Musa Cavendishii*, *Lamb. in Paxt. Mag.* iii. p. 51, with col. fig.; *Baker, in Ann. Bot.* vii. p. 209; *Kew Bulletin*, 1894, p. 244; *Seem. Fl. Vit.* p. 289.

*Musa chinensis*, *Sweet, Hort. Brit.* ed. 2, p. 596 (*nomen*).

*Musa sinensis*, *Sagot, ex Baker, in Ann. Bot.* vii. p. 209.

SOUTHERN CHINA (*ex Sweet*).

3. *Musa coccinea*, *Andr. Bot. Rep.* t. 47; *Red. Lil.* tt. 307-8; *Bot. Mag.* t. 1559; *Fl. des Serres*, tt. 722-3; *Walp. Ann.* vi. p. 38; *Baker, in Ann. Bot.* vii. p. 220; *Kew Bulletin*, 1894, p. 258.

*Musa Uranoscopos*, *Lour. Fl. Cochinch.* p. 645; *Bretschn. Early Res.* p. 176.

SOUTHERN CHINA (*ex Andrews*).

Cochinchina.

4. *Musa lasiocarpa*, *Franch. in Journ. de Bot.* iii. (1889) p. 329; *Baker, in Ann. Bot.* vii. p. 208; *Kew Bulletin*, 1894, p. 243.

YUNNAN: near Tapintze (*Delavay ex Franchet*).

Native name, Ngaptsiao.

5. *Musa sapientum*, *Linn. Syst.* ed. x. p. 1303; *Sp. Pl.* ed. 2, p. 1477; *Rumph. Hort. Amboin.* v. p. 130, t. 60; *Roxb. Pl. Corom.* t. 275; *Horan. Monogr. Scit.* p. 42; *Benth. Fl. Hongk.*

p. 348; *Bretschn. Early Res.* p. 175; *Hook. f. Fl. Brit. Ind.* vi. p. 262; *Baker, in Ann. Bot.* vii. p. 211; *Kew Bulletin*, 1894, p. 250.

*Musa corniculata*, *M. odorata*, *M. seminifera*, and *M. nana*, *Lour. Fl. Cochinch.* p. 644; *Desv. in Journ. de Bot.* 1814, pp. 28, 30, 38–39, 43

HONGKONG (*Champion ex Benth*am); Mount Parker (*Ford*, 60!).  
Herb. Kew.

India and Malaya.

6. *Musa Wilsoni*, *W. J. Tutcher, in Gard. Chron.* ser. iii. xxxii. (1902) p. 450, fig. 151.

YUNNAN (*Wilson ex Tutcher*).

Young plants are in cultivation at Kew.

7. *Musa* sp.

YUNNAN: Szemao (*A. Henry*, 12325!).

This is cultivated and the enlarged base of the stem is eaten as a vegetable. It agrees with the description of *M. nepalensis*, *Wall.* (in *Roxb. Fl. Ind.* ii. 492 [1824]), of which no specimens are known to exist, and which is, as suggested by *Wallich* (*l. c.*), doubtfully distinct from the widely cultivated *M. superba*, *Roxb.* (*Hort. Beng.* 19).

### 137. HÆMODORACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Aletris foliosa*, var. *sikkimensis*, *Franch. in Journ. de Bot.* x. (1896) p. 198.

*Aletris sikkimensis*, *Hook. f. Fl. Brit. Ind.* vi. p. 265.

SZETCHUEN: Tachienlu (*Soulié*, 525!), Tchenkéoutin (*Farges ex Franchet*). Herb. Kew.

Sikkim.

2. *Aletris glabra*, *Bureau et Franch. in Journ. de Bot.* v. (1891) p. 156; *Franch. l. c.* x. (1896) p. 197.

SZETCHUEN: Tachienlu (*Bonvalot ex Franchet, Pratt*, 675, 691!), Tongolo (*Soulié*, 309!). Herb. Kew.

3. *Aletris glandulifera*, *Bureau et Franch. in Journ. de Bot.* v. (1891) p. 156, x. (1896) p. 200.

SZETCHUEN: Tachienlu (*Bonvalot ex Franchet*).

4. *Aletris japonica*, *Lamb. in Trans. Linn. Soc.* x. (1811) p. 407; *A. Gray, in Mem. Amer. Acad. new ser.* vi. (1859) p. 417; *S. Moore, in Journ. Bot.* xvi. (1878) p. 138; *Hance, in Journ. Bot.* xx. (1882) p. 295.

*Aletris spicata*, *Franch. in Journ. de Bot.* x. (1896) p. 199.

*Hypoxis spicata*, *Thunb. Fl. Jap.* p. 136.

*Hypoxis farinosa*, *Thunb. in Trans. Linn. Soc.* ii. (1794) p. 334.

ANWHEI: Wuhu (*Bullock*!); CHEKIANG: Ningpo (*Everard*!); KIANGSI: Kiukiang (*Shearer*! *Carles*, 460!); FORMOSA: Tamsui (*Oldham*, 563!), Kelung (*Ford*, 13!); HUPEH: Ichang (*Henry*, 269!), Patung (*Henry*, 1748!), Yangtsekiang (*Faber*, 895! *Gill*! *Stronach*!); KANSUH: Lidshapu (*Potanin*!); SZECHUEN: Chunching (*Bourne*!); COREA: Tsusima (*Wilford*, 866!); LUCHU ARCHIPELAGO (*Wright*, 346!): Oosima Islands (*Carpenter*, 12!). Mus. Brit., Herb. Kew.

Japan.

Var. *Fargesii*, *C. H. Wright*.

*Aletris spicata*, var. *Fargesii*, *Franch. in Journ. de Bot.* [x. (1896) p. 200.

SZECHUEN: Tsoueuitsao (*Farges*, 437, ex *Franchet*).

5. *Aletris lactiflora*, *Franch. in Journ. de Bot.* x. (1896) p. 200.

SZECHUEN: Tachienlu (*Soulié* ex *Franchet*).

6. *Aletris lanuginosa*, *Bureau et Franch. in Journ. de Bot.* v. (1891) p. 155, x. (1896) p. 201.

*Stachyopogon pauciflorus* et *S. spicatus*, *Klotzsch, in Bot. Ergeb. Reise Pr. Waldem.* p. 49, t. 94.

*Aletris nepalensis*, *Hook. f. Fl. Brit. Ind.* vi. p. 264, ex parte.

SZECHUEN (*Faber*!), Litang (*Bonvalot* ex *Franchet*), Tachienlu (*Pratt*, 13, 182! *Soulié*, 445, 562!), Tongola (*Soulié* ex *Franchet*). Herb. Kew.

Kumaon.

7. *Aletris laxiflora*, *Bureau et Franch. in Journ. de Bot.* v. (1891) p. 155, x. (1896) p. 199.

SZECHUEN: Tachienlu (*Bonvalot* ex *Franchet*, *Soulié*, 12, 82, 102! *Pratt*, 86!). Herb. Kew.

8. *Aletris nepalensis*, var. *Delavayi*, *Franch. in Journ. de Bot.* x. (1896) p. 198.

*Aletris Delavayi*, *Franch. in Journ. de Bot.* v. (1891) p. 157 (*name only*).

YUNNAN : Tsang-chan (*Delavay*, 981, 2420, ex *Franchet*).

9. *Aletris revoluta*, *Franch. in Journ. de Bot.* x. (1896) p. 202.

SZECHUEN : Tchenkéoutin (*Farges* ex *Franchet*).

10. *Aletris stenoloba*, [*Franch. in Journ. de Bot.* x. (1896) p. 203.

SZECHUEN : Souifou (*Delavay*, 2282! ex *Franchet*).

1. *Peliosanthes Delavayi*, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 43.

YUNNAN : Longki (*Delavay*, 3141, 4989! ex *Franchet*).

2. *Peliosanthes macrostegia*, *Hance, in Journ. Bot.* xxiii. (1885) p. 328.

KWANGTUNG : Lofaushan (*Ford*, 72!). Herb. Kew.

1. *Ophiopogon clavatus*, *C. H. Wright, in Kew Bulletin* (1895) p. 116, and *Hook. Ic. Pl.* t. 2382.

HUPEH : Patung (*A. Henry*, 6065!), Kuei (*A. Henry*, 6065 A!).

Allied to *O. dracænoides*, *Hook. f.*, but differing in the shape of the leaves and larger, less numerous flowers.

2. *Ophiopogon Griffithii*, *Hook. f. Fl. Brit. India*, vi. p. 270.

*Flueggea Griffithii*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 502.

SZECHUEN : Mount Omei (*Faber*, 890!). Herb. Kew.

Eastern Himalaya.

3. *Ophiopogon Jaburan*, *Lodd. Bot. Cab.* t. 1876; *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 86; *Mél. Biol.* vii. p. 324; *Engler, Bot. Jahrb.* vi. (1885) p. 54.

*Convallaria japonica*, var. *major*, *Thunb. Fl. Jap.* p. 139.

*Slateria Jaburan*, *Sieb. in Verh. Batav. Gen.* xii. (1830) p. 15.

*Flueggea Jaburan*, *Kunth. Enum.* v. p. 303; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 502.

LUCHU ARCHIPELAGO : Amami Osima (*Döderlein* ex *Engler*).

Japan.

4. *Ophiopogon japonicus*, *Ker*, in *Bot. Mag.* t. 1063 ; *Maxim.* in *Bull. Acad. St. Pétersb.* xv. (1871) p. 87 ; *Mél. Biol.* vii. p. 325.

*Convallaria japonica*, *Linn. f. Suppl.* 204 ; *Red. Lil.* t. 80.

*Convallaria japonica*, var. *minor*, *Thunb. Fl. Jap.* p. 139.

*Flüggea japonica*, *Rich. in Schrad. N. Journ.* II. i. (1807) p. 9, t. 1 A ; *Kunth, Enum.* v. p. 301 ; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 501 ; *A. Gray, in Bot. Perry's Exped.* p. 322.

*Slateria japonica*, *Desv. in Verh. Batav. Gen.* xii. (1830) p. 15.

*Ophiopogon japonicus*, var. *genuinus*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 87 ; *Mél. Biol.* 1870, p. 327.

KIANGSI: Kiukiang (*Shearer* !); FORMOSA: Ape's Hill (*A. Henry*, 1098 A !), South Cape (*A. Henry*, 1703 A !); HUPEH (*A. Henry*, 6438 !), Patung (*A. Henry*, 3705 !); SZECHUEN: Tachienlu (*Soulié*, 177 !); COREAN ARCHIPELAGO (*Oldham*, 883 !). *Mus. Brit.*; *Herb. Kew.*

Japan.

Var. *umbraticola*, *C. H. Wright.*

*Ophiopogon* (*Flüggea*) *umbraticola*, *Hance, in Journ. Bot.* vi. (1868) p. 115 ; *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) 88 ; *Mél. Biol.* vii. p. 328.

*Flüggea japonica*, var. *umbraticola*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 501.

FORMOSA: Takao (*Playfair*, 85 !); HUPEH: Ichang (*Henry*, 4344 A !); KWANGTUNG: Canton (*Hance et Sampson*, 13759 !), Lofaushan (*Ford* !). *Herb. Kew.*

Japan.

This is not, as stated by Maximowicz, synonymous with *O. japonicus* var. *intermedius*, Maxim., which has much larger, cernuous flowers ; in this they are small and erect.

Var. *umbrosus*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 88 ; *Mél. Biol.* vii. p. 327.

CHIHLI: Peking (*Shaczkoﬀ ex Maximowicz*).

Japan.

Var. *Wallichianus*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 89 ; *Mél. Biol.* vii. p. 328.

*Chloopsis acaulis*, *Blume, Enum.* i. p. 14 ?

*Flüggea Wallichiana*, *Kunth, Enum.* v. p. 303.

*Flüggea japonica*, var. *Wallichiana*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 501, *excl. syns.*

*Ophiopogon Wallichianus*, *Hook. f. Fl. Brit. Ind.* vi. p. 268 (*excl. syn. Ophiopogon longifolius, Decne.*).

SZECHUEN: Mount Omei (*Faber*, 310 !); KIANGSU: Woosung (*Home* !); KWANGTUNG: Canton (*Hance*, 10086 !). Mus. Brit.; Herb. Kew.

Japan, Temp. and Trop. Himalayas and Khasia.

1. *Liriope kansuensis*, *C. H. Wright*.

*Ophiopogon* (*Liriope*) *kansuensis*, *Batalin*, in *Act. Hort. Petrop.* xiii. (1893) p. 103.

KANSUH: by the River Heiho, near Sapa (*Potanin ex Batalin*).

2. *Liriope spicata*, *Lour. Fl. Cochinch.* p. 201.

*Dracæna graminifolia*, *Linn. Syst.* ed. 12, p. 246.

*Convallaria spicata*, *Thunb. Fl. Jap.* p. 141.

*Fluggea spicata*, *Schult. f. Syst.* vii. p. 309.

*Ophiopogon japonicus*, *Wall. Cat.* n. 5139, *ex parte*.

*Ophiopogon spicatus*, *Ker*, in *Bot. Reg.* t. 593; *conf. Bot. Mag.* sub t. 1063; *Kunth, Enum.* v. p. 299; *Benth. Fl. Hongk.* p. 371; *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 83; *Mél. Biol.* vii. p. 321; *Hance, in Journ. Bot.* xii. (1874) p. 262; *Debeaux, Fl. Tchéf.* p. 135; *Maxim. Fl. As. Or. Fragm.* p. 63; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 83.

*Ophiopogon spicatus*, var. *gracilis*, *Miq. Prol. Fl. Jap.* p. 143; *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 85; *Mél. Biol.* vii. p. 323; *Franch. Plantes de Tchéfou*, p. 258.

*Ophiopogon gracilis*, *Kunth, Enum.* v. p. 298.

*Ophiopogon longifolius*, *Decne. in Fl. des Serres*, xvii. (1867-68) p. 182.

*Ophiopogon spicatus*, var. *Kunthianus*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 84; *Mél. Biol.* vii. p. 322.

*Liriope graminifolia*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 499.

SHINGKING: Moukden (*Webster*, 20 !); SHANTUNG: Chefoo (*Hancock*, 11 !); FORMOSA (*Oldham*, 575 !); HUPEH: Ichang (*Henry*, 353, 498, 911, 1044, 1093, 1329 !), Patung (*Henry*, 2550, 2806 !), Tunghu (*Henry*, 6439 !); KIANGSI: Kiukiang (*Shearer* ! *Möllendorff ex Hance*); KWANGTUNG: Macao (*Calléry*, 272 ! *Maingay*, 391 !); HONGKONG (*Champion*, 286 ! *Tate* ! *Hance*, 550, 703 ! *Wright*, 530 ! *Urquhart* !); KWANGSI (*Wenyon* !); HAINAN (*Henry*, 8385 !). Mus. Brit.; Herb. Kew.

Var. *densiflora*, *C. H. Wright*.

*Ophiopogon spicatus*, *Hook. f. in Bot. Mag.* t. 5348.

*Ophiopogon Muscari*, *Decne. in Fl. des Serres*, xvii. (1867-68) p. 181.

*Ophiopogon spicatus*, var. *communis*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 85; *Mél. Biol.* vii. p. 323; *Hance, in Journ. Bot.* xxi. (1883) p. 358.

*Liriope graminifolia*, var. *densiflora*, *Baker*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 500.

KIANGSU : Chinkiang (*Bullock ex Hance*); FORMOSA (ex *Baker*); LUCHU ARCHIPELAGO (*Wright*, 330!). *Mus. Brit.*; *Herb. Kew.*

Japan.

Var. *intermedia*, *Maxim.* fide *Franch. Pl. David.* p. 296.

KIANGSI : Kiukiang Mountains (*David*, 809, ex *Franchet*).

This may possibly be an error for *Ophiopogon japonicus*, var. *intermedius*, *Maxim.* in *Bull. Acad. St. Pétersb.* xv. (1871) p. 88; *Mél. Biol.* vii. p. 327.

Var. *koreana*, *C. H. Wright*.

*Ophiopogon spicatus*, var. *koreanus*, *Palibin*, in *Act. Hort. Petrop.* xix. (1901) p. 106.

COREA : Kyongkwi, Seoul (*Kalinowsky ex Palibin*).

Var. *latifolia*, *Franch. Pl. David.* p. 296.

KIANGSI : Kiukiang Mountains (*David*, 854, ex *Franchet*).

Var. *minor*, *C. H. Wright*.

*Ophiopogon spicatus*, *Lodd. Bot. Cab.* t. 694.

*Ophiopogon spicatus*, var. *minor*, *Maxim.* in *Bull. Acad. St. Pétersb.* xv. (1871) p. 85; *Mél. Biol.* vii. p. 327.

CHINA (ex *Loddiges*).

Japan.

## 138. IRIDEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Iris Bungei*, *Maxim.* in *Bull. Acad. St. Pétersb.* xxvi. (1880) p. 509; *Mél. Biol.* x. p. 695; *Franch. Pl. David.* i. p. 296; *Baker, Handb. Irideæ*, p. 5.

*Iris ventricosa*, *Maxim.* *Ind. Fl. Mongol. in Prim. Fl. Amur. Suppl.* p. 485, non *Pall.*

KIANGSI : Kiukiang (*Carles*, 447 !); KANSUH *occidentalis* (*Potanin* !), Tangut (*Przewalski* !). *Herb. Kew.*

Mongolia.

2. *Iris Delavayi*, *Micheli*, in *Rev. Hort.* (1895) p. 398, figs. 128-9; *Hook. f. Bot. Mag.* t. 7661.

YUNNAN (*Delavay ex Micheli*).



3. *Iris dichotoma*, *Pall. Reise*, iii. p. 712 (1776); *Linn. fl. Suppl.* p. 97 (1781); *Bot. Reg.* t. 246; *Sweet, Brit. Flow. Gard.* t. 96; *Bunge, Enum. Pl. Chin. Bor.* p. 63; *Hance, in Journ. Bot.* xiii. (1875) p. 104; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1877) p. 142; *Bot. Mag.* t. 6428; *Franch. Pl. David.* p. 298; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 540; *Mél. Biol.* x. p. 738.

*Iris pomeridiana*, *Fisch. ex Ledeb. Fl. Ross.* iv. p. 106.

*Pardanthus dichotoma*, *Ledeb. Fl. Ross.* iv. p. 106.

*Evansia dichotoma et E. vespertina*, *Decne. in Bull. Soc. Bot. France*, xx. (1873) p. 302.

CHIHLI: Peking (*Bushell! Williams! Hance, 12632! David, 442, ex Franchet, Bretschneider!*), Jehol (*David, 2165, ex Franchet*), Siaowutaishan (*Hancock ex Maximowicz*); SHINGKING: Moukden (*Webster, 52!*); SHANTUNG: Chefoo (*Forbes ex Hance*). *Mus. Brit.*; *Herb. Kew.*

*Dahuria*, Mandshuria, Mongolia.

4. *Iris ensata*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 328; *Baker, in Journ. Linn. Soc., Bot.* xvi. (1877) p. 139, *et* xvii. (1879) p. 387; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 512; *Mél. Biol.* x. p. 699; *Hance, in Journ. Bot.* xi. (1883) p. 358; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 258; *Baker, Handb. Irideæ*, p. 8; *Hook. f. Fl. Brit. Ind.* vi. p. 272; *Kanitz, in Bot. Res. Széch. Centr. As. Exped.* p. 58.

*Iris graminea*, *Thunb. Fl. Jap.* p. 34, *non Linn.*

*Iris triflora*, *Balbis, Misc. Bot.* p. 6, t. 1.

*Iris biglumis*, *Vahl, Enum.* ii. p. 149; *Sweet, Brit. Flow. Gard.* ser. 2, t. 187.

*Iris Pallasii*, *Fisch. in Trev. Ind. Sem. Hort. Vrat.* 1821.

*Iris Pallasii*, var. *chinensis*, *Fisch. in Bot. Mag.* t. 2331.

*Iris longispatha*, *Fisch. in Bot. Mag.* t. 2528.

*Iris Moorcroftiana*, *Wall. Cat.* no. 5021.

*Iris oxypetala*, *Bunge, Enum. Pl. Chin. Bor.* p. 63, *et in C. A. Meyer, Ind. Pl. Mus. Petrop.* (1834) p. 8, *non Hance.*

*Iris caricifolia*, *Pallas ex Link, Jahrb.* i. III. (1820) p. 72.

*Iris longifolia*, *Royle, Ill. Bot. Himal.* p. 372, t. 91. f. 2.

*Iris fragrans*, *Lindl. in Bot. Reg.* (1840) t. 1.

*Iris Doniana*, *Spach, Hist. Phan.* xiii. p. 34.

*Iris ensata*, var. *chinensis*, *Maxim. in Gartenfl.* t. 1011; *Bull. Acad. St. Pétersb.* xxvi. (1880) p. 514; *Mél. Biol.* x. p. 701.

CHIHLI: Peking (*Bullock, 193! Bretschneider, 723! Bushell! Hancock, 22!*), Jehol (*David, 1744!*); SHINGKING: Moukden

(*James*!), Tieling (*Webster*, 71!), Fooling (*Ross*, 332!); SHANTUNG: Chefoo, Tungtschaufu (*Hancock* ex *Maximowicz*); FOKIEN: Foochow (*Wykeham Perry*!); SHENSI (*David* ex *Franchet*, *Piasezki* ex *Maximowicz*); KANSUH (*Piasezki*, *Przewalski* ex *Maximowicz*), Kanessoufu (*Loczy*, 74, ex *Kanitz*); COREA (*Wykeham Perry*!). Mus. Brit.; Herb. Kew.

North Asia, Japan, Cashmir.

The variety *chinensis* differs only in having lighter flowers with narrower perianth-segments.

5. *Iris gracilis*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 527; *Mél. Biol.* x. p. 720; *Baker, Handb. Irideæ*, p. 28.

KANSUH: Tangut (*Przewalski*, 3, 10!); SZECHUEN: Tachienlu (*Pratt*, 262!), Tongola (*Soulié*, 876!). Herb. Kew.

Tibet.

6. *Iris Grijsi*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 515; *Mél. Biol.* x. p. 702; *Hance, in Journ. Bot.* xx. (1882) p. 295; *Baker, Handb. Irideæ*, p. 6; *Franch. in Bull. Soc. Bot. France*, xlvi. (1899) p. 213.

*Iris oxypetala*, *Hance, in Journ. Bot.* viii. (1870) p. 314, non *Bunge*.

ANWHEI: Wuhu (*Bullock*!); FOKIEN (*De Grijs* in Herb. *Hance*, 8583! & 21351!), Kuatun (*De la Touche* ex *Franchet*). Mus. Brit.; Herb. Kew.

7. *Iris Henryi*, *Baker, Handb. Irideæ*, p. 6.

HUPEH: Nanto (*Henry*, 6372!). Mus. Brit.; Herb. Kew.

8. *Iris japonica*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 327; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 141; *Baker, in Gard. Chron.* ser. ii. vi. (1876) p. 37; *Journ. Linn. Soc., Bot.* xvi. (1877) p. 143; *Handb. Irideæ*, p. 24; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 41; *Franch. Pl. David.* i. p. 298, et ii. p. 127; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 538; *Mél. Biol.* x. p. 736.

*Iris squalens*, *Thunb. Fl. Jap.* p. 33, non *Linn.*

*Iris chinensis*, *Curt. in Bot. Mag.* t. 373; *Regel, in Gartenfl.* t. 511. fig. 2-3.

*Iris fimbriata*, *Vent. Hort. Cels.* t. 9; *Red. Lil.* t. 152.

*Isis fimbriata*, *Tratt. Arch. Gewächsk.* p. 668.

*Evansia chinensis*, *Salisb. in Trans. Hort. Soc. Lond.* i. (1812) p. 303.

*Evansia fimbriata*, *Decne. in Bull. Soc. Bot. France*, xx. (1873) p. 302.

CHIHLI: Peking (*David ex Franchet*); HUPEH: Ichang (*Henry*, 786, 1161, 1254!), Nanto (*Henry*, 1967!), Yangtsekiang (*Gill*!); KIANGSI: Kiukiang (*Maries! Shearer*!); KWANGTUNG: Canton (*Hance*, 16441! cult.); CHINA (*Fortune*, 41!). Mus. Brit.; Herb. Kew.

Japan, Tibet.

9. *Iris lævigata*, *Fisch. in Ind. Sem. Hort. Petrop.* v. (1839) p. 36; *Gartenfl.* t. 442. fig. 1; *Bot. Mag.* t. 6132; *Baker, in Journ. Linn. Soc., Bot.* xvi. (1877) p. 142; *Baker, Handb. Irideæ*, p. 13.

*Iris Kämpferi*, *Sieb. in Ill. Hort.* (1858) t. 157; *Fl. des Serres*, tt. 2073-4.

*Iris Itsihatsi*, *Hassk. Cat. Hort. Bogor.* p. 35.

*Iris violacea*, *Klatt, in Linnæa*, xxxv. (1867-68) p. 384.

*Iris lævigata*, var. *Kämpferi*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 521; *Mél. Biol.* x. p. 712.

SHINGKING: Moukden (*James! Ross!*), Toonghwahien (*Ross!*); COREA: Port Chusan (*Wilford*, 973!). Mus. Brit.; Herb. Kew.  
Amur, Japan.

10. *Iris Loczyi*, *Kanitz, Bot. Res. Széch. Centr. As. Exped.* p. 58, t. 6. fig. ii. 5.

KANSUH: Nanszan (*Széchenyi ex Kanitz*).

Probably not distinct from *Iris tenuifolia*, Pall.

11. *Iris pandurata*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 529; *Mél. Biol.* x. p. 723; *Baker, Handb. Irideæ*, p. 29.

KANSUH: Tangut (*Przewalski!*). Herb. Kew.

12. *Iris Potanini*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 528; *Mél. Biol.* x. p. 721; *Baker, Handb. Irideæ*, p. 27.

*Iris flavissima*, var. *rupestris*, *Ledeb. Fl. Alt.* i. p. 60.

KANSUH (*Potanin!*). Herb. Kew.

Dahuria, Altai, Mongolia.

13. *Iris Rossii*, *Baker, in Gard. Chron.* (1877) ii. p. 809; *Journ. Linn. Soc., Bot.* xvii. (1879) p. 387; *Handb. Irideæ*, p. 3; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 523; *Mél. Biol.* x. p. 714.

SHINGKING: Changdien (*Ross*, 160, 162!); ANWHEI: Nanking (*Faber*, 1542!); KIANGSI: Kiukiang (*Maries!*); COREA (*Carles*, 5!); Chemulpo (*Carles!*), Seoul (*Carles!*). Herb. Kew.

14. *Iris ruthenica*, *Dryand. in Ait. Hort. Kew.* ed. 2, i. p. 117; *Gawl. in Bot. Mag.* tt. 1123 et 1393; *Ledeb. Fl. Ross.* i. p. 55; *Maxim. Prim. Fl. Amur. Suppl.* pp. 477 et 485; *Bull. Acad. St. Pétersb.* xxvi. (1880) p. 516; *Mél. Biol.* x. p. 704; *Hance, in Journ. Bot.* xiii. (1875) p. 139; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 387; *Franch. Pl. David.* i. p. 297; *Baker, Handb. Irideæ*, p. 4.

*Iris cæspitosa*, *Pallas, in Spreng. et Link, Jahrb.* i. III. (1820) p. 71.

*Ioniris ruthenica*, *Klatt, in Bot. Zeit.* xxx. (1872) p. 502.

CHIHLI: Peking (*Bullock*, 234!), Jehol (*David*, 1741, ex *Franchet*), Sangyu (*David*, 2235, ex *Franchet*), Pohuashan (*Bretschneider* ex *Hance*), Kalgan (*Przewalski* ex *Maximowicz*); SHINGKING: Ngaiyang (*Ross*, 228!). Mus. Brit.; Herb. Kew. North Asia.

Var. *nana*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 516; *Mél. Biol.* x. p. 705.

CHIHLI: Peking, Lunzuanssy (*Tatarinow* ex *Maximowicz*), Pohuashan (*Bretschneider* ex *Maximowicz*); SHANSI (*Potanin*!). Herb. Kew.

South Mongolia.

15. *Iris sibirica*, var. *orientalis*, *Baker, in Journ. Linn. Soc., Bot.* xvi. (1877) p. 139; *Handb. Irideæ*, p. 9; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 519; *Mél. Biol.* x. p. 709.

*Iris orientalis*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 328.

*Iris sibirica*, var. *sanguinea*, *Gawl. in Bot. Mag.* t. 1604.

*Iris hæmatophylla*, *Fisch. in Sweet, Brit. Flow. Gard.* t. 118.

*Iris nertchinskia*, *Lodd. Bot. Cab.* t. 1843.

SHINGKING: Moukden (*James! Webster!*), Toonghwahien (*Ross*, 48!); SZECHUEN (*Pratt*, 247!); COREA: Tsusima (*Wilsford*, 752!), Chemulpo (*Carles*, 177!). Mus. Brit.; Herb. Kew. Japan, Eastern Siberia.

16. *Iris songarica*, var. *gracilis*, *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 510; *Mél. Biol.* x. p. 696; *Baker, Handb. Irideæ*, p. 5.

KANSUH: Tangut (*Przewalski*!). Herb. Kew.

17. *Iris speculatrix*, *Hance, in Journ. Bot.* xiii. (1875) p. 196, and xiv. (1876) p. 75; *Baker, in Gard. Chron.* ser. ii. vi. (1876)

p. 36; and *Bot. Mag.* t. 6306; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 538; *Mél. Biol.* x. p. 736.

HONGKONG: between Victoria Peak and Mount Davis (*Ford* in *Herb. Hance*, 783 A! 18465!); KWANGTUNG: Canton, Lienchau (*Lamont!*), Lantao (*Ford*, 53!). *Mus. Brit.*; *Herb. Kew.*

18. *Iris tectorum*, *Maxim. in Bull. Acad. St. Pétersb.* xv. (1871) p. 380; *Mél. Biol.* vii. p. 563, et x. p. 736; *Gartenfl.* t. 716; *Hook. f. Bot. Mag.* t. 6118; *Fl. des Serres*, t. 2287; *Baker, in Gard. Chron.* ser. ii. vi. (1876) p. 37; *Handb. Irideæ*, p. 23; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 41; *Franch. Pl. David.* i. p. 297; *Kanitz, Bot. Res. Széch. Centr. As. Exped.* p. 59.

*Iris chinensis*, *Bunge, Enum. Pl. Chin. Bor.* p. 64, non *Curt.*

*Iris fimbriata*, *Klatt, in Linnæa*, xxxiv. (1865-66) p. 619, *partim.*

*Iris cristata*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 141, non *Soland.*

*Iris tomiolopha*, *Hance, in Journ. Bot.* x. (1872) p. 229.

SHANTUNG (*Maingay*, 742!); HUPEH: Changlo (*Henry*, 6330!), Ichang (*Henry*, 3970!), Patung (*Henry*, 3781!); HUNAN (*Hancock*, 80!); SHENSI (*Piasezki ex Maximowicz, David ex Franchet*); KANSUH (*Piasezki ex Maximowicz*); SZECHUEN (*Lóczy*, 122, ex *Kanitz*). *Mus. Brit.*; *Herb. Kew.*

Japan.

This species has been cultivated about Peking and Canton.

19. *Iris tenuifolia*, *Pall. Reise*, iv. p. 683, t. 7. fig. 2; *Maxim. Ind. Fl. Mongol. in Prim. Fl. Amur.* p. 485; in *Bull. Acad. St. Pétersb.* xxvi. (1880) p. 511; *Mél. Biol.* x. p. 697; *Baker, in Gard. Chron.* ser. ii. vi. (1876) p. 144; *Journ. Linn. Soc., Bot.* xvi. (1877) p. 138; *Handb. Irideæ*, p. 4.

*Neubeckia tenuifolia*, *Alefeld, in Bot. Zeit.* xxi. (1863) p. 297.

*Joniris tenuifolia*, *Klatt, in Bot. Zeit.* xxx. (1872) p. 502.

KANSUH (*Potanin! Przewalski ex Maximowicz*). *Herb. Kew.*  
Central Asia.

20. *Iris Tigridia*, *Bunge, in Ledeb. Fl. Alt.* i. p. 60; *Ledeb. Icon. Pl. Ross.* iv. p. 342; *Maxim. in Bull. Acad. St. Pétersb.* xxvi. (1880) p. 530; *Mél. Biol.* x. p. 722; *Franch. Pl. David.* i. p. 297; *Baker, Handb. Irideæ*, p. 28.

CHIHLI: Jehol (*David*, 1696, ex *Franchet*); KANSUH (*Potanin!*). *Herb. Kew.*

Altai, Eastern Siberia, and Mongolia.

1. **Belamcanda punctata**, *Moench, Meth.* p. 529 (1794); *Franch. Pl. David.* i. p. 299.

*Ixia chinensis*, *Linn. Sp. Pl.* ed. 1, p. 36; *Curt. Bot. Mag.* t. 171; *Lour. Fl. Cochinch.* p. 36.

*Belamcanda chinensis*, *Leman, in Red. Lil.* t. 121 (1807); *Baker, in Journ. Linn. Soc., Bot.* xvi. (1877) p. 113; *Baker, Handb. Irideæ*, p. 117; *Hook. f. Fl. Brit. Ind.* vi. p. 277.

*Moræa chinensis*, *Thunb. Fl. Jap.* p. 34.

*Pardanthus chinensis*, *Ker, in Kon. Ann. Bot.* i. (1805) p. 246; *Benth. Fl. Hongk.* p. 365; *Fl. des Serres*, t. 1632; *Hance, in Journ. Bot.* xii. (1874) p. 262; *Debeaux, Fl. Tchéf.* p. 133; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 256.

SHANTUNG: Chefoo (*Fauvel ex Franchet*); FORMOSA (*Oldham*, 642! *Watters*, 70! *Wilford*, 550! *A. Henry*, 968!); Kiangsi: Kiukiang (*Shearer! Möllendorf ex Hance, David ex Franchet*); HUPEH: Ichang (*Henry*, 96!), Patung (*Henry*, 4905!), Nanto (*Henry*, 2112 A!); SHENSI (*David ex Franchet*); SZECHUEN: Chung Ching (*Bourne*, 207!); HONGKONG (*Champion*, 284! *Hance*, 718! *Lamont*, 783!); COREA: Chemulpo (*Carles*, 193! *Wykeham Perry!*). *Mus. Brit.*; *Herb. Kew.*

Mandshuria, Japan, North India.

### 139. AMARYLLIDEÆ.

(By C. H. WRIGHT, A.L.S.)

1. **Hypoxis aurea**, *Lour. Fl. Cochinch.* p. 200; *Hance, in Journ. Bot.* viii. (1870) p. 276, et 1878, p. 111; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1877) p. 108; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 75; *Hook. f. Fl. Brit. Ind.* vi. p. 277; *Franch. in Bull. Soc. Bot. France*, xlvi. (1901) p. 214.

*Hypoxis curculigoides*, *Wall. Cat.* n. 5164.

*Hypoxis minor*, *D. Don, Prodr. Fl. Nepal.* p. 53; *Royle, Ill. Bot. Himal.* p. 376, t. 91. fig. 3.

*Curculigo graminifolia*, *Nimmo, in J. Graham, Bomb. Cat.* p. 215; *Dalzell & Gibson, Bombay Flora*, p. 276.

*Hypoxis Franquevillei*, *Miq. Fl. Ind. Bat.* iii. p. 586.

HUPEH: South Patung (*Henry*, 6242!); FOKIEN: Kuantun (*De la Touche ex Franchet*); KWANGTUNG: Canton (*Sampson*, 1624! et in hb. *Hance*, 16516!); LUCHU ARCHIPELAGO (*Wright*, 344!). *Mus. Brit.*; *Herb. Kew.*

North India, Cochin China, Java, Japan.

1. **Curculigo orchioides**, *Gaertn. Fruct. et Sem.* i. p. 63, fig. 11; *Roxb. Pl. Corom.* i. p. 14, t. 13; *Ait. Hort. Kew.* ed. 2, ii. p. 253; *Roem. et Schult. Syst. Veget.* vii. p. 756.

*Curculigo ensifolia*, *R. Br. Prodr.* p. 290; *Maxim. Fl. As. Or. Fragm.* p. 61; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 75.

*Curculigo stans*, *Labill. Sert. Austr. Caled.* i. p. 18, t. 24.

*Curculigo malabarica*, *Wight, Ic. Pl. Ind. Or.* t. 2043.

*Curculigo orchioides*, var. *minor*, *Benth. Fl. Hongk.* p. 366; *Bot. Mag.* t. 1076.

*Hypoxis minor*, *Seem. Bot. Voy. 'Herald,'* p. 419.

*Hypoxis orchioides*, *Kurz, in Ann. Mus. Bot. Lugd.-Bat.* iv. p. 177.

CHEKIANG: Ningpo (*Hancock ex Maximowicz*); SZECHUEN: Min River (*Faber, 959!*); KWANGTUNG: Lofaushan (*Ford!*), Canton (*Hance ex Maximowicz*); HONGKONG (*Champion, 533!*); LUCHU ARCHIPELAGO (*Wright, 343!*). *Herb. Kew.*

Japan, Northern India to Australia.

A very variable species, with extreme forms, differing much in size and in the breadth of the leaves, yet presenting every stage of transition, so that no satisfactory line can be drawn between them.

2. **Curculigo recurvata**, *Dryand. in Ait. Hort. Kew.* ed. 2, ii. p. 253; *Wall. Cat.* n. 5159; *Bot. Reg.* t. 770; *Roxb. Fl. Ind.* ii. p. 145; *Miq. Fl. Ind. Bat.* iii. p. 585; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 75; *Hook. f. Fl. Brit. Ind.* vi. p. 278.

*Molineria recurvata*, *Herb. Amaryll.* p. 84; *Kurz, in Ann. Mus. Bot. Lugd.-Bat.* iv. p. 175; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 120; *Hance, in Journ. Bot.* xxi. (1883) p. 358.

*Molineria capitulata*, *Herb. Amaryll.* p. 84.

*Molineria plicata*, *Colla, Hort. Rip. App.* ii. p. 333, t. 18.

*Leucojum capitulatum*, *Lour. Pl. Cochinch.* p. 199.

CHEKIANG: Ningpo (*Hancock ex Maximowicz*); FORMOSA: Tamsui (*Oldham, 645!*); KWANGTUNG: Lofaushan (*Ford, 61!*), Siushi-Lienchan (*B. C. Henry!*), Canton (*Hance ex Maximowicz*); HONGKONG: Mount Parker (*Ford, 56!*). *Mus. Brit.; Herb. Kew.*

India, Ceylon, Malaya, Borneo, Queensland.

1. **Narcissus Tazetta**, var. *chinensis*, *M. Roem. Syn. Monog.* fasc. iv. p. 223; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 140; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 45; *Franch. Fl. Tchéf.*

p. 256; *Debeaux, Fl. Tchéf.* p. 133; *Fl. Shangh.* p. 63; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 75.

CHIHLI: Peking (*Bretschneider ex Maximowicz*); SHANTUNG: Chefoo (*Debeaux, Fauvel ex Franchet*); KIANGSU: Shanghai (*Debeaux*), Saddle Island (*Weyrich ex Maximowicz*).

Japan.

1. *Crinum asiaticum*, var. *declinatum*, *Herb. Amaryll.* p. 244; *Kunth, Enum.* v. p. 549; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 139; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 45; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 77; *Baker, Handb. Amaryll.* p. 75.

*Crinum declinatum*, *Herb. in Bot. Mag.* t. 2231.

*Crinum latifolium*, *Thunb. Fl. Jap.* p. 131.

*Crinum asiaticum*, *Benth. Fl. Hongk.* p. 366.

HONGKONG: Saywan (*Champion ex Benth.*), Skeko (*Hance, 1332!*). Mus. Brit.

Var. *sinicum*, *Baker, Handb. Amaryll.* p. 75.

*Crinum sinicum*, *Roxb. ex Herb. in Bot. Mag.* sub t. 2121, p. 7; *Herb. Amaryll.* p. 244; *Kunth, Enum.* v. p. 550; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 77.

FORMOSA: Tamsui (*Hancock, 20!*); HAINAN: Hoihow (*Hancock, 19!*). Herb. Kew.

Var. *anomalum*, *Herb. Amaryll.* p. 244; *Kunth, Enum.* v. p. 549; *Baker, Handb. Amaryll.* p. 75.

*Crinum anomalum*, *Herb. in Bot. Mag.* sub t. 2120, p. 5; *Herb. App.* p. 22.

*Crinum plicatum*, *Livingst. ex Hook. in Bot. Mag.* t. 2908.

CHINA (*Livingstone ex Hooker*).

2. *Crinum Loureirii*, *M. Roem. Syn. Ensat.* fasc. iv. p. 85; *Kunth, Enum.* v. p. 581; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 77.

*Crinum zeylanicum*, *Lour. Fl. Cochinch.* p. 198, *nec Linn.*

CHINA (ex *Loureiro*).

Cochinchina.

This species has not been found by recent collectors. From the figure in Rumphius's *Hort. Amboin.* viii. p. 306, t. 105 (quoted by Loureiro), it would appear to be closely allied to *Crinum zeylanicum*, Linn.



1. **Lycoris aurea**, *Herb. Append.* p. 20; *Kunth, Enum.* v. p. 545; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 139; *Hance, in Journ. Bot.* xxi. (1874) p. 262; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 44; *Franch. Pl. David.* i. p. 299; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 79; *Baker, Handb. Amaryll.* p. 40.

*Amaryllis aurea*, *L'Hérit. Sert. Angl.* p. 14, t. 15 b; *Jacq. Hort. Schoenb.* t. 73; *Bot. Mag.* t. 409; *Redouté, Lil.* t. 61; *Bot. Reg.* t. 611; *Roxb. Fl. Ind.* ii. p. 140.

*Nerine aurea*, *Bury, Hevand.* t. 3.

KIANGSU: Shanghai (*Maingay*, 426!); Kiangsi: Kiukiang (*Bullock*, 251! *Carles*, 499, 505! *Moellendorff*! *David*, 804, ex *Franchet*); FORMOSA (*Hancock*, 3!), Tamsui (*Oldham*, 643!); HUPEH: Ichang (*Henry*, 2326, 2326 B!). *Mus. Brit.*; *Herb. Kew.*  
Cultivated in Japan.

2. **Lycoris radiata**, *Herb. App.* p. 20; *Kunth, Enum.* v. p. 546; *Hance, in Journ. Bot.* xxi. (1874) p. 262; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 44; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 78; *Baker, Handb. Amaryll.* p. 40.

*Amaryllis radiata*, *L'Hérit. Sert. Angl.* p. 16; *Bot. Reg.* t. 596; *Andr. Bot. Rep.* t. 95.

*Amaryllis sarniensis*, *Thunb. Fl. Jap.* p. 131, non *Linn.*; *Lour. Fl. Cochinch.* p. 200.

*Nerine japonica*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 139; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 44.

KIANGSU: Fengwangshan (*Poli ex Franchet*); CHEKIANG: Ningpo (*Hancock*, 33!); Kiangsi: Kiukiang (*Shearer*! *Bullock*, 88!); FOKIEN: Chinchew (*Fortune*, 148!); HUPEH: Ichang (*A. Henry*, 428, 1094, 2326 A, 2326 C!); KWANGSI (*Graves*!). *Mus. Brit.*; *Herb. Kew.*

Japan.

3. **Lycoris sanguinea**, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 80; *Baker, Handb. Amaryll.* p. 40.

CHEKIANG: Ningpo (*Forbes ex Maximowicz*). Cultivated.  
Japan.

4. **Lycoris squamigera**, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 79; *Baker, Handb. Amaryll.* p. 40.

CHIHLI: Peking (*Bretschneider ex Maximowicz*); Kiangsu: Shanghai (*Carles*, 417!); CHEKIANG: Ningpo (*Hancock*, 34!). *Herb. Kew.*

Japan.

Perhaps occurring in China only as a cultivated plant.

5. *Lycoris straminea*, *Lindl. in Journ. Hort. Soc. Lond.* iii. (1848) p. 76; *Walp. Ann.* i. p. 834 et iii. p. 616; *Kunth, Enum.* v. p. 546; *Baker, Handb. Amaryll.* p. 40.

CHINA (*Fortune ex Lindley*).

1. *Pancratium biflorum*, *Roxb. Fl. Ind.* ii. p. 125; *Kunth, Enum.* v. p. 661; *Benth. Fl. Hongk.* p. 367; *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 81; *Hook. f. Fl. Brit. Ind.* vi. p. 285; *Baker, Handb. Amaryll.* p. 119.

HONGKONG (*Champion, Hance ex Benth*).

Northern India.

*Agave americana*, Linn., *Agave lurida*, Jacq., and *Pancratium zeylanicum*, Linn., have been cultivated in China, but not found in a wild state.

#### 140. TACCACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Tacca pinnatifida*, *Forst. Char. Gen.* p. 70, t. 35; *Lour. Fl. Cochinch.* p. 300, *excl. syn. Tacca sativa, Rumph.*

CHINA *ex Loureiro*.

This has not been gathered by any recent collector, and its sole claim to be regarded as a Chinese plant rests on the statement of Loureiro.

Tropical Asia, Africa, Australia, and Polynesia.

1. *Schizocapsa plantaginea*, *Hance, in Journ. Bot.* xix. (1881) p. 292.

KWANGTUNG: North River, Canton (*Faber ex Hance, Ford, 376! Gerlash!*), Lienchau (*B. C. Henry!*). Mus. Brit.; Herb. Kew.

#### 141. DIOSCOREACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Dioscorea alata*, *Linn. Sp. Pl.* ed. 1, p. 1033; *Lour. Fl. Cochinch.* p. 623; *Kunth, Enum.* v. p. 387; *Wight, Ic. Pl. Ind. Or.* t. 810; *Hook. f. Fl. Brit. Ind.* vi. p. 296; *Bretschneider, Early Res.* p. 177.

CHINA (*ex Loureiro*).

Tropical India.

2. *Dioscorea Batatas*, *Decne. in Rev. Hort.* (1854) p. 243, and (1855) p. 69 cum icone; *Benth. Fl. Hongk.* p. 368; *Makino, Phan. Jap. Ic. Ill.* tt. 93-95.

CHIHLI: Peking (*Bretschneider!*); KIANGSU: Shanghai (*Faber!*); KIANGSI: Kiukiang (*Shearer! Bullock, 246!*); KWANGTUNG: Canton (*Sampson, 13760!*); HONGKONG (*Wright!*), Lantao Island (*Lamont, 788!*). *Mus. Brit.*; *Herb. Kew.*

3. *Dioscorea bulbifera*, *Linn. Sp. Pl.* ed. 1, p. 1033; *Miq. Journ. Bot. Néerl.* i. p. 90.

KWANGTUNG (*Krone ex Miquel*).

Tropical Asia.

4. *Dioscorea dæmona*, var. *reticulata*, *Hook. f. Fl. Brit. Ind.* vi. p. 289.

*Dioscorea macrocarpa*, *Wall. Cat.* n. 5100.

FORMOSA: Takow (*A. Henry, 1875!*); HAINAN (*A. Henry, 8106, 8288, 8725!*). *Herb. Kew.*

Malay Peninsula.

5. *Dioscorea Decaisneana*, *Carr. in Rev. Hort.* (1865) p. 111, cum icone; *Chappellier, in Bull. Soc. Acclim.* v. (1888) pp. 373-5; *Bretsch. Hist. Europ. Bot. Disc.* p. 832.

CHINA: without locality (*ex Carrière*).

6. *Dioscorea doryophora*, *Hance, in Ann. Sc. Nat.* 5<sup>me</sup> sér. v. (1866) p. 244.

*Dioscorea Swinhoei*, *Rolfe, in Journ. Bot.* xx. (1882) p. 359.

FORMOSA (*Swinhoe, 33! 7573 in Herb. Hance! Playfair!*), Takow (*A. Henry, 1922!*), Bankinsing (*A. Henry, 878, 1672!*). *Herb. Kew.*

7. *Dioscorea glabra*, *Roxb. Hort. Beng.* p. 72; *Fl. Ind.* iii. p. 804; *Kunth, Enum. Pl.* v. p. 383; *Hook. f. Fl. Brit. Ind.* vi. p. 294.

*Dioscorea sagittata*, *Royle, ex Hook. t. l. c.*

KIANGSI: Lushan Mountains (*Bullock, 245!*); KWANGTUNG: Lantao (*Ford, 6!*), mainland opposite Hongkong (*Ford, 183!*); HAINAN (*A. Henry, 8407, 8690!*).

From the sub-tropical Himalayas southwards to the Concan and Malay Peninsula.

8. *Dioscorea japonica*, *Thunb. Fl. Jap.* p. 151; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 159 excl. syn.; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 47 excl. syn.; *Somoku Zusetsu*, xx. tt. 56-7.

CHEKIANG : Ningpo (*Faber*, 80 !); KIANGSI : Kiukiang (*Shearer* !); FORMOSA (*Oldham*, 573 ! *Wilford*, 561 !), Tamsui (*Watters*, 37 !); KWANGTUNG : Lofaushan (*Ford*, 330 !). *Mus. Brit.*; *Herb. Kew.*

“Shanyotsai, a valuable medicine.”—*Watters.*

Japan.

9. *Dioscorea oppositifolia*, *Linn. Sp. Pl.* ed. 1, p. 1033; *Lour. Fl. Cochinch.* p. 624; *Wight, Ic. Pl. Ind. Or.* t. 813; *Kunth, Enum. Pl.* v. p. 390; *Benth. Fl. Hongk.* p. 367; *Hook. f. Fl. Brit. Ind.* vi. p. 292.

FORMOSA : Bankinsing (*A. Henry*, 1673 !); HONGKONG (*Wright*, 528 ! *Ford*, 24 ! *Lamont*, 787 !), Tytam (*Hance*, 458 !). *Mus. Brit.*; *Herb. Kew.*

Tropical India.

10. *Dioscorea polystachya*, *Turcz. in Bull. Soc. Nat. Mosc.* (1837) no. 7, p. 158; *Kunth, Enum. Pl.* v. p. 403; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478.

N. CHINA (*Kirilow*, 198, ex *Turczaninow*); CHIHLI : Peking (ex *Maximowicz*).

I have seen no specimen of this species, which, from *Turczaninow*'s brief description, must be very closely allied to *D. doryophora*, *Hance*. The following may belong here:—*Ichang* (*A. Henry*, 1538, 2023 !).

11. *Dioscorea quinqueloba*, *Thunb. Fl. Jap.* p. 150, non *Vell.*; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Kunth, Enum. Pl.* v. p. 350; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 46; *Franch. Pl. David.* i. p. 300; *Banks, Icon. Kæmpf.* t. 15; *Houtt. Handl.* xi. p. 358, t. 74. figg. 1-2; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 159.

CHIHLI : Peking (*Bretschneider*, 732 ! *David*, 14560 !); SHINGKING : between Tungcheshien (*James* !); HUPEH : Chien-shih (*A. Henry*, 5870, 5870 B !), Patung (*A. Henry*, 4769, 7358 !), *Ichang* (*A. Henry*, 183 !). *Mus. Brit.*; *Herb. Kew.*

Japan and Amur.

12. *Dioscorea rhipogonioides*, *Oliver*, in *Hook. Ic. Pl.* t. 1868.

FORMOSA: South Cape (*A. Henry*, 589, 970, 1297!), Bankinsing (*A. Henry*, 105!); HONGKONG (*Ford*, 10, 54/88!). Herb. Kew.

“The root of this climber is called Shulang, and is much used for dyeing and tanning nets, &c.”—*A. Henry*.

13. *Dioscorea sativa*, *Linn. Sp. Pl.* ed. 1, p. 1033, & *Hort. Cliff.* t. 28; *Benth. Fl. Hongk.* p. 368; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 47; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Debeaux, Fl. Shangh.* p. 65, & *Flor. Tchef.* p. 140; *Duthie, Field Crops*, t. 80; *Hook. f. Fl. Brit. Ind.* vi. p. 295.

SHANTUNG: Chefoo, cult. (ex *Debeaux*); KIANGSU: Shanghai, cult. (ex *Debeaux*); FORMOSA: Tamsui (*Oldham*, 574!), Takow (*A. Henry*, 1871 A & B!), South Cape (*A. Henry*, 1210!), Apes Hill (*Playfair*, 188!); HUPEH: Ichang (*A. Henry*, 3641, 3641 A, 4193!), Nanto (*A. Henry*, 2068!); SZECHUEN: Mount Omei (*Faber*, 281!); KWANGTUNG: Lofaushan (*Ford*!); HONGKONG (*Hance ex Bentham, Lamont*, 789!). Mus. Brit.; Herb. Kew.

Widely dispersed through the tropics of both hemispheres.

14. *Dioscorea spinosa*, *Roxb. ex Wall. Cat.* n. 5103; *Hook. f. Fl. Brit. Ind.* vi. p. 291.

*Dioscorea tiliæfolia*, *Kunth, Enum. Pl.* v. p. 401.

FORMOSA: Bankinsing (*A. Henry*, 856!). Herb. Kew.

Tropical India, Malaya, and Tropical America.

15. *Dioscorea zingiberensis*, *Wright*, n. sp.

*Caulis* suffruticosus, tenuis, levis, inerimis. *Folia* alterna, 2 poll. longa, 5-7-nervia, 3-(rarissime 5-)loba; lobus centralis triangularis, mucronatus; lobi laterales rotundati plus minusve deflexi; petiolus 1 poll. longus. *Spicæ masculæ* solitariae petiolis paulo longiores. *Perianthium* 1 lin. diam., lobis 6 ovatis, rarius emarginatis. *Stamina* 6; antheræ extrorsæ, latæ. *Racemi feminei* solitarii. *Perianthii* lobi 6, late ovati, imbricati. *Staminodia* minuta. *Stylus* 3-fidus. *Ovarium* oblongum, trilobulare. *Fructus* 1 poll. diam., æqualiter trilobatus. *Semen* ala membranacea circumdatum.

HUPEH: Ichang (*A. Henry*, 407, 1520, 1621!). Mus. Brit.; Herb. Kew.

This differs from *D. japonica*, Thunb., in its alternate leaves, and from *D. doryophora*, Hance, in its solitary male spikes.

“Creeper, with dark red flowers and green fruit, growing many feet in length. The root is eaten in winter. The local name is ‘Hjang Chiang,’ *i. e.* yellow ginger.”—*A. Henry*.

16. *Dioscorea* sp.

COREA: Chemulpo (*Carles*, 178!). Herb. Kew.

This is probably a new species, but the material is insufficient for specific description. The lower leaves are verticillate, with slender petioles as long as the cordate acute laminae. The male spikes are solitary in the axils of the alternate upper leaves.

17. *Dioscorea* sp.

HUPEH: Tunghu (*A. Henry*, 6419!), Changyang (*A. Henry*, 7501!), Nanto (*A. Henry*, 2666! 4486!); SZECHUEN: North Wushan (*A. Henry*, 7103!). Herb. Kew.

Resembling *D. pentaphylla*, Linn., and *D. tomentosa*, Linn., but not determinable, the specimens bearing leaves and very young fruit only.

142. ROXBURGHACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Stemona acuta*, C. H. Wright, in *Journ. Linn. Soc., Bot.* xxxii. (1896) p. 493.

KWANGTUNG (*Ford*, 283!). Herb. Kew.

Differs from *Stemona japonica*, Franch. & Savat., in having very acute leaves with cordate bases, and in the peduncles not being adnate to the petiole.

2. *Stemona erecta*, C. H. Wright, in *Kew Bulletin* (1895), p. 117; *Hook. Ic. Pl.* t. 2389; *Journ. Linn. Soc., Bot.* xxxii. (1896) p. 493.

ANWHEI: Nanking (*Faber*, 1541!). Herb. Kew.

A species with erect stems and verticillate leaves, somewhat resembling *Stemona sessilifolia*, Miq., figured in the ‘Somoku Zusetsu,’ vol. ii. t. 55, but with more flowers on shorter pedicels. The anthers are appendiculate and the ovules erect, as in the other species of *Stemona*.

3. *Stemona japonica*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 92; *C. H. Wright, in Journ. Linn. Soc., Bot.* xxxii. (1896) p. 493.

*Roxburghia japonica*, *Blume, Enum. Pl. Jap.* i. p. 9, non *Miq. Somoku Zusetsu*, ii. t. 56.

*Roxburghia ruscifolia*, *Zucc. ex Schnitzl. Icon.* t. 56 b. fig. 17.

KIANGSU: Hills near Shanghai (*Carles*, 357!). *Herb. Kew.*  
Japan.

4. *Stemona parviflora*, *C. H. Wright, in Journ. Linn. Soc., Bot.* xxxii. (1896) p. 496.

HAINAN (*A. Henry*, 8698!). *Herb. Kew.*

This somewhat resembles *Roxburghia javanica*, var. ? *australiana*, *Benth. (Fl. Austral.* vii. p. 1), but has much smaller, shortly pedicellate, flowers, with broadly ovate-lanceolate perianth-segments.

5. *Stemona tuberosa*, *Lour. Fl. Cochinch.* p. 404; *Hook. f. Fl. Brit. Ind.* vi. p. 298.

*Roxburghia gloriosoides*, *Roxb. Pl. Corom.* i. p. 29, t. 32, non *Zoll.*; *Schnitzl. Icon.* t. 56 b. fig. 1; *C. H. Wright, in Journ. Linn. Soc., Bot.* xxxii. (1896) p. 494.

*Roxburghia viridiflora*, *Smith, Exot. Bot.* i. p. 111, t. 57; *Griff. Notul.* i. p. 29, *Icon.* tt. 12-16 & 62. fig. 2; *Wall. Pl. As. Rar.* iii. p. 49 t. 282.

*Roxburghia Stemona*, *Steud. Nom.* ed. 2, ii. p. 475.

FOKIEN: Amoy (*Swinhoe*!); FORMOSA: Bankinsing (*A. Henry*, 816!); HUPEH: Ichang (*A. Henry*, 566!). *Mus. Brit.*; *Herb. Kew.*

Northern India.

## 143. LILIACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Heterosmilax Gaudichaudiana*, *Maxim. in Bull. Acad. St. Pétersb.* xvii. (1872) p. 176 in obs.; *Mél. Biol.* viii. p. 415; *A. DC. Monogr. Phan.* i. p. 44, cum. var. *hongkongensis*, *A. DC.*

*Smilax Gaudichaudiana*, *Kunth, Enum. Pl.* v. p. 252; *Benth. Fl. Hongk.* p. 370.

*Smilax hongkongensis*, *Seem. Bot. Voy. 'Herald,'* p. 420.

HONGKONG (*Hance*, 858! *Wright*, 535! *Urquhart*! *Seemann*,

2401 !); SOUTH CHINA (*Gaudichaud, Ivan, Calléry, 104, ex A. DeCandolle*). *Herb. Kew.*

2. *Heterosmilax japonica*, *Kunth, Enum. Pl. v. p. 270; Maxim. in Bull. Acad. St. Péterb. xvii. (1872) p. 175; Mél. Biol. viii. p. 415; Franch. et Savat. Enum. Pl. Jap. ii. p. 50; A. DC. Monogr. Phan. i. p. 43.*

This species, which is cultivated in Japan, is said to have been introduced from China, but no Chinese specimens are known to exist.

1. *Smilax calophylla*, *Wall., var. concolor, C. H. Wright*. Folia oblongo-lanceolata, acuminata, trinervia, 4 poll. longa, 1 poll. lata, concoloria.

KIANGSI: Kiukiang (*Shearer! Bullock, 89!*).

This differs from the type in the leaves not being glaucous beneath and in the less prominent veins.

2. *Smilax China*, *Linn. Sp. Pl. ed. 1, p. 1029; Thunb. Fl. Jap. p. 152; Lour. Fl. Cochinch. p. 622; Hance, in Journ. Linn. Soc., Bot. xiii. (1872) p. 130; Franch. et Savat. Enum. Pl. Jap. ii. p. 49; Franchet, in Mém. Soc. Sc. Nat. Cherbourg, xxiv. (1882) p. 258; Swinhoe, List of Formosa Plants, no. 191; Maxim. Mél. Biol. viii. p. 408; A. DC. Monogr. Phan. i. p. 46.*

*Smilax ferox*, *Wall. Cat. no. 5119; Benth. Fl. Hongk. p. 370; Walp. Ann. vi. p. 145; S. Moore, in Journ. Bot. xvi. (1878) p. 138; Hook. f. Fl. Brit. Ind. vi. p. 307.*

*Smilax japonica*, *A. Gray, in Narrative of Perry's Exped. ii. p. 320, et Mem. Amer. Acad. vi. (1857) p. 412.*

*Coprosmanthus japonicus*, *Kunth, Enum. Pl. v. p. 268.*

SHANTUNG (*Maingay, 130!*), Chefoo (*Faber! Fauvel ex Franchet*); CHEKIANG: Taihoo Lake (*Carles, 209!*), Chusan (*Horne!*), Ningpo (*Horne!*); KANGSI: Kiukiang (*Bullock, 254! Shearer! Maries!*); FOKIEN: Amoy (*Fortune, 71!*), Foochow (*Popow!*); FORMOSA (*Swinhoe*), Tamsui (*Oldham, 571!*); HUPEH: Ichang (*Henry, 1267, 466, 1177, 1197, 1208, 1219, 1222, 2943!*), Patung (*Henry, 1417, 1775!*); SZECHUEN: Mount Omei (*Faber, 125!*); KWANGTUNG (*Krone ex Miquel*), Macao (*Abel!*), Kowlung Hills (*Lamont, 783!*); HONGKONG (*Wilford, 351! Hance, 1025! Wright, 534! Harland, 629!*); KWANGSI (*Playfair!*); HAINAN



(*Henry*, 8520, 8538 !); COREA (*Wilford*, 684! *Wykeham Perry*, 58, 108 !); LUCHU ARCHIPELAGO (*Wright*!). Mus. Brit.; Herb. Kew.

3. *Smilax Davidiana*, *A. DC. Monogr. Phan.* i. p. 104; *Franchet, Pl. David.* p. 300.

KIANGSI (*David ex De Candolle*).

4. *Smilax* (§ *Eusmilax*) *flaccida*, *C. H. Wright, in Kew Bulletin* (1895), p. 118.

HUPEH: Ichang (*A. Henry*, 3630, 3630 A, 3630 B !). Herb. Kew.

The leaves of this plant somewhat resemble those of *S. parvifolia*, Wall., but the stem has not the same zigzag habit.

5. *Smilax glabra*, *Roxb. Fl. Ind.* iii. p. 792; *Seem. Bot. Voy. 'Herald,'* p. 420, t. 100; *Hance, in Journ. Bot.* x. (1872) p. 102; *Benth. Fl. Hongk.* p. 369; *A. DC. Monogr. Phan.* i. p. 60; *Hook. f. Fl. Brit. Ind.* vi. p. 302.

KWANGTUNG: Canton (*Hance*, 10535 ! *Tate* !); HONGKONG (*Wright*, 532 ! *Seemann*, 2401 ! *Hinds* ! *Millett* !). Herb. Kew. Northern India.

This is recorded by Hance as the source of the "China-root" of commerce.

6. *Smilax herbacea*, var. *acuminata*, *C. H. Wright. Folia* membranacea, ovato-lanceolata, acuminata, 4-5 poll. longa, 1½-2 poll. lata, 5-7-nervia, subtus tenuiter puberula; petiolus 1 poll. longus; stipulæ 6 lin. longæ, apice cirrhiferæ. *Pedunculus* 4 poll. longus; pedicelli 8 lin. longi.

HUPEH: Patung (*Henry*, 4002, 5600 B !). Herb. Kew.

Var. *angusta*, *C. H. Wright. Folia* membranacea, lanceolata, acuminata, pellucide lineolata, quinquenervia, 3 poll. longa, 1 poll. lata, glabra; petiolus 9 lin. longus; stipulæ cirrhiferæ. *Pedunculus* 1-1½ poll. longus.

HUPEH: Nanto (*A. Henry*, 5600 F !). Herb. Kew.

Var. *intermedia*, *C. H. Wright. Caulis* erectus, herbaceus. *Folia* ovata, acuta, basi cordata, 7-9-nervia, 4 poll. longa, 2 poll. lata; petiolus 1 poll. longus basi vaginatus; stipulæ cirrhiferæ.

*Pedunculus* 3 poll. longus, flores plures gerens. *Stamina* 6.  
*Ovula* geminata.

HUPEH: Chiensih (*A. Henry*, 5600 C!), Nanto (*Henry*, 4509!); SZECHUEN: South Wushan (*A. Henry*, 5600 A & B!).  
Herb. Kew.

This appears to be intermediate between the American varieties *peduncularis* and *lasioneuron*. Dr. Henry's numbers 939, 2810, and 5600 H from Hupeh are probably mere broad-leaved forms of this variety.

Var. *nipponica*, *Maxim. in Bull. Phys.-math. Acad. Pétersb.* xvii. (1872) p. 174; *A. DC. Monogr. Phan.* i. p. 52.

*Smilax nipponica*, *Miq. in Versl. en Med. K. Akad. Wetensch.* ser. 2, ii. (1868) p. 87, et *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 150.

SHINGKING: Kwandien Mountains (*Ross*, 79!), Moukden (*James*!); KIANGSI: Kiukiang (*Maries! Bullock*, 252!); HUPEH: Changyang (*Henry*, 5600 K!); SZECHUEN: South Wushan (*Henry*, 5600!); KWANGTUNG (*Ford*, 341!).  
Herb. Kew.

Var. *oblonga*, *C. H. Wright. Folia* oblonga, acuta,  $4\frac{1}{2}$  poll. longa,  $1\frac{1}{2}$  poll. lata, septemnervia, glabra; petiolus 6 lin. longus, basi cirrhiferus. *Pedunculus* 3 poll. longus, glaber; pedicelli plures, 6 lin. longi. *Bacca* globosa, 6 lin. diam.; semina geminata.

HUPEH: Ichang (*Henry*, 5216!). Herb. Kew.

Var. *pubescens*, *C. H. Wright. Caulis* suffrutescens, pilis rufescentibus pubescens. *Folia* ovato-lanceolata, 4 poll. longa, 2 poll. lata, quinquenervia; pagina superiora glabra, inferiora et petioli pubescens; petiolus 9 lin. longus, basi bicirrhiferus. *Pedunculus* 2 poll. longus; pedicelli 6 lin. longi, dense pubescentes. *Bacca* globosa, 6 poll. diam., nigra. *Semina* geminata.

HUPEH: Tunghu (*Henry*, 5600 K bis!). Herb. Kew.

7. *Smilax hypoglauca*, *Benth. Fl. Hongk.* p. 369; *A. DC. Monogr. Phan.* i. p. 61.

HONGKONG (*Wright*, 533! *Champion*, 2! *Wilford*, 103! *Urquhart*, 123! *Hinds*!); HAINAN (*A. Henry*, 8443, 8544!).  
Herb. Kew.

8. *Smilax lævis*, var. *ophirensis*, *A. DC. Monogr. Phan.* i. p. 56.  
KIANGSI: Kiukiang (*Shearer!*). Herb. Kew.

Var. *Parkii*, *A. DC. Monogr. Phan.* i. p. 57.

CHINA (*Park ex A. DC.*).

The type occurs in the Malay Peninsula and var. *ophirensis* in Sumatra.

9. *Smilax lanceæfolia*, var. *opaca*, *A. DC. Monogr. Phan.* i. p. 57.

*Smilax lanceæfolia*, *Seem. Bot. Voy. 'Herald,'* p. 420, t. 99: *Benth. Fl. Hongk.* p. 370; *Franchet, in Bull. Soc. Bot. France*, xlv. (1901) p. 214.

FOKIEN: Kuatun (*De la Touche ex Franchet*); FORMOSA: Tamsui (*Oldham, 329!*), Taiwan (*Playfair!*); HONGKONG (all collectors); HAINAN (*Swinhoe*). Mus. Brit.; Herb. Kew.

The type occurs in Northern India.

10. *Smilax* (§ *Eusmilax*) *megalantha*, *C. H. Wright, in Kew Bulletin* (1895), p. 118.

SZECHUEN (*Pratt, 811!*), Mount Omei (*Faber, 241!*). Herb. Kew.

This species somewhat resembles *Smilax stenopetala*, *A. Gr.*, but differs in the inflorescence, which, instead of arising directly from the axil of a mature leaf, consists of a subumbellate raceme, borne in the axil of a very young leaf (rarely an inch long) situated on an axillary branch about four inches long, at the base of which a large bud-scale persists opposite to the leaf-sheath.

11. *Smilax menispermoidea*, *A. DC. Monogr. Phan.* i. p. 108.

HUPEH: Patung (*Henry, 4679!*). Herb. Kew.

Sikkim.

12. *Smilax microphylla*, *C. H. Wright, in Kew Bulletin* (1895), p. 117.

HUPEH: Ichang (*Henry, 1521, 3089, 3089 a, 3980, 3996, 4410!*). Herb. Kew.

This appears to be related to *Smilax obtusa*, *Benth.*, from Mexico. It can be distinguished from the other Chinese species by its small, glaucous, somewhat coriaceous leaves.

13. **Smilax Oldhami**, *Miq. in Versl. en Med. K. Akad. Wetensch.* ser. 2, ii. (1868) p. 86; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 150; *A. DC. Monogr. Phan.* i. p. 53.

*Smilax herbacea*, var. *Oldhami*, *Maxim. in Bull. Acad. St. Pétersb.* xvii. (1872) p. 174; *Mél. Biol.* x. p. 413; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 48.

COREAN ARCHIPELAGO (*Oldham*, 887, 997! *Perry*, 53!).  
Herb. Kew.

Japan.

Var. *ussuriensis*, *A. DC. Monogr. Phan.* i. p. 54; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 257.

SHANTUNG: Chefoo (*Fauvel ex Franchet*).

14. **Smilax ovalifolia**, *Roxb. Fl. Ind.* iii. p. 794; *Seem. Bot. Voy. 'Herald,'* p. 421; *Benth. Fl. Hongk.* p. 370.

KWANGTUNG (*Ford*, 369!); HONGKONG (*Hance ex Seemann*).  
Herb. Kew.

India, Ceylon.

15. **Smilax pekingensis**, *A. DC. Monogr. Phan.* i. p. 108.

CHILI: Peking (*Wawra*, 1059, ex *A. DC.*).

16. **Smilax perfoliata**, *Lour. Fl. Cochinch.* p. 622; *Hance, in Journ. Bot.* xvii. (1879) p. 15; *A. DC. Monogr. Phan.* i. p. 211.

HAINAN (*Swinhoe ex Hance*).

17. **Smilax riparia**, *A. DC. Monogr. Phan.* i. p. 55.

KWANGTUNG: Canton (*Sampson ex Hance*).

18. **Smilax scobinicaulis**, *C. H. Wright, in Kew Bulletin* (1895), p. 117.

HUPEH (*Henry*, 6554!). Herb. Kew.

Easily distinguished from the other Chinese species by the numerous black fragile prickles which clothe the stem, as in some forms of *Smilax californica*, *A. Gray* (*Wats. Bot. Calif.* ii. p. 186).

19. **Smilax Sieboldi**, *Miq. in Versl. en Med. K. Akad. Wetensch.* ser. 2, ii. (1868) p. 87; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 150; *Maxim. in Bull. Acad. St. Pétersb.* xvii. (1872) p. 169; *Mél. Biol.* viii. p. 406; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 49; *A. DC. Monogr. Phan.* i. p. 48.

SHANTUNG: Chefoo (*Wawra* ex *A. DC.*); COREA (*Carles*, 172!). Herb. Kew.

Japan.

20. *Smilax stans*, *Maxim.* in *Bull. Acad. St. Pétersb.* xvii. (1872) p. 170; *Mél. Biol.* viii. p. 407; *A. DC. Monogr. Phan.* i. p. 49.

HUPEH: South Patung (*Henry*, 5436 partly!), Chiensih (*Henry*, 5941!), Changlo (*Henry*, 6276, 6334!); SZECHUEN: Wushan (*Henry*, 7906!). Mus. Brit.; Herb. Kew.

Japan.

21. *Smilax stenopetala*, *A. Gray*, in *Mem. Amer. Acad.* n. s. vi. (1859) p. 412; *Maxim.* in *Bull. Acad. St. Pétersb.* xvii. (1872) p. 169; *Mél. Biol.* viii. p. 405; *A. DC. Monogr. Phan.* i. p. 189.

FORMOSA (*Maries!* *Playfair*, 89!); LUCHU ARCHIPELAGO (*Wright*, 312!). Herb. Kew.

Japan.

22. *Smilax vaginata*, *Decne.* in *Jacquem. Voy.* iv. *Bot.* p. 169, t. 169; *Hook. f. Fl. Brit. Ind.* vi. p. 305.

HUPEH: South Patung (*Henry*, 5436 partly!). Herb. Kew.  
North India.

In addition to the specimens of this genus enumerated above, there are many others, both at the British Museum and Kew, which are not referable to any of the above mentioned species, but are too imperfect for description.

1. *Asparagus brachyphyllus*, *Turcz.* in *Bull. Soc. Nat. Mosc.* (1840) p. 78; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 602; *Bretschn. Hist. Europ. Bot. Disc.* p. 352.

*Asparagus maritimus*, *Pall. Reise*, ii. p. 329, non *Mill.*; *Bunge, Enum. Pl. Chin. Bor.* p. 65, no. 368; *Franch. Pl. David.* i. p. 301.

*Asparagus trichophyllus*, var. *trachyphyllus*, *Kunth, Enum. Pl.* v. p. 63.

CHIHLI: Jehol (*David*, 1800! 1639, ex *Franchet*), Nankau (*Williams*, hb. *Hance*, 14887!). Mus. Brit.; Herb. Kew.

Songaria.

2. *Asparagus davuricus*, *Fisch.* ex *Link*, *Enum. Hort. Berol.* i. p. 340; *Kunth, Enum. Pl.* v. p. 64; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 599, et xvii. (1879) p. 387; *Franch. Pl. David.* i. p. 301; *Bretschn. Hist. Europ. Bot. Disc.* p. 181.

CHIHLI: Jehol (*David*, 1800bis, ex *Franchet*), Peking (*Bushell!*),

Tsien Mountains (*Faber*, 1536!); SHINGKING: Sarhoo (*Ross*, 341!), Hingjing (*Ross*, 263!); SHANTUNG: Chefoo (*Faber*, 76! *Forbes*!); COREA (*Perry*, 44! *Carles*, 41!). Herb. Kew.

Dahuria and Mongolia.

3. *Asparagus filicinus*, *Ham. in D. Don, Prodr. Fl. Nepal.* p. 49; *Kunth, Enum. Pl.* v. p. 73; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 605; *Hook. f. Fl. Brit. Ind.* vi. p. 314; *Baroni, in Bull. Soc. Bot. Ital.* (1898) p. 185.

HUPEH: South Patung (*Henry*, 5712 B!); SZECHUEN: South Wushan (*Henry*, 5712!); SHENSI: Mount Tuikiotsau (*Giraldi*!). Mus. Brit.; Herb. Kew.

Northern India.

4. *Asparagus gibbus*, *Bunge, Enum. Pl. Chin. Bor.* p. 65; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 600, et xvii. (1879) p. 387; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 257.

*Asparagus Sieboldi*, *Debeaux, Fl. Tchef.* p. 134, non *Maxim.*

CHIHLI: Peking (*Williams*, 12635 in herb. *Hance*!); SHINGKING: Foolong (*Ross*, 330!), Jaoling (*Ross*, 385!), Liaoyang (*Ross*, 9!); SHANTUNG: Chefoo (*Perry*! *Fauvel ex Franchet*). Herb. Kew.

5. *Asparagus longiflorus*, *Franch. Pl. David.* i. p. 300.

CHIHLI: Jehol (*David*, 1766, ex *Franchet*).

6. *Asparagus lucidus*, *Lindl. in Bot. Reg.* 1844, *Misc.* p. 29, no. 36; *Benth. Fl. Hongk.* p. 371; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 151; *Kunth, Enum. Pl.* v. p. 72; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 605; *Debeaux, Fl. Shangh.* p. 63; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 58.

*Asparagus falcatus*, *Benth. in Hook. Lond. Journ. Bot.* i. (1842) p. 493; *Seem. Bot. Voy. 'Herald,'* p. 421, non *Linn.*

*Asparagus insularis*, *Hance, in Ann. Sc. Nat.* 5<sup>me</sup> sér. v. (1866) p. 245.

*Melanthium cochinchinense*, *Lour. Fl. Cochinch.* p. 216?

KIANGSU: Shanghai (ex *Debeaux*); FOKIEN: Amoy (*Swinhoe*!); FORMOSA: Tamsui (*Oldham*, 567, 568, 569!); HUPEH: Ichang (*Henry*, 192, 1009, 1124!), Nanto (*Henry*, 2093!); SZECHUEN: Mount Omei (*Faber*, 36!); KWANGTUNG: Macao (*Staunton*! *Nelson*!); HONGKONG (*Hinds*! *Wright*, 538! *Hance*, 274!),

Mount Victoria (*Lamont*, 796!); HAINAN (*Henry*, 8163\*); COREA: Port Hamilton (*Wilford*, 696! 1005!), West Corea (*Perry*, 46!). Mus. Brit.; Herb. Kew.

Japan.

7. *Asparagus officinalis*, *Linn. Sp. Pl.* ed. 1, p. 313; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 598; *Baker & S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 387.

SHINGKING: Fungwangshan (*Ross*, 196!), Jaoling (*Ross*, 375!), between Moukden and Tungcheshien (*James*!). Herb. Kew.

Europe, Caucasus, North Persia, and Siberia.

8. *Asparagus schoberioides*, *Kunth, Enum. Pl.* v. p. 70; *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 151; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 604; *Baker & S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 387; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 58.

*Asparagus Wrightii*, *A. Gr. in Mem. Amer. Acad.* vi. (1858-59) p. 413.

*Asparagus micranthus*, *Sieb. et Zucc. ex Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 151.

*Asparagus Sieboldi*, *Maxim. Prim. Fl. Amur.* p. 287.

*Asparagus parviflorus*, *Turcz. Fl. Baic.-Dahur.* ii. pt. i. p. 226.

SHINGKING: Sarhoo (*Ross*, 342!), Chienshan (*Ross*, 449), Changpeishan (*James*!), between Moukden and Tungcheshien (*James*!), Jaoling (*Webster*, 21!); SHANTUNG: Chefoo (*Faber*, 76 bis!); COREA (*Carles*, 79!). Herb. Kew.

Japan and Northern Asia.

Var. *subsetaceus*, *Franch. Pl. David.* i. p. 302.

CHIHLI: Jehol (*David*, 1779, ex *Franchet*).

9. *Asparagus sinicus*, *C. H. Wright*.

*Asparagopsis sinica*, *Miq. in Journ. Bot. Néerl.* i. (1861) p. 90.

HONGKONG (*Krone ex Miquel*).

10. *Asparagus trichophyllus*, *Bunge, Enum. Pl. Chin. Bor.* p. 65, no. 369; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 600; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 257, et *Bull. Soc. Bot. France*, xxix. (1882) p. 12; *Pl. David*, i. p. 301.

CHIHLI: Jehol (*David* ex *Franchet*), Peking (*Incarville* ex *Franchet*); SHANTUNG: Chefoo (*Fauvel* ex *Franchet*).

Central Siberia.

1. ***Polygonatum agglutinatum***, *Hua*, in *Journ. de Bot.* vi. (1892) p. 448.

YUNNAN: Lo-Kochan (*Delavay*, 1935, ex *Hua*), Santchaho (*Delavay*, 2880 p. p., ex *Hua*), Che-Tong (*Delavay*, 2550, ex *Hua*).

*Delavay's* no. 4332, collected at Kimise, is said by *Hua* to be a form of this with a stipitate perianth, the inner lobes of which have their margins densely ciliate.

2. ***Polygonatum anomalum***, *Hua*, in *Journ. de Bot.* vi. (1892) p. 420.

SZECHUEN (*Pratt*, 41!). Herb. Kew.

This species bears the aspect of *P. officinale*, var. *humile*, Baker, but can be distinguished by its smaller solitary flowers, borne on slender erect pedicels, more than half an inch long. The lower leaves are opposite, the upper alternate, and every one bears a flower in its axil.

3. ***Polygonatum cirrhifolium***, *Royle*, *Ill. Him. Bot.* i. p. 380; *Kunth*, *Enum. Pl.* v. p. 145; *Maxim.* in *Bull. Acad. St. Pétersb.* xxix. (1884) p. 212; *Mél. Biol.* xi. p. 854; *Hook. f.* in *Fl. Brit. Ind.* vi. p. 322; *Hua*, in *Journ. de Bot.* vi. (1892) p. 428.

*Convallaria cirrhifolia*, *Wall.* in *Asiat. Research.* xiii. (1820) p. 382, cum tab.

*Polygonatum chinense*, *Kunth*, *Enum. Pl.* v. p. 146.

*Polygonatum sibericum*, *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 561 p. p.

SHINGKING: Moukden (*Webster*!); HUPEH: South Patung (*Henry*, 5501 A); KANSU (fide *Hua*); SZECHUEN: Tatsienlou (*Bonvalot* ex *Hua*).

Tibet and Temperate Himalayas.

4. ***Polygonatum curvistylum***, *Hua*, in *Journ. de Bot.* vi. (1892) p. 424.

YUNNAN: Yentzehay (*Delavay* ex *Hua*).

5. ***Polygonatum cyrtonema***, *Hua*, in *Journ. de Bot.* vi. (1892) p. 393.

SZECHUEN (*Farges* ex *Hua*).

6. ***Polygonatum Delavayi***, *Hua*, in *Journ. de Bot.* vi. (1892) p. 422.

YUNNAN: Piouse (*Delavay*, 3024, ex *Hua*).



7. *Polygonatum erythrocarpum*, *Hua*, in *Journ. de Bot.* vi. (1892) p. 424.

SZECHUEN (*Farges ex Hua*).

8. *Polygonatum Fargesi*, *Hua*, in *Journ. de Bot.* vi. (1892) p. 446.

SZECHUEN (*Farges, 538, ex Hua*).

Sikkim.

The native name is "Lao-fou-Kiang." It is doubtful whether this is specifically distinct from *P. sibiricum*, Red.

9. *Polygonatum Franchetii*, *Hua*, in *Journ. de Bot.* vi. (1892) p. 392.

SZECHUEN (*Farges, 395, 529, 671, ex Hua*).

10. *Polygonatum fuscum*, *Hua*, in *Journ. de Bot.* vi. (1892) p. 444.

SZECHUEN: Fangyangtchong (*Delavay, 3037, ex Hua*).

11. *Polygonatum Hookeri*, *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 558; *Hook. Ic. Pl.* t. 2218; *Hook. f. Fl. Brit. Ind.* vi. p. 320.

SZECHUEN: Tachienlu (*Pratt, 867! Bonvalot ex Hua*). Mus. Brit.; Herb. Kew.

Sikkim.

12. *Polygonatum kansuense*, *Maxim.* in *Act. Hort. Petrop.* xi. (1891) p. 493; *Hua*, in *Journ. de Bot.* vi. (1892) p. 426.

*Polygonatum roseum*, *Maxim.* in *Bull. Acad. St. Pétersb.* xxix. (1884) p. 211; *Mél. Biol.* xi. p. 853, non *Kunth*; *Hua*, in *Journ. de Bot.* vi. (1892) p. 423 ex parte.

KANSUH: Tetung River (*Przewalski!*); Dshoni (*Potanin!*); Myndansha River (*Grum-Grshimailo ex Maximowicz*). Herb. Kew.

13. *Polygonatum Kingianum*, *Collett et Hemsl.*, in *Journ. Linn. Soc., Bot.* xxviii. (1890) p. 138, t. 21; *Hua*, in *Journ. de Bot.* vi. (1892) p. 449.

YUNNAN: Tapintze (*Delavay, 2044, ex Hua*), Talongtan (*Delavay, 3048, ex Hua*), Mengtze (*Tanant ex Hua*).

Burmah, Tonquin.

14. *Polygonatum latifolium* var. *commutatum*, *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 555.

*Convallaria commutata*, *Schult. f. Syst.* vii. p. 1671.

*Polygonatum giganteum*, *Dietr. in Otto, Gartenz.* 1835, p. 222; *Kunth, Enum. Pl.* v. p. 136; var. *falcatum*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 210; *Mél. Biol.* xi. p. 851.

*Polygonatum commutatum*, *Dietr. in Otto, Gartenz.* 1835, p. 223; *Kunth, Enum. Pl.* v. p. 135.

*Polygonatum Thunbergii*, *Morren et Decne. in Ann. Sc. Nat.* 2<sup>me</sup> sér. ii. (1834) p. 312; *Kunth, Enum. Pl.* v. p. 133.

*Polygonatum canaliculatum*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 148, non *Pursh*.

COREAN ARCHIPELAGO (*Oldham*, 879!), *Tsusima* (*Wilford*!). *Herb. Kew.*

South and Central Europe, North America, and Japan.

15. *Polygonatum macropodum*, *Turcz. in Bull. Soc. Nat. Mosc.* 1832, pt. ii. p. 205; *l. c.* 1837, fasc. vii. p. 158; *l. c.* 1879, p. 61; *Kunth, Enum. Pl.* v. p. 140; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 208; *Mél. Biol.* xi. p. 848; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 557; *Franch. Pl. David.* p. 302; *Hua, in Journ. de Bot.* vi. (1892) p. 420.

*Polygonatum umbellatum*, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 553.

*Convallaria umbellata*, *Bunge, ex Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 553.

CHIHLI: *Peking* (*Bullock*, 230!, *Skatschkow*!, *Bretschneider*, 737!), *Jehol* (*David*, 1842!), *Kalgan* (*Kirilow ex Turczaninow*); SHANTUNG: *Chefoo* (*Hancock ex Maximowicz, Fauvel ex Franchet*). *Herb. Kew.*

16. *Polygonatum multiflorum*, *Allioni, Fl. Ped.* i. p. 131; *Kunth, Enum. Pl.* v. p. 138; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 555; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 55.

*Convallaria multiflora*, *Linn. Sp. Pl.* ed. 1, p. 315.

*Convallaria govaniana*, *Wall. Cat.* n. 5137.

*Polygonatum govanianum*, *Royle, Ill. Him. Bot.* p. 380.

HUPEH: *Ichang* (*Henry*, 2020, 2682, 3849, 4087, 5460 B!); KWANGTUNG (*Ford*, 365!). *Mus. Brit.*; *Herb. Kew.*

Europe, Temperate Asia, Japan.

This is a very variable species, between which and *P. latifolium* many intermediate forms occur. The extreme states of these two species can be distinguished thus:—

*P. multiflorum*.—Leaves sessile, sheathing, acute, margins more or less undulate, nerves prominent beneath. Perianth greenish; tube constricted above the ovary, pilose inside; segments spreading.

*P. latifolium*.—Leaves broader, shortly petiolate, acuminate. Perianth yellowish; tube not constricted above the ovary, glabrous inside; segments more or less incurved.

The differences in the length of the peduncle, number of flowers and number of seeds appear to be of little diagnostic value.

17. **Polygonatum nodosum**, *Hua*, in *Journ. de Bot.* vi. (1892) p. 394.

SZECHUEN (*Farges ex Hua*).

18. **Polygonatum officinale**, *Allioni*, *Fl. Ped.* i. p. 131 (1785); *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 554; *Hance*, in *Journ. Bot.* xiii. (1875) p. 139, et xx. (1882) p. 39; *Maxim.* in *Bull. Acad. St. Pétersb.* xxix. (1884) p. 206; *Mél. Biol.* xi. p. 846; *Debeaux*, *Fl. Tchef.* p. 134; *Hua*, in *Journ. de Bot.* vi. (1892) p. 395.

*Polygonatum vulgare*, *Desf.* in *Ann. Mus. Par.* ix. (1807) p. 49; *Red. Lil.* t. 258; *Kunth*, *Enum. Pl.* v. p. 132; *Baker & S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 387; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 54.

*Convallaria Polygonatum*, *Linn. Sp. Pl.* ed. 1, p. 315; *Bunge*, *Enum. Pl. Chin. Bor.* p. 65.

CHIHLI: Peking (*Bullock*, 231! *Möllendorff ex Maximowicz*), Pohuashan (*Bretschneider ex Hance*); SHINGKING: Moukden (*James! Webster*, 116!); Kwandien Mountains (*Ross*, 78!), Hingjing (*Ross*, 249, 296!); SHANTUNG: Chefoo (*Fauvel ex Franchet*); SHANSI (*Potanin!*); KIANGSU: Chinkiang (*Bullock!*); KANSUH: Tangut (*Przewalski!*); SZECHUEN: South Wushan (*Henry*, 556!); COREA: Chemulpo (*Carles*, 59!). *Mus. Brit.*; *Herb. Kew.*

Europe, Siberia, and Dahuria.

Var. **japonicum**, *Maxim.* in *Bull. Acad. St. Pétersb.* xxix. (1884) p. 207; *Mél. Biol.* xi. p. 847.

*Polygonatum japonicum*, *Morren et Decne.* in *Ann. Sc. Nat.* 2<sup>m</sup>e sér. ii. (1834) p. 311; *Kunth*, *Enum. Pl.* v. p. 133; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 554.

*Polygonatum vulgare*, var. *japonicum*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 54.

CHIHLI: Peking (*Bullock*, 232!); SHANTUNG: Chefoo (*Hancock*, 28!); COREA (*Carles*!). Herb. Kew.

Japan.

Var. *macranthum*, *Hua, in Journ. de Bot.* vi. (1892) p. 395.

*Polygonatum vulgare*, var. *macranthum*, *Hook. f. in Bot. Mag.* t. 6133.

SZETCHUEN: Tchenkeoutin (*Farges*, 513, 586, ex *Hua*).

Japan.

Var. *Maximowiczii*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 207; *Mél. Biol.* xi. p. 847.

*Polygonatum Maximowiczii*, *Fr. Schmidt, Reisen in Amur-Lande*, p. 185.

*Polygonatum* sp., *Maxim. Prim. Fl. Amur.* p. 275.

NORTH CHINA (*Kirilow ex Maximowicz*); CHIHLI: Peking (*Tatarinow ex Maximowicz*).

Var. *papillosum*, *Franch. Pl. David.* i. p. 302.

CHIHLI: Jehol (*David*, 1767, ex *Franchet*).

19. *Polygonatum platyphyllum*, *Franch. in Journ. de Bot.* iv. (1890) p. 318.

CHIHLI: Peking, Pekeou, St. Michel (*Bodinier ex Franchet*).

20. *Polygonatum Prattii*, *Baker, in Hook. Ic. Pl.* t. 2217; *Hua, in Journ. de Bot.* vi. (1892) p. 420.

SZETCHUEN: Tatsienlou (*Pratt*, 28\*!). Herb. Kew.

21. *Polygonatum pumilum*, *Hua, in Journ. de Bot.* vi. (1892) p. 423.

SZETCHUEN (*Farges*, 797, ex *Hua*).

22. *Polygonatum roseum*, *Kunth, Enum. Pl.* v. p. 144, non *Maxim.*; *Hook. in Bot. Mag.* t. 5049; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 560.

*Convallaria rosea*, *Ledeb. Ic.* t. 1.

HUPEH: Fang (*A. Henry*, 6497 B!). Mus. Brit.; Herb. Kew. Southern Siberia.

Great confusion prevails as to this species. In the 'Flora of British India,' vi. p. 321, it is doubtfully united with *P. verticillatum*, All.

23. *Polygonatum sibiricum*, *Red. Lil.* t. 315; *Kunth, Enum. Pl.* v. p. 145; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 561, pro parte; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 212; *Mél. Biol.* xi. p. 854; *Hua, in Journ. de Bot.* vi. (1892) p. 447; *Franch. Pl. David.* i. p. 302; *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1882) p. 257.

*Polygonatum chinense*, *Kunth, Enum. Pl.* v. p. 146.

CHIHLI: Jehol (*David*, 1784!); Pohuashan (*Bretschneider!*), Peking (*Bretschneider*, hb. *Hance*, 17036!); SHANTUNG: Chefoo (*Fauvel ex Franchet*); HUPEH: Nanto (*A. Henry*, 1984!). *Mus. Brit.*; *Herb. Kew.*

Siberia.

This species is very closely allied to *P. cirrhifolium*, from which it differs in having the filaments papillose and the style twice as long as the ovary.

24. *Polygonatum Souliei*, *Hua, in Journ. de Bot.* vi. (1892) p. 427.

SZECHUEN: Tachienlu (*Soulié*, 306, ex *Hua*).

25. *Polygonatum trinerve*, *Hua, in Journ. de Bot.* vi. (1892) p. 445.

YUNNAN: Santchaho (*Delavay*, 2879, ex *Hua*).

26. *Polygonatum verticillatum*, *Allioni, Fl. Pedem.* i. p. 131; *Kunth, Enum. Pl.* v. p. 142; *Red. Lil.* t. 244; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 560, and xvii. (1879) p. 387; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 210; *Mél. Biol.* xi. p. 852; *Hook. f. Fl. Brit. Ind.* vi. p. 321, pro parte.

*Convallaria verticillata*, *Linn. Sp. Pl.* ed. 1, p. 315.

SHINGKING: Moukden (*James! Webster*, 4, 32, 323!), Sarhoo (*Ross*, 346!); SZECHUEN: Mount Omei (*Faber*, 969!). *Herb. Kew.*

Europe, Caucasus, Temperate Himalayas.

1. *Oligobotrya Henryi*, *Baker, in Hook. Ic. Pl.* t. 1537.

HUPEH: Patung (*A. Henry*, 914, 5583 B!), Hsingshan (*A. Henry*, 7535 A!); HUNAN: Shihmen (*A. Henry*, 7535!); SZECHUEN: South Wushan (*A. Henry*, 5583!), Mount Omei (*Faber*, 899, 964!), Tatsienlou (*Pratt*, 54, 826! *Soulié*, 181!). *Herb. Kew.*

1. **Disporopsis fusco-picta**, *Hance*, in *Journ. Bot.* xxi. (1883) p. 278; *Kew Bulletin* (1893), p. 344.

*Aulisconema aspersa*, *Hua*, in *Journ. de Bot.* vi. (1892) p. 471, pl. xiv. fig. 1.

HUPEH: Kuei (*A. Henry*, 6168!), Fang (*A. Henry*, 6705!); KWANGTUNG: Lofaushan at 3100 feet (*B. C. Henry* in hb. *Hance*, 22186!). Mus. Brit.; Herb. Kew.

1. **Streptopus amplexifolius**, *DC. Fl. Franç.* iii. p. 174; *Red. Lil.* t. 259; *Kunth, Enum. Pl.* iv. p. 205; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 212; *Mél. Biol.* xi. p. 855; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 146; *A. Gray, Bot. Jap. in Mem. Amer. Acad.* vi. (1859) p. 415; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 51.

*Streptopus amplexicaulis*, *Poir. Encycl.* vii. p. 467; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 591.

*Uvularia amplexifolia*, *Linn. Sp. Pl.* ed. 1, p. 304.

KANSUH (*Przewalski!*); SZECHUEN: Tachienlu (*Pratt*, 52!). Mus. Brit.; Herb. Kew.

Central Europe, North Asia, North America.

2. **Streptopus paniculatus**, *Baker, in Hook. Ic. Pl.* t. 1932.

HUPEH: Chienshih (*A. Henry*, 5723 A and B!); SZECHUEN: South Wushan (*A. Henry*, 5723!). Mus. Brit.; Herb. Kew.

1. **Tovaria atropurpurea**, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 45.

YUNNAN: Mount Tsangchan (*Delavay*, 1824, ex *Franchet*).

2. **Tovaria Delavayi**, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 47.

YUNNAN: Longki (*Delavay* ex *Franchet*).

3. **Tovaria Fargesii**, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 45.

SZECHUEN: Tchenkeoutin (*Farges*, 422, ex *Franchet*).

4. **Tovaria japonica**, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 570, and xvii. (1879) p. 387; *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 46.

*Smilacina japonica*, *A. Gray, in Rep. Perry's Exped.* p. 321; *Bot.*

Japan, in *Mem. Amer. Acad.* vi. (1859) p. 414; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 53; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 214; *Mél. Biol.* xi. p. 857.

*Smilacina japonica*, var. *mandschurica*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 214; *Mél. Biol.* xi. p. 857.

*Smilacina hirta*, *Maxim. Prim. Fl. Amur.* p. 276.

*Smilacina Rossii*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 214; *Mél. Biol.* xi. p. 857.

*Tovaria Rossii*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 387.

CHIHLI: Peking (*Tatarinow and Bretschneider ex Maximowicz*), Jehol (*David*, 1769, ex *Franchet*), Tatchiaochen (*David*, 2224, ex *Franchet*); SHINGKING: Kwandien Mountains (*Ross*, 86!), Soongwahien (*Ross*, 44!), Moukden (*Webster*, 42!); HUPEH: South Patung (*A. Henry*, 5366!); SZECHUEN: Tchenkeoutin (*Farges*, 777 bis, ex *Franchet*). Mus. Brit.; Herb. Kew.

Japan and E. Siberia.

5. ***Tovaria Prattii***, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 46.

Var. ***quadrifolia***, *Franch. l. c.*

SZECHUEN: Tachienlu (*Pratt*, 847!). Herb. Kew.

Var. ***robusta***, *Franch. l. c.*

SZECHUEN: Tachienlu (*Pratt*, 106!). Herb. Kew.

6. ***Tovaria Souliei***, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 45.

SZECHUEN: Tachienlu (*Soulié*, 806, ex *Franchet*).

7. ***Tovaria stenoloba***, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 47.

SZECHUEN: Tchenkeoutin (*Farges*, 593 bis, ex *Franchet*).

8. ***Tovaria tatsienensis***, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 47.

SZECHUEN: Tachienlu (*Pratt*, 34, ex *Franchet*).

9. ***Tovaria tubifera***, *C. H. Wright*.

*Smilacina tubifera*, *Batalin, in Act. Hort. Petrop.* xiii. i. (1893) p. 104.

HUPEH: Fag (*A. Henry*, 6845, 6858!), Hsingshan (*A. Henry*, 6845 A!), Chienshih (*A. Henry*, 5791!); Changlo (*A. Henry*, 6252!); KANSUH: Dshoni, Mount Idshuschan

(*Potantin*!); SZECHUEN: South Wushan (*Henry*, 7527!), Mount Omei (*Faber*, 961!). Herb. Kew.

10. *Tovaria yunnanensis*, *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 48.

YUNNAN: Koutoui and Lyenyn (*Delavay*, 3556, 4160, ex *Franchet*).

1. *Maianthemum Convallaria*, *Wigg. Prim. Fl. Holsat.* p. 15.

*Maianthemum bifolium*, *DC. Fl. Franç.* iii. p. 177; *Desf. in Ann. Mus. Par.* ix. (1807) p. 54; *Red. Lil.* t. 216. fig. 2; *Kunth, Enum. Pl.* v. p. 147; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 563.

*Smilacina bifolia*, *Schult. f. Syst.* vii. p. 307; *Turcz. in Bull. Soc. Nat. Mosc.* (1837) fasc. vii. p. 158; *A. Gray, in Rep. Perry's Exped.* p. 321; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 53; *Franch. Pl. David.* i. p. 303.

N. CHINA (*Kirilow ex Turczaninow*); CHIHLI: Jehol (*David*, 689! 1814, ex *Franchet*), Puhuashan (*David*, 2257!); SHINGKING: Changpeishan (*James*!); COREA (*Bushnell*!). Mus. Brit.; Herb. Kew.

Widely spread through the north temperate zone.

1. *Convallaria majalis*, *Linn. Sp. Pl.* ed. 1, p. 314; *Red. Lil.* t. 227; *Kunth, Enum. Pl.* v. p. 131; *A. Gray, in Rep. Perry's Exped.* p. 321; *Maxim. Fl. As. Or. Fragm.* p. 61; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 552; *Hance, in Journ. Bot.* xiii. (1875) p. 139; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Maxim. in Bull. Soc. Nat. Mosc.* (1879) p. 61; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 54; *Franch. Pl. David.* i. p. 303.

*Polygonatum maiale*, *Allioni, Fl. Pedem.* i. p. 130.

CHIHLI: Puhuashan (*David*, 1838! *Bretschneider*!), Peking (*David*, 1120!); SHINGKING: Ngaiyang (*Ross*, 245!). Hingjing (*Ross*, 295!), Toonghwahien (*Ross*, 4!), Moukden (*James*!), Tungling (*Webster*, 113, 124!); SHANTUNG: Chefoo (*Hancock*, 6!). Mus. Brit.; Herb. Kew.

Europe, Northern Asia, Japan, Eastern United States.

1. *Speirantha convallarioides*, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 563, t. 17.

Albuca? *Gardeni, Hook. Bot. Mag.* t. 4842.

KIANGSU: Shanghai (*Baines*!). Mus. Brit.; Herb. Kew.



1. **Reineckia carnea**, *Kunth, in Abh. Akad. Wiss. Berl.* (1842) p. 29; *Kunth, Enum. Pl.* v. p. 22; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 57; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 551.

*Sansevieria carnea*, *Andr. Bot. Rep.* t. 361; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478.

*Sansevieria sessiliflora*, *Ker-Gawl. in Bot. Mag.* t. 739.

HUPEH: Patung (*A. Henry*, 1000, 5933 A!), Chienshih (*A. Henry*, 5933!), Hsingshan (*A. Henry*, 6562!), Ichang (*A. Henry*, 3267, 4283!); SZECHUEN: Mount Omei (*Faber*, 889!), Tachienlu (*Pratt*, 163!); KWANGTUNG: Canton (hb. *Hance*, cult.); HONGKONG (hb. *Hance*, cult.). Mus. Brit.; Herb. Kew. Japan.

1. **Aspidistra** \* *elatior*, *Blume, in Hœv. et De Vriese, Tijdschrift voor Nat.* i. (1834) p. 76, tt. 4 & 3 d; *Useful Plants of Japan*, iii. p. 962.

*Aspidistra punctata*, *Carrière, in Rev. Hort.* (1875) pp. 36-37, figs. 4-8.

*Aspidistra punctata*, var. *albo-maculata*, *Hook. Bot. Mag.* t. 5386.

*Aspidistra variegata*, *Hort. Turic. ex Regel, in Flora*, xxxiii. (1850) p. 354.

*Aspidistra* sp., *Baill. Dict. Bot.* i. p. 294; *J. H. Wilson, in Gard. Chron.* ser. III. xxvi. (1899) p. 155.

*Plectogyne variegata*, *Link, in Otto et Dietr. Gartenz.* (1834) p. 265; *Kunth, Enum. Pl.* v. p. 313; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 85; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 580, t. 18; *Yasuda, in Bot. Mag. Tokyo*, viii. (1894) p. 75.

HONGKONG: Mount Parker (*A. B. Westland*!). Herb. Kew. Japan.

2. **Aspidistra lurida**, *Ker-Gawl. in Bot. Reg.* t. 628; *Bot. Mag.* t. 2499; *Morr. & Decne. in Ann. Sc. Nat.* 2<sup>me</sup> sér. ii. (1834) p. 311; *Lodd. Bot. Cab.* t. 1468; *Kunth, Enum. Pl.* v. p. 311; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 579, ex parte; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 85; *Baill. in Bull. Soc. Linn. Paris*, ii. p. 1129, non *Sieb.*

*Macroglyne convallariæfolia*, *Link et Otto, Ic. Select.* p. 69, t. 31.

CHINA (ex *Kunth*); HONGKONG: Wongneichung (*Ford* in hb. *Hance*, 18984!). Mus. Brit.

3. **Aspidistra minutiflora**, *Stapf*, sp. n.

*Perigonium urceolatum*, dentibus parvis triangularibus por-

\* By Dr. O. STAPF.

rectis, circiter 4 lin. longum. Stigma stylo paulo latius, haud pileiforme.

HONGKONG: Wongneichung, *Ford*, 626, 627. Mus. Brit.

4. *Aspidistra punctata*, *Lindl. in Bot. Reg. t. 977, et in Trans. Hort. Soc. Lond. vii. (1826) p. 66; Kunth, Enum. Pl. v. p. 312; Bretschn. Hist. Europ. Bot. Disc. p. 273; non Durand.*

CHINA (*Parks ex Lindley*); HONGKONG (*Lamont!*). Mus. Brit.

1. *Tupistra chinensis*, *Baker, in Hook. Ic. Pl. t. 1867; Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 40.*

*Tupistra Fargesii*, *Baill. in Bull. Soc. Linn. Paris, i. p. 1114.*

HUPEH: Patung (*A. Henry, 5023, 5023 A!*), Changyang (*A. Henry, 5231, 5692 C!*), Chiensih (*A. Henry, 5692 A!*), Ichang (*A. Henry, 2044!*); SZECHUEN: South Wushan (*A. Henry, 5692*), Tchenkeoutin (*Farges, 114, ex Baillon*). Mus. Brit.; Herb. Kew.

2. *Tupistra chlorantha*, *Baill. in Bull. Soc. Linn. Par. ii. (1893) p. 1115; Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 42.*

SZECHUEN (*Farges, 923, ex Franchet*).

3. *Tupistra Delavayi*, *Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 40.*

HUPEH (*A. Henry, 5231 A, ex Franchet*); YUNNAN: Outchay (*Delavay ex Franchet*).

4. *Tupistra lorifolia*, *Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 41.*

SZECHUEN: Tchenkeou (*Farges ex Franchet*).

5. *Tupistra viridiflora*, *Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 41.*

SZECHUEN: Heoupin (*Farges, 933, ex Franchet*).

6. *Tupistra Wattii*, *Hook. f. Fl. Brit. Ind. vi. p. 325; Franch. in Bull. Soc. Bot. France, xliii. (1896) p. 42.*

*Campylandra Wattii*, *C. B. Clarke, in Journ. Linn. Soc., Bot. xxv. (1889) p. 78, t. 32.*

YUNNAN: Longki (*Delavay, 5109, 5139, ex Franchet*).

1. *Rohdea japonica*, *Roth, Nov. Pl. Sp.* p. 197; *Kunth, Enum. Pl.* v. p. 321; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 582; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 86; *Franch. in Bull. Soc. Bot. France*, xliii. (1896) p. 40.

*Orontium japonicum*, *Thunb. Flor. Jap.* p. 144; *Bot. Mag.* t. 898.

KIANGSU: Shanghai (*Faber*, "wild?"); HUPEH: Nanto (*A. Henry*, 4601!), Patung (*A. Henry*, 5023 B!); SZECHUEN: Mount Omei (*Faber*, 34!), Tchenkeoutin (*Farges*, 424, ex *Franchet*); COREA: Tsusima (*Wilford*, 898!). *Herb. Kew.*

Japan.

"Named K'ai-K'ou Chien: root is used as drug in cases of sore mouth."—*Dr. A. Henry.*

1. *Hemerocallis citrina*, *Baroni, in Nuovo Giorn. Bot. Ital.* iv. (1897) p. 305, t. 9; *Bull. Soc. Tosc. Ort.* (1897) p. 160, cum ic. color.

SHENSI: Luntun (*Giraldi, ex Baroni*).

2. *Hemerocallis flava*, *Linn. Sp. Pl.* ed. 2, p. 462; *Jacq. Hort. Vind.* ii. t. 139; *Curt. Bot. Mag.* t. 19; *Red. Lil.* t. 15; *Kunth, Enum. Pl.* iv. p. 588; *Baker, in Journ. Linn. Soc., Bot.* xi. (1870) p. 358; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 79.

*Hemerocallis graminea*, *Kew Bulletin*, 1889, p. 116, non *Andr.*

SHINGKING: Moukden (*James*!); KANGSU: Chinkiang (*Carles*, 448, 449!); HUPEH: Ichang (*A. Henry*, 195!), Patung (*A. Henry*, 1851, 6063!); COREA: Soul (*Veitch*, 68 a pro parte!). *Herb. Kew.*

Southern Europe, Northern Asia, Japan.

*Dr. A. Henry* states that the flowers are eaten with meat.

3. *Hemerocallis fulva*, *Linn. Sp. Pl.* ed. 2, p. 462; *Bot. Mag.* t. 64; *Red. Lil.* t. 16; *Lour. Fl. Cochinch.* p. 205; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Bull. Soc. Nat. Mosc.* (1879) p. 63; *Kunth, Enum. Pl.* iv. p. 588; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 80; *Baker, in Journ. Linn. Soc., Bot.* xi. (1870) p. 359; *Hook. f. Fl. Brit. Ind.* vi. p. 326.

*Hemerocallis disticha*, *Donn, Hort. Cantab.* ed. 6, p. 93; *Ker-Gawl. in Bot. Mag.* sub t. 1433; *Sweet, Brit. Flow. Gard.* t. 28.

*Hemerocallis fulva*, var. *Kwanso*, *Regel, Gartenfl.* (1866) p. 66, t. 500.

CHIHLI: Peking (*Bretschneider*!); FOKIEN: Amoy (*Hance*, 1502!); HUPEH: South Patung (*Henry*, 6142!); KWANGTUNG:

Canton, Pakwan (*Sampson* in hb. *Hance*); COREA: Soul (*Veitch*, 68, 68 a pro parte!). Mus. Brit.; Herb. Kew.

South Europe and Japan to the Himalayas.

Var. *maculata*, *Baroni*, in *Nuovo Giorn. Bot. Ital.* iv. (1897) p. 306.

SHENSI (*Giraldi* ex *Baroni*).

4. *Hemerocallis Middendorffii*, *Trautv. et Meyer*, in *Middend. Reise*, i. pt. 2, *Fl. Ochot.* p. 94; *Maxim. Prim. Fl. Amur.* p. 285; *Baker*, in *Journ. Linn. Soc., Bot.* xi. (1870) p. 359; *Regel*, *Gartenfl.* t. 522.

SHINGKING: Laoyehling (*James*!), Changpeishan (*James*!). Herb. Kew.

Amur.

5. *Hemerocallis minor*, *Mill. Dict.* ed. 8, no. 2; *Baker*, in *Journ. Linn. Soc., Bot.* xi. (1870) p. 358; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 79; *Franch. Pl. David.* i. p. 303.

*Hemerocallis graminea*, *Andr. Bot. Rep.* t. 244; *Kunth, Enum. Pl.* iv. p. 588; *Bot. Mag.* t. 873; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Debeaux, Fl. Tchef.* p. 140.

CHIHLI: Peking (*Bretschneider*, 755! *David* in hb. *Hance*, 13430!), Jehol (*David*, 1863!), Pohuashan (*Möllendorff*!); CHEKIANG: Chusan (*Wilford*!); HUPEH: Ichang (*A. Henry*, 1618!), Kuei (*A. Henry*, 6182!); HUNAN (*Hancock*, 123!); COREA (*Veitch*, 69, 70! *Wykeham Perry*!), Chemulpo (*Carles*, 170, 181, 182!). Mus. Brit.; Herb. Kew.

Eastern Siberia and Japan.

1. *Funkia lancifolia*, *Spreng. Syst.* ii. p. 41; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 83; *Baker*, in *Journ. Linn. Soc., Bot.* xi. (1870) p. 368.

*Funkia ovata*, var.  $\beta$ , *Kunth, Enum. Pl.* iv. p. 592.

*Hemerocallis lancifolia*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 335.

KIANGSI: Kiukiang (*Shearer*!); COREA: Waisan (*Veitch*, 82!). Herb. Kew.

Japan.

2. *Funkia ovata*, *Spreng. Syst.* ii. p. 40; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 81; *Baker*, in *Journ. Linn. Soc., Bot.* xi. (1870) p. 367.

*Funkia ovata*, var. *a*, *Kunth, Enum. Pl.* iv. p. 592.

*Hemerocallis cœrulea*, *Andr. Bot. Rep.* t. 6; *Red. Lil.* t. 106; *Bot. Mag.* t. 894.

*Hemerocallis japonica*, *Thunb. Fl. Jap.* p. 142, non *Trans. Linn. Soc.* ii. (1787) p. 142; *Bot. Mag.* t. 1433.

HUPEH (*A. Henry*, 6209 A !), Ichang (*A. Henry*, 287 !); SZECHUEN: Tatsienlou (*Pratt*, 407 !), Mount Omei (*Faber*, 898 !).

Var. *minor*, *Baker, in Journ. Linn. Soc., Bot.* xi. (1870) p. 368.

SHINGKING: Moukden (*James* !); SZECHUEN: Patung (*A. Henry*, 2577 !); COREAN ARCHIPELAGO (*Oldham*, 865 !). Mus. Brit.; Herb. Kew.

Japan and Eastern Siberia.

3. *Funkia Sieboldiana*, *Hook. in Bot. Mag.* t. 3663; *Miq. in Arch. Néerl.* iv. (1869) p. 226; *Kunth, Enum. Pl.* iv. p. 592; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 81; *Baker, in Journ. Linn. Soc., Bot.* xi. (1870) p. 367.

*Funkia Sieboldi*, *Lindl. in Bot. Reg.* 1839, t. 50.

*Hemerocallis Sieboldtiana*, *Lodd. Bot. Cab.* t. 1869.

KIANGSI: Kiukiang (*Bullock*, 247 !). Herb. Kew.

Japan.

4. *Funkia subcordata*, *Spreng. Syst.* ii. p. 41; *Kunth, Enum. Pl.* iv. p. 591; *Miq. in Arch. Néerl.* iv. (1869) p. 229; *Baker, in Journ. Linn. Soc., Bot.* xi. (1870) p. 367; *Franch. in Bull. Soc. Bot. France*, xxix. (1882) p. 12.

*Hosta japonica*, *Tratt. Thes.* p. 10, t. 52.

*Hemerocallis japonica*, *Red. Lil.* t. 3; *Bot. Mag.* t. 1433.

*Hemerocallis alba*, *Andr. Bot. Rep.* t. 194.

NORTH CHINA (*Staunton ex Baker*); CHIHLI: Peking (*Incarville ex Franchet*); HUPEH: Ichang (*A. Henry*, 526, 2356 !), Nanto (*A. Henry*, 2480 A !); SZECHUEN (*A. Henry*, 8892 !), Tatsienlou (*Pratt*, 414 !); KWANGTUNG: Canton, Fukwing (*Krone in hb. Hance*, 1506 !). Mus. Brit.; Herb. Kew.

Japan.

[1. *Aloe chinensis*, *Baker, in Bot. Mag.* t. 6301. *Aloë barbadensis*, var. *chinensis*, *Haw. Suppl.* p. 45. Originally recorded from China, where, however, it is most unlikely to be indigenous.]

1. *Cordyline terminalis*, *Kunth*, in *Abh. Akad. Berl.* (1820) p. 30, et *Enum. Pl.* v. p. 25; *Baker*, in *Journ. Bot.* xi. (1873) p. 265, et in *Journ. Linn. Soc., Bot.* xiv. (1875) p. 539; *Hook. f. Fl. Brit. Ind.* vi. p. 331.

*Dracæna terminalis*, *Linn. Syst.* ed. 12, p. 246; *Hook. et Arn. Bot. Beech. Voy.* p. 97.

*Aletris chinensis*, *Lam. Encycl.* i. p. 79.

CHINA (ex *Kunth*); FORMOSA: Bankinsing (*A. Henry*, 813!). *Herb. Kew.*

India, Malaya, and Pacific Islands.

Var. *ferrea*, *Baker*, in *Journ. Bot.* xi. (1873) p. 265, et *Journ. Linn. Soc., Bot.* xiv. (1875) p. 540; *Hook. f. Fl. Brit. Ind.* vi. p. 331.

*Cordyline Jacquini*, *Kunth*, in *Abh. Akad. Berl.* (1842) p. 30, et *Enum. Pl.* v. p. 23.

*Dracæna ferrea*, *Linn. Syst.* ed. 12, p. 246; *Bot. Mag.* t. 2053; *Lour. Fl. Cochinch.* p. 196.

*Dracæna terminalis*, *Jacq. Ic.* t. 448; *Red. Lil.* t. 91; *Lodd. Bot. Cab.* t. 1224.

CHINA (ex *Kunth*); KWANGTUNG: Saikeung (*Krone* in hb. *Hance*, 412!). *Mus. Brit.*

India, Malaya.

1. *Anemarrhena asphodeloides*, *Bunge*, *Enum. Pl. Chin. Bor.* no. 373; *Maxim. Ind. Fl. Pekin. in Prim. Fl. Amur.* p. 478; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 83; *Franch. Pl. David.* i. p. 303; *Baker*, in *Journ. Linn. Soc., Bot.* xv. (1876) p. 268; *Kunth, Enum. Pl.* iv. p. 600.

CHIHLI: Peking (*Williams* in hb. *Hance*, 13471! *Bretschneider*, 759!). *Mus. Brit.*; *Herb. Kew.*

1. *Diuranthera major*, *Hemsl. in Hook. Ic. Pl.* t. 2734.

WESTERN CHINA (*Wilson*!). *Herb. Kew.*

2. *Diuranthera minor*, *C. H. Wright*, ex *Hemsl. l. c.*

*Paradisea minor*, *C. H. Wright*, in *Kew Bulletin* (1895), p. 118.

YUNNAN: Mengtze, 6000 ft. (*Hancock*, 94!). *Herb. Kew.*

1. *Chlorophytum chinense*, *Franch. in Journ. de Bot.* (1891) v. p. 154.

SZECHUEN: Tachienlu (*Bonvalot* ex *Franchet*).

2. *Chlorophytum laxum*, *R. Br. Prodr.* p. 277; *Kunth, Enum. Pl.* iv. p. 603; *Hook. f. Fl. Brit. Ind.* vi. p. 336.

*Chlorophytum laxiflorum*, *Baker, in Journ. Linn. Soc., Bot.* xv. (1876) p. 328.

*Chlorophytum falcatum*, *Baker, in Ref. Bot.* t. 333.

*Phalangium parviflorum*, *Wight, Ic. Pl. Ind. Or.* t. 2039.

*Anthericum parviflorum*, *Benth. Fl. Hongk.* p. 373.

KWANGTUNG: Pakhoi (*Playfair*, 66!); HONGKONG (*Wright*, 536! *Hance*, 1264! *Lamont*, 800!); HAINAN (*A. Henry*, 8364!). *Mus. Brit.*; *Herb. Kew.*

Southern India, Ceylon, Borneo, Tropical Africa, and Australia.

1. *Thysanotus chinensis*, *Benth. Fl. Hongk.* p. 372; *Hance, in Ann. Sc. Nat.* 5<sup>me</sup> sér. v. (1866) p. 245; *Journ. Bot.* ix. (1871) p. 202; *Baker, in Journ. Linn. Soc., Bot.* xv. (1876) p. 337.

*Thysanotus chrysantherus*, *F. Muell. Fragm.* v. p. 202.

*Thysanotus chrysantherus*, *Benth. Fl. Austral.* vii. p. 40.

KWANGTUNG: Canton (*Sampson*, 535! *Hance*, 3882!); HONGKONG (*Hance*); SOUTH CHINA (*Krone*!). *Mus. Brit.*; *Herb. Kew.*

Philippine Islands, Tonkin, Northern Australia.

1. *Dianella nemorosa*, *Lam. Encycl.* ii. p. 276 (1786); *Jacq. Hort. Schœnb.* i. t. 94.

*Dianella ensifolia*, *Red. Lil.* t. 1 (1802); *Bot. Mag.* t. 1404; *Hook. f. Fl. Brit. Ind.* vi. p. 337; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 576.

*Dianella odorata*, *Blume, Enum. Pl. Jav.* i. p. 13; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 58; *Engl. Bot. Jahrb.* vi. (1885) p. 53.

*Dianella javanica*, *Kunth, Enum. Pl.* v. p. 52.

*Dracæna ensifolia*, *Linn. Mant.* p. 63; *Ait. Hort. Kew.* i. p. 454.

*Dracæna ensata*, *Thunb. et Dallm. Diss.* p. 3.

FORMOSA: Tamsui (*Oldham*, 566!), Bankingsing (*A. Henry*, 66!), Apes Hill (*A. Henry*, 792! *Swinhoe*!); SZECHUEN: Chungching (*Faber*, 963!); KWANGTUNG: Pakoi (*Playfair*, 240!), Macao (*Vachell*!); HONGKONG (*Wilford*, 87, 287! *Wright*, 537! *Hinds*! *Hance*, 450! *Lamont*, 798!); HAINAN (*A. Henry*, 8663!); LUCHU ARCHIPELAGO (*Wright*, 332!). *Mus. Brit.*; *Herb. Kew.*

Tropical Asia to Tropical Australia, Mascarene Islands.

1. *Nothoscordum nerinifolium*, *Benth. et Hook. f. Gen. Pl.* iii. p. 802.

*Caloscordum nerinifolium*, *Herb. in Bot. Reg.* (1844), *Misc.* p. 66; *Walp. Ann.* i. p. 859; *Lindl. in Bot. Reg.* (1847) t. 5.

*Allium nerinifolium*, *Baker, in Journ. Bot.* xii. (1874) p. 290.

*Allium Thunbergii*, *Franch. Pl. David.* i. p. 306 ex parte, non *G. Don*.

*Allium Thunbergii*, var. *Davidii*, *Franch. in David, Exs.* n. 2124.

CHIHLI: Peking (*Bretschneider*, 157! *Cantor*!), Jehol (*David*, 2124!); SHANSI (*Pierson*!). *Herb. Kew.*

1. *Allium Bakeri*, *Regel, Alliorum Monogr.* p. 141.

*Allium splendens*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 154.

*Caloscordum exsertum*, *Herb. in Bot. Reg.* (1847), sub t. 5.

*Allium exsertum*, *Baker, in Journ. Bot.* xii. (1874), p. 294, non *G. Don*.

SHINGKING: Moukden (*Webster*, 138, 233!); SHANTUNG (*Maingay*, 222!), Chefoo Mountains (*Faber*, 79!); CHEKIANG: Chusan (*Fortune*, 102!); KIANGSI (*Staunton*!), Kiukiang (*Shearer*!); HUPEH: Fang (*A. Henry*, 6739!), Tunghu (*A. Henry*, 7493!), Nanto (*A. Henry*, 3071, 4413!); SZECHUEN: North Wushan (*A. Henry*, 7038!). *Mus. Brit.*; *Herb. Kew.*

Japan.

2. *Allium Bouddhæ*, *Debeaux, Fl. Tchef.* p. 137; *Fl. Tients.* p. 44; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 259.

CHIHLI: Tientsin (ex *Debeaux*); SHANTUNG: Chefoo (ex *Debeaux*).

This is said by Franchet to greatly resemble *Allium ascalonicum*, Linn. *Debeaux* states that this species is extensively cultivated in Northern China and used, in the fresh state, in Chinese cookery.

3. *Allium Cepa*, *Linn. Sp. Pl.* ed. 1, p. 300; *Lour. Fl. Cochinch.* p. 201; *Bretschn. Early Res.* p. 178.

Extensively cultivated in China.

4. *Allium chinense*, *G. Don, Monogr. All.* p. 83; *Kunth, Enum. Pl.* iv. p. 454; *Baker, in Journ. Bot.* xii. (1874) p. 291.

*Allium Thunbergii*, *G. Don, Monogr. All.* p. 84, non *Franch.*; *Kunth, Enum. Pl.* iv. p. 454; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 154;



*A. Gray*, *Rep. Perry's Exp.* p. 322; *Debeaux*, *Fl. Tchef.* p. 139; *Regel*, *Alliorum Monogr.* p. 234, ex parte; *Act. Hort. Petrop.* x. (1887) p. 355; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 388.

*Allium triquetrum*, *Lour. Fl. Cochinch.* p. 202.

*Allium nereidum*, *Hance*, in *Ann. Sc. Nat.* 5<sup>me</sup> sér. v. (1866) p. 244; *Franch. in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1882) p. 259

*Allium Grayi*, *Regel*, *Alliorum Monogr.* p. 125; *Franch. et Savat. Enum. Pl. Jap.* p. 76; *Franch. Pl. David.* i. p. 304; *Regel*, in *Act. Hort. Petrop.* x. (1889) p. 325; *Baroni*, in *Bull. Soc. Bot. Ital.* (1898) p. 185.

*Allium odorum*, *Lour. Fl. Cochinch.* p. 203, non *Linn.*

CHIHLI: Jehol (*David ex Franchet*); SHINGKING: Chienshan (*Ross!*), Moukden (*James!*); SHANTUNG: Chefoo (*Wykeham Perry! F. B. Forbes! Hance, 7487! King!*) KIANGSU: Shanghai (*Maingay, 410!*); CHEKIANG: Ningpo (*Oldham, 82!*); KIANGSI: Kiukiang (*Shearer!*); FOKIEN: Lamyit Island (*Swinhoe!*); FORMOSA: Tamsui (*Oldham, 564!*); HUPEH: Hsingshan (*A. Henry, 6926!*), Ichang (*A. Henry, 563! Watters!*); SHENSI: Mount Sikutzuisan (*Giraldi ex Baroni*); COREA (*Wykeham Perry, 45!*); COREAN ARCHIPELAGO (*Oldham, 875!*); LUCHU ARCHIPELAGO (*Beechey! Wright, 333!*). Mus. Brit.; Herb. Kew.

Japan.

Although *Allium Grayi* is separated from *A. chinense*, G. Don, by *Regel* and placed in another section of the genus, it does not appear to differ even specifically from the latter.

5. *Allium chrysocephalum*, *Regel*, in *Act. Hort. Petrop.* x. (1887) pp. 291 et 335, t. 3. fig. 1.

KANSUH: Tangut (*Przewalski*). Herb. Kew.

6. *Allium condensatum*, *Turcz. Fl. Baic.-Dahur.* n. 1151, absque descr.; *Bull. Soc. Nat. Mosc.* xxvii. (1854), ii. p. 121; *Kunth*, *Enum. Pl.* iv. p. 456; *Maxim. Prim. Fl. Amur.* pp. 281 et 478; *Regel*, in *Act. Hort. Petrop.* iii. (1875) p. 105; *Franch. Pl. David.* i. p. 304.

*Allium Stevenii*, *Willd.*, var.  $\epsilon$ , *Ledeb. Fl. Ross.* iv. p. 177.

CHIHLI: Jehol (*David, 1964, 2817, 2818, ex Franchet*), Peking (ex *Maximowicz*).

Dahuria, Amur, and Mandshuria.

7. *Allium cyaneum*, *Regel, in Act. Hort. Petrop.* iii. (1875) p. 174, et x. p. 345, t. 4. fig. 3 c (excl. var.  $\beta$ . *brachystemon*); *Kanitz, Bot. Res. Széch. Centr. As. Exped.* p. 59.

KANSUH: Tangut (*Przewalski*!); SZECHUEN: Tachienlu (*Lóczy*, 257, ex *Kanitz*). Herb. Kew.

8. *Allium cyathophorum*, *Franch. in Journ. de Bot.* v. (1891) p. 154.

SZECHUEN: Tachienlu (*Bonvalot ex Franchet*).

9. *Allium Fetisowii*, *Regel, in Act. Hort. Petrop.* v. (1877) p. 631; *Gartenfl.* t. 971. figs. a-f; *Kanitz, Bot. Res. Széch. Centr. As. Exped.* p. 59.

KANSUH: Czingeson (*Lóczy*, 177, ex *Kanitz*).

Eastern Turkestan.

10. *Allium fistulosum*, *Linn. Sp. Pl.* ed. 1, p. 301; *Bretschn. Early Res.* p. 178.

Cultivated in Northern China.

11. *Allium flavovirens*, *Regel, in Act. Hort. Petrop.* x. (1887) pp. 294 et 344, t. 8. fig. 1.

KANSUH (*Przewalski ex Regel*).

12. *Allium* (§ *Rhiziridium*) *Henryi*, *C. H. Wright, in Kew Bulletin* (1895), p. 119.

HUPEH: Hsingshan (*A. Henry*, 6924!). Mus. Brit.; Herb. Kew.

This is allied to *A. Przewalskianum*, var. *planifolium*, *Regel*, from which it differs in having longer leaves and pedicels, a shorter broader spathe, and fewer flowers.

13. *Allium Herderianum*, *Regel, in Act. Hort. Petrop.* x. (1887) pp. 289 et 324, t. 8. fig. 2.

KANSUH (ex *Regel*).

14. *Allium japonicum*, *Regel, Alliorum Monogr.* p. 133; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 77; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 259.

SHANTUNG: Chefoo (*Fauvel ex Franchet*).

Japan.

15. *Allium jeholense*, *Franch. Pl. David. i. p. 305.*

CHIHLI: Jehol (*David, 2157, ex Franchet*).

16. *Allium kansuense*, *Regel, in Act. Hort. Petrop. x. (1887) pt. II. p. 690; Gartenfl. t. 1317; Bot. Mag. t. 7290.*

*Allium cyaneum*,  $\beta$ . *brachystemon*, *Regel, in Act. Hort. Petrop. x. (1887) p. 346.*

KANSUH: Tangut (*Przewalski*); SZECHUEN: Tachienlu (*Soulié, 434!*), Tongolo (*Soulié, 256!*). *Herb. Kew.*

17. *Allium lineare*, *Linn. Sp. Pl. ed. 1, p. 295; Baker & S. Moore, in Journ. Linn. Soc., Bot. xvii. (1879) p. 388; Regel, Alliorum Monogr. p. 166; Allii Sp. Asiæ Cent. p. 66.*

SHINGKING: Moukden (*James! Ross, 348, 444!*), Chienshan (*Ross, 477, 558!*). *Herb. Kew.*

Caucasus, Central and Northern Asia, Sachalin Islands.

18. *Allium macrostemon*, *Bunge, in Mém. Sav. Etr. Acad. Pétersb. ii. (1835) p. 139; Enum. Pl. Chin. Bor. p. 65; Franch. et Savat. Enum. Pl. Jap. ii. p. 76; Regel, Alliorum Monogr. p. 105.*

CHIHLI: Peking (*Bunge, Bretschneider, 783!*). *Herb. Kew.*  
Mandshuria and Japan.

19. *Allium monadelphum*, *Turcz., var. tibeticum, Regel, in Act. Hort. Petrop. x. (1887) p. 311.*

KANSUH: Dshachar-Dsargyn (*Przewalski ex Regel*).

20. *Allium odorum*, *Linn. Mant. p. 62; Bot. Mag. t. 1142; Kunth, Enum. Pl. iv. p. 432; Franch. in Mém. Soc. Sc. Nat. Cherbourg, xxiv. (1882) p. 260; Franch. et Savat. Enum. Pl. Jap. ii. p. 78; Baker, in Journ. Bot. xii. (1874) p. 291; Regel, Alliorum Monogr. p. 175; Act. Hort. Petrop. x. (1887) p. 346; Hook. f. Fl. Brit. Ind. vi. p. 343; Franch. Pl. David. i. p. 306.*

*Allium uliginosum*, *G. Don, in Mem. Wern. Soc. vi. (1832) p. 60; Hance, in Ann. Sci. Nat. 5<sup>me</sup> sér. v. (1866) p. 245.*

*Allium tuberosum*, *Roxb. Hort. Beng. p. 24, et Fl. Ind. ii. p. 141; Kunth, Enum. Pl. iv. p. 454; Baker, in Journ. Bot. xii. (1874) p. 291; Hook. f. Fl. Brit. Ind. vi. p. 343.*

*Allium senescens*, *Miq. in Ann. Mus. Bot. Lugd.-Bat. iii. p. 154.*

CHIHLI: Peking (*Bullock! Bushell! Bretschneider! Cantor! Williams! David!*), Mount Conolly (*Bullock!*); SHINGKING:

Moukden (*Webster*, 199!); SHANTUNG: Chefoo (*Swinhoe*, 4! *Fauvel* ex *Franchet*); KIANGSU: Shanghai (*Faber*! *Maingay*!); HUPEH: Patung (*A. Henry*, 2884, 4656, 5109, 6929 A!), Ichang (*A. Henry*, 443!), Hsingshan (*A. Henry*, 6929!); HUNAN (*A. Henry*, 7537, 7940!); KWANGTUNG (*Ford*, 366!); HONG-KONG (*Hance*!); FORMOSA: Takow (*A. Henry*, 1856!). Mus. Brit.; Herb. Kew.

Siberia.

21. *Allium polyrhizum*, *Turcz. ex Regel, Alliorum Monogr.* p. 162, et in *Act. Hort. Petrop.* x. (1887) p. 339.

KANSUH: Tangut (*Przewalski ex Regel*).

22. *Allium Prattii*, *C. H. Wright*, n. sp.

Glabrum. *Bulbus* cylindricus, fibris reticulatis involutus. *Folia* ligulata, basi apiceque attenuata, 9 poll. longa, 6 lin. lata. *Pedunculus* foliis æquilongus, cylindricus. *Spatha* scariosa, 6 lin. longa, acuminata. *Pedicelli* 6–7 lin. longi, apice non incrassati. *Perianthii* segmenta oblonga, acuta. *Filamenta* perianthio paullo longiora, basi breviter connata, alterna lanceolata. *Ovarium* profunde trilobatum; stylus brevis.

SZECHUEN: Tachienlu (*Pratt*, 576!), Tongola (*Soulié*, 669!). Herb. Kew.

Allied to *Allium Stoliczki*, *Regel*.

23. *Allium Przewalskianum*, *Regel, Alliorum Monogr.* p. 164, et in *Act. Hort. Petrop.* x. (1887) p. 343, t. 4. fig. 2; *Kanitz, Bot. Res. Szech. Centr. As. Exped.* p. 59.

KANSUH: River Tetung (*Przewalski ex Regel*), Kunesangfu (*Lóczy*, 132 b, ex *Kanitz*).

24. *Allium sacculiferum*, *Maxim. Prim. Fl. Amur.* p. 281; *Regel, Alliorum Monogr.* p. 171.

*Allium longistylum*, *Baker, in Journ. Bot.* xii. (1874) p. 294.

NORTH CHINA: between Peking and Moukden (*Staunton*!); FOKIEN: Amoy, Namtaiwu (without collector's name, 1431!). Mus. Brit.

Central and Northern Asia.

25. *Allium sativum*, *Linn. Sp. Pl.* ed. 1, p. 296; *Lour. Fl. Cochinch.* p. 201; *Bretschn. Early Res.* p. 178.

Cultivated throughout China.

26. *Allium Semenovi*, Regel, in *Bull. Soc. Nat. Mosc.* xli. (1868) i. p. 449; *Alliorum Monogr.* p. 85; Baker, in *Journ. Bot.* xii. (1874) p. 293.

SZECHUEN: Tachienlu (Pratt, 733!). Herb. Kew.

Mongolia, Temperate and Alpine Himalayas.

27. *Allium senescens*, Linn. *Sp. Pl.* ed. 1, p. 299; Kunth, *Enum. Pl.* iv. p. 421; *Bot. Mag.* t. 1150; Regel, *Alliorum Monogr.* p. 137; *Allii Sp. Asiæ Centr.* p. 58.

*Allium angulosum*, Ker-Gawl. in *Bot. Mag.* t. 1149, non Linn.; *Red. Lil.* t. 281.

*Allium baicalense*, Willd. *Enum. Hort. Berol.* p. 360.

SHINGKING: Changpeishan (James!). Herb. Kew.

Central Europe, Central and Northern Asia, Japan.

28. *Allium subangulatum*, Regel, in *Act. Hort. Petrop.* x. (1887) pp. 292 et 340, t. 5. fig. 1.

KANSUH: Tangut (Przewalski!). Herb. Kew.

29. *Allium tanguticum*, Regel, in *Act. Hort. Petrop.* x. (1887) pp. 287 et 317, t. 2. fig. 1.

KANSUH: Tangut (Przewalski). Herb. Kew.

30. *Allium tchefouense*, Debeaux, *Fl. Tchef.* p. 138; Franch. in *Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 259.

SHANTUNG: Chefoo (Debeaux).

31. *Allium tenuissimum*, Linn. *Sp. Pl.* p. 301; Baker, in *Journ. Bot.* xii. (1874) p. 292; Regel, *Alliorum Monogr.* p. 157; *Allii Sp. Asiæ Cent.* p. 63; Debeaux, *Fl. Tchef.* p. 138; Franch. *Pl. David.* i. p. 305; Franch. in *Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 259.

CHIHLI: Peking (Staunton! Bushell! Cantor! Williams in hb. Hance, 11463! Bretschneider! Bullock!); SHANTUNG: Chefoo (Faber, 80! Forbes!); KIANGSI: Kiukiang (Bullock, 27!); HUPEH: Fang (A. Henry, 6860!). Mus. Brit.; Herb. Kew.

Siberia.

Var. *anisopodium*, Regel, *Alliorum Monogr.* p. 157; *Allii Sp. Asiæ Cent.* p. 63; Baker, in *Journ. Bot.* xii. (1874) p. 292.

*Allium anisopodium*, *Ledeb. Fl. Ross.* iv. p. 183.

CHIHLI: Peking (*Bretschneider*!); KANSU: Tangut (*Przewalski*!); FORMOSA (ex *Regel*). Mus. Brit.; Herb. Kew.  
Siberia and Mongolia.

32. *Allium uratense*, *Franch. Pl. David.* i. p. 304.

CHIHLI: Jehol (*David ex Franchet*).  
Mongolia.

33. *Allium venustum*, *C. H. Wright*, n. sp.

Gracilis. *Bulbus* ignotus. *Folia* linearia, plana, glabra, obtusa, 10 poll. longa, 1 lin. lata. *Scapus* erectus, foliis æquilongus, 8-12-florus; spatha membranacea, suborbicularis. *Perianthii* segmenta ovata, subacuta, cæruleo-purpurea. *Stamina* inclusa; filamenta in tubo alte connata, perianthio breviter adnata; antheræ oblongæ. *Ovarium* oblongum; stylus brevis.

SZECHUEN: Tachienlu (*Soulié*, 609!). Herb. Kew.

This species much resembles *Allium Henryi*, C. H. Wright, which differs in having broader leaves and free stamens, with the alternate filaments toothed. It is also allied to *Allium Griffithianum*, Boiss.

34. *Allium Victorialis*, *Linn. Sp. Pl.* ed. 1, p. 295; *Kunth, Enum. Pl.* iv. p. 432; *Red. Lil.* t. 265; *Bot. Mag.* t. 1222; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 154; *Maxim. Prim. Fl. Amur.* p. 283; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 78; *Franch. Pl. David.* p. 306; *Regel, Alliorum Monogr.* p. 170; *Allii Sp. Asiæ Cent.* p. 68; *Hook. f. Fl. Brit. Ind.* vi. p. 342.

*Cepa victorialis*, *Mænoch, Meth.* p. 243.

*Allium ellipticum*, *Wall. Cat.* n. 5069; *Kunth, Enum. Pl.* iv. p. 456.

*Allium plantagineum*, *Lam. Fl. Franç.* iii. p. 262.

CHIHLI: Jehol (*David ex Franchet*), Puhuashan (*David*, 1843, ex *Franchet*); HUPEH: Hsingshan (*A. Henry*, 5590 F!), Tunghu (*A. Henry*, 5590 E!), South Patung (*A. Henry*, 5590 C!); KANSUH: Tangut (*Przewalski*!); SZECHUEN: South Wushan (*A. Henry*, 5590!). Mus. Brit.; Herb. Kew.

Central Europe, Central and Northern Asia, Temperate Himalayas, Japan, North-west America.

Var. *angustifolia*, *Hook. f. Fl. Brit. Ind.* vi. p. 342.

SZECHUEN: Tachienlu (*Soulié*, 623!). Herb. Kew.

Sikkim, Western Nepal.

35. *Allium weichanicum*, Palibin, in *Act. Hort. Petrop.* xiv. (1895) p. 142.

CHIHLI: Weichang (ex *Palibin*).

36. *Allium* sp. *A. tangutico*, Regel, affine.

SZECHUEN: Tachienlu (*Soulié*, 129!). *Herb. Kew.*

1. *Scilla chinensis*, Benth. *Fl. Hongk.* p. 373; Baker, in *Journ. Linn. Soc., Bot.* xiii. (1872) p. 233; Franch. in *Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 259; Franch. *Pl. David.* i. p. 306; Debeaux, *Fl. Tchef.* p. 136.

*Barnardia scilloides*, Lindl. in *Bot. Reg.* t. 1029; Hook. *Bot. Mag.* t. 3788; Maxim. *Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; Kunth, *Enum. Pl.* iv. p. 337.

CHIHLI: Peking (*Bretschneider*, 754! *Bushell*! *Williams*, 291!), Chihli Plains (*David*, 398, ex *Franchet*); SHINGKING: Chienshan (*Ross*, 317!); SHANTUNG: Chefoo (*F. B. Forbes*! *Maingay*, 110! *Fauvel* ex *Franchet*); KIANGSU: near Shanghai (*Carles*, 211!); KIANGSI: Kiukiang (*F. B. Forbes*! *Shearer*! *David*, 918, ex *Franchet*); HUPEH: Ichang (*A. Henry*, 126, 2927, 3264!), Nanto (*A. Henry*, 4411, 2927, 3264!); KWANGTUNG: Lappa Island (*Vachell*!); HONGKONG: Chuckchew (*Champion*!); COREA: Chemaulpo (*Carles*, 186!). *Mus. Brit.*; *Herb. Kew.*

2. *Scilla japonica*, Baker, in *Journ. Linn. Soc., Bot.* xiii. (1872) p. 233, non *Thunb.*

*Ornithogalum japonicum*, *Thunb. Fl. Jap.* p. 137.

*Barnardia japonica*, *Rœm. et Schult. Syst.* vii. p. 555; *Kunth, Enum. Pl.* iv. p. 337.

COREA (*Wykeham Perry*, 68!); LUCHU ARCHIPELAGO (*Wright*, 331!). *Herb. Kew.*

Japan.

1. *Lilium amabile*, Palibin, in *Act. Hort. Petrop.* xix. (1901) p. 113.

COREA: Kyongkwi, Soul (*Kalinowski* ex *Palibin*).

2. *Lilium apertum*, Franch. in *Journ. de Bot.* xii. (1898) p. 220.

*Lilium oxypetalum*, Franch. in *Journ. de Bot.* vi. (1892) p. 320, non *Baker*.

YUNNAN: Tchangephan (*Delavay*, 4178, ex *Franchet*), Maeulchan (*Delavay* ex *Franchet*).

Var. *thibetica*, *Franch. in Journ. de Bot.* xii. (1898) p. 221.

SZECHUEN: Séla near the Mékong (*Soulié*, 1031, ex *Franchet*).

3. *Lilium Bakerianum*, *Collett et Hemsl. in Journ. Linn. Soc., Bot.* xxviii. (1890) p. 138, t. 22.

— YUNNAN: Szemao, 6000 ft. (*A. Henry*, 13026!). Herb. Kew.

This species, previously known only from the Shan States, supports the theory advanced by Dr. A. Henry in the Kew Bulletin (1898), p. 294, that the flora of the Szemao district is closely allied to that of Upper Burma.

4. *Lilium Brownii*, *F. E. Brown*, ex *Spae* in *Ann. Soc. Agric. de Gand*, i. (1845) p. 437, t. 41, et *Mém. Couronn. Bruxelles*, xix. (1845-46) p. 11; non *Rev. Hort.* 2<sup>me</sup> sér. ii. p. 496; *Fl. des Serres*, t. 46; *Elwes, Monogr. Lilium*, t. 8; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 230, sub *L. japonico*, *Thunb.*; *Baker, in Gard. Chron.* (1885) ii. p. 134 & (1891) ii. p. 225; *Franch. in Journ. de Bot.* vi. (1892) p. 312; *Baroni, in Nuovo Giorn. Bot. Ital.* iv. (1897) p. 303.

*Lilium longiflorum*, *Franch. Pl. David.* i. p. 307, non *Thunb.*

*Lilium japonicum*, *Sims, Bot. Mag.* t. 1591, non *Thunb.*; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 230, ex parte.

*Lilium odorum*, *Planch. in Fl. des Serres*, tt. 876-7.

CHUSAN ARCHIPELAGO (*Fortune*, 66!); KIANGSI: Kiukiang (*Bullock*, 87! *Shearer!* *David*, 855, ex *Franchet*); FOKIEN: Amoy (*Maclagan*, 111!); HUPEH: Ichang (*A. Henry*, 514, 4160!), Nanto (*A. Henry*, 2047, 2047 A!); SHENSI (*Giraldi* ex *Baroni*); SZECHUEN: Tchenkeoutin (*Farges*, 174, ex *Franchet*); YUNNAN: Mengtze, 4600-7000 ft. (*A. Henry*, 9453, 9453 A, 9453 B, 9453 C!); HONGKONG (*Champion*, 23!); COREAN ARCHIPELAGO (*Oldham*, 869!), Herschel Island (*Oldham*, 1025!); KWANGSI: Lungchow Hills (*Morse*, 602!). Mus. Brit.; Herb. Kew.

Var. *chloraster*, *Baker, in Gard. Chron.* (1891) ii. p. 225.

*Lilium longiflorum*, var. *chloraster*, *Baker, in Gard. Chron.* (1891) ii. p. 66.

HUPEH: Ichang (*A. Henry*, 4162!). Herb. Kew.

Var. *leucanthum*, *Baker, in Gard. Chron.* (1894) ii. p. 180; *Bot. Mag.* t. 7722.

HUPEH: Ichang (*A. Henry!*).



5. **Lilium callosum**, *Sieb. et Zucc. Fl. Jap.* p. 86, t. 41; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 156; *Kunth, Enum. Pl.* iv. pp. 262, 674; *Baker, in Journ. Hort. Soc. n. s. iv.* (1877) p. 46; *in Journ. Linn. Soc., Bot.* xiv. (1874) p. 250; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 65; *Elwes, Monogr. Lilium*, t. 41.

*Lilium pomponicum*, *Thunb. Fl. Jap.* p. 134, excl. syn.

HUPEH: Ichang (*A. Henry*, 2327!); COREA: Soul (*J. H. Veitch*, 87!); LUCHU ARCHIPELAGO (ex *Kunth*). *Herb. Kew.*  
Japan, Mandshuria, Dahuria.

6. **Lilium concolor**, *Salisb. Parad.* t. 47; *Lindl. & Paxt. Flow. Gard.* ii. p. 115, fig. 193; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 236; *Baker, in Journ. Hort. Soc. n. s. iv.* (1877) p. 44; *Kunth, Enum. Pl.* iv. p. 259; *Baroni, in Nuovo Giorn. Bot. Ital.* iv. (1897) p. 304. Typus bulbis solitariis, foliis latis, floribus immaculatis.

*Lilium sinicum*, *Fl. des Serres*, t. 1206; *Ill. Hort.* 1856, t. 100.

Var. **sinicum**, *Hook. f. in Bot. Mag.* t. 6005; bulbis solitariis, foliis latis, floribus maculatis.

*Lilium concolor*, *Bot. Mag.* t. 1165.

Var. **Buschianum**, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 236; bulbis solitariis, foliis angustis, floribus maculatis.

*Lilium Buschianum*, *Lodd. Bot. Cab.* t. 1628.

*Lilium pulchellum*, *Fisch. in Fisch., Meyer et Avé-Lall., Ind. Sem. Hort. Petrop.* vi. p. 56; *Gartenfl.* t. 284. fig. 2; *Rev. Hort.* (1862) p. 131.

*Lilium concolor*, var. *pulchellum*, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 237.

*Lilium concolor*, *Elwes, Monogr. Lilium*, t. 18. fig. 1.

*Lilium partheneion*, *Sieb. et De Vriese, in Tuinb. Flora*, ii. p. 341.

CHIHLI: Peking (*Bushell!*), Jehol (*David*, 1864!), Sylinchan (*Provost*, 139, ex *Franchet*); SHINGKING: Moukden (*James!*), Changpeishan (*James!*), Peilung (*Webster!*), Chienshan (*Ross*, 370!), Toonghuahien (*Ross!*); SHANTUNG: Chefoo (*Wykeham Perry! Hancock, 7! Fauvel ex Franchet*); SHANSI? between Taiming and Liuchiachung (*Pierson in hb. Hance*, 13273!); CHEKIANG: Ningpo (*Cooper!*), Tinghae (*Sandilands!*); SHENSI: Mount Tunusse (*Giraldi ex Baroni*), Peishan (*Maximowicz!*). *Mus. Brit.*; *Herb. Kew.*

Eastern Siberia, Japan.

This species has long been in cultivation in China and Japan, whence it was introduced into Europe. It presents a series of forms, which so gradually merge into each other, that any attempt at grouping into varieties is a matter of great difficulty. The segments of the perianth vary much in direction and also in spotting, the type being quite destitute of spots.

7. *Lilium cordifolium*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 332, non *D. Don*; *Kunth, Enum. Pl.* iv. p. 268; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 227; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 72; *Franch. in Journ. de Bot.* vi. (1892) p. 309; *Gard. Chron.* (1877) ii. p. 304, fig. 61; *Baker, in Bot. Mag.* t. 6337; *Kew Bulletin*, 1889, p. 118.

*Hemerocallis cordata*, *Thunb. Fl. Jap.* p. 143.

KIANGSI: Kiukiang (*David*, 846, ex *Franchet*).

Japan.

The bulb of this species is eaten by the Japanese.

8. *Lilium Delavayi*, *Franch. in Journ. de Bot.* vi. (1892) p. 314.

YUNNAN: Lankong (*Delavay*, 4310, ex *Franchet*), Koutoui (*Delavay* ex *Franchet*), Yanginchan (*Delavay*, 4309, ex *Franchet*), Houanglipin (*Delavay*, 3576, ex *Franchet*).

9. *Lilium Duchartrei*, *Franch. Pl. David.* ii. p. 128, et in *Journ. de Bot.* vi. (1892) p. 316.

KANSUH: Tshagon (*Potanin*!); SZECHUEN: Tachienlu (*Prince Henri d'Orléans* ex *Franchet*); YUNNAN: Koutoui (*Delavay*, 3983, ex *Franchet*). *Herb. Kew.*

10. *Lilium Fargesii*, *Franch. in Journ. de Bot.* vi. (1892) p. 317.

SZECHUEN: Tchenkeoutin (*Farges* ex *Franchet*).

11. *Lilium formosum*, *Franch. in Journ. de Bot.* vi. (1892) p. 313.

SZECHUEN: Tchenkeoutin (*Farges*, 231, ex *Franchet*).

12. *Lilium giganteum*, *Wall. Tent. Fl. Nep.* 21, tt. 12-13; *Bot. Mag.* t. 4673; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874)

p. 227; *Kunth, Enum. Pl.* iv. p. 268; *Hook. f. Fl. Brit. Ind.* vi. p. 349; *Franch. in Journ. de Bot.* vi. (1892) p. 310.

*Lilium cordifolium*, *D. Don, Prodr. Fl. Nepal.* p. 52, non *Thunb.*

HUPEH: Patung (*A. Henry*, 3141, 4085!), Nanto (*A. Henry*, 2681, 4545, 6197 B!), Changyang (*A. Henry*, 6197 A!), Chien-shih (*A. Henry*, 5376!); YUNNAN: Mengtze, 6000-9000 ft. (*Hancock*, 300! *A. Henry*, 10805!). Mus. Brit.; Herb. Kew.

Temperate Himalayas.

Dr. A. Henry says of the Patung plant that his collector "calls this *Peh Wa*, and says that the root is used in surgery. This name I have not seen mentioned in books. Large lilies with edible bulbs are named *Peh Ho* at Patung and Ichang."

13. *Lilium Henrici*, *Franch. in Journ. de Bot.* xii. (1898) p. 220.

YUNNAN: on the banks of the Mékong, near Loukou (*Prince Henri d'Orléans ex Franchet*).

14. *Lilium Henryi*, *Baker, in Gard. Chron.* (1888) ii. p. 660, et (1890) ii. p. 380, fig. 75; *Bot. Mag.* t. 7177.

HUPEH: Ichang (*A. Henry*, 4205, 4245, 4378!). Mus. Brit.; Herb. Kew.

15. *Lilium lankongense*, *Franch. in Journ. de Bot.* vi. (1892) p. 317.

YUNNAN: Lankong (*Delavay*, 4437, ex *Franchet*), Heechanmen (*Delavay*, 2559, ex *Franchet*), Mosoyn (*Delavay*, 3795, ex *Franchet*).

16. *Lilium longiflorum*, *Thunb. in Trans. Linn. Soc.* ii. (1797) p. 333; *Kunth, Enum. Pl.* iv. p. 266; *Walp. Ann.* iii. p. 628; *Lodd. Bot. Cab.* t. 985; *Fl. des Serres*, t. 270; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 157; *Benth. Fl. Hongk.* p. 374; *Swinhoe, List of Formosa Plants*, no. 189; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 229; *Hance, in Journ. Bot.* xii. (1874) p. 262; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 71; *Franch. in Journ. de Bot.* vi. (1892) p. 311 (non *Pl. David.* i. p. 307); *Baker, in Gard. Chron.* (1891) ii. p. 225.

*Lilium candidum*, *Thunb. Fl. Jap.* p. 133.

*Lilium longiflorum*, var. *suaveolens*, *Bot. Reg.* t. 560.

*Lilium Jama Juri*, *Sieb. et De Vriese, in Tuinbouw Fl.* i. p. 319, t. 11.

FOKIEN: Amoy (*Fortune*, 57!); FORMOSA: Tamsui (*Oldham*,

565! *Swinhoe!* *Hancock*, 24! *Maries!*); KWANGTUNG: Lofaushan (*Ford!*); HONGKONG (*Urquhart!* *Hance*, 626! *Lamont*, 802!); LUCHU ARCHIPELAGO (*Beechey!* *Wright*, 334!), Oosima (*Carpenter!*). *Herb. Kew.*

Japan.

17. *Lilium lophophorum*, *Franch. in Journ. de Bot.* xii. (1898) p. 221.

*Fritillaria lophophora*, *Franch. in Journ. de Bot.* v. (1891) p. 153; *Hook. Ic. Pl.* t. 2219.

SZECHUEN: between Batang and Litang (*Bonvalot ex Franchet*), Tachienlu (*Pratt*, 261, 568!), Tongola (*Soulié*, 296, 326, 725!) between Litang and Tatsien (*Prince Henri d'Orléans ex Franchet*); YUNNAN: near Likiang (*Delavay*, 28, 237 ex *Franchet*), Fangyangtchang (*Delavay ex Franchet*). *Herb. Kew.*

18. *Lilium Lowii*, *Baker, in Bot. Mag.* t. 7232; *Hook. f. Fl. Brit. Ind.* vi. p. 350; *Gard. Chron.* (1893) i. p. 121, fig. 29; *Journ. Hort.* ser. 3, xxvii. (1893) p. 79, fig. 13.

*Lilium nepalense*, *Collett et Hemsl. in Journ. Linn. Soc., Bot.* xxviii. (1890) p. 138, non *D. Don*.

↑  
1 — YUNNAN: Mengtze, 5000-7000 ft. (*Hancock*, 214, 309, 568! (*A. Henry*, 10774, 10774 A!)). *Herb. Kew.*

Burma.

19. *Lilium medeoloides*, *A. Gray, in Mem. Amer. Acad.* vi. (1859) p. 415; *Oliver, in Journ. Linn. Soc., Bot.* ix. (1865) p. 169; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 63, excl. syn.; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 156; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 236; *Elwes, Monogr. Lilium*, t. 35, fig. sinist.

CHEKIANG: Tsien Mountains (*Faber*, 1543!); COREAN ARCHIPELAGO (*Oldham*, 873, 1024!). *Herb. Kew.*

Japan.

20. *Lilium mirabile*, *Franch. in Journ. de Bot.* vi. (1892) p. 310.

SZECHUEN: Tchenkeoutin (*Farges ex Franchet*).

21. *Lilium myriophyllum*, *Franch. in Journ. de Bot.* vi. (1892) p. 313.

YUNNAN: Mosoyn (*Delavay*, 3273, 3817, 4784, ex *Franchet*).

22. *Lilium nepalense*, *D. Don*, in *Mem. Wern. Soc.* iii. (1820) p. 412, et *Prodr. Fl. Nepal.* p. 52; *Wall. Pl. As. Rar.* iii. p. 67, t. 291; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 231; *Elwes*, *Monogr. Lilium*, t. 5; *Bot. Mag.* t. 7043; *Hook. f. Fl. Brit. Ind.* vi. p. 350.

YUNNAN: Mengtze, 5000–7000 ft. (*A. Henry*, 9320! *Hancock*, 392!). *Herb. Kew.*

Himalayas.

A species with numerous forms, several of which are in cultivation.

23. *Lilium ochraceum*, *Franch.* in *Journ. de Bot.* vi. (1892) p. 319.

YUNNAN: Tchangchan (*Delavay*), Tapintze (*Delavay*, 369), Kanhaydze (*Delavay*, 3274, ex *Franchet*).

24. *Lilium papilliferum*, *Franch.* in *Journ. de Bot.* vi. (1892) p. 316.

*Lilium Biondii*, *Baroni*, in *Nuovo Giorn. Bot. Ital.* n. s. ii. (1895) p. 337, tt. 8–9, et iv. (1897) p. 304.

YUNNAN: Chouimaouan (*Delavay*, 4592, ex *Franchet*); SHENSI: Mount Thaepechan (*Giraldi* ex *Baroni*).

25. *Lilium pseudotigrinum*, *Carrière*, in *Rev. Hort.* (1867) p. 411; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 248; *Franch.* in *Journ. de Bot.* vi. (1892) p. 319.

SZECHUEN: Tchenkeoutin (*Farges* ex *Franchet*); YUNNAN: Mengtze, 6000 ft. (*A. Henry*, 10774 B!). *Herb. Kew.*

The leaves of the Mengtze plant are much shorter than those of the cultivated plant, upon which the species was founded, and resemble those of *L. papilliferum*, *Franch.*

26. *Lilium roseum*, *Wall. Cat.* n. 5077; *Bot. Mag.* t. 4725; *Belg. Hort.* (1854) p. 130, pl. 21. figs. 5–9; *Hook. f. Fl. Brit. Ind.* vi. p. 352.

*Lilium Thomsonianum*, *Lindl.* in *Bot. Reg.* (1845) t. 1; *Hook. f. Fl. Brit. Ind.* vi. p. 352; *Franch.* in *Journ. de Bot.* vi. (1892) p. 320.

*Fritillaria macrophylla*, *D. Don*, *Prodr. Fl. Nepal.* p. 51; *Kunth, Enum. Pl.* iv. p. 253; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 268.

*Fritillaria Thomsoniana*, *D. Don*, in *Royle, Ill. Him. Bot.* p. 388, t. 92. fig. 1; *Kunth, Enum. Pl.* iv. p. 672.

SZECHUEN: Tachienlu (*Pratt*, 480, 670! *Soulié*, 336! *Prince Henri d'Orléans* ex *Franchet*); YUNNAN: Yentzehay (*Delavay*, 3061, ex *Franchet*), Tchangchan (*Delavay*, 1098, ex *Franchet*).  
Herb. Kew.

Northern India, Afghanistan.

27. *Lilium speciosum*, *Thunb.* in *Trans. Linn. Soc.* ii. (1797) p. 332, non *Andr.*; *Kunth*, *Enum. Pl.* iv. p. 259; *Sieb. et Zucc. Fl. Jap.* p. 31, tt. 12-13. fig. 1; *Bot. Reg.* t. 2000; *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 234; *Hance*, in *Journ. Bot.* xii. (1874) p. 262; *Franch. Pl. David.* p. 307; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 67; *Franch. in Journ. de Bot.* vi. (1892) p. 315.

*Lilium superbum*, *Thunb. Fl. Jap.* p. 134, non *Linn.*

KIANGSI: Kiukiang (*Shearer*! *Möllendorff* in hb. *Hance*, 18422! *David*, 813, ex *Franchet*). Mus. Brit.; Herb. Kew.

Japan.

Var. *gloriosoides*, *Baker*, in *Gard. Chron.* (1880) ii. p. 198.

*Lilium lancifolium*, var. *gloriosoides* (*sic*), *Bretsch. Hist. Europ. Bot. Disc.* p. 742.

KIANGSI: Lushan, Dragon's Pool (*Maries* ex *Bretschneider*).

28. *Lilium sutchuenense*, *Franch. in Journ. de Bot.* vi. (1892) p. 318; *Baroni*, in *Nuovo Giorn. Bot. Ital.* n. s. iv. (1897) p. 304; *Bot. Mag.* t. 7715.

*Lilium tenuifolium*, var. *punctatum*, *Bur. et Franch. ex Franch. l. c.*

*Lilium chinense*, *Baroni*, in *Nuovo Giorn. Bot. Ital.* n. s. ii. (1895) p. 333, tt. 6-7.

SHENSI: Inkiapo (*Giraldi* ex *Baroni*); SZECHUEN: Tachienlu (*Prince Henri d'Orléans* ex *Franchet*), Tchenkeoutin (*Farges* ex *Franchet*).

Var. *atropurpureum*, *Baroni*, in *Nuovo Giorn. Bot. Ital.* n. s. iv. (1897) p. 304.

*Lilium chinense*, var. *atropurpureum*, *Baroni*, *l. c.* ii. (1895) p. 335.

SHENSI (*Giraldi* ex *Baroni*).

29. *Lilium taliense*, *Franch. in Journ. de Bot.* vi. (1892) p. 319.

YUNNAN: Koualapo (*Delavay* ex *Franchet*), Mosoyu (*Delavay* ex *Franchet*).

30. *Lilium tenuifolium*, *Fisch. Hort. Gorenk.* ed. 2 (1812) p. 8; *Schrank, Pl. Rar. Hort. Monac.* t. 91; *Schult. fil., Syst.* vii. p. 409; *Kunth, Enum. Pl.* iv. p. 263; *Bot. Mag.* t. 3140; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 478; *Bunge, Enum. Pl. Chin. Bor.* p. 65; *Ledeb. Fl. Ross.* iv. p. 151; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 250; *Debeaux, Fl. Tchef.* p. 136; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 258; *in Bull. Soc. Bot. France*, xxix. (1882) p. 12; *et in Pl. David.* i. p. 307; *Kanitz, Bot. Res. Széch. Centr. As. Exped.* p. 59; *Franch. in Journ. de Bot.* vi. (1892) p. 318; *Baroni, in Nuovo Giorn. Bot. Ital.* iv. (1897) p. 305.

CHIHLI: Peking (*Bretschneider!* *Bushell!* *Provost*, 11, ex *Franchet, Williams* in hb. *Hance*, 6504! *David*, 401, ex *Franchet*), Jehol (*David*, 1886, ex *Franchet*), Yaloo River (*James!*); NORTHERN CHINA: Guangou (*Bunge*, 371); SHENSI: Mounts Thaepeisan, Gniuju, and Lunsanhuo (*Giraldi* ex *Baroni*); HUPEH: Ichang (*A. Henry*, 511!), Chienshih (*A. Henry*, 5917!), Fang (*A. Henry*, 6786!); KANSUH: Tangut (*Przewalski!*), Tonkerr (*Lóczy*, 15, ex *Kanitz*); COREA: Chemulpo (*Carles*, 129! *Veitch*, 86!) Mus. Brit.; Herb. Kew.

Dahuria.

31. *Lilium tigrinum*, *Ker-Gawl. in Bot. Mag.* t. 1237; *Kunth, Enum. Pl.* iv. p. 259; *Red. Lil.* t. 475; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* p. 156; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 233; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 66; *Franch. Pl. David.* i. p. 301, *et Journ. de Bot.* vi. (1892) p. 319; *Maxim. in Bull. Soc. Nat. Mosc.* (1879) p. 62; *Elwes, Monogr. Lilium*, t. 38.

*Lilium speciosum*, *Andr. Bot. Rep.* t. 586.

CHIHLI: Peking (*David*, 2310, ex *Franchet*); HUPEH: Ichang (*A. Henry*, 4140!), Patung (*A. Henry*, 2397!); SZECHUEN: Tchenkeoutiu (*Farges*, 229, ex *Franchet*); COREA (*Wykeham Perry!*), Chemulpo (*Carles*, 180!); COREAN ARCHIPELAGO (*Oldham*, 871!). Mus. Brit.; Herb. Kew.

Japan.

Dr. A. Henry states that in Ichang this species is cultivated and the bulb eaten. Dried lily flowers are used by the Chinese for flavouring soups and as a remedy for pulmonary diseases. See *Kew Bulletin*, 1889, p. 116.

32. *Lilium yunnanense*, *Franch. in Journ. de Bot.* vi. (1892) p. 314.

YUNNAN: Tapintze (*Delavay*, 1837, 2737, 3577, ex *Franchet*), Mengtze, 6000 ft. (*A. Henry*, 10743! *Hancock*, 174!). *Herb. Kew.*

1. *Nomocharis meleagrina*, *Franch. in Journ. de Bot.* xii. (1898) p. 196.

SZECHUEN: Sela, on the Mékong (*Soulié*, 1032, ex *Franchet*).

2. *Nomocharis pardanthina*, *Franch. in Journ. de Bot.* iii. (1889) p. 113, t. 3.

YUNNAN: Mount Koualapo (*Delavay*, 257, ex *Franchet*).

This genus is intermediate between *Lilium* and *Fritillaria*, from both of which it differs in the following characters:—Lower part of the filaments inflated and hollow, foveola multifid and partly free, inner lobes of the perianth differing from the outer.

1. *Fritillaria camschatcensis*, *Ker-Gawl. in Bot. Mag.* sub t. 1216; *Lamb. in Trans. Linn. Soc.* x. (1811) p. 265, t. 11; *Kunth, Enum. Pl.* iv. p. 254.

*Lilium camschatcense*, *Linn. Sp. Pl.* ed. 1, p. 303; *Lour. Fl. Cochinch.* p. 207.

*Fritillaria kamtschatcensis*, *Fisch. ex Hook. Fl. Bor.-Amer.* ii. p. 181, t. 193a; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 273.

CHINA (ex *Loureiro*).

America north of the United States, Kamtschatka and Japan.

2. *Fritillaria cantoniensis*, *Lour. Fl. Cochinch.* p. 206; *Ker-Gawl. in Bot. Mag.* sub t. 1537; *Kunth, Enum. Pl.* iv. p. 255.

KWANGTUNG: Canton (ex *Loureiro*).

A doubtful plant, supposed by Gawler to be the same as *Uvularia chinensis*, which is now reduced to *Disporum pullum*, *Salisb.*

3. *Fritillaria cirrhosa*, *D. Don, Prodr. Fl. Nepal.* p. 51; *Kunth, Enum. Pl.* iv. p. 253; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 257; *Hook. f. Fl. Brit. Ind.* vi. p. 253.



## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they would appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

Part I. for 1901, consisting of 378 pages, price 21 shillings, was issued on July 10, 1902; Part II., completing the volume, is now in course of compilation.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.



VOL. XXXV. is still in course of issue, the Parts already published being Nos. 242-245. Those still unpublished are Parts 246-248 inclusive.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."

THE JOURNAL  
OF  
THE LINNEAN SOCIETY.

VOL. XXXVI. BOTANY. No. 251.

CONTENTS.

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.—	
PART XVI. ....	137

See Notice on last page of Wrapper.

LONDON:  
SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;  
AND BY  
LONGMANS, GREEN, AND CO.,  
AND  
WILLIAMS AND NORGATE.  
1903.

F 12

# LINNEAN SOCIETY OF LONDON.

---

## LIST OF THE OFFICERS AND COUNCIL.

Elected 24th May, 1902.

---

### PRESIDENT.

Prof. Sydney Howard Vines, M.A., D.Sc., F.R.S.

### VICE-PRESIDENTS.

Frank Crisp, LL.B., B.A.	Prof. J. B. Farmer, F.R.S.
C. B. Clarke, Esq., M.A., F.R.S.	Mr. A. O. Walker.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

Prof. G. B. Howes, D.Sc., LL.D., F.R.S. | D. H. Scott, M.A., Ph.D., F.R.S.

### COUNCIL.

Rev. R. A. Bullen, B.A., F.G.S.	W. G. Ridewood, D.Sc., F.Z.S.
Charles Baron Clarke, M.A., F.R.S.	G. S. Saunders, F.E.S.
Frank Crisp, LL.B., B.A.	Dukinfield H. Scott, M.A., Ph.D., F.R.S.
Prof. J. B. Farmer, M.A., F.R.S.	A. C. Seward, M.A., F.R.S.
Prof. J. Reynolds Green, M.A., D.Sc., F.R.S.	Col. C. Swinhoe, F.Z.S., F.E.S.
Prof. G. B. Howes, D.Sc., LL.D., F.R.S.	Prof. Sydney H. Vines, D.Sc., F.R.S.
B. Daydon Jackson, Esq.	Alfred O. Walker, Esq.
George Masee, Esq.	

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

---

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

---

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*. The former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meets at 4 p.m., as required during the Session.

*Fritillaria verticillata*, *Wall. Cat.* 5079 A, non *Willd.*

SZECHUEN: near Tachienlu (*Pratt*, 609, 845, 852!), Tongolo (*Soulié*, 707!), Tcheto near Tongolo (*Soulié*, 578, ex *Franchet*).  
Herb. Kew.

Var. *ecirrhusa*, *Franch. in Journ. de Bot.* xii. (1898) p. 233.

SZECHUEN: Tachienlu (*Pratt*, 739, ex *Franchet*), Tizon (*Soulié*, 620, 707, ex *Franchet*).

Central and Eastern Himalayas.

4. *Fritillaria dagana*, *Turcz. in Flora*, xvii. (1834) i. p. 25; *Ledeb. Fl. Ross.* iv. p. 148; *Franch. Pl. David.* i. p. 308; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 271.

*Imperialis dagana*, *Turcz. Bull. Soc. Nat. Mosc.* (1837) i. p. 62, et 1838, p. 102.

CHIHLI: Jehol, Mount Nantachan (*David*, 1816, ex *Franchet*), Peishan (*David* in hb. *Hance*, 13188!). Mus. Brit.

Siberia.

5. *Fritillaria Delavayi*, *Franch. in Journ. de Bot.* xii. (1898) p. 222.

YUNNAN: at the base of the glacier of Likiang (*Delavay*, 27, ex *Franchet*).

6. *Fritillaria Przewalskii*, *Maxim. ex Batalin, in Act. Hort. Petrop.* xiii. (1893) i. p. 105.

KANSUH: Tangut, Mudshik (*Przewalski*!), River Myndanscha (*Grum-Grschimailo* ex *Batalin*).

7. *Fritillaria Souliei*, *Franch. in Journ. de Bot.* xii. (1898) p. 221.

SZECHUEN: Dongchingtchratsong, near Tsékong on the Mékong (*Soulié*, 1029, ex *Franchet*).

8. *Fritillaria verticillata*, *Willd.*, var. *Thunbergii*, *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 258; *Hance, in Journ. Bot.* xxi. (1883) p. 358.

*Uvularia cirrhosa*, *Thunb. Fl. Jap.* p. 136; *Kunth, Enum. Pl.* iv. p. 202.

*Fritillaria Thunbergii*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 157.

*Fritillaria collicola*, *Hance, in Journ. Bot.* viii. (1870) p. 76.

CHINA (*Fortune*, 27!); CHEKIANG (*Tarrant* in hb. *Hance*, 1665!), Ningpo (*Maries*!), Meichi (*Forbes & Carles*!); YUNNAN:

summit of Lopinchan (*Delavay*, 2066, ex *Franchet*). Mus. Brit.; Herb. Kew.

Var. *ecirrhusa*, *Franch.* in *Journ de Bot.* xii. (1898) p. 223.

SZECHUEN: Heoupin, near Tchenkeou (*Farges*, 984, ex *Franchet*).

Japan.

1. *Tulipa edulis*, *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 295; *Maxim.* in *Bull. Soc. Nat. Mosc.* (1879) p. 62.

*Orithyia edulis*, *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 158; *Franch.* in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1882) p. 258; *Debeaux*, *Fl. Shanghai*, p. 64.

*Tulipa graminifolia*, *Baker*, in *Journ. Bot.* xiii. (1875) p. 230, et xvi. (1878) p. 138.

SHANTUNG: Chefoo (*Hancock!* *Fauvel* ex *Franchet*); CHEKIANG: Ningpo (*Everard!* *Hancock*, 13!); KIANGSI: Kiukiang (*Bullock*, 244! *Maries!* *Shearer!* *Carles*, 500!); HUPEH: Ichang (*A. Henry*, 3323, 3323 A!). Mus. Brit.; Herb. Kew.

Japan.

2. *Tulipa erythronioides*, *Baker*, in *Journ. Bot.* xiii. (1875) p. 292.

CHEKIANG: Snowy Valley (*Quekett!*), Tientsi Mountains (*Faber*, 96!). Herb. Kew.

1. *Gagea lutea*, *Ker-Gawl.* in *Bot. Mag.* t. 1200; *Roem. et Schult. Syst.* vii. pp. 538 & 1703; *Kunth*, *Enum. Pl.* iv. p. 236; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1874) p. 388; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 61; *Hook. f. Fl. Brit. Ind.* vi. p. 355.

*Gagea fascicularis*, *Salisb.* in *Kon. & Sims, Ann. Bot.* ii. (1806) p. 555, ex parte.

*Ornithogalum luteum*, *Linn. Sp. Pl.* ed. 1, p. 306.

SHINGKING: Fungshan (*Ross*, 30!), Muchi (*Webster*, 141!); COREA: Soul Mountains (*Carles!*). Herb. Kew.

Northern India and Japan.

2. *Gagea pauciflora*, *Turcz.* in *Bull. Soc. Nat. Mosc.* (1838) p. 102; *Ledeb. Fl. Ross.* iv. p. 143; *Franch. Pl. David.* i. p. 308.

*Plecostigma pauciflorum*, *Turcz.* in *Trautv. Pl. Fl. Ross. Imag.* p. 8,

t. 2, et *Fl. Baic.-Dahur.* ii. p. 209; *Maxim. Ind. Fl. Mongol. in Prim. Fl. Amur.* p. 485.

*Ornithogalum pauciflorum*, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. p. 113.

*Tulipa ornithogaloïdes*, *Fisch. ex Ledeb. Fl. Ross.* iv. p. 143.

*Szechenyia lloydioïdes*, *Kanitz, in Bot. Res. Szech. Centr. As. Exped.* p. 60, t. 7. figs. 1-3.

KANSUH: Tangut (*Przewalski! Loczy ex Kanitz*). Herb. Kew.

Central and Northern Asia.

1. *Lloydia alpina*, *Salisb. in Trans. Hort. Soc.* i. (1812) p. 328.

*Lloydia serotina*, *Sweet, Hort. Brit.* ed. 2, p. 527; *Reichb. Fl. Germ. Excurs.* p. 102; *Kunth, Enum. Pl.* iv. p. 244; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 300; *Hance, in Journ. Bot.* xvii. (1879) p. 15; *Hook. f. Fl. Brit. Ind.* vi. p. 354.

*Anthericum serotinum*, *Linn. Sp. Pl.* ed. 2, p. 444.

*Gagea serotina*, *Ker-Gawl. in Journ. Roy. Inst.* i. (1816) p. 180.

NORTH CHINA: Siaowutaishan (*Hancock in hb. Hance!*); SZECHUEN: near Tachienlu (*Pratt, 532!*). Mus. Brit.; Herb. Kew.

Widely distributed through the mountainous regions of the North Temperate Zone.

2. *Lloydia Delavayi*, *Franch. in Journ. de Bot.* xii. (1898) p. 193.

YUNNAN: Mount Tsangchan (*Delavay, 95, ex Franchet*).

3. *Lloydia filiformis*, *Franch. in Journ. de Bot.* xii. (1898) p. 192.

SZECHUEN: Tongola and Mount Dzeura (ex *Franchet*; collector's name not stated).

4. *Lloydia ixiolirioides*, *Baker, in Hook. Ic. Pl.* t. 2215.

SZECHUEN: Tachienlu (*Pratt, 533!*). Herb. Kew.

5. *Lloydia oxycarpa*, *Franch. in Journ. de Bot.* xii. (1898) p. 192.

YUNNAN: Heechanmen (*Delavay, 1554, ex Franchet*), Maeulchan (*Delavay ex Franchet*), Likiang (*Delavay ex Franchet*), Tsangchan, near Tali (*Delavay, 93, 294, ex Franchet*).

6. *Lloydia tibetica*, *Baker*, in *Hook. Ic. Pl.* t. 2216.

SZECHUEN: near Tachienlu (*Pratt*, 857 !). Herb. Kew.

Var. *lutescens*, *Franch.* in *Journ. de Bot.* xii. (1898) p. 193.

SZECHUEN: Tchenkeoutin (*Farges*, 429, ex *Franchet*), Tongolo (*Soulié*, 301, 805, ex *Franchet*), between Litang and Batang (*Prince Henri d'Orléans* ex *Franchet*).

Var. *purpurascens*, *Franch.* in *Journ. de Bot.* xii. (1898) p. 193.

YUNNAN: Likiang (*Delavay*, 25, ex *Franchet*).

7. *Lloydia triflora*, *Baker*, in *Journ. Linn. Soc., Bot.* xiv. (1874) p. 300, et xvii. (1879) p. 388.

*Gagea triflora*, *Schultes f. Syst.* vii. p. 551; *Kunth, Enum. Pl.* iv. p. 242; *Ledeb. Fl. Ross.* iv. p. 141; *Maxim. Prim. Fl. Amur.* p. 278.

*Ornithogalum triflorum*, *Ledeb. in Mém. Acad. Sc. Pétersb.* v. (1812) p. 529; *Ic.* t. 379.

SHINGKING: Kwandien (*Ross*, 115 !), Fungwangchung (*Ross*, 119 !). Herb. Kew.

North-east Asia, Kurile Is., Japan.

1. *Heloniopsis japonica*, *Maxim.* in *Bull. Acad. St. Pétersb.* xi. (1867) p. 436; *Mél. Biol.* vi. p. 211; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 87; *Baker*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 460.

*Scilla japonica*, *Thunb. Fl. Jap.* p. 137; *Ic.* dec. iv. t. 4; *Kunth, Enum. Pl.* iv. p. 330.

*Sugerokia japonica*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 145.

COREA: near Phyong Kang (*Carles* !), Soul Mountains (*Carles*, 136 !). Herb. Kew.

Japan.

2. *Heloniopsis umbellata*, *Baker*, in *Journ. Bot.* xii. (1874) p. 278, et *Journ. Linn. Soc., Bot.* xvii. (1879) p. 460.

FORMOSA (*Swinhoe* !).

1. *Ypsilandra thibetica*, *Franch. Pl. David.* ii. p. 132, t. 17; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 240.

SZECHUEN: near Tachienlu (*Pratt*, 766 !). Herb. Kew.

Tibet.



1. *Tofieldia brevistyla*, *Franch. in Journ. de Bot.* xii. (1898) p. 223.

SZECHUEN: Tachienlu (*Soulié*, 833, ex *Franchet*).

2. *Tofieldia divergens*, *Bureau et Franch. in Journ. de Bot.* v. (1891) p. 157.

SZECHUEN: Fouli (*Bonvalot* ex *Bureau et Franchet*).

3. *Tofieldia iridacea*, *Franch. in Journ. de Bot.* xii. (1898) p. 224; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 239.

YUNNAN: Tchenfongchan (*Delavay*, 4906, ex *Franchet*), near Kulongtchang (*Delavay* ex *Franchet*).

4. *Tofieldia nuda*, *Maxim. in Bull. Acad. St. Pétersb.* xvii. (1872) p. 176; *Mél. Biol.* x. p. 416, non *Franch. et Savat. Enum. Pl. Jap.* ii. p. 89; *Somoko Zusetsu*, vii. t. 29; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 239.

*Tofieldia thibetica*, *Franch. Pl. David.* ii. p. 133.

HUPEH (*A. Henry*, 7627 A!), Hsingshan (*A. Henry*, 6561!); HUNAN (*Hancock*, 40!); SZECHUEN: North Wushan (*A. Henry*, 7627!), Mount Omei (*Faber*, 893!), near Tachienlu (*Pratt*, 9!). Mus. Brit.; Herb. Kew.

Tibet, Japan.

5. *Tofieldia setchuenensis*, *Franch. in Journ. de Bot.* xii. (1898) p. 224; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 239.

SZECHUEN: near Tchenkeoutin (*Farges*, 238, ex *Franchet*).

6. *Tofieldia yunnanensis*, *Franch. in Journ. de Bot.* xii. (1898) p. 225.

YUNNAN: Tsangchan (*Delavay*, 79, ex *Franchet*).

1. *Tricyrtis formosana*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 465.

FORMOSA (*Oldham*, 570! *Hancock*, 11! *A. Henry*, 1267, 1393a!). Mus. Brit.; Herb. Kew.

2. *Tricyrtis latifolia*, *Maxim. in Bull. Acad. St. Pétersb.* xi. (1867) p. 435; *Mél. Biol.* vi. p. 208; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 74; *Baker, in Journ. Linn. Soc., Bot.* xvii. p. 464.

FORMOSA: Bankinsing Mountains (*A. Henry*, 1485!); HUPEH:

Ichang (*A. Henry*, 953 !), Chienshih (*A. Henry*, 5871 !); Patung (*A. Henry*, 4690 !), Nanto (*A. Henry*, 4619 a !). Mus. Brit.; Herb. Kew.

Japan.

3. *Tricyrtis macropoda*, *Miq. in Versl. en Med. Kon. Akad. Wetensch.* ii. (1868) p. 86, et in *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 155; *Gartenfl.* t. 613; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 74; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 464; *Hemsl. in Journ. Bot.* xiv. (1876) p. 210.

KIANGSI: Kiukiang (*Forbes* !). Mus. Brit.

Japan.

4. *Tricyrtis pilosa*, *Wall. Tent. Fl. Nep.* p. 62; *Kunth, Enum. Pl.* iv. p. 279; *Bot. Mag.* t. 4955; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 464; *Franch. in Journ. de Bot.* iv. (1890) p. 319 (*sphalmate T. villosa*); *Hook. f. Fl. Brit. Ind.* vi. p. 359.

*Tricyrtis elegans*, *Wall. Tent. Fl. Nep.* t. 46.

CHIHLI: Tongkeou (*Bodinier ex Franchet*); CHEKIANG: Ningpo (*Faber*, 357 ! *Hancock*, 37 !); KANGSI: Kiukiang (*Shearer! Forbes ex Hemsley*), Lushan Mountains (*Bullock*, 86 !); HUPEH: Patung (*A. Henry*, 4780 !); SZECHUEN: Mount Omei (*Faber*, 222 !). Herb. Kew.

Temperate Himalayas.

1. *Disporum calcaratum*, *D. Don, in Proc. Linn. Soc.* i. (1839) p. 45, et in *Trans. Linn. Soc.* xviii. (1841) p. 516; *Kunth, Enum. Pl.* iv. p. 207; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1874) p. 588; *Hook. f. Fl. Brit. Ind.* vi. p. 359.

*Disporum Wallichii*, *D. Don, in Proc. Linn. Soc.* i. (1839) p. 45, et in *Trans. Linn. Soc.* xviii. (1841) p. 516; *Kunth, Enum. Pl.* iv. p. 207.

*Uvularia Wallichii*, *Wall. Cat.* n. 5087.

YUNNAN: Mengtze (*Hancock*, 46 !). Herb. Kew.

Temperate Himalayas, Khasia Hills, Birma.

2. *Disporum Leschenaultianum*, *D. Don?*, *Hance, in Journ. Bot.* xxv. (1887) p. 13.

KWANGTUNG: Lofaushan (*B. C. Henry*).

3. *Disporum pullum*, *Salisb. in Trans. Hort. Soc.* i. (1812) p. 331; *D. Don, in Trans. Linn. Soc.* xviii. (1841) p. 521; *Kunth, Enum. Pl.* iv. p. 208; *Baker, in Journ. Linn. Soc., Bot.* xiv.

(1875) p. 589; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 216; *Mél. Biol.* xi. p. 860; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 52; *Hook. f. Fl. Brit. Ind.* vi. p. 360.

*Uvularia chinensis*, *Ker-Gawl. in Bot. Mag.* t. 916.

*Streptopus chinensis*, *Sm. in Rees, Cycl.* xxxvii. sub *Uvularia*, n. 4.

FORMOSA: Kelung (*Ford!*); HUPEH (*A. Henry*, 5459, 5545!), Ichang (*A. Henry*, 577, 919, 1179, 2960, 2960 A, 3321, 3782! *Watters* in hb. *Hance*, 21206!); KWANGTUNG: Whampoa (*Hance*, 19079! cult.). Mus. Brit.; Herb. Kew.

Japan, Temperate Himalayas, Java, Sumatra.

4. *Disporum sessile*, *D. Don, Prodr. Fl. Nepal.* p. 50; *Kunth, Enum. Pl.* iv. p. 208; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 216; *Mél. Biol.* xi. p. 860; *A. Gray, in Rep. Perry's Exped.* p. 321; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 589; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 51; *Hance, in Journ. Bot.* xxi. (1883) p. 358, et xxv. (1887) p. 13; *Franch. in Bull. Soc. Bot. France*, xlvi. (1899) p. 214.

*Uvularia sessilis*, *Thunb. Fl. Jap.* p. 135, excl. syn.

CHEKIANG: Meichi (*Carles*, 212! *Maries!*); KIANGSI: Kiu-kiang (*Faber*, 896!); FOKIEN: Mountains of Kuatun (*De la Touche* ex *Franchet*); HUPEH: Nanto (*A. Henry*, 7635!), Patung (*A. Henry*, 247, 1474!); KWANGTUNG: Taimoshan (*Westland!*). Mus. Brit.; Herb. Kew.

Japan.

5. *Disporum smilacinum*, *A. Gray, in Proc. Amer. Acad.* (1858) p. 321, et *Bot. Jap.* p. 414; *Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 147; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 590 excl. syn.; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 387; *Maxim. in Engl. Bot. Jahrb.* vi. p. 53; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 52.

SHINGKING: Chienshan (*Ross*, 525!), Toonghwahien (*Ross*, 26!), Moukden (*James! Webster*, 18!); COREA: Chemulpo (*Carles*, 66!); LUCHU ARCHIPELAGO (*Döderlein* ex *Maximowicz*). Mus. Brit.; Herb. Kew.

Japan, Eastern Siberia.

Var. *viridescens*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 216; *Mél. Biol.* xi. p. 859; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 114.

*Uvularia*? *viridescens*, *Maxim. Prim. Fl. Amur.* p. 273.

*Prosartes viridescens*, *Regel, Flor. Ussur.* p. 162, n. 493.

CHIHLI: Mountains west of Peking (*Tatarinow ex Maximowicz*).  
 COREA: Kyong kwi, Soul (*Gortsche ex Palibin*), Huttschaimeo,  
 Pauckhan, Tap Tong, and Mt. Yisan (*Sontag ex Palibin*).  
 Manschuria.

6. *Disporum uniflorum*, *Baker, in Journ. Bot.* xiii. (1875)  
 p. 230; *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 216;  
*Mél. Biol.* xi. p. 860.

KIANGSI: Kiukiang (*Shearer*!). *Herb. Kew.*

1. *Clintonia udensis*, *Trautv. et Mey. Fl. Ochot.* p. 92; *Maxim. Prim. Fl. Amur.* p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 50; *Maxim. in Bull. Soc. Nat. Mosc.* (1879) p. 62; *Franch. Pl. David.* ii. p. 309; *Baker, in Journ. Linn. Soc., Bot.* xiv. (1875) p. 585.

CHIHLI: Pohuashan (*Bretschneider ex Maximowicz*); SHINGKING: Changpeishan (*James*!); HUPEH: South Patung (*A. Henry, 6085*!), Fang (*A. Henry, 6085 a*!), Kuei (*A. Henry, 6085 b*!); SZECHUEN (*A. Henry, 8832! Potanin*!), Mount Omei (*Faber, 891*!), Tachienlu (*Pratt, 319, 685*!). *Mus. Brit.*; *Herb. Kew.*

Japan, Eastern Siberia.

1. *Trillium Tschonoskii*, *Maxim. in Bull. Acad. St. Pétersb.* xxix. (1884) p. 218; *Mél. Biol.* xi. p. 836.

SHINGKING: Moukden (*James*!); HUPEH: (*A. Henry, 6067, 6067 B*!); SZECHUEN: near Tachienlu (*Pratt, 840*!), Mount Omei (*Faber, 980*!). *Mus. Brit.*; *Herb. Kew.*

Japan.

This is very near to *T. erectum*, *Linn.*, of which it may possibly be a variety.

1. *Paris chinensis*, *Franch. Pl. David.* ii. p. 135, *et in Mém. Soc. Philom. Paris, Cent.* (1888) p. 287.

HUPEH: Changyang (*A. Henry, 5230*!); KWEICHAU (*Perny ex Franchet*). *Herb. Kew.*

Tibet.

2. *Paris Delavayi*, *Franch. in Journ. de Bot.* xii. (1898) p. 190.

YUNNAN: Longki (*Delavay ex Franchet*).

3. *Paris Fargesii*, *Franch. in Journ. de Bot.* xii. (1898) p. 190.  
SZECHUEN: Tchenkeoutin (*Farges ex Franchet*).

4. *Paris incompleta*, *Bieb. Fl. Taur.-Cauc.* i. p. 306, et iii. p. 287; *Bieb. Cent. Pl. Rar. Ross.* t. 74; *Kunth, Enum. Pl.* v. p. 119; *Ledeb. Fl. Ross.* iv. p. 121; *Franch. in Mém. Soc. Philom. Paris, Cent.* (1888) p. 283.

COREA (*Wykeham Perry*, 43!). Herb. Kew.

Caucasus.

5. *Paris petiolata*, *Baker*, MSS. in Herb. Kew.

*Herba* erecta. *Folia* circa 7, verticillata, herbacea, ovata, breviter acuminata, basi subcordata, sæpe inæqualia, 3 poll. longa, 2 poll. lata; petiolus 1-1½ poll. longus. *Pedunculus* 7 poll. longus. *Perianthii* segmenta exteriora 5, ovato-elliptica breviter acuminata, basi brevissime et late unguiculata, 1¼ poll. longa, 9 lin. lata; interiora exterioribus multo breviora, anguste oblanceolata, 10 lin. longa, 2 lin. lata. *Stamina* 10, 6 lin. longa; connectivum supra antheram longe productum. *Ovarium* 5-angulatum; stigmata 5, crassiuscula, recurva.

SZECHUEN: near Tachienlu (*Pratt*, 572\*!). Herb. Kew.

Var. *membranacea*, *Wright* (an sp. distincta?). A typo differt foliis majoribus basi cordatis, 4-6 poll. longis, 2-4 poll. latis, perianthii segmentis exterioribus angustioribus, interioribus linearibus, staminum connectivo apice in globulo (non mucrone) producto.

HUPEH: South Patung (*A. Henry*, 5385, 5385 A!). Herb. Kew.

6. *Paris polyphylla*, *Sm. in Rees's Cyclop.* xxvi. n. 2; *D. Don, Prodr. Fl. Nepal.* p. 49; *Kunth, Enum. Pl.* v. p. 118; *Wall. Pl. As. Rar.* t. 126; *Hook. f. Illust. Himal. Pl.* t. 24; *Hook. f. Fl. Brit. Ind.* vi. p. 362; *S. Moore, in Journ. Bot.* xvi. (1878) p. 138, et xxi. (1883) p. 358; *Franch. in Mém. Soc. Philom. Paris, Cent.* (1888) p. 287, et *Bull. Soc. Bot. France*, xlvi. (1899) p. 214.

SHINGKING: Laoling (*Webster*, 10a!); CHEKIANG: Niugpo (*Everard*!); KIANGSI: Kiukiang (*Shearer*! *Maries*! *Faber*, 305!), Lushan Mountains (*Styan*!); FOKIEN: Mountains of Kuantun (*De la Touche ex Franchet*); HUPEH: Patung (*A. Henry*, 206, 2432, 2831, 4083, 4088!); SZECHUEN: South Wushan (*A. Henry*, 5387 a), Mount Omei (*Faber*, 306!), near Tachienlu (*Pratt*, 56! *Soulié*, 185, 213!); KWEICHAU (*Perny ex Franchet*);

YUNNAN (*Delavay ex Franchet*); KWANGTUNG: Lofaushan (*Faber* in hb. *Hance*, 22122!). Mus. Brit.; Herb. Kew.

Himalaya.

Var. *brachystemon*, *Franch. in Journ. de Bot.* xii. (1898) p. 191.

SZECHUEN: Tchenkeoutin (*Farges ex Franchet*).

Var. *platypetala*, *Franch. in Journ. de Bot.* xii. (1898) p. 191.

SZECHUEN: Tchenkeoutin (*Farges*, 573, ex *Franchet*).

7. *Paris quadrifolia*, *Linn. Sp. Pl.* ed. 1, p. 367; *Kunth, Enum. Pl.* v. p. 116; *Franch. in Mém. Soc. Philom. Paris, Cent.* (1888) p. 278.

SZECHUEN: Mount Omei (*Faber*, 307!). Herb. Kew.

Var. *setchuenensis*, *Franch. in Journ. de Bot.* xii. (1898) p. 191.

SZECHUEN: Tchenkeoutin (*Farges*, 414, ex *Franchet*).

Europe, Northern Asia.

Var. *obovata*, *Regel et Til. Fl. Ajan.* p. 121; *Maxim. Prim. Fl. Amur.* p. 272; *Regel, Fl. Ussur.* p. 162; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 56; *Franch. Pl. David.* ii. p. 134; *Franch. in Mém. Soc. Philom. Paris, Cent.* (1888) p. 282.

*Paris obovata*, *Ledeb. Ic. Pl. Ross.* i. t. 16; *Kunth, Enum. Pl.* v. p. 117.

*Paris hexaphylla*, *Cham. in Linnæa*, vi. (1831) p. 586; *Ledeb. Fl. Ross.* iv. p. 120; *Kunth, Enum. Pl.* v. p. 117; *Franch. Pl. David.* i. p. 309.

SHINGKING: Tungcheshien (*James*!), near Moukden (*James*!), Chienshan (*Ross*, 448!), Ngai River (*Ross*, 226!). Herb. Kew.

Northern Asia, Japan.

8. *Paris yunnanensis*, *Franch. in Mém. Soc. Philom. Paris, Cent.* (1888) p. 290.

YUNNAN: Chetchotze (*Delavay ex Franchet*).

Said to be allied to *P. polyphylla*, Sm., but to differ in the leaves being almost coriaceous and the anthers elongated.

[*Melanthium cochinchinese*, Lour. *Fl. Cochinch.* p. 216, is a doubtful plant. *Hance* suggests that it may be *Asparagus lucidus*, Lindl., with which the description agrees, except that the fruit is said to be capsular. *Vide Benth. Fl. Hongk.* p. 371.]

1. *Veratrum album*, *Linn. Sp. Pl.* p. 1044; *Maxim. Prim. Fl. Amur.* p. 289; *Red. Lil.* t. 447; *Kunth, Enum. Pl.* iv. p. 186; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 470.

CHIHLI: Peking (*Tatarinow ex Maximowicz*); HUPEH: South Patung (*A. Henry, 6060!*); SZECHUEN: Mount Omei (*Faber, 888!*). *Mus. Brit.*; *Herb. Kew.*

Mountains of Europe and Northern Asia.

*Var. viride*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 471. *Veratrum viride*, *Ait. Hort. Kew.* iii. p. 422; *Kunth, Enum. Pl.* iv. p. 188.

*Helonias viridis*, *Ker-Gawl. in Bot. Mag.* t. 1096.

*Melanthium viride*, *Thunb. Diss. Melanth.* pp. 3 & 6.

SHINGKING: between Moukden and Kirin (*James*). *Herb. Kew.*

North America.

2. *Veratrum bracteatum*, *Batalin, in Act. Hort. Petrop.* xiii. (1893) i. p. 106.

KANSUH: between Morping and Wuping (*Potanin ex Batalin*), Mount Tshagola (*Potanin ex Batalin*).

3. *Veratrum Maackii*, *Regel, Fl. Ussur.* p. 169, t. xi. figg. 8-14; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 473.

*Zygadenus japonicus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 146.

SHINGKING: between Moukden and Kirin (*James!*); YUNNAN: Mengtze (*Hancock, 81!*); COREA: Chemulpo (*Carles, 86!*). *Herb. Kew.*

Northern Asia, Japan.

An imperfect specimen at Kew collected in Ichang may belong to this species.

4. *Veratrum Maximowiczii*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 472; *Franch. Pl. David.* i. p. 309.

*Veratrum parviflorum*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 91, non *Michx.*

KIANGSI: Kiukiang (*David, 842, ex Franchet*); HUPEH: Nanto (*A. Henry, 4610 A!*), Ichang (*A. Henry, 4610!*). *Mus. Brit.*; *Herb. Kew.*

Japan.

5. *Veratrum nigrum*, *Linn. Sp. Pl.* ed. 1, p. 1044; *Red. Lil.* t. 416; *Bot. Mag.* t. 963; *Kunth, Enum. Pl.* iv. p. 186; *Franch.*

*et Savat. Enum. Pl. Jap.* ii. p. 90; *Maxim. Prim. Fl. Amur.* p. 289; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 472; *Franch. Pl. David.* i. p. 310.

*Melanthium nigrum*, *Thunb. Diss. Melanth.* pp. 4 & 8.

CHIHLI: Jehol (*David, 2037, ex Franchet*); SHINGKING: Tanghoko (*James!*); HUPEH: Hsingshan and Fang (*A. Henry, 6082 a!*), South Patung (*A. Henry, 6082 b!*); SZECHUEN: Tachienlu (*Pratt, 718!*), Tongolo (*Soulié, 931!*). *Mus. Brit.*; *Herb. Kew.*

Central Europe, Northern Asia, Japan.

Var. *japonicum*, *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 472.

FORMOSA: Tamsui (*Hancock, 30!*); SZECHUEN: Mount Omei (*Faber, 937, 960!*); KWANGTUNG: Lofaushan (*Ford!*). *Herb. Kew.*

Japan.

This variety connects *V. nigrum*, Linn., with *V. Mackii*, Regel.

1. *Zygadenus sibiricus*, *A. Gray, in Ann. Lyc. New York*, iv. (1837) p. 112.

*Melanthium sibiricum*, *Linn. Amœn. Acad.* ii. p. 348, t. 4, n. 11; *Linn. Sp. Pl.* ed. 1, p. 339.

*Anthericum?* *Gmelinianum*, *Schultes f. Syst.* vii. p. 481.

*Leimanthium?* *sibiricum*, *Schultes f. Syst.* vii. p. 1551.

*Anticlea sibirica*, *Kunth, Enum. Pl.* iv. p. 191; *Ledeb. Fl. Ross.* iv. p. 207; *Baker, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 483.

HUPEH: Fang (*A. Henry, 6889!*); SZECHUEN: one specimen from North Wushan (ex *A. Henry*). *Herb. Kew.*

Siberia.

#### 144. PONTEDERIACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Monochoria hastæfolia*, *Presl, Reliq. Haenk.* i. p. 128; *Kunth, Enum. Pl.* iv. p. 133; *Miquel, Fl. Ind. Bat.* iii. p. 548; *Thwaites, Enum. Pl. Zeyl.* p. 321; *Hook. f. Fl. Brit. Ind.* vi. p. 362.

*Monochoria hastata*, *Solms-Laubach, in DC. Monogr. Phan.* iv. p. 523.

*Monochoria dilatata*, *Kunth, Enum. Pl.* iv. p. 134.

*Monochoria sagittata*, *Kunth, Enum. Pl.* iv. p. 134.



*Pontederia hastata*, *Linn. Sp. Pl.* p. 288; *Linn. f. Suppl.* p. 192; *Loureiro, Fl. Cochinch.* p. 199; *Roxburgh, Pl. Corom.* ii. p. 6, t. 111.

*Pontederia dilatata*, *Buchanan, in Symes, Embassy to Ava*, p. 475 cum fig., et ed. 2, iii. p. 309, t. 23; *Andr. Bot. Rep.* vii. t. 490.

*Pontederia sagittata*, *Roxb. Fl. Ind.* ii. p. 124.

KWANGTUNG: Lofaushan (*Ford*, 331!), opposite Hongkong (*Ford*, 161!), Fatshan (*Wenyon*!), Whampoa (*Hance*, 1372, ex *Solms-Laubach*); HAINAN (*A. Henry*, 8238! a form with more racemose flowers). *Herb. Kew.*

Cambodia, India, Malay Archipelago, New Guinea.

2. *Monochoria Korsakowii*, *Regel & Maack, in Mém. Acad. Pétersb.* iv. 1861, p. 155; *Regel, Opuit Florui Usuriuskoi*, p. 170, t. 12. figs. 1-7; *Gartenfl.* (1862) p. 313, t. 374; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 88; *Franch. Pl. David.* i. p. 310.

*Monochoria vaginalis*, var. *Korsakowii*, *Solms-Laubach, in DC. Monogr. Phan.* iv. p. 525; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 115.

CHIH LI: Peking (*Bretschneider*, 768!; *Williams* in herb. *Hance*, 13463!, *David*!); SHINGKING (*Ross*, 40! *Webster*, 262!); SHANTUNG (*Faber*, 92!); COREA: Chemulpo (*Carles*!), Soul (*Gottsche, Kalinowsky*). *Mus. Brit.*; *Herb. Kew.*

Mandshuria, Japan.

3. *Monochoria ovata*, *Kunth, Enum. Pl.* iv. p. 665.

*Pontederia ovata*, *Hook. & Arn. Bot. Beechey's Voy.* p. 218.

KWANGTUNG: Canton (*Millett*).

I have seen no specimen of this imperfectly described species, which may, perhaps, be the same as *M. Korsakowii*, *Regel & Maack*.

4. *Monochoria vaginalis*, *Presl, Reliq. Haenk.* i. p. 128; *Kunth, Enum. Pl.* iv. p. 134; *Miq. Fl. Ind. Bat.* iii. p. 548; *Hook. f. Fl. Brit. Ind.* vi. p. 363; *Benth. Fl. Hongk.* p. 374; *Solms-Laubach, in DC. Monogr. Phan.* iv. p. 524.

*Pontederia vaginalis*, *Burm. f. Fl. Ind.* p. 80; *Linn. Mant.* p. 222; *Roxb. Fl. Ind.* ii. p. 121; et *Pl. Coromand.* ii. p. 6, t. 110.

CHIH LI (*Ross*, 288!); KIANGSI: Kiukiang (*Shearer*!); FORMOSA (*Wilford*, 150! *Oldham*, 644! *Swinhoe*, 5! *Playfair*, 170!); HUPEH: Ichang (*A. Henry*, 129! 2357!); SZECHUEN: Mount Omei (*Faber*, 243!); HONGKONG: Happy Valley (*Wilford*, 150!); HAINAN (*A. Henry*, 8368! 8780!). *Mus. Brit.*; *Herb. Kew.*

Japan, India, and Malay Archipelago.

Var. *plantaginea*, *Solms-Laubach*, in *DC. Monogr. Phan.* iv. p. 524.

*Monochoria plantaginea*, *Kunth*, *Enum. Pl.* iv. p. 135; *Benth. Fl. Hongk.* p. 375.

*Monochoria linearis*, *Miq. Fl. Ind. Bat.* iii. p. 549.

*Monochoria Junghuhniana*, *Hassk.*, in *Flora*, xxxv. (1852) p. 115.

*Pontederia plantaginea*, *Roxb. Fl. Ind.* ii. p. 123.

*Pontederia linearis*, *Hassk. Cat. Hort. Bogor.* p. 28, et in *Flora*, xxv. (1842) ii. *Beibl.* p. 4.

FOKIEN: Amoy (*Hance*, 1426!); FORMOSA: Takow (*Henry*, 718!), South Cape (*Henry*, 718 A!); HONGKONG (*Wright*, 540!); HAINAN (*A. Henry*, 8386!). *Mus. Brit.*; *Herb. Kew.*

Japan, Tonquin, India, and Malay Archipelago.

#### 145. PHILYDRACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Philydrum lanuginosum*, *Banks*, in *Gaertn. Fruct.* i. p. 62, t. 16. fig. 10; *Bot. Mag.* t. 783; *Kunth*, *Enum. Pl.* iii. p. 380; *Griffith*, *Notul.* iii. p. 231, *Icon.* t. 269; *Benth. Fl. Hongk.* p. 380; *Hance*, in *Journ. Bot.* ix. (1871) p. 202; *Caruel*, in *DC. Monogr. Phan.* iii. p. 3; *Hook. f. Fl. Brit. Ind.* vi. p. 363.

*Garciana cochinchinensis*, *Lour. Fl. Cochinch.* p. 15.

KWANGTUNG: Pakhoi (*Playfair*, 199!), Canton (*Loureiro*); HONGKONG, Happy Valley and Saywan (*Wilford*, 315! *Wright*, 539! *Champion*, 290! *Seemann*, and *Leclancher ex Caruel*). *Mus. Brit.*; *Herb. Kew.*

Cochinchina, Tonquin, Malay Peninsula and Archipelago, Australia.

#### 146. XYRIDEÆ.

(By N. E. BROWN, A.L.S.)

1. *Xyris bancana*, *Miq. Fl. Ind. Bat., Suppl.* i. p. 608; *Nilsson*, in *Kongl. Svensk. Vet.-Akad. Handl.* xxiv. no. 14 (1892), p. 37; *Hook. f. Fl. Brit. Ind.* vi. p. 366.

*Xyris schœnoides*, *Benth. Fl. Hongk.* p. 379, non *Martius*.

HONGKONG (*Wright*, 548!). *Herb. Kew.*

The material of this plant at Kew is too meagre to permit of

an accurate comparison by dissection, but it is either *X. bancana* or a new species, and is perfectly distinct from *X. schænoïdes*, Mart., with which Bentham identified it.

2. *Xyris pauciflora*, Willd. *Phytog.* p. 2, t. 1. fig. 1; et *Sp. Pl.* i. p. 255; *Wall. Pl. Asiat. Rar.* iii. p. 29; *Kunth, Enum. Pl.* iv. p. 17; *Benth. Fl. Hongk.* p. 379; *Nilsson, in Kongl. Svensk. Vet.-Akad. Handl.* xxiv. no. 14 (1892), p. 36; *Hook. f. Fl. Brit. Ind.* vi. p. 365.

KWANGTUNG: Pakhoi (*Playfair*, 2!); HONGKONG: Happy Valley (*Wilford*, 57! *Wright*, 547!). Herb. Kew.

Tonquin, India.

## 147. COMMELINACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Aneilema angustifolium*, N. E. Brown, n. sp.

*Herba A. sinico* simillima sed minor, 6–12 poll. alta. *Caules* floriferi erecti, graciles, glabri. *Folia* radicalia 4–9 poll. longa,  $1\frac{1}{2}$ –3 lin. lata, linearia, acuta, glabra; caulina simillima sed minora, 1–3 poll. longa, vaginis ciliatis. *Paniculæ* terminales, pauciramosæ, glabræ; rami 1–3 poll. longi, parte florifera cicatricibus approximatis notata. *Bracteæ*  $1\frac{1}{2}$  lin. longæ, ovatæ, acutæ vel subacutæ, caducæ. *Pedicelli*  $1\frac{1}{2}$  lin. longi. *Sepala*  $1\frac{3}{4}$ –2 lin. longa, ovata, subobtusata. *Petala* circa 2 lin. longa, cærulea. *Stamina* 2 fertilia, 3 sterilia, filamentis barbatis. *Capsula* trigona,  $1\frac{3}{4}$  lin. longa, obtusa, apiculata. *Semina* foveolato-rugosa, minutissime furfuracea.

*Aneilema nudiflorum*, R. Br., var. *rigidior*, *Benth. Fl. Hongk.* p. 377.

FORMOSA: Takow (*A. Henry*, 1897!), and without precise locality (*Oldham*, 576 partly!); SZECHUEN: above Chungking (*Faber*, 106!); KWANGTUNG: Lantao Island (*Ford*, 398!), Pakhoi (*Playfair*, 135!, 225!); HONGKONG (*Wright*, 546!). Mus. Brit.; Herb. Kew.

2. *Aneilema divergens*, C. B. Clarke, *Commel. & Cyrt. Beng.* t. 16; et in *DC. Monogr. Phan.* iii. p. 203; *Hook. f. Fl. Brit. Ind.* vi. p. 376.

*Aneilema herbaceum*, Wall., var. *divergens*, C. B. Clarke, in *Journ. Linn. Soc., Bot.* xi. (1870) p. 448.

FORMOSA: Bankinsing Mountains (*A. Henry*, 899!); YUNNAN: Hotha (*Anderson*!). Herb. Kew.

India.

3. *Aneilema formosanum*, *N. E. Brown*, n. sp.

*Herba* 12–15 poll. alta. *Radices* fibrosæ, crassæ. *Folia* radicalia 8–11 poll. longa,  $\frac{1}{2}$ –1 poll. lata, ensiformia, acuminata, basi angustata, utrinque parce puberula. *Scapi* erecti, 7–10 poll. longi, aphylli, vaginis duabus 5–8 lin. longis instructi, puberuli, fasciculos 1–4 gerentes, fasciculis e ramulis 2–3 erectis,  $\frac{1}{4}$ –2 poll. longis puberulis compositis. *Bracteæ* 3–6 lin. longæ et bracteolæ 2–3 lin. longæ, ovatæ, acutæ, amplexicaules, submembranaceæ. *Flores*? *Capsula* trigono-oblonga, obtusa, apiculata. *Semina* obscure et minute tuberculata.

FORMOSA: Bankinsing (*A. Henry*, 872!). Herb. Kew.

Allied to *A. scapiflorum*, Wight.

4. *Aneilema japonicum*, *Maxim. in Engl. Bot. Jahrb.* vi. (1885) p. 53, vix *Kunth*.

LUCHU ARCHIPELAGO: Amami-O-sima (*Döderlein*).

This can scarcely be the *Commelina japonica*, Thunb., in *Trans. Linn. Soc.* ii. (1794) p. 332, with which Maximowicz identified it, since the description differs in several important particulars. Mr. C. B. Clarke appears to have seen no specimen that he could identify with Thunberg's description. (See DC. *Monogr. Phan.* iii. p. 231.)

5. *Aneilema Keisak*, *Hassk. Commel. Ind.* p. 32; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 209; *Franch. Pl. David.* i. p. 310.

*Aneilema oliganthum*, *Franch. et Savat. Enum. Fl. Jap.* ii. pp. 94 et 532.

SHANTUNG: at the foot of Chefoo Mountains (*Faber*, 280!); KIANGSU: Hills near Shanghai (*Carles*, 420!); KIANGSI: Kiu-kiang (*Shearer! David*, 922); FOKIEN: Amoy (*Swinhoe*!); HUPEH: Ichang (*A. Henry*, 128, 2757!). Mus. Brit.; Herb. Kew.

6. *Aneilema Loureirii*, *Hance*, in *Journ. Bot.* vi. (1868) p. 250; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 201; *Hook. f. Fl. Brit. Ind.* vi. p. 375.

*Aneilema spectabile*, *Kurz*, in *Journ. As. Soc. Beng.* xl. (1871) pt. 2, p. 77; *C. B. Clarke*, *Commel. & Cyrt. Beng.* t. 20.

*Commelina tuberosa*, *Lour. Fl. Cochinch.* p. 40.

KWANGTUNG: Foot of White Cloud Hill or Pakwan Mountain, Canton (*Sampson*, 252! hb. *Hance*, 13842!). Mus. Brit.; Herb. Kew.

Burma, Siam.

7. *Aneilema medicum*, *R. Br. Prodr.* p. 271 in adn.; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 202.

*Commelina medica*, *Lour. Fl. Cochinch.* p. 40.

CHINA: without locality (*Loureiro*).

Cochinchina, *ex Loureiro*.

Bretschneider, in his 'Early European Researches into the Flora of China,' p. 179, states that the Chinese name quoted by Loureiro for this plant, "in China as well as in Japan is applied to *Ophiopogon japonicus*, Ker." But Loureiro's description of the flowers and fruit in no way agrees with those of *Ophiopogon*.

8. *Aneilema nudiflorum*, *R. Br. Prodr.* p. 271 in adn.; *Benth. Fl. Hongk.* p. 376; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 21; *et in DC. Monogr. Phan.* iii. p. 210, excl. var.  $\gamma$ ; *Hook. f. Fl. Brit. Ind.* vi. p. 378.

*Commelina nudiflora*, *Linn. Mant.* ii. p. 177 (nec *Sp. Pl.*); *Reichenb. Icon. Bot. Exot.* cent. 2, p. 15, t. 136.

FOKIEN: Amoy (*Swinhoe*!); FORMOSA (*Oldham*, 576 partly! *Wilford*, 551! *A. Henry*, 882!); HUPEH: Ichang (*A. Henry*, 2252!); SZECHUEN: Mount Omei, lower regions (*Faber*, 245!); KWANGTUNG: Lofaushan (*Ford*, 138!), Canton (*Sampson*, 253! *Lamont*!); HONGKONG (*Wilford*, 49! *Hance*, 256! *Lamont*!); HAINAN (*A. Henry*, 8334!); LUCHU ARCHIPELAGO (*Wright*, 353!). Mus. Brit.; Herb. Kew.

India, Siam, Malay Archipelago.

9. *Aneilema paucifolium*, *N. E. Brown*, n. sp.

*Herba* 6-8 poll. alta, basi ramosa. *Radices* fibrosæ, crassæ, tomentosæ. *Caules* erecti vel basi subdecumbentes, simplices vel subsimplices, glabri, basi unifoliati rarius bifoliati, florum fasciculos 3-4 racemosim dispositos gerentes. *Folia*  $\frac{1}{2}$ - $2\frac{1}{4}$  poll. longa,  $1\frac{1}{2}$ - $2\frac{1}{2}$  lin. lata, linearia, acuta, canaliculata, arcuato-recurva. *Bracteæ* vaginiformes,  $\frac{1}{2}$  poll. longæ, erectæ, submembranaceæ. *Fasciculi* 4-8-flori. *Pedicelli* erecti, 5-8 lin. longi, medio bracteolati, supra bracteolas articulati et pubescentes, infra articulos glabri. *Bracteolæ* vaginiformes, membranaceæ,

parvæ. *Sepala*  $2\frac{1}{2}$  lin. longa, 1 lin. lata, oblonga vel elliptico-oblonga, subobtusata, apice cærulea, parce pubescentia. *Petala* obovata, cærulea. *Stamina* 2 fertilia, 4 sterilia, filamentis barbatis. *Capsula* immatura trigono-oblonga, loculis 3-4-spermis, seminibus uniseriatis.

KWANGTUNG: Pakhoi (*Playfair*, 131!). Herb. Kew.

A very distinct and remarkable species, belonging to the same group as *A. Keisak*, Hassk.; but the stems appear usually to bear only one developed leaf at their base, the others being reduced to submembranous sheaths, which are erect and rather closely embrace the stem and fascicles of pedicels.

10. *Aneilema scapiflorum*, *Wight*, var. *latifolium*, *N. E. Brown*, n. var.; folia quam ea typi latiora et minus acuminata, 4-7 poll. longa,  $\frac{3}{4}$ - $1\frac{1}{4}$  poll. lata.

HAINAN: Lingmen (*A. Henry*!). Herb. Kew.

The type extends from the Temperate and Tropical Himalayas southwards to Ceylon and Tenasserim.

11. *Aneilema sinicum*, *Lindl. Bot. Reg.* t. 659; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 212; *Hook. f. Fl. Brit. Ind.* vi. p. 379.

*Aneilema secunda*, *Wight, Ic. Pl. Ind. Or.* vi. p. 31, t. 2075.

*Commelina sinica*, *Roem. et Schult. Syst.* i., *Mant.* i., *Additament.* i., p. 376.

KWANGTUNG: West River (*Ford*!), at the foot of Pakwan Mountains (*Sampson* in hb. *Hance*, 14736!); Lantao Island (*Ford*, 402!). Mus. Brit.; Herb. Kew.

India, Malay Archipelago, Tropical and South Africa?, Madagascar?

I doubt the identity of the African plant with this.

12. *Aneilema spiratum*, *R. Br. Prodr.* p. 271 in adn.; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 207; *Hook. f. Fl. Brit. Ind.* vi. p. 377.

*Aneilema nanum*, *Kunth, Enum. Pl.* iv. p. 65; *Wight, Ic. Pl. Ind. Or.* vi. p. 31, t. 2077; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 18.

*Aneilema melanostictum*, *Hance, in Journ. Bot.* vii. (1869) p. 167.

HAINAN (*Swinhoe*! herb. *Hance*, 14412!). Mus. Brit.; Herb. Kew.

India, Malay Archipelago.

*A. melanostictum*, Hance, was doubtfully placed by C. B. Clarke, *DC. Monogr. Phan.* iii. p. 206, as a synonym of *A. esculentum*, Wall., and the incorrect locality "Hongkong" quoted for it; but Swinhoe's specimen from Hainan has been named (in my opinion correctly) *A. spiratum* by Mr. Clarke.

13. *Aneilema vaginatum*, *R. Br. Prodr.* p. 271 in adn.; *Wight, Ic. Pl. Ind. Or.* vi. p. 31, t. 2076; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 23; *et in DC. Monogr. Phan.* iii. p. 216; *Hook. f. Fl. Brit. Ind.* vi. p. 381.

HONGKONG (*Lamont!*). Mus. Brit.

India, Cambodia.

1. *Commelina auriculata*, *Blume, Enum. Pl. Jav.* i. p. 2; *C. B. Clarke, in DC. Monogr. Phan.* iii. p. 186.

FORMOSA: Tamsui (*Oldham, 578!*); KWANGTUNG: Macao (*Meyen*); HONGKONG (*Wright, 544!*). Mus. Brit.; Herb. Kew.  
Malay Archipelago, Australia.

2. *Commelina benghalensis*, *Linn. Sp. Pl.* p. 41; *Wight, Ic. Pl. Ind. Or.* vi. p. 29, t. 2065; *Benth. Fl. Hongk.* p. 376; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 4, *et in DC. Monogr. Phan.* iii. p. 159; *Hook. f. Fl. Brit. Ind.* vi. p. 370.

*Commelina procurrens*, *Schlecht. in Linnæa*, xxiv. (1851) p. 656, *et* xxv. (1852) p. 183; *cf. Gartenfl.* iv. (1855) p. 44.

*Commelina delicatula*, *Schlecht. in Gartenfl.* iii. p. 349, t. 104. fig. a.

FORMOSA (*Oldham, 577!*); HUPEH: Ichang (*A. Henry, 2353!*); SZECHUEN: Hokiang and Nankee rice-fields (*Faber, 965!*); KWANGTUNG: Macao (*Tate!*); HONGKONG (*Wright, 542! Hance, 407! Lamont!*). Mus. Brit.; Herb. Kew.

Widely distributed through the tropics of the eastern hemisphere.

3. *Commelina communis*, *Linn. Sp. Pl.* p. 40; *Benth. Fl. Hongk.* p. 376; *Debeaux, in Act. Soc. Linn. Bordeaux*, xxx. (1874) p. 117, & xxxii. (1875) p. 27; *Fl. Shangh.* p. 65, *et Fl. Tchef.* p. 140; *Hance, in Journ. Bot.* xi. (1874) p. 262; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 260, *et Pl. David.* i. p. 310; *C. B. Clarke, in DC. Monogr. Phan.* iii. p. 170; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 115; *nec Kunth, nec plur. auct.*

*Commelina polygama*, Roth, in Roemer et Usteri, *Mag. Bot.* iv. (1790) p. 14; *et Cat. Bot.* i. p. 1.

*Commelina vulgaris*, Schmidel,  *Ic. Pl.* ed. 2, p. 159, tt. 40, 41.

CHINA (*Fortune*, 24!); CHIHLI: Peking (*Bretschneider*, 769! *Bushell! David*, 471); SHINGKING: Moukden (*Webster*, 310!); Kaichow to Kinchow (*Ross*, 246!); SHANTUNG: Chefoo (*Ford! Debeaux, Fauvel*); without locality (*Maingay*, 133!); KIANGSU: Shanghai (*Maingay*, 389, 439!); CHEKIANG: Ningpo Mountains (*Faber*, 18!); KIANGSI: Kiukiang (*Shearer! Moellendorf, David*, 929); HUPEH: Patung (*A. Henry*, 1800, 4927!), Ichang (*A. Henry*, 76!); HONGKONG (*Lamont*); COREA: Inchon (*Gottsche*), Soul (*Kalinowsky, Sontag*); Chemulpo (*Bunge*). Mus. Brit.; Herb. Kew.

Japan, Mandshuria, Ussuri.

Var. *ludens*, C. B. Clarke, in DC. *Monogr. Phan.* iii. p. 171.

*Commelina ludens*, *Miq. Journ. Bot. Néerl.* i. (1861) p. 88.

KWANGTUNG: Fukwing (*Krone ex Miquel*).

I have seen no specimen of this variety, although it is said in Mr. Clarke's monograph to be frequent in South China.

4. *Commelina? Loureirii*, *Kunth, Enum. Pl.* iv. p. 60; C. B. Clarke, in DC. *Monogr. Phan.* iii. p. 191.

*Lechea chinensis*, *Lour. Fl. Cochinch.* p. 60.

KWANGTUNG: near Canton (ex *Loureiro*).

An altogether doubtful plant.

5. *Commelina nudiflora*, *Linn. Sp. Pl.* p. 41, neque *Mant.* 177, nec *auct.*; C. B. Clarke, in DC. *Monogr. Phan.* iii. p. 144; *Hook. f. Fl. Brit. Ind.* vi. p. 369.

*Commelina longicaulis*, *Jacq. Ic. Pl. Rar.* ii. p. 8, t. 294.

*Commelina ochreatea*, *Schauer, in Nov. Act. Acad. Nat. Cur.* xix., *Suppl.* i. (1843) p. 447.

*Commelina salicifolia*, *Benth. Fl. Hongk.* p. 376, non *Roxb.*

*Commelina communis*, C. B. Clarke, *Commel. & Cyrt. Beng.* t. 1, non *Linn.*

FORMOSA: Takow Plain (*A. Henry*, 2041!), South Cape (*A. Henry*, 357!); KWANGTUNG: Lappas Island (*Vachell*, 83!), Macao and Lintin (*Meyen*); HONGKONG (*Wright*, 543! *Wilford*, 362! *Hance*, 406!); LUCHU (*Wright*, 347!). Mus. Brit.; Herb. Kew.

Widely distributed throughout the warmer regions of both hemispheres.



6. *Commelina obliqua*, *Buch.-Ham., in D. Don, Prodr. Fl. Nep.* p. 45; *C. B. Clarke, Commel. & Cyrt. Beng.* tt. 9, 10, 11, et in *DC. Monogr. Phan.* vi. p. 178; *Hook. f. Fl. Brit. Ind.* vi. p. 372 (non *Vahl*).

*Commelina polyspatha*, *Wight, Ic. Pl. Ind. Or.* vi. p. 29, t. 2066.

FORMOSA: Tamsui (*A. Henry, 1447!*); KWANGTUNG: Lofau-shan (*Ford!*), Lantao Island (*Ford, 406!*). *Herb. Kew.*

India, Malay Archipelago.

7. *Commelina undulata*, *R. Br. Prodr.* p. 270; *C. B. Clarke, in DC. Monogr. Phan.* iii. p. 179; *Hook. f. Fl. Brit. Ind.* vi. p. 373.

CHINA (*Millet! Lay!*); CHUSAN ARCHIPELAGO (*Yvan ex C. B. Clarke*); FORMOSA: Takow (*Playfair, 62!*); KWANGTUNG: Canton (*Hance, 21055!*), Macao (*Gaudichaud, 62, ex C. B. Clarke*); COREA: N.E. coast (*Perry, 105!*); LUCHU (*Wright, 347, 348!*). *Mus. Brit.*; *Herb. Kew.*

India, Australia, Philippines.

1. *Cyanotis arachnoidea*, *C. B. Clarke, in DC. Monogr. Phan.* iii. p. 250; *Hook. f. Fl. Brit. Ind.* vi. p. 386.

*Cyanotis pilosa*, *Wight, Ic. Pl. Ind. Or.* vi. p. 32, t. 2083, non *Roem. et Schult.*

FORMOSA: Bankinsing Mountains (*A. Henry, 900!*). *Herb. Kew.*

Indian Peninsula and Ceylon.

2. *Cyanotis axillaris*, *Roem. & Schult. Syst.* vii. p. 1154; *Benth. Fl. Hongk.* p. 378; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 35, et in *DC. Monogr. Phan.* iii. p. 244; *Hook. f. Fl. Brit. Ind.* vi. p. 388.

*Tradescantia axillaris*, *Roxb. Pl. Corom.* ii. p. 5, t. 107.

KWANGTUNG: Pakhoi (*Playfair, 216!*); HONGKONG (*Hance, 427!*). *Mus. Brit.*; *Herb. Kew.*

India, Malay Archipelago, Australia.

3. *Cyanotis barbata*, *D. Don, Prodr. Fl. Nep.* p. 46; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 37, et in *DC. Monogr. Phan.* iii. p. 248; *Hook. f. Fl. Brit. Ind.* vi. p. 385.

*Cyanotis nobilis*, *Hassk. Commel. Ind.* p. 148; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 39.

*Cyanotis fasciculata*, *Benth. Fl. Hongk.* p. 378, non *Roem. & Schult.*

KWANGTUNG: Mountains of the province of Pakwan, near

Canton (*Sampson*, hb. *Hance*, 428); HONGKONG (*Harland!* *Lamont!*). Mus. Brit.; Herb. Kew.

India, Burma.

4. *Cyanotis cristata*, *Roem. & Schult.*, var. *Griffithii*, *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 248; *Hook. f. Fl. Brit. Ind.* vi. p. 385.

HAINAN (*A. Henry*, 8413!). Herb. Kew.

Malay Peninsula.

Mr. Clarke does not describe the corolla, which is gamopetalous.

1. *Floscopa scandens*, *Lour. Fl. Cochinch.* p. 193; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 265; *Hook. f. Fl. Brit. Ind.* vi. p. 390.

*Floscopa paniculata*, *Benth. Fl. Hongk.* p. 377 (excl. synonym. Afr. et Amer.).

*Dithyrocarpus paniculatus*, *Kunth, Enum. Pl.* iv. p. 79; *C. B. Clarke, Commel. & Cyrt. Beng.* t. 34.

*Dithyrocarpus petiolatus*, *D. undulatus*, & *D. Rothii*, *Wight, Ic. Pl. Ind. Or.* vi. p. 32, tt. 2079, 2080.

KIANGSI (*Staunton!*); FOKIEN: Amoy (*Swinhoe!*); KWANG-TUNG: Lappas Island (*Vachell*, 109!), Macao (*Lay!*), Lofaushan (*Faber*, 4!); HONGKONG (*Harland*, 544! *Lamont!*). Mus. Brit.; Herb. Kew.

India, Burma, Malay Archipelago, Australia.

1. *Forrestia chinensis*, *N. E. Brown*, n. sp.

*Herba* perennis, caule robusto, elato, glabro. *Folia* 7–12 poll. longa,  $1\frac{3}{4}$ – $2\frac{3}{4}$  poll. lata, lanceolata vel elliptico-lanceolata, apice acuminata, basi longe attenuata, supra glabra, subtus minute puberula, interdum costis hirsutis; vaginæ puberulæ vel longe et appresse villosæ, ore villosa-ciliatæ. *Flores* dense capitati, sessiles; bracteæ ovatæ, acutæ, carinatæ, glabræ, marginibus carinisque ciliatis; sepala  $4\frac{1}{2}$ – $5\frac{1}{2}$  lin. longa, oblonga, cucullata, obtusa, acute carinata, carinis dense vel sparse fulvo-hirtis, marginibus ciliatis; petala oblanceolato-oblonga, obtusa; filamenta apice barbata; ovarium superne hirsutum. *Capsula* 3 lin. longa,  $2\frac{1}{2}$  lin. lata, ellipsoideo-trigona, superne hirsuta; semina  $1\frac{1}{4}$ – $1\frac{1}{2}$  lin. longa,  $1\frac{1}{4}$  lin. lata, rugosa.

FORMOSA: Tamsui (*Oldham*, 578/1!), Bankinsing Mountains

(*A. Henry*, 149!, 1618 A!); KWANGTUNG: without locality (*Ford*, 256!); HAINAN: without locality (*A. Henry*, 8670!). Mus. Brit.; Herb. Kew.

This plant was united with *F. hispida*, Less. & Rich., by Mr. Clarke in *DC. Monogr. Phan.* iii. p. 236, but that species is a native of New Guinea and quite distinct.

1. *Pollia sorzogonensis*, *Endl. ex Miq. Fl. Ind. Bat.* iii. p. 541; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 126; *Benth. Fl. Hongk.* p. 377; *Hook. f. Fl. Brit. Ind.* vi. p. 367.

*Aclisia sorzogonensis*, *E. Meyer*, in *Presl, Rel. Haenk.* i. p. 138, t. 25.

FORMOSA: Bankinsing (*A. Henry*, 133, 575!); HUPEH: Ichang (*A. Henry*, 4163!); KWANGTUNG: Lofaushan (*Ford*!); HONGKONG (*Wright*, 545! *Lamont*! *Forbes*! *Hance*, 1035!). Mus. Brit.; Herb. Kew.

Var. *gigantea*, *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 127; *Hook. f. Fl. Brit. Ind.* vi. p. 368.

*Pollia japonica*, *Hance*, in *Journ. Bot.* xvi. (1878) p. 233, not of *Thunb.*

*Aneilema didymum*, *Seem. Bot. Voy. 'Herald,'* p. 422.

HUPEH: Changyang (*A. Henry*, 6231 A!); KWANGTUNG: Filoitsz, North River (*Sampson* in hb. *Hance*, 11388!). Mus. Brit.; Herb. Kew.

India, Malay Archipelago.

2. *Pollia Zollingeri*, *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 127.

FORMOSA (*Wilford*, 521!); SZECHUEN: Mount Omei, 4000 ft. (*Faber*, 309!). Herb. Kew.

Malay Archipelago.

1. *Streptolirion volubile*, *Edgew. in Proc. Linn. Soc.* i. (1845) p. 254; *et Trans. Linn. Soc.* xx. (1846) p. 90, t. 2; *Wight, Ic. Pl. Ind. Or.* vi. p. 32, t. 2081; *C. B. Clarke*, in *DC. Monogr. Phan.* iii. p. 261; *Franch. Pl. David.* i. p. 311; *Hook. f. Fl. Brit. Ind.* vi. p. 389.

CHIHLI: Peking Mountains (*Bretschneider*, 154! *David*, 403, *Bullock*, 98!); HUPEH: Patung district (*A. Henry*, 2428, 2482, 4170!), Nanto (*A. Henry*, 2587!); KWANGTUNG (*Ford*, 261, 310!). Mus. Brit.; Herb. Kew.

Himalaya, Khasia, Burma.

2. *Streptolirion longifolium*, Gagnep. in *Bull. Soc. Bot. France*, xlvii. (1900) p. 334, t. 11.

YUNNAN: near Yunnan-sen (*Ducloux ex Gagnepain*).

1. *Tradescentia vaga*, *Lour. Fl. Cochinch.* p. 193, non *Blume*.

KWANGTUNG: Canton (*Loureiro*).

This is altogether a doubtful plant. It was combined by Roemer & Schultes in their *Systema*, vii. p. 1153, with a Javan plant (*T. vaga*, *Blume*) under the name of *Cyanotis vaga*, *Roem. & Schult.*; but their description is based upon a specimen from Java sent to them by *Blume*, and does not at all accord with that of *Loureiro*. The plant described by *C. B. Clarke* in *DC. Monogr. Phan.* iii. p. 252 as *Cyanotis vaga* is probably that which *Roemer & Schultes* had in view, and cannot be the *T. vaga*, *Lour.*, therefore the name *Cyanotis vaga* must be restricted to the Javan plant.

#### 148. FLAGELLARIACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Flagellaria indica*, *Linn. Sp. Pl.* 333; *Lam. Encycl.* ii. p. 502, t. 266; *Gaertn. Fruct.* i. p. 61, t. 16. f. 9; *Lour. Fl. Cochinch.* p. 211; *Willd. Sp. Pl.* ii. p. 263; *Redouté, Lil.* v. t. 257; *Roem. & Schult. Syst.* vii. pt. 2, p. 1492; *Roxb. Fl. Ind.* ii. p. 154; *Kunth, Enum. Pl.* iii. p. 370; *Hook. f. Fl. Brit. Ind.* vi. p. 391.

FORMOSA: South Cape (*A. Henry*, 1355!); HAINAN (*A. Henry*, 8657!). *Herb. Kew.*

India, Malay Archipelago, Polynesia, and Australia.

#### 149. JUNCACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Luzula campestris*, *DC. Fl. Franç.* iii. p. 161; *Kunth, Enum. Pl.* iii. p. 307; *Hook. f. Fl. Brit. Ind.* vi. p. 401; *Reichb. Ic. Fl. Germ.* ix. p. 14, tt. 375-376; *Syme, English Bot.* ed. 3, x. p. 8, t. 1551; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 155.

*Luzula multiflora*, *Lej. Fl. Envir. Spa*, i. p. 169.

*Luzula campestris*, *DC.*, var. *multiflora*, *Celakov. Prodr. Fl. Böhmen*, 1869, p. 85, & 1881, p. 749; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 161; *Franch. Pl. David.* i. p. 312.

*Luzula campestris*, *DC.*, var. *congesta*, *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 162.

*Luzula multiflora*, *Syme, English Bot. ed. 3, x. t. 1550.*

*Luzula campestris*, *DC., var. capitata, Buchenau, in Engl. Bot. Jahrb. xii. p. 160; Palibin, in Act. Hort. Petrop. xix. (1901) p. 117.*

CHIHLI: Jehol (*David ex Franchet*); KIANGSU: Chingkiang (*Maries!*), Shanghai (*Maingay, 279!*); CHEKIANG: Tientai Mountain, 3000 ft. (*Faber, 74!*); KIANGSI: Kiukiang (*Maries!*); HUPEH: Patung District (*A. Henry, 1793, 3739!*); SZECHUEN: near Tachienlu, 9000–13,000 ft. (*Pratt, 12! Soulié, 753*); COREA (*Perry*), Soul (*Kalinowsky ex Palibin*), Mount Yisan (*Sontag ex Palibin*). Mus. Brit.; Herb. Kew.

Generally distributed in temperate regions.

2. *Luzula chinensis*, *N. E. Brown, n. sp.*

*Herba* perennis, cæspitosa, 2–2½ ped. alta. *Folia* 4–8 poll. longa, 4–9 lin. lata, late linearia, acuminata, glabra, juniora sparsim ciliata, adulta glabra. *Inflorescentia* elongata, interrupta, subcontracta, e cymis 3–4 composita; cymæ corymbosæ, erectæ, compactæ, multifloræ, 2 inferiores distantes; rami filiformes, erecti. *Bracteæ* late ovatæ, acutæ vel breviter acuminatæ. *Perianthii* segmenta subæqualia, 1 lin. longa, acuminata, rubescentia. *Stamina* 6, inclusa. *Pistillum* inclusum; stigmata 3, ovario æquilonga.—*L. effusa*, *Buchenau, in Engl. Bot. Jahrb. xii. (1890) p. 463, nec p. 106.*

SZECHUEN (*A. Henry, 8829! Pratt, 360!*). Mus. Brit.; Herb. Kew.

This more closely resembles the Tropical American *L. gigantea*, *Desv.*, than any other species, but the perianth-segments do not taper to such fine awn-like points, and are reddish instead of dark brown. From *L. effusa*, *Buchen.*, it is at once distinguished by the more numerous flowers and compact, not diffuse, cymes.

3. *Luzula effusa*, *Buchen. Kritisch. Verz. Junc. p. 88; et in Engl. Bot. Jahrb. vi. (1885) p. 196, & xii. (1890) p. 106.*

HUPEH: Fang, 6000–9500 ft. (*A. Henry, 6809!*); KANSU (*Potanin*); SZECHUEN: Mount Omei (*Faber, 1175!*). Mus. Brit.; Herb. Kew.

India.

4. *Luzula pilosa*, *Willd., var. plumosa, Franch. Pl. David. ii. p. 138.*

*Luzula plumosa*, *E. Mey. in Linnæa, xxii. (1849) p. 387; Buchenau, in Engl. Bot. Jahrb. xii. (1890) p. 85.*

SZECHUEN: Moupin (*David ex Franchet*).  
North India to Japan.

5. *Luzula vernalis*, *DC. Fl. Franç.* iii. p. 160; *Desv. in Journ. Bot.* (1808) i. p. 138; *Kunth. Enum. Pl.* iii. p. 297; *E. Mey. in Linnæa*, xxii. (1849) p. 383.

*Luzula pilosa*, *Willd. Enum. Pl. Hort. Berol.* p. 393; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 83, t. 1. fig. 18; *Syme, Engl. Bot.* ed. 3, x. p. 5, t. 1548; *Reichb. Ic. Fl. Germ.* ix. t. 381.

HUPEH: Patung district (*A. Henry*, 3515, 3515 A, 3515 B, 5206!), Changlo (*A. Henry*, 6316!). Mus. Brit.; Herb. Kew.  
North Temperate Zone.

1. *Juncus alatus*, *Franch. & Savat. Enum. Pl. Jap.* ii. pp. 98 & 534; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 304, & xxix. (1900) p. 238.

CHEKIANG: Ningpo Mountains (*Faber*, 6!); KIANGSI: Kiu-kiang (*Shearer*); HUPEH: Patung district (*A. Henry*, 1794!), Chienshih (*A. Henry*, 5785!). Herb. Kew.  
Japan.

2. *Juncus allioides*, *Franch. Pl. David.* ii. p. 137; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 399.

*Juncus macranthus*, *Buchenau, l. c.* xii. (1890) p. 398, & xxix. (1900) p. 238.

HUPEH: Hsingshan (*A. Henry*, 6927!); KANSUH (*Potantin*!); SZECHUEN (*A. Henry*, 8916!), near Tachienlu, 9000–13,000 ft. (*Pratt*, 398! *Soulié*, 827!), Moupin (*David*), Tongolo (*Soulié*, 720, 961!). Herb. Kew.

3. *Juncus bufonius*, *Linn. Sp. Pl.* ed. 1, p. 328; *Kunth, Enum. Pl.* iii. pp. 353 & 597; *Bunge, Enum. Pl. Chin. Bor.* p. 67; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 174, t. 1. figs. 4–5, t. 2. fig. 3, & t. 3. fig. 5; *Syme, English Bot.* ed. 3, x. p. 34, tt. 1572–1573; *Reichb. Ic. Fl. Germ.* ix. p. 18, t. 395; *Franch. Pl. David.* i. p. 311; *Hook. f. Fl. Brit. Ind.* vi. p. 392.

CHIHLI: Jehol (*David*, 1896, ex *Franchet*), near Kantai and Tanshan (ex *Bunge*); SZECHUEN (*A. Henry*, 8843! *Pratt*, 390!). Mus. Brit.; Herb. Kew.

Generally distributed except in the hottest and coldest regions.

4. *Juncus bulbosus*, *Linn. Sp. Pl.* ed. 1, p. 327; *Kunth, Enum. Pl.* iii. p. 351.

*Juncus compressus*, *Jacq. Enum. Stirp. Vind.* pp. 60 & 235; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 261; *et Pl. David.* i. p. 311; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 185; *Reichb. Ic. Fl. Germ.* ix. p. 19, t. 399; *Syme, English Bot.* ed. 3, x. p. 37, t. 1575; *Hook. f. Fl. Brit. Ind.* vi. p. 393; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 116.

*Juncus bottnicus*, *Wahlb. Fl. Lapp.* p. 82, t. 5; *Bunge, Enum. Pl. Chin. Bor.* p. 67.

CHIHLI: Jehol (*David, 1907!*), near Kantai (ex *Bunge*), Tanshan (ex *Bunge*), and without precise locality (*Potanin*); SHINGKING: Laoling (*Ross, 380!*); SHANTUNG: Chefoo (*Fauvel, Perry!*); COREAN ARCHIPELAGO: Port Hamilton (*Oldham, 896!*). *Herb. Kew.*

Europe and North Asia, Kashmir.

5. *Juncus diastrophanthus*, *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 309, & xxix. (1900) p. 238.

SZECHUEN: Nanchuan (*Bock von Rosthorn, 2102, ex Buchenau*).

6. *Juncus effusus*, *Linn. Sp. Pl.* ed. 1, p. 326; *Kunth, Enum. Pl.* iii. p. 320; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 228; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1882) p. 261; *et Pl. David.* ii. p. 137; *Hance, in Journ. Bot.* xiii. (1875) p. 107; *Kurz, in Journ. Bot.* xxxi. (1893) p. 193; *Syme, English Bot.* ed. 3, x. p. 21, t. 1561; *Reichb. Ic. Fl. Germ.* ix. p. 21, t. 413; *Hook. f. Fl. Brit. Ind.* vi. p. 392; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 116.

*Juncus Leersii*, *Buchenau, in Engl. Bot. Jahrb.* xxix. (1900) p. 238, non *Marsson*.

*Scirpus capsularis*, *Lour. Fl. Cochinch.* p. 44.

SHINGKING: Liautong Peninsula (*Swinhoe!*); SHANTUNG: Chefoo (*Fauvel ex Franchet*); KIANGSU: Shanghai (*Maingay, 421!*); CHEKIANG: Ningpo (*Cooper!, Swinhoe!*); FOKIEN: Amoy (*Sampson!*); FORMOSA: Tamsui (*Oldham, 580!*); HUPEH: Patung district (*A. Henry, 1781, 4049!*), Nanto and mountains to the northward (*A. Henry, 1953, 3855!*); SZECHUEN: Mount Omei, 3400 ft. (*Faber, 1136!*), Kinshan (*Bock von Rosthorn, 78, ex Buchenau*), Moupin (*David ex Franchet*); YUNNAN (*Anderson!*); KWANGTUNG: Canton, near the White Cloud Monastery

(*Sampson*, 843, 2157 !); COREA, Port Chusan (*Wilford*, 935 !), Soul (*Kalinowsky ex Palibin*); COREAN ARCHIPELAGO (*Oldham*, 894 !); LUCHU (*Wright*, 328 !). Mus. Brit.; Herb. Kew.

Europe, Asia, America, Africa, and Australia.

7. *Juncus glaucus*, *Sibth. Fl. Oxon.* p. 113; *Kunth, Enum. Pl.* iii. p. 316; *Syme, English Bot.* ed. 3, x. p. 25, t. 1563; *Reichb. Ic. Fl. Germ.* ix. p. 22, t. 415; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 243, t. 1. figs. 2-3, et t. 3. fig. 8; *Hook. f. Fl. Brit. Ind.* vi. p. 393.

*Juncus inflexus*, *Huds. Fl. Angl.* ed. 1, p. 130 (1762)! and perhaps also of *Linn. Sp. Pl.* ed. 1, p. 326.

KANSUH (*Potanin*!). Herb. Kew.

Europe, Asia, Africa, and New Zealand.

8. *Juncus himalensis*, *Klotzsch, in Klotzsch & Garcke, Bot. Ergebn. Reise Prinz Waldemar*, p. 60, t. 97; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 405; *Hook. f. Fl. Brit. Ind.* vi. p. 398.

SZECHUEN: near Tachienlu, 9000-13,500 ft. (*Pratt*, 96, 844 !). Mus. Brit.; Herb. Kew.

India.

*Pratt's* no. 844 has broader leaves than is usual in this species, but in other characters seems not to differ.

9. *Juncus khasiensis*, *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 407; *Hook. f. Fl. Brit. Ind.* vi. p. 399.

SZECHUEN: Tachienlu (*Soulié*, 652 !). Herb. Kew.

Also in Assam.

10. *Juncus luzuliformis*, *Franch. Pl. David.* ii. p. 137; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 204, et xxix. (1900) p. 238.

SZECHUEN: Moupin (*David ex Franchet*).

11. *Juncus modestus*, *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 203.

HUPEH: Fang (*A. Henry*, 6846, 6868 !); SZECHUEN: Valley of the River Heiho, above Hunneiku (*Potanin ex Buchenau*), near Tachienlu, 9000-10,000 ft. (*Pratt*, 162 !). Mus. Brit.; Herb. Kew.



12. *Juncus modicus*, *N. E. Brown*, n. sp.

Dense cæspitosus. *Caules* 3-5 poll. longi, filiformes vel setacei. *Cataphylla* basilaria truncata, setaceo-mucronata. *Folia* caulina 2, setacea vel filiformia, inferius  $2\frac{1}{2}$ - $4\frac{1}{2}$  poll. longum, superius  $\frac{1}{4}$ -1 poll. longum. *Capitula* 3-8-flora. *Bracteæ* flores subæquantes vel breviores, lanceolatae, acutæ, pallidæ. *Perianthii* segmenta  $1\frac{1}{2}$ - $2\frac{1}{2}$  lin. longa,  $\frac{1}{4}$  lin. lata, linearia vel lineari-lanceolata, acuta, pallida. *Stamina* 6, exserta; filamenta filiformia; antheræ lineares. *Fructus* 2- $2\frac{1}{2}$  lin. longus, perianthium æquans; stylus exsertus  $\frac{1}{2}$  lin. longus.

HUPEH: Fang, 8000-9000 ft. (*A. Henry*, 6834, 6854, 6868 A!). Mus. Brit.; Herb. Kew.

Allied to *J. sphenostemon*, Buchenau.

13. *Juncus pauciflorus*, *R. Br. Prodr.* p. 259; *Kunth, Enum. Pl.* iii. p. 320; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 238; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 116.

CHEKIANG: in running streams near Ningpo (*Oldham*, 76!); KIANGSI: Kiukiang (*Shearer*!); HUPEH: Ichang (*A. Henry*, 222, 592, 3653!); SZECHUEN: Mount Omei, 3400 ft. (*Faber*, 1137!); COREAN ARCHIPELAGO (*Oldham*, 895!), Tracey Island, in a marshy place near the beach (*Oldham*, 1067!). Herb. Kew.

Japan, New Caledonia, Australia, and New Zealand.

14. *Juncus Potanini*, *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 394, et xxix. (1900) p. 238.

KANSUH: by the Lumbu River and Jombunon River (*Potanin ex Buchenau*); SZECHUEN: Kzeruzo (*Potanin ex Buchenau*).

I have not seen an authentic example of this plant, but probably a specimen from the following locality should be referred to it:—

HUPEH: Kuei, on wooded precipices, 6000 ft. (*A. Henry*, 6169!). Herb. Kew.

15. *Juncus prismatocarpus*, *R. Br. Prodr.* p. 259; *Kunth, Enum. Pl.* iii. p. 337; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 310; *Hook. f. Fl. Brit. Ind.* vi. p. 395.

*Juncus Leschenaultii*, *J. Gay, in Laharpe, Monogr. Jonc.* p. 137; *Kunth, Enum. Pl.* iii. p. 336; *Benth. Fl. Hongk.* p. 380.

*Juncus sinensis*, *J. Gay, in Laharpe, Monogr. Jonc.* p. 137; *Kunth, Enum. Pl.* iii. p. 336; *Buchenau, in Engl. Bot. Jahrb.* xii. (1890) p. 313.

*Juncus prismatocarpus*, *R. Br.*, var. *Leschenaultii*, *Buchenau*, in *Engl. Bot. Jahrb.* xii. (1890) p. 311, & xxix. (1900) p. 238.

*Juncus lamprocarpus*, *Buchenau*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 238, non *Ehrh.*

SHANTUNG: at the foot of Chefoo Mountains (*Faber*, 75! *Fauvel*); FORMOSA (*Oldham*, 579! *Swinhoe*!); HUPEH: Patung district (*A. Henry*, 2471, 4902!), Ichang and neighbourhood (*A. Henry*, 3504, 3518, 3984!), Nanto (*A. Henry*, 3863!); YUNNAN (*Anderson*!); KWANGTUNG: Pakhoi (*Playfair*, 50!); HONGKONG (*Hance*, 599! *Faber*, 240! *Wright*, 554! *Wilford*, 269!); STRAITS OF COREA: Tsusima Island (*Wilford*, 809!); LUCHU ISLANDS (*Wright*, 329!). Mus. Brit.; Herb. Kew.

Japan, India, Australia, New Zealand.

16. *Juncus Przewalskii*, *Buchenau*, in *Engl. Bot. Jahrb.* xii. (1890) p. 401.

KANSUH: alpine heights near the river Tetung, a tributary of the Hoangho (*Przewalski* ex *Buchenau*).

17. *Juncus setchuenensis*, *Buchenau*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 238, name only.

SZETCHUEN: Nanchuan (*Bock von Rosthorn*, 2101, ex *Buchenau*).

18. *Juncus Thomsoni*, *Buchenau*, in *Bot. Zeit.* xxv. (1867) p. 148; et in *Engl. Bot. Jahrb.* xii. (1890) p. 390.

KANSUH: Tangut region (*Przewalski*!); SZETCHUEN: Tongolo (*Soulié*, 960!). Herb. Kew.

Himalaya.

This is treated as a form of *J. leucomelas*, *Royle*, in *Hook. f. Fl. Brit. Ind.* vi. p. 397, and it may be, but it seems constantly to differ in wanting the leafy prolongation to the lowest bract which is so characteristic of *J. leucomelas*, *Royle*.

19. *Juncus Hancockii*, *Hance*, in *Journ. Bot.* xvi. (1878) p. 111.

NORTH CHINA: Siao Wutaishan (*Hancock*). Mus. Brit.

Only one specimen of this is at present known, which is referred to *J. triglumis*, *Linn.*, by *Buchenau* in *Engl. Bot. Jahrb.* xii. (1890) p. 388; but it differs from that species in having a short sheathing leaf near the top of the stem, and when better known will probably prove distinct.

## 150. PALMÆ.

(By C. H. WRIGHT, A.L.S.)

1. **Areca Catechu**, *Willd. Sp. Pl.* iv. i. p. 594; *Lour. Fl. Cochinch.* p. 567; *Kunth, Enum. Pl.* iii. p. 184; *Mart. Nat. Hist. Palm.* iii. p. 169, t. 102; *Hook. f. Fl. Brit. Ind.* vi. p. 405; *Bretschn. Hist. Europ. Bot. Disc.* p. 11.

*Areca Cathecu*, *Linn. Sp. Pl.* ed. 1, p. 1189; *Burm. f. Fl. Ind.* p. 241.

*Areca hortensis*, *Lour. Fl. Cochinch.* p. 568 in obs.

SOUTH CHINA (rare, ex *Loureiro*); FORMOSA: Takow (*A. Henry*!); HAINAN (*A. Henry*, 8406!). Mus. Brit.; Herb. Kew. India and Malaya.

1. **Arenga Engleri**, *Becc. Malesia*, iii. p. 184; *Kew Bulletin*, 1895, p. 19.

FORMOSA: Tamsui (*Oldham*, 626, 627!), Kelung (*Ford*, 26!), Takow, Ape's Hill (*A. Henry*, 798!), Bankinsing (*A. Henry*, 798 A!). Mus. Brit.; Herb. Kew.

"This palm is called in Formosa *Ts'ung*, the name given in North [Formosa?] to *Trachycarpus Fortunei*."—*A. Henry*.

2. **Arenga saccharifera**, *Labill. in Mém. Inst. Par.* iv. p. 209; *Griff. Palms Brit. Ind.* p. 164, t. 235 A; *Mart. Nat. Hist. Palm.* iii. p. 191, t. 108; *Kunth, Enum. Pl.* iii. p. 197; *Hook. f. Fl. Brit. Ind.* vi. p. 421; *Engl. Bot. Jahrb.* vi. (1885) p. 52.

LUCHU ARCHIPELAGO (*Döderlein ex Engler*).

South-east Asia and Malaya.

1. **Caryota mitis**, *Lour. Fl. Cochinch.* p. 569; *Mart. Nat. Hist. Palm.* iii. p. 191; *Hook. f. Fl. Brit. Ind.* vi. p. 423.

*Caryota urens*, *Jacq. Fragm.* p. 20, t. 12. fig. 1, non *Linn.*

HAINAN: Kuanglang (*A. Henry*, 8373!). Herb. Kew.

2. **Caryota ochlandra**, *Hance, in Journ. Bot.* xvii. (1879) p. 176.

KWANGTUNG: Canton, and KWANGSI (*Hance*, 20332!). Mus. Brit.

1. **Phoenix Hanceana**, *Naudin, in Kerch., Palm.* p. 252, et *Man. Acclim.* p. 407; *Hance, in Journ. Bot.* xvii. (1879) p. 174.

*Phoenix acaulis?*, *Benth. Flor. Hongk.* p. 340; *Hance, in Ann. Sc. Nat., Bot.* 5<sup>me</sup> sér. v. (1866) p. 247, et *Journ. Linn. Soc., Bot.* xiii. (1872) p. 129; *Seemann, Bot. Voy. 'Herald,'* p. 416.

*Phoenix farinifera*, *Hance, in Journ. Bot.* vii. (1869) p. 15, et *Journ. Linn. Soc., Bot.* xiii. (1872) p. 129, non *Roxb.*

*Phoenix humilis*, *Royle*, var. *Hanceana*, *Becc. Malesia*, iii. pp. 348 et 392.

FORMOSA: Takow (*Swinhoe*, 17!), Taiwan (*Playfair*, 374!); KWANGTUNG: Macao (*Vachell*!); HONGKONG (*Wright*, 507! *Hance*, 1192! *Ford* in hb. *Forbes*, 608!); HAINAN (*A. Henry*, 8595!). *Mus. Brit.*; *Herb. Kew.*

1. **Licuala Fordiana**, *Becc. Malesia*, iii. p. 198.

KWANGTUNG: Lofaushan (*Ford*, 139! 22267 in hb. *Hance*!). *Mus. Brit.*; *Herb. Kew.*

1. **Livistona chinensis**, *Mart. Nat. Hist. Palm.* iii. p. 240, t. 146. figs. 1-3; *Walp. Ann.* v. p. 816; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 2; *Hance, in Journ. Bot.* xvii. (1879) p. 174; *Hook. f. Fl. Brit. Ind.* vi. p. 434; *Bull. Soc. Bot. France*, viii. (1861) p. 426.

*Livistona sinensis*, *Griff. Palms Brit. Ind.* p. 131, t. 226 D.

*Latania chinensis*, *Jacq. Fragm.* p. 16, t. 11. fig. 1; *Bretschn. Hist. Europ. Bot. Disc.* pp. 120 et 712.

SOUTH CHINA (ex *Martius*); HONGKONG: cultivated (*Hance*, 21013!). *Mus. Brit.*

Cultivated in India and Japan.

1. **Trachycarpus excelsus**, *H. Wendl. ex C. Gay, in Bull. Soc. Bot. France*, viii. (1861) p. 429; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 233.

*Trachycarpus Fortunei*, *H. Wendl. in Bull. Soc. Bot. France*, viii. (1861) p. 429; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 233.

*Chamærops excelsa*, *Thunb. Fl. Jap.* p. 130; *Mart. Nat. Hist. Palm.* iii. p. 251, t. 125; *Kunth, Enum. Pl.* iii. p. 250; *Walp. Ann.* v. p. 818; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 1.

*Chamærops Fortunei*, *Hook. in Bot. Mag.* t. 5221; *Debeaux, Fl. Shangh.* p. 59.

KIANGSU: Shanghai (*Montigny ex Gay*), Kiangnan (*ex Gay*);  
 CHEKIANG: Chusan (*Fortune*), Hangtcheoufou (*ex Gay*);  
 KIANGSI (*Staunton*!); FORMOSA: Tamsui (*Oldham*, 628!);  
 SZECHUEN: Chung Ching (*Mesny in hb. Hance*, 21707!);  
 COREAN ARCHIPELAGO: Tsusima (*Wilford*, 750!). Mus. Brit.;  
 Herb. Kew.

Japan.

This plant has been cultivated in many parts of China for a considerable time, and it is doubtful in what provinces it is really wild. C. Gay in the above-mentioned paper sums up the evidence without, however, coming to any definite conclusion.

2. *Trachycarpus khasianus*, *H. Wendl. ex C. Gay, in Bull. Soc. Bot. France*, viii. (1861) p. 429.

*Chamærops khasiana*, *Griff. in Calc. Journ. Nat. Hist.* v. (1845) p. 341; *Kurz, in Journ. Bot.* xi. (1873) p. 193; *Bot. Mag.* t. 7128; *Hook. f. Fl. Brit. Ind.* vi. p. 436.

YUNNAN: Momyen (*Anderson*!). Herb. Kew.

Eastern Bengal and Burmah.

This species is inserted here on the faith of Kurz's determination. The specimen at Kew is very imperfect.

1. *Rhapis flabelliformis*, *Ait. Hort. Kew.* ed. 1, iii. p. 473; *Mart. Nat. Hist. Palm.* iii. p. 253, t. 144; *Kunth, Enum. Pl.* iii. p. 251; *Jacq. Hort. Schænbr.* t. 316; *Bot. Mag.* t. 1371; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 2; *Benth. Fl. Hongk.* p. 340; *Bretschn. Hist. Europ. Bot. Disc.* p. 126.

*Rhapis Kwanwortsik*, *H. Wendl. Ind. Palm.* p. 34; *Seem. Bot. Voy. 'Herald,'* p. 416.

CHINA (*Macartney*! *Staunton*!); HONGKONG (*Hance*, 10164! *Lamont*! *Urquhart*! *Wilford*, 138!); LUCHU ARCHIPELAGO (*Wright*!). Mus. Brit.; Herb. Kew.

Japan.

2. *Rhapis humilis*, *Blume, Rumph.* ii. p. 54; *Mart. Nat. Hist. Palm.* iii. p. 254; *Walp. Ann.* v. p. 818.

CHINA (*ex Blume*).

3. *Rhapis major*, *Blume, Rumph.* ii. p. 55; *Mart. Nat. Hist. Palm.* iii. p. 254; *Walp. Ann.* v. p. 818.

CHINA (*ex Blume*).

1. **Calamus formosanus**, *Becc. in Records Bot. Survey India*, ii. (1902) p. 211.

FORMOSA: Kelung (*Oldham*, 629!), Bankinsing (*A. Henry*, 522, 522 A!), South Cape (*A. Henry*, 587!). Mus. Brit.; Herb. Kew.

2. **Calamus Henryanus**, *Becc. in Records Bot. Survey India*, ii. (1902) p. 199.

YUNNAN: Szemao Mountains (*A. Henry*, 12239!). Herb. Kew.

3. **Calamus Margaritæ**, *Hance, in Journ. Bot.* xii. (1874) p. 266.

FORMOSA: Bankinsing (*A. Henry*, 521, 521 A!); HONGKONG (*Mrs. Dods* in hb. *Hance*, 18407!), Happy Valley (*Ford*!), Washerman's Ravine (*Urquhart*!). Mus. Brit.; Herb. Kew.

4. **Calamus tetradactylus**, *Hance, in Journ. Bot.* xii. (1875) p. 289.

HONGKONG: Wongneichung (*Dods* in hb. *Hance*, 18979! *Ford*!), Victoria (*Lamont*!), Little Hongkong (*Lamont*!); HAINAN: Hoihere (*A. Henry*, 8213!). Mus. Brit.; Herb. Kew. Saigon.

5. **Calamus thysanolepis**, *Hance, in Journ. Bot.* xi. (1874) p. 265.

HONGKONG (*Dods* in hb. *Hance*, 18373! *Lamont*!). Mus. Brit.; Herb. Kew.

6. **Calamus Walkeri**, *Hance, in Journ. Bot.* xi. (1874) p. 266.

HONGKONG (*Walker* in hb. *Hance*, 18225!). Mus. Brit.; Herb. Kew.

1. **Cocos nucifera**, *Linn. Sp. Pl.* ed. 1, p. 1188; *Lour. Fl. Cochinch.* p. 566; *Seem. Bot. Voy. 'Herald,'* p. 416; *Williams, The Middle Kingdom*, ed. 2, i. p. 360.

HONGKONG: planted on the sea-shore (ex *Seemann*); HAINAN: cultivated (ex *Loureiro*).

All tropical shores.

## 151. PANDANACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Pandanus furcatus*, *Roxb. Hort. Beng.* p. 71, *et Fl. Ind.* iii. p. 744; *Hook. f. Fl. Brit. Ind.* vi. p. 484; *Balf. f. in Journ. Linn. Soc.* xvii. (1878) p. 47; *Warburg, in Engl. Pflanzenr., Pandanaceæ*, p. 75.

*Pandanus urophyllus*, *Hance, in Journ. Bot.* xii. (1875) p. 68.

*Pandani* sp., *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 129.

KWANGTUNG: Canton, Lienchan (*Lamont*!); HONGKONG (*Harland and Lamont* in hb. *Hance*, 18394!). Mus. Brit.

India and Malaya.

2. *Pandanus odoratissimus*, *Linn. f. Suppl.* p. 424; *Lour. Fl. Cochinch.* p. 603; *Balf. f. in Journ. Linn. Soc.* xvii. (1878) p. 54; *Hance, in Journ. Linn. Soc.* xiii. (1872) p. 129; *Warburg, in Engl. Pflanzenr., Pandanaceæ*, p. 46.

*Pandanus fascicularis*, *Lam. Encycl.* i. p. 372 (excl. var.  $\beta$ ); *Seem. Bot. Voy. 'Herald,'* p. 416; *Hook. f. Fl. Brit. Ind.* vi. p. 485.

FORMOSA: Tamsui (*Oldham*, 631!); KWANGTUNG: Macao (*Vachell*!); HONGKONG (*Hance*, 715!); HAINAN: Hoihow (*A. Henry*, 8290!). Mus. Brit.; Herb. Kew.

Indian Archipelago and North Australia to Sandwich Islands.

1. *Freycinetia formosana*, *Hemsl. in Kew Bulletin*, 1896, p. 166; *Warburg, in Engl. Pflanzenr., Pandanaceæ*, p. 41.

FORMOSA: without locality (*Oldham*, 630!), Kelung (*Ford*, 45!). Herb. Kew.

## 152. TYPHACEÆ.

(By N. E. BROWN, A.L.S.)

1. *Typha angustata*, *Bory & Chaub. Expéd. Sc. de Morée*, ii. pt. 2, p. 338; *Boiss. Fl. Orient.* v. p. 50; *Rohrb. in Verhandl. Bot. Ver. Prov. Brandenb.* xi. (1869) p. 87, figs. 4-6; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 159, t. 4. fig. 6, t. 5. fig. 1; *Hook. f. Fl. Brit. Ind.* vi. p. 489.

CHINA (*Karolkow* ex *Kronfeld*); CHIHLI: near Peking (*Bretschneider, Fischer* ex *Kronfeld*).

Throughout the warmer parts of Asia, South-eastern Europe, and the north-eastern part of Africa.

2. *Typha angustifolia*, *Linn. Sp. Pl.* ed. 1, p. 971; *Bunge, Enum. Pl. Chin. Bor.* p. 67; *Syme, English Bot.* ed. 3, ix. p. 4, t. 1886; *Reichb. Ic. Fl. Germ.* ix. p. 2, t. 321; *Rohrb. in Verhandl. Bot. Ver. Prov. Brandenb.* xi. (1869) p. 81, fig. 3; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 150, t. 5. fig. 2; *Debeaux, Fl. Tchef.* p. 132; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 260.

CHIHLI: near Kantai (ex *Bunge*); SHANTUNG: Chefoo and vicinity (ex *Debeaux, Fauvel* ex *Franchet*); FORMOSA: Takow (*A. Henry*, 1815!). *Herb. Kew.*

Widely dispersed in temperate regions.

3. *Typha latifolia*, *Linn. Sp. Pl.* ed. 1, p. 971; *Kunth, Enum. Pl.* iii. p. 90; *Syme, English Bot.* ed. 3, ix. p. 2, t. 1385; *Reichb. Ic. Fl. Germ.* ix. p. 2, t. 323; *Rohrh. in Verhandl. Bot. Ver. Prov. Brandenb.* xi. (1869) p. 75, fig. 1; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 176, t. 5. fig. 11; *Lour. Fl. Cochinch.* p. 552?; *Bretschn. Early Res.* p. 181.

CHINA (ex *Loureiro*). Frequent in North China (ex *Bretschneider*).

Widely distributed in north temperate and subtropical regions.

4. *Typha Laxmanni*, *Lepechin, in Nov. Act. Acad. Petrop.* xii. (1801) pp. 84, 335, t. 4; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 167, t. 4. fig. 3, et t. 5. fig. 15; *Hook. f. Fl. Brit. Ind.* vi. p. 489.

CHIHLI: near Peking (*Bretschneider*, 771!). *Herb. Kew.*

Widely dispersed in temperate Asia.

5. *Typha Martini*, *Jord.*, var. *Davidiana*, *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 149.

*Typha Laxmanni*, *Franch. Pl. David.* p. 312, non *Lepechin* (ex *Kronfeld*).

CHIHLI: Jehol (*David*, 1860, ex *Franchet*).

6. *Typha minima*, *Funck, in Hoppe, Bot. Taschenb.* (1794) pp. 118, 187; *Hoffm. Deutschl. Fl.* ed. 2, ii. p. 251; *Braune, Salz. Fl.* ii. p. 624; *Willd. Sp. Pl.* iv. p. 198; *Franch. Pl. David.* p. 313; *Reichb. Ic. Fl. Germ.* ix. p. 1, t. 319. figs. 742, 743; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889,



p. 144, t. 4. fig. 2, & t. 5. fig. 7, and incl. var. *Regelii*, *Kronf.*  
p. 147, t. 4. fig. 4, & t. 5. fig. 9.

*Typha Laxmanni*, *Ledeb. Fl. Ross.* iv. p. 3; *Rohrb. in Verhandl. Bot. Ver. Prov. Brandenb.* xi. (1869) p. 91, non *Lepechin*.

*Typha minor*, *Smith, Fl. Brit.* iii. p. 960, & in *Engl. Bot.* ed. 1, t. 1457; *Bunge, Enum. Pl. Chin. Bor.* p. 67.

*Typha Bungeana*, *Presl, Epimel. Bot.* p. 239; *Walp. Ann.* iii. p. 495; *Kurz, in Journ. Bot.* v. (1867) p. 96.

CHIHLI: Jehol (*David, 2723, ex Franchet*), near Kantai (ex *Bunge*); KANSUH: Tangut (*Przewalski!*); CHINA (*Karolkow & Krause ex Kronfeld*). *Herb. Kew.*

Temperate Asia and Europe.

7. *Typha orientalis*, *Presl, Epimel. Bot.* p. 239; *Walp. Ann.* iii. p. 495; *Kronfeld, in Verhandl. zool.-bot. Gesellsch. Wien*, 1889, p. 175, t. 4. fig. 1, & t. 5. fig. 14.

*Typha Shuttleworthii*, *Koch & Sond.*, var. *orientalis*, *Rohrb. in Verhandl. Bot. Ver. Prov. Brandenb.* xi. (1869) p. 80.

*Typha latifolia*, *Bretschn. Early Res.* p. 181, non *Linn.* (ex *Kronfeld*).

*Typha japonica*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 160; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 11.

NORTH CHINA (ex *Bretschneider*); CHIHLI: north-west of Peking (*Bretschneider, Moellendorf!*); HUPEH: Nanto (*A. Henry, 2105!*), Patung (*A. Henry, 5118, 7396!*), Hsingshan (*A. Henry, 6501!*). *Mus. Brit.*; *Herb. Kew.*

Japan.

### 153. AROIDEÆ.

(By N. E. BROWN, A.L.S.)

1. *Pinellia cordata*, *N. E. Br.*, n. sp.

*Tuber* globosum,  $\frac{1}{2}$ – $\frac{3}{4}$  poll. diam. *Folia* subhastato-cordata, acuminata,  $2\frac{1}{2}$ – $3\frac{1}{2}$  poll. longa,  $1\frac{1}{4}$ – $1\frac{3}{4}$  poll. lata, petiolo 5–6 poll. longo, basi et apice tuberifero. *Scapus*  $1\frac{1}{2}$ –2 poll. longus, quam petiolus dimidio brevior. *Spathæ* tubus ovoideus,  $\frac{1}{4}$  poll. longus,  $\frac{1}{8}$  poll. latus; lamina  $\frac{3}{4}$ –1 poll. longa, cucullata, obtusa, marginibus recurvis? *Spadicis appendix* filiformis, sigmoideo-erecta,  $2\frac{1}{2}$  poll. longa.

CHEKIANG: Foot of Tientai Mountain (*Faber, 82!*); KIANGSI: Kiukiang (*Shearer!*). *Herb. Kew.*

The leaves of this species somewhat resemble the cordate ones of the juvenile state of *P. tuberifera*, Tenore, but are not so constricted above the basal lobes. It is readily distinguished by its short scape and broader and more hooded limb of the smaller spathe, which has a much shorter and more ovoid tube; the bulbils on the petioles are also smaller and more pointed than in *P. tuberifera*, Tenore.

2. *Pinellia integrifolia*, *N. E. Br. in Hook. Ic. Plant.* xix. t. 1875.

HUPEH: Ichang (*A. Henry*, 663, 4323!). Mus. Brit.; Herb. Kew.

3. *Pinellia pedatisecta*, *Schott, in Oest. Bot. Wochenbl.* vii. 1857, p. 341; *Schott, Prodr. Aroid.* p. 20.

*Pinellia tuberifera*, Tenore, var. *pedatisecta*, *Engl. in DC. Monogr. Phan.* ii. p. 567.

*Pinellia Wawræ*, *Engl. in DC. Monogr. Phan.* ii. p. 568; *Franch. Pl. David.* i. p. 313.

CHIHLI: Peking, Nankow Pass (*Wawra*, 1037, ex *Engler*); Sang-yu (*David*, 2236, ex *Franchet*); SZECHUEN: Yangtze-kiang (*Faber*, 979!). Herb. Kew.

4. *Pinellia tripartita*, *Schott, Synop. Aroid.* p. 5; *Prodr. Aroid.* p. 20; *Engl. in DC. Monogr. Phan.* ii. p. 566; *N. E. Brown, in Journ. Linn. Soc., Bot.* xviii. (1881) p. 246.

*Atherurus tripartitus*, *Blume, Rumphia*, i. p. 137, tt. 31 et 37 F; *Kunth, Enum. Pl.* iii. p. 54.

*Arisæma tripartitum*, *Engl. in DC. Monogr. Phan.* ii. p. 538.

LUCHU ARCHIPELAGO (*C. Wright*, 319!). Herb. Kew.  
Japan.

5. *Pinellia tuberifera*, Tenore, in "Atti IIIa. riun. Scienz. Ital." p. 522; *Sem. Cat. Hort. Neap.* 1830," et in *Atti Reale Accad. Scienze*, iv. (1839) p. 70, t. x.; *Schott, Prodr. Aroid.* p. 20; *Gen. Aroid.* t. 4; *Engl. in DC. Monogr. Phan.* ii. p. 566; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 88; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 260; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 117.

*Arum macrourum*, *Bunge, Enum. Pl. Chin. Bor.* p. 67.

*Arum ternatum*, *Thunb. Fl. Jap.* p. 233.

*Atherurus ternatus*, *Blume, Rumphia*, i. p. 136; *Kunth, Enum. Pl.* iii. p. 54.

CHIHLI: Peking (*Williams! Bushell! Bullock, 94! Bunge*), Nankow Pass (*Wawra, 987, ex Engler*); HUPEH: Ichang (*A. Henry, 3646!*); SHANTUNG: Chefoo (*Fauvel ex Franchet*); PUTOY ISLAND (*Wright, 508*); COREA: Soul (*Kalinowsky ex Palibin*). Mus. Brit.; Herb. Kew.

Japan.

Var. *subpandurata*, *Engl. Bot. Jahrb.* i. (1881) p. 488; *Franch. Pl. David.* i. p. 313.

CHIHLI: Peking (*David, 2330, ex Engler*). Herb. Mus. Paris.

I have not seen a specimen of this variety, but from the description it would appear to be merely an abnormal form.

1. *Pistia Stratiotes*, *Linn. Sp. Pl.* ed. 1, p. 963; *Kunth, Enum. Pl.* iii. p. 8; *Engl. in DC. Monogr. Phan.* ii. p. 634; *Hook. f. Fl. Brit. Ind.* vi. p. 497.

KWANGTUNG: Whampoa (*Hance, 6065!*). Mus. Brit.; Herb. Kew.

Throughout the tropics.

1. *Arisæma amurense*, *Maxim. Prim. Fl. Amur.* p. 264; *Engl. in DC. Monogr. Phan.* ii. p. 549.

*Arisæma amurense*, *Maxim.*, var. *robustum*, *Engl. in DC. Monogr. Phan.* ii. p. 550; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 117.

SHINGKING: Laoling (*Webster, 31!*); Black Bear Valley, Toonghwahien (*Ross, 32!*); COREA: Port Hamilton (*Wilford!*). Herb. Kew.

Mandshuria, Amur.

Var. *Sazensoo*, *Engl. in DC. Monogr. Phan.* ii. p. 550.

*Arisæma japonicum*, var. *Sazensoo*, *Blume, Rumphia*, i. p. 107?

*Arisæma serratum*, *Schott*, var., *Baker et S. Moore, in Journ. Linn. Soc. Bot.* xvii. (1879) p. 387.

SHINGKING: Chienshan and south of Hingjing (*Ross, 250!*); CHEKIANG: Tientai Mountain, 3000 ft. (*Faber, 84!*); KIANGSI: Chinkiang and Kiukiang (*Maries!*); HUPEH: South Patung (*A. Henry, 5394!*), Nanto (*A. Henry, 5394 B, 4571!* and probably 3030). Mus. Brit.; Herb. Kew.

Var. *magnidens*, *N. E. Br.*, n. var.; foliolis grosse dentatis, dentibus  $\frac{1}{4}$ – $\frac{1}{2}$  poll. longis, cum minoribus interjectis.

HUPEH (*A. Henry*, without locality, 6363 A! 6730 A!), South Patang (*A. Henry*, 5393!). Mus. Brit.; Herb. Kew.

The variety *Sazensoo* is very variable in the size and form of the leaflets, which sometimes arise almost directly from the apex of the petiole, at others have long stalks as in the typical form. The more southern forms have two leaves to each plant, with leaflets attaining 9 inches in length and 6 inches in breadth, and occasionally six or seven in number instead of five. The variety *magnidens* is a very striking form, but I can find nothing to distinguish it except the remarkably large teeth of the leaflets.

2. *Arisæma asperatum*, *N. E. Br.*, n. sp.

*Petiolus* 12 poll. longus, toto processibus brevibus dense obtectus; lamina tripartita, segmentis omnibus sessilibus, laterilibus 10–10 $\frac{1}{2}$  poll. longis, 9 poll. latis, subrhomboideo-ellipticis, acutis, intermedio paullo brevior, latissime obovato, apice retuso apiculato, basi cuneato, costis subtus asperatis. *Scapus* 10 poll. longus, ut in petiolo processibus brevibus asperatus. *Spathæ* tubus 2–2 $\frac{1}{4}$  poll. longus, cylindricus; lamina 4 $\frac{1}{2}$  poll. longa, abrupte procurva (vel decurva?), oblanceolata, acuminata. *Spadicis appendix* 2 $\frac{1}{2}$  poll. longa, anguste cylindrica, obtusa,  $\frac{1}{10}$ – $\frac{1}{8}$  poll. crassa, basi abrupte incrassata, truncata, apice e fauce longe procurva.

HUPEH: Patung district, cultivated, the tuber used as a drug (*A. Henry*, 3776!). Herb. Kew.

More nearly allied to *A. verrucosum*, Schott, than to any other species, but perfectly distinct in its inflorescence and in the petiole being much more densely covered with short cylindric or conical processes.

3. *Arisæma Bockii*, *Engl. in Engl. Bot. Jahrb.* xxix. (1900) p. 235.

SZECHUEN: Nanchuan (*Bock von Rosthorn*, 882, ex *Engler*).

4. *Arisæma consanguineum*, *Schott, in Bonplandia*, vii. (1859) p. 27, et *Prodr. Aroid.* p. 52; *Hook. f. Fl. Brit. Ind.* vi. p. 505.

*Arisæma erubescens*, Schott, var. *consanguineum*, *Engl. in. DC. Monogr. Phan.* ii. p. 558.

*Arisæma Giraldii*, Baroni, in *Bull. Soc. Tosc. Ort.* xviii. (1893) p. 356, t. 12; et *Bull. Soc. Bot. Ital.* (1893) p. 515; *Engl. in Engl. Jahrb.* xxix. (1900) p. 236.

*Arisæma Tatarinowii*, Schott, in *Bonplandia*, vii. (1859) p. 27; et *Prodr. Aroid.* p. 53; *Engl. in DC. Monogr. Phan.* ii. p. 559, et in *Engl. Bot. Jahrb.* xxix. (1900) p. 236, incl. var. *latisecta*, *Engl.*

CHIHLI: Peking (*Tatarinow ex Schott*); HUPEH: Patung (*A. Henry*, 3780, 4653, 5303, 5303 A, 5397, 5379 A & B!); HONGKONG: Lantao Island, 2000–3000 feet (*Westland*, 93!); SZECHUEN: Nanchuan (*Bock von Rosthorn*, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, ex *Engler*), Changlingkang (*Bock von Rosthorn*, 644, 650, ex *Engler*), Tsakulao (*Bock von Rosthorn*, 2562, ex *Engler*); SHENSI: Tsinlingshan (*Giraldi!*). *Mus. Brit.*; *Herb. Kew.*

Sikkim, Khasia, Kumaon.

5. *Arisæma cordatum*, *N. E. Br.*, n. sp.

*Petiolus* 18 poll. longus, basi breviter vaginatus, brunneus, atroviride maculatus, lamina pedatipartita, segmentis 5–6½ poll. longis, 2–2½ poll. latis, oblongo-ellipticis utrinque acutis, sessilibus. *Scapus* 2–4 poll. longus. *Spathæ* tubus 1½–1¼ poll. longus, cylindricus; lamina ampla, explanata triangulari-cordata, acuta, 2–2½ poll. longa et lata, auriculis rotundatis. *Spadix* unisexualis, appendice in filum tenuem quam spatha duplo longiorem attenuata.

KWANGTUNG: Tai Mo Mountain opposite Hongkong, 2300 feet (*Westland*, 92!). *Herb. Kew.*

A very distinct species, unlike any other known to me, readily distinguished by the triangular-cordate limb of the spathe, when flattened out, and long filiform appendix. In the living state the form of the spathe would probably be something like that of *A. nepenthoides*, Mart., but with proportionately broader auricles. The scapes may be longer than I have described them, as they are detached, and there is no tuber.

6. *Arisæma Franchetianum*, *Engl. in Engl. Bot. Jahrb.* i. (1881) p. 487, et xxix. (1900) p. 236; *Franch. Pl. David.* ii. p. 139.

SZECHUEN: Moupin (*David ex Engler*).

7. *Arisæma heterophyllum*, *Blume, Rumphia*, i. p. 110; *Kunth, Enum. Pl.* iii. p. 20; *Schott, Prodr. Aroid.* p. 35; *N. E. Brown, in Journ. Linn. Soc., Bot.* xviii. (1881) p. 250.

*Arisæma Thunbergii*, *Blume*, var. *heterophyllum*, *Engl. in DC. Monogr. Phan.* ii. (1882) p. 546.

*Arisæma curvatum*, *Hance, in Journ. Bot.* xviii. (1880) p. 302, non *Kunth*.

KIANGSU: Fengwangshan near Shanghai (*Forbes* in hb. *Hance*, 8139!); CHEKIANG: Tientai Mountain, 3000 feet (*Faber*, 85!); Kiangsi: Kiukiang (*Shearer*!); HUPEH: Ichang (*A. Henry*, 212, 3574!), South Patung (*A. Henry*, 5508!); COREA: Soul Mountains (*Carles*!), West coast (*Wykeham Perry*, 50!); COREAN ARCHIPELAGO: Surly Island (*Oldham*, 817!). *Mus. Brit.*; *Herb. Kew.*

Japan, according to *Blume*, but I have seen no Japanese specimen.

8. *Arisæma japonicum*, *Blume, Rumphia*, i. p. 106; *Kunth, Enum.* iii. p. 19; *Schott, Prodr. Aroid.* p. 40; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 5; *Engl. in DC. Monogr. Phan.* ii. p. 549.

*Arisæma latisectum*, *Blume, Rumphia*, i. p. 110.

*Arum dracunculus*, *Thunb. Fl. Jap.* p. 233, non *Linn.*

CHEKIANG: Tientai Mountain (*Faber*, 86!); FORMOSA: Kelung (*Oldham*, 634? leaf only); HUPEH: Patung (*A. Henry*, 5371!); SZECHUEN: South Wushan (*A. Henry*, 5536!). *Herb. Kew.*

Japan.

9. *Arisæma lobatum*, *Engl. Bot. Jahrb.* i. (1881) p. 487; *Franch. Pl. David.* ii. p. 138.

SZECHUEN: Moupin (*David ex Engler*).

Var. *Rosthornianum*, *Engl. in Engl. Bot. Jahrb.* xxix. (1900) p. 235.

SZECHUEN: Nanchuan (*Bock von Rosthorn*, 651, 1237, ex *Engler*).

10. *Arisæma parvum*, *N. E. Brown, in Journ. Linn. Soc., Bot.* xxix. (1892) p. 320.

SZECHUEN: near Tachienlu (*Pratt*, 841!). *Herb. Kew.*

11. *Arisæma penicillatum*, *N. E. Brown*, in *Journ. Linn. Soc., Bot.* xviii. (1881) p. 248, t. 5.

*Arisæma laminatum*, *Benth. Fl. Hongk.* p. 342, non *Blume*.

HONGKONG (*Harland*, 1026! *Hance*, 1238!), top of Ridge (*Urquhart*, 182!), Victoria Peak (*Ford*, 1! *Lamont*!). Mus. Brit.; Herb. Kew.

12. *Arisæma pictum*, *N. E. Brown*, in *Journ. Linn. Soc., Bot.* xxix. (1892) p. 321.

SZECHUEN: near Tachienlu (*Pratt*, 61!). Herb. Kew.

Probably specimens collected by Dr. Henry in the province of HUPEH (5381 A & 5381 B) and in the province of SZECHUEN at S. Wushan (5381 C) belong to this species, but the tuber bears two leaves, the leaflets are all much broader, the middle one being elliptic-rhomboid and nearly as broad as long, and the spathe appears to have a very short tail. Henry's 5381, however, from the province of HUPEH, has the middle leaflet oblong in form and more like that of *A. pictum*, but the lateral leaflets are stalked. All these may be forms of one variable species, but more material with perfect spathes are needed to decide. Mus. Brit.; Herb. Kew.

13. *Arisæma ringens*, *Schott, Melet.* p. 17; *Prodr. Aroid.* p. 31; *Kunth, Enum. Pl.* iii. p. 17; *Gartenfl.* 1861, p. 1, t. 313; *Flore des Serres*, xii. (1857) p. 167, tt. 1269, 1270; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 4; *Hance, in Journ. Bot.* xvii. (1879) p. 15; *Engl. in DC. Monogr. Phan.* ii. p. 534.

*Arisæma præcox*, *De Vriese, Hortus Spurnbergensis*, ex *K. Koch*, in *Allgem. Gartenzeit.* 1857, p. 87; *Schott, Prodr. Aroid.* p. 32; *Bot. Mag.* t. 5267; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 4.

*Arisæma Sieboldii*, *De Vriese*, ex *K. Koch*, in *Allgem. Gartenzeit.* 1857, p. 85.

*Arisæma ringens*, *Schott*, var. *Sieboldii*, *Engl. in DC. Monogr. Phan.* ii. p. 534; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 117.

*Arisæma ringens*, *Schott*, var. *præcox*, *Engl. in DC. Monogr. Phan.* ii. p. 535.

KIANGSU: Gutzlaff Island, near Shanghai (*Herb. Hongkong Bot. Gard.* 101!); FORMOSA (*Oldham*, 633! & 118! [hb. *Hance*, 11066!], *Swinhoe*! *Hancock*, 10, a form with narrower leaflets!); COREA: Port Hamilton (*Wilford*, 662!); COREAN ARCHIPELAGO (*Oldham*, 816!); LUCHU ARCHIPELAGO (*Wright*, 314!). Mus. Brit.; Herb. Kew.

Japan.

14. *Arisæma Thunbergii*, *Blume, Rumphia*, i. p. 105; *Kunth, Enum. Pl.* iii. p. 19; *Schott, Prodr. Aroid.* p. 38; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 4; *Engl. in DC. Monogr. Phan.* ii. p. 546, et in *Engl. Bot. Jahrb.* xxix. (1900) p. 236.

*Arum dracontium*, *Thunb. Fl. Jap.* p. 233, non *Linn.*

HUPEH: Chienshih, South Patung (*A. Henry*, 5370, 5370 A!); COREA: Port Hamilton (*Wilford*, 660!); COREAN ARCHIPELAGO: Tsusima Island (*Wilford*, 660!). *Mus. Brit.*; *Herb. Kew.*

Japan.

15. *Arisæma pentaphyllum*, *Schott, Melet.* p. 17, et *Prodr. Aroid.* p. 59; *Blume, Rumphia*, i. p. 109; *Kunth, Enum. Pl.* iii. p. 20; *Engl. in DC. Monogr. Phan.* ii. p. 560.

*Arum pentaphyllum*, *Linn. Sp. Pl.* ed. 1, p. 964; *Lour. Fl. Coch.* p. 533.

CHINA (ex *Loureiro*).

This is altogether a doubtful plant, and is only known from the figure given by Zanoni in his 'Istoria Botanica,' p. 170, t. 78 (1675), under the name "Rumphal pianta orientale," which has been copied in later works by other authors. Zanoni gives India as its habitat, and Louriero's Chinese plant may be different. The plant figured by Zanoni is certainly not an *Arisæma*, but probably some species of *Typhonium*, badly drawn.

1. *Typhonium divaricatum*, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) p. 367 [*Herb. Timor.* p. 39]; *Blume, Rumphia*, i. p. 130, t. 36 a; *Kunth, Enum. Pl.* iii. pp. 26 et 581; *Schott, Aroid.* i. p. 12, t. 18, *Gen. Aroid.* t. 17, et *Prodr. Aroid.* p. 106; *Somoku Zusetsu*, ed. 2, xix. t. 4; *Seem. Bot. Voy. 'Herald,'* p. 417; *Engl. in DC. Monogr. Phan.* ii. p. 611; *Hook. f. Fl. Brit. Ind.* vi. p. 510.

*Arum divaricatum*, *Linn. Sp. Pl.* ed. 1, p. 966; *Roxb. Fl. Ind.* iii. p. 503; *Wight, Ic. Pl. Or.* iii. pt. 1, p. 6, t. 790; *Lodd. Bot. Cab.* t. 516; *Benth. Fl. Hongk.* p. 342.

*Arum trilobatum*, *Curt. Bot. Mag.* tt. 339 et 2324, non *Linn.*

*Arum diversifolium*, *Blume, Cat. Gewass. Buitenz.* p. 102.

CHINA (*Hance* ex *Seemann*); KWANGTUNG: Tingushan, Canton (*Lamont*, 1076!); HONGKONG (*Champion*, 289! *Hance*!). *Mus. Brit.*; *Herb. Kew.*

Ceylon, Malay Archipelago, and Japan, according to authors; but I have seen no native specimens of this plant except those from



Hongkong above quoted ; all the others I have seen are cultivated specimens. Possibly *T. Roxburghii*, Schott (*T. javanicum*, Miq.), or *T. Motleyanum*, Schott, may have been mistaken for this species, as they are very similar, but the neuter organs are reflexed or decurved at their tips, whilst in *T. divaricatum* they are erect.

2. **Typhonium giganteum**, *Engl. in Bot. Jahrb.* iv. (1883) p. 66, t. 1.

*Typhonium giganteum*, *Engl.*, var. *Giraldii*, *Baroni*, in *Nuovo Giorn. Bot. Ital.* ser. 2, iv. (1897) p. 189, t. 6 ; *Sprenger*, in *Gard. Chron.* ser. 3, xxxii. (1902) p. 150, fig. 51.

CHIHLI : Peking Mountains (*Skatschkow ex Engler, Bretschneider*, 132 !); HUPEH : Hsingshan, 4500 feet (*A. Henry*, 6537 !); SHENSI (*Giraldi* !). Mus. Brit. ; Herb. Kew.

1. **Amorphophallus campanulatus**, *Decne. in Nouv. Ann. Mus. Par.* iii. (1834) p. 366 [*Herb. Timor.* p. 38] ; *Blume, Rumphia*, i. p. 139, excl. tt. 32, 33, et descriptione ; *Wight, Ic.* iii. pt. 1, p. 5, t. 785 ; *Engl. in DC. Monogr. Phan.* ii. p. 309, excl. planta javanica ; *Hance, in Journ. Bot.* xvii. (1879) p. 15 ; *Hook. f. Fl. Brit. Ind.* vi. p. 513.

*Amorphophallus virosus*, *N. E. Brown*, in *Gard. Chron.* ser. 2, xxiii. (1885) p. 759 ; *Bot. Mag.* t. 6978.

*Arum campanulatum*, *Roxb. Fl. Ind.* iii. p. 509.

SOUTH CHINA (*Hance*, 17601 !). Mus. Brit.

Tropical Asia.

2. **Amorphophallus Henryi**, *N. E. Brown*, n. sp.

*Folia*  $1\frac{1}{2}$ –2 ped. diam., glabra ; *petiolus* levis ; *segmenta terminalia*,  $3\frac{1}{2}$ –5 poll. longa, 1 poll. lata, lanceolata, acuminata, cetera minora. *Pedunculus* 2–4 poll. longus, levis. *Spatha* 4–5 poll. longa, inferne convoluta, intus basi verrucosa ; *lamina* late ovata, acuta, atropurpurea. *Spadix* sessilis, quam *spatha* duplo longior ; *pars feminea* 6–10 lin. longa, 4–6 lin. crassa, cylindrica, *mascula*  $\frac{3}{4}$ –1 poll. longa, obconica ; *appendix* 5–7 poll. longa, fusiformis, acuta. *Ovarium* globosum, 2-loculare ; *stylus*  $\frac{1}{2}$ – $\frac{2}{3}$  lin. longus ; *stigma* magnum, capitatum.

FORMOSA : Takow, on Ape's Hill in rocky situations (*A. Henry*, 776 !). Herb. Kew.

3. **Amorphophallus hirtus**, *N. E. Brown*, n. sp.

*Folii* immaturi glabri *segmenta* lineari-lanceolata, attenuato-

acuminata, petiolo levi. *Pedunculus* 15 poll. vel ultra longus, levis. *Spatha* 6 poll. longa, glabra, inferne convoluta, intus basi verrucosa; lamina ovata, subacuta. *Spadix* sessilis, quam spatha duplo longior, inter partes masculæ et femineæ pilis paucis instructus, pars feminea  $\frac{3}{4}$ -1 poll. longa,  $\frac{1}{2}$  poll. crassa, mascula 1-1 $\frac{1}{4}$  poll. longa, basi angustata; appendix 7-10 poll. longa,  $\frac{3}{4}$ -1 poll. crassa, tereto-attenuata, pilis longis flexuosis laxè oblecta. *Ovarium* ellipsoideum vel subglobosum, 2-loculare; stylus  $\frac{1}{2}$  lin. longus, crassus; stigma 2-lobum.

FORMOSA: Takow, on Ape's Hill (*A. Henry*, 1914!). Herb. Kew.

This differs from all the other species of *Amorphophallus* by the presence of hairs upon the appendix, resembling *Helicodicerus crinitus*, Schott, in this respect, and the few that occur between the male and female parts of the spadix represent the only approach to neuter organs hitherto found in the genus.

4. *Amorphophallus Konjac*, *K. Koch*, in *Berlin. Allgem. Gartenz.* 1858, pp. 166, 411.

*Amorphophallus Rivieri*, "*Durieu*, in *Cat. des graines du jard. Bordeaux*, 1869, p. 12"; *Rev. Hort.* 1870, p. 573, figs. 75-76; 1872, p. 19; 1873, pp. 305 & 344; *Gard. Chron.* 1873, p. 610, figs. 117-119; *Fl. des Serres*, xix. (1873) p. 43, tt. 1958-1961; *Engl. in DC. Monogr. Phan.* ii. p. 312; *Illustr. Hort.* xix. (1872) p. 246; *Monats. Ver. Beförd. Gartenb. Berlin*, 1880, p. 187, t. 7.

*Brachyspatha Konjac*, *K. Koch*, in *Berlin. Allgem. Gartenz.* 1858, pp. 166, 412.

*Conophallus Konjac*, *Schott*, ex *Miq. Annal. Mus. Bot. Lugd.-Bat.* ii. p. 202; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 7.

*Arisæma Konjac*, *Siebold*, ex *K. Koch*, in *Berlin. Allgem. Gartenz.* 1858, pp. 166, 412.

*Dracontium polyphyllum*, *Thunb. Fl. Jap.* p. 234, non *Linn.*

*Proteinophallus Rivieri*, *Hook. f. in Bot. Mag.* t. 6195.

*Tapeinophallus Rivieri*, *Baill. Dict. Bot.* i. plate issued in part 5, June 1877, et iv. p. 153.

*Hydrosme Rivieri*, *Engl. Bot. Jahrb.* i. (1881) p. 188.

SZECHUEN: South Wusban, cultivated (*A. Henry*, 5544!). Herb. Kew.

Cochinchina, Japan (cultivated).

Probably leafy specimens from Ichang (*Henry*, 4223), distributed from Kew, also belong here. Dr. Henry states of it, that "The same plant is cultivated by the name of *Mo-yii*; the root-stock is deprived of its acrid qualities by washing with water, and

is made into what is sold as *Mo-yii tou-fu*. *Tou-fu* is a term applied to cheesy-looking stuff extracted in China from beans (Soja)."

5. *Amorphophallus variabilis*, *Blume, Rumphia*, i. p. 146, tt. 35 & 37 H; *Kunth, Enum. Pl.* iii. p. 33; *Benth. Fl. Hongk.* p. 343; *Engl. in DC. Monogr. Phan.* ii. p. 313.

*Brachyspatha variabilis*, *Schott, Synop. Aroid.* p. 35, *Gen. Aroid.* t. 29, et *Prodr. Aroid.* p. 127; *Miq. Fl. Ind. Bat.* iii. p. 200; *Gard. Chron.* ser. 2, vi. (1876) p. 680, figs. 129, 130.

KWANGTUNG: Taimoshan (*Westland* in hb. *Hance*, 22422!); HONGKONG (*Harland*, 300!), Mount Gough (*Ford*, hb. *Hance*, 21368!). *Mus. Brit.*; *Herb. Kew.*

Java.

1. *Colocasia antiquorum*, *Schott, Melet.* p. 18, *Gen. Aroid.* t. 37, et *Prodr. Aroid.* p. 138; *Miq. Fl. Ind. Bat.* iii. p. 202; *Kunth, Enum. Pl.* iii. p. 37; *Engl. in DC. Monogr. Phan.* ii. p. 491, cum synonymis varietatibusque, var. *illustris*, *Engl.*, exclusa; *Hook. f. Fl. Brit. Ind.* vi. p. 523.

*Colocasia antiquorum* et *C. nymphæfolia*, *Wight, Ic. Pl. Ind. Or.* iii. pt. 1, p. 5, t. 786.

*Colocasia prunipes*, *K. Koch, in Ind. Sem. Hort. Berol.* 1854, *App.* p. 4; *Walp. Ann.* v. p. 871.

*Arum esculentum*, *Linn. Sp. Pl.* ed. 1, p. 965; *Lour. Fl. Cochinch.* p. 535.

KIANGSI: Kiukiang (*Shearer*!); SZECHUEN (*A. Henry*, 8830!). *Herb. Kew.*

India, Malay Archipelago, widely cultivated in the tropics.

1. *Alocasia cucullata*, *Schott, in Oestr. Bot. Wochenbl.* iv. (1854) p. 410; *Prodr. Aroid.* p. 155; *Engl. in DC. Monogr. Phan.* ii. p. 498, cum synonym.; *Hook. f. Fl. Brit. Ind.* vi. p. 525.

*Arum cucullatum*, *Lour. Fl. Cochinch.* p. 536; *Roxb. Fl. Ind.* iii. p. 501; *Wight, Ic. Pl. Ind. Or.* iii. pt. 1, p. 6, t. 787.

SZECHUEN: below Chungking (*Faber*, 29!); KWANGTUNG: Canton (ex *Loureiro*): HAINAN (*A. Henry*, 8626!). *Herb. Kew.*  
Khasia.

2. *Alocasia hainanica*, *N. E. Br.*, n. sp.

*Petiolus* 7-10 poll. longus, usque ad medium vaginatus; lamina 7-8 poll. longa, 4-5 poll. lata, peltata, ovata, basi cordato-sagittata, apice acuta, lobis posticis late ovatis vel triangulari-ovatis obtusis ad  $\frac{1}{2}$  poll. connatis, sinu triangulari  $1\frac{1}{2}$ -2 poll. profundo

sejunctis, nervis lateralibus 5 utrinque costa media abeuntibus, rectis, patentibus. *Pedunculus* solitarius, 4–5 poll. longus, in vaginem petioli longe inclusus. *Spathæ* tubus  $\frac{3}{4}$ –1 poll. longus, subcylindricus vel anguste ovoideus; lamina 2–2 $\frac{3}{4}$  poll. longa, lanceolata acuminata. *Spadix* 1 $\frac{1}{2}$ –1 $\frac{3}{4}$  poll. longus, parte feminea  $\frac{1}{4}$  poll. longa, parte mascula  $\frac{1}{2}$  poll. longa, parte intermedia organis neutris dense oblecta; appendix 3 lin. longa, interdum deest; ovaria ovoidea; ovula 4–5.

HAINAN (*A. Henry*, 8679!). Herb. Kew.

3. *Alocasia macrorrhiza*, *Schott*, in *Oestr. Bot. Wochenbl.* iv. (1854) p. 409, *Gen. Aroid.* t. 40, et *Prodr. Aroid.* p. 146; *Miq. Fl. Ind. Bat.* iii. p. 205; *Engl. in DC. Monogr. Phan.* ii. p. 502; *Hook. f. Fl. Brit. Ind.* vi. p. 526.

*Alocasia commutata*, *Schott*, in *Oestr. Bot. Wochenbl.* iv. (1854) p. 409; *Seemann, Bot. Voy. 'Herald.'* p. 416.

*Alocasia odora*, *C. Koch*, in *Ind. Sem. Hort. Berol.* 1854, *App.* p. 5; *Walp. Ann.* v. p. 874; *Engl. in DC. Monogr. Phan.* ii. p. 503.

*Arum macrorrhizon*, *Linn. Sp. Pl.* ed. 1, p. 965; *Lour. Fl. Cochinch.* p. 535.

*Arum odorum*, *Roxb. Fl. Ind.* iii. p. 499; *Wight, Ic. Pl. Ind. Or.* iii. pt. 1, p. 6, t. 797; *Lodd. Bot. Cab.* t. 416.

*Caladium odorum*, *Lindl. Bot. Reg.* t. 641.

*Caladium odoratum*, *Roxb. ex Lindl. Bot. Reg.* sub t. 641.

*Colocasia odora*, *Brongn. in Nouv. Ann. Mus. Par.* iii. (1834) p. 145, pl. 7; *Benth. Fl. Hongk.* p. 343.

*Colocasia indica*, *Benth. Fl. Hongk.* p. 343, non *Kunth*.

KWANGTUNG: Macao (*Gaudichaud ex Engler*); HONGKONG: Tank Ravine (*Urquhart! Harland*, 749! *Hance*, 1189!), Mount Gough (*Lamont*, 736!). Mus. Brit.; Herb. Kew.

India, and cultivated in the Eastern Tropics.

[*Arum sagittifolium*, *Lour. Fl. Cochinch.* p. 534, not of *Linn.*, and excluding synonymy, stated to be “cultivated and wild in Cochinchina and China,” is probably a species of *Alocasia*, but I know of no species to which *Loureiro's* description will apply.]

1. *Aglaonema modestum*, *Schott, ex Engl. in DC. Monogr. Phan.* ii. p. 442.

*Aglaonema acutispathum*, *N. E. Brown, in Gard. Chron.* ser. 2, xxiv (1885) p. 39.

“Cultivated in South China” (*Hance*, 11459!).

KWANGTUNG? (*Knaggs*!); KIANGSU? (*Knaggs*!). Mus. Brit.;  
Herb. Kew.

Philippines.

There is some uncertainty about the locality of this; the plant which I described as *A. acutispathum* was bought in Hongkong by Dr. Knaggs and thought to come from the vicinity of Canton. But there is another specimen in the Kew Herbarium from a plant flowered by him labelled “From Shanghai. Brought home by Dr. H. V. Knaggs.” From this, and the statement by Hance on his label that it is cultivated, it seems probable that it may prove not to be a native of China. In my description I stated that it was distinct from *A. modestum*, but Engler’s reproduction of Schott’s figure of this species (Engler, *Araceæ*, no. 74), which I had not then seen, demonstrates that the two are identical. The specimens distributed by Engler as *A. marantifolium* (Engler, *Araceæ* exsicc. et illustratæ No. 258) also belong to *A. modestum*.

According to a statement with Engler’s figure of the type, *A. modestum* was collected at Manila by Gaudichaud and is in the Paris Herbarium.

1. *Scindapsus sinensis*, *Engl. in Engl. Jahrb.* xxix. (1900)  
p. 234.

SZECHUEN: Nanchuan (*Bock von Rosthorn*, 758, ex Engler).

1. *Rhaphidophora angustifolia*, *Schott, in Bonplandia*, v.  
(1857) p. 45, et *Prodr. Aroid.* p. 378.

*Rhaphidophora hongkongensis*, *Schott, in Bonplandia*, v. (1857) p. 45;  
*Prodr. Aroid.* p. 378; *Seem. Bot. Voy. ‘Herald,’* p. 417; *Engl. in DC.*  
*Monogr. Phan.* ii. p. 240.

*Rhaphidophora sylvestris*, *Engl. in DC. Monogr. Phan.* ii. p. 239.

*Rhaphidophora Peepla*, *Benth. Fl. Hongk.* p. 344, non Schott.

*Scindapsus sylvestris*, *Kunth, Enum. Pl.* iii. p. 64; *Miq. Fl. Ind. Bat.*  
iii. p. 187.

HONGKONG: trailing over rocks below Buddhist temple E.  
Point (*Lamont! Champion*, 535! *Hance* ex *Seemann*). Mus. Brit.;  
Herb. Kew.

Java, Borneo.

Bentham, *Fl. Hongk.* p. 344, unites this with *R. lancifolia*,  
Schott, and *R. Peepla*, Schott, two quite distinct species, and

describes from Indian specimens of *R. Peepla*, so that his description does not apply to this plant at all. *R. hongkongensis* was founded on a single leaf collected by Champion, now in the Kew Herbarium, which does not differ in any way from the leaves of *R. angustifolia*. Some leaves of Lamont's specimen in the British Museum Herbarium are 14 inches long, exclusive of the petiole, and  $3\frac{1}{2}$  inches broad.

1. **Epipremnum mirabile**, *Schott, Gen. Aroid. p. 79; Prodr. Aroid. p. 388; Engl. in DC. Monogr. Phan. ii. p. 249; N. E. Br. in Gard. Chron. ser. 2, xvii. (1882) p. 180; Journ. Bot. xx. (1882) p. 332; Hook. f. Fl. Brit. Ind. vi. p. 549.*

*Rhaphidophora lacera*, *Hassk. in Tijdschr. Nat. Gesch. Phys. ix. (1842) p. 168; in Flora, xxv. (1842) II. Beibl. p. 11; et Pl. Jav. Rar. p. 155, excl. synonym.*

*Rhaphidophora pinnata*, *Schott, in Bonplandia, v. (1857) p. 45, et Prodr. Aroid. p. 384; Hance, in Journ. Bot. xvii. (1879) p. 16.*

KWANGTUNG: near Whampoa (*Hance, 15,600!* and at Lien-fashan, along the banks of Canton River, ex *Hance*). Mus. Brit.; Herb. Kew.

This is the Tonga plant, of which a full account will be found in the 'Journal of Botany,' *l. c.* It is widely distributed from South China and the Malay Peninsula, through the Malay Archipelago and Polynesia, to tropical Australia.

1. **Pothos Loureiri**, *Hook. & Arn. Bot. Beech. Voy. p. 220, Schott, Aroid. p. 23, t. 49, et Prodr. Aroid. p. 569; Engl. in DC. Monogr. Phan. ii. p. 87.*

*Pothos terminalis*, *Hance, in Ann. Sc. Nat. Bot. 5<sup>me</sup> sér. v. (1866) p. 247.*

*Pothos microphyllus*, *Schott, Aroid. p. 23, t. 40. fig. B (quoad plantam sinensem et figuram), non Presl.*

*Flagellaria repens*, *Lour. Fl. Cochinch. p. 212.*

CHINA (*Millett!*); KWANGTUNG: Tingushan, on the West River (*Sampson* in hb. *Hance, 11,168! Lamont, 1064! Ford!*); HAINAN (*A. Henry, 112! 7980! & 7980 A! Swinhoe!*). Mus. Brit.; Herb. Kew.

Tonquin, Cochinchina.

2. **Pothos Seemanni**, *Schott, in Bonplandia, v. (1857) p. 45; Aroid. p. 22, t. 43; Prodr. Aroid. p. 564; Engl. in DC. Monogr. Phan. ii. p. 83.*

*Pothos scandens*, *Benth. Fl. Hongk.* p. 344; *Lindl. Bot. Reg.* t. 1337, non *Linn.*

FORMOSA (*Oldham*, 308, 632! *Walters*, 765!), Ape's Hill, Takow (*Playfair*, 438!); HUPEH: Ichang (*A. Henry*, 4395!); SZECHUEN: Chungking (*Bourne*!); KWANGTUNG: Macao (*Cal- lery*, 295!), Lofaushan, without collector's name; HONGKONG (*Champion*, 288! *Wilford*, 280! *Wright*, 509! *Hance*, 765! *Lamont*, 738!). *Mus. Brit.*; *Herb. Kew.*

1. *Acorus Calamus*, *Linn. Sp. Pl.* ed. 1, p. 324; *Schott, Prodr. Aroid.* p. 578; *Engl. in DC. Monogr. Phan.* ii. p. 216; *Bunge, Enum. Pl. Chin. Bor.* p. 67; *Debeaux, Fl. Shangh.* p. 62; *Franch. Pl. David.* i. p. 313; *Hook. f. Fl. Brit. Ind.* vi. p. 555.

*Acorus cochinchinensis*, *Schott, Prodr. Aroid.* p. 580.

*Orontium cochinchinense*, *Lour. Fl. Cochinch.* p. 208.

KIANGSU: Shanghai (*Maingay*, 405!); HUPEH: Ichang, Nanto and mountains to northward (*A. Henry*, 4506!). *Mus. Brit.*; *Herb. Kew.*

North Temperate zone.

2. *Acorus gramineus*, *Ait. Hort. Kew.* i. p. 474; *Schott, Prodr. Aroid.* p. 580; *Engl. in DC. Monogr. Phan.* ii. p. 218; *Hook. f. Fl. Brit. Ind.* vi. p. 556; *Diels, in Engl. Jahrb.* xxix. (1900) p. 234.

*Acorus Calamus*, *Lour. Fl. Cochinch.* p. 208; *Benth. Fl. Hongk.* p. 345, non *Linn.*

*Acorus terrestris*, *Spreng. Syst. Veg.* ii. p. 118; *Schultes, Syst. Veg.* vii. p. 174; *Schott, Prodr. Aroid.* p. 579.

*Acorus Tatarinowii*, *Schott, in Oest. Bot. Zeitschr.* ix. (1859) p. 101.

*Acorus Calamus*, var. *terrestris*, *Engl. in DC. Monogr. Phan.* ii. p. 217.

CHEKIANG: Tienta Mountain, 3000 ft. (*Faber*, 87!); KIANGSI: Kiukiang (*Shearer! Forbes*, 59!); HUPEH: Ichang (*A. Henry* 727, 3370, 7857!), Nanto (*A. Henry*, 1904!), Changyang (*A. Henry*, 5246!); SZECHUEN: Nanchuan (*Bock von Rosthorn*, 1253, ex *Engler*), Mount Omei, 2500 ft. (*Faber*, 1193!), near Tachienlu (*Pratt*, 792!); KWANGTUNG: West River, Canton (*Lamont*, 739!); HONGKONG (*Urquhart*, 112! *Eyre! Wright*, 510! *Wilford*, 314! *Lamont*, 739! *Hance*, 973!); HAINAN (*Swinhoe*!). *Mus. Brit.*; *Herb. Kew.*

Japan, Tonquin.

This is the common *Acorus* of China, and is very variable in size. The larger forms have been confused with *A. Calamus*, Linn., from which they are readily distinguished by the absence of a distinctly thickened midrib to the leaves and spathe, a more slender spadix, and a shorter spathe. *A. Calamus* seems to be rare in China, as I have seen but two specimens. Owing to the confusion of names, only specimens actually seen are here quoted, except one on the authority of Dr. Engler.

## 154. LEMNACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. **Lemna minor**, *Linn. Sp. Pl.* ed. 1, p. 970; *Lour. Fl. Cochinch.* p. 550; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 129; *Bretschneider, Early Res.* p. 181; *Hegelmaier, Die Lemnaceen*, p. 142; *Debeaux, Fl. Shangh.* p. 62; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 12; *Hook. f. Fl. Brit. Ind.* vi. p. 556.

CHIHLI: Peking (*Bretschneider*, 777!); KIANGSU: Shaughai (ex *Debeaux*); CHEKIANG (ex *Debeaux*); HONGKONG (*Hance*). Herb. Kew.

“Frequent in S. China. Common also in N. China” (*Bretschneider*).

Temperate and warmer regions of the world.

2. **Lemna paucicostata**, *Hegelmaier, Die Lemnaceen*, p. 139; *Mart. Fl. Bras.* iii. II. p. 20, t. i. fig. 5; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 12; *Hook. f. Fl. Brit. Ind.* vi. p. 556.

LUCHU ARCHIPELAGO (*Wright*, 324!). Herb. Kew.

Throughout the tropics.

This has been confused by many authors with *Lemna minor*, Linn., from which it differs in its less symmetrical frond and erect ovule.

3. **Lemna polyrrhiza**, *Linn. Sp. Pl.* ed. 1, p. 970; *Flor. Dan.* t. 1589; *Syme, Engl. Bot.* ed. 3, t. 1397; *Kurz, in Journ. Linn. Soc.* ix. p. 267; *Hook. f. Fl. Brit. Ind.* vi. p. 557.

*Spirodela polyrrhiza*, *Schleid. in Linnæa*, xiii. p. 392, et in *Ann. Sci. Nat.* 2<sup>me</sup> sér. xiii. p. 148; *Maxim. Prim. Fl. Amur.* p. 266; *Mart. Flor. Bras.* iii. II. p. 21, t. i. fig. 6; *Hegelmaier, Die Lemnaceen*, p. 151, t. xiii. figs. 10–16, tt. xiv.–xv.; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 13.



*Lemna bannatica*, *Waldst. et Kit. ex Schleid. in Linnæa*, xiii. p. 392.

*Lemna orbiculata*, *Roxb. Fl. Ind.* iii. p. 565.

*Lemna major*, *Griff. Notulæ*, iii. p. 216.

KIANGSU: Shanghai (*Maingay*, 674, ex *Kurz*); LUCHU ARCHIPELAGO (*Wright*, 325!). *Herb. Kew.*

Warm and temperate regions of the Old World, Northern United States to Venezuela and Cuba.

### 155. ALISMACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Alisma Plantago*, *Linn. Sp. Pl.* ed. 1, p. 342; *Kunth, Enum. Pl.* iii. p. 148; *Reichb. Ic. Fl. Germ.* vii. t. 57; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 16; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 256; *Hook. f. Fl. Brit. Ind.* vi. p. 559; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 477; *Kurz, in Journ. Bot.* xi. (1873) p. 193; *Micheli, in DC. Monogr. Phan.* iii. p. 32.

*Alisma flava* et *A. cordifolia*, *Thunb. Fl. Jap.* p. 153, non *Linn.*

Var. *latifolium*, *Kunth, Enum. Pl.* iii. p. 148; *Gren. et Godr. Fl. France*, iii. p. 165; *Franch. Pl. David.* i. p. 134.

CHIHLI: Jehol (*David*, 2009, ex *Franchet*); SHINGKING (*Ross*, 39!); SHANTUNG: Chefoo (*Stuhlmann! Forbes! Fauvel ex Franchet*). *Mus. Brit.*; *Herb. Kew.*

Var. *angustifolium*, *Kunth, Enum. Pl.* iii. p. 148.

HUPEH: Patung (*A. Henry*, 1852, 5117!); KWANGTUNG (*Ford*, 278!); COREAN ARCHIPELAGO: Long Reach (*Oldham*, 1062!). *Mus. Brit.*; *Herb. Kew.*

Widely distributed through the north and south temperate regions.

This species is also stated by *Kurz* to have been collected by *Anderson* in Yunnan, but without any indication of which variety is intended.

1. *Lophotocarpus guyanensis*, *Durand & Schinz, Conspect. Fl. Afr.* v. p. 487; *C. H. Wright, in Dyer, Fl. Trop. Afr.* viii. p. 210.

*Sagittaria guyanensis*, *Humb., Bonpl. et Kunth, Nov. Gen. et Spec.* i. p. 250; *Kunth, Enum. Pl.* iii. p. 161; *Hook. f. Fl. Brit. Ind.* vi. p. 561.

*Sagittaria cordifolia*, *Roxb. Fl. Ind.* iii. p. 647; *Kunth, Enum. Pl.* iii. p. 161; *Benth. Fl. Hongk.* p. 346.

*Echinodorus guianensis*, *Griseb. Fl. Brit. W. Ind.* p. 505.

*Lophiocarpus guyanensis*, *Micheli, in DC. Monogr. Phan.* iii. p. 62.

HONGKONG (*Harland ex Bentham*).

Tropical Asia, Africa, Madagascar, and America.

1. *Sagittaria sagittifolia*, *Linn. Sp. Pl.* ed. 1, p. 993; *Kunth, Enum. Pl.* iii. p. 156; *Red. Lil.* tt. 279, 280; *Lour. Fl. Cochinch.* p. 570; *Kurz, in Journ. Bot.* xi. (1873) p. 193; *Franch. Pl. David.* i. p. 314; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 16; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 256; *Micheli, in DC. Monogr. Phan.* iii. p. 66; *Hook. f. Fl. Brit. Ind.* vi. p. 561.

*Sagittaria sagittata* et *S. obtusa*, *Thunb. Fl. Jap.* p. 242.

*Sagittaria chinensis*, *Sims, Bot. Mag.* t. 1631.

*Sagittaria macrophylla*, *Bunge, in Mém. Sav. Étrang. Acad. Pétersb.* ii. (1834) p. 137; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 477.

*Sagittaria sagittifolia*, var. *diversifolia*, *Micheli, in DC. Monogr. Phan.* iii. p. 67.

*Sagittaria sagittifolia*, var. *longiloba*, *Turcz. in Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. p. 57; *Regel, Fl. Ussur.* p. 154; *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 138, iii. p. 192; *Debeaux, Fl. Shangh.* p. 61.

CHIHLI: Peking (*Bretschneider, 1972*, in hb. *Forbes!*), Jehol (*David, 1917*, ex *Franchet*); SHINGKING (*Ross, 38!*); KIANGSU: Shanghai and Woosong (ex *Debeaux*); KIANGSI (*Staunton!*), Kiukiang (*Shearer! Forbes!*); FORMOSA (*Oldham, 368, 640! Wilford, 500!*), Takow (*Playfair, 182!*); HUPEH: Ichang (*A. Henry, 105, 300, 2324!*), South Patung (*A. Henry, 7387!*), Nanto (*A. Henry, 2664!*); YUNNAN (*Anderson!*); KWANGTUNG: Canton (*Seemann! Hance, 5344!*), Macao? (*Lay!*); HAINAN (*A. Henry, 8064, 8390!*); COREA: Port Chusan (*Wilford, 903!*). *Mus. Brit.; Herb. Kew.*

Widely dispersed in Europe and Temperate Asia.

The leaves of this species vary greatly in shape, the one extreme being represented by *Bretschneider's* Peking specimen, which has the lobes 7 in. long by 5½ in. broad, the other by *Wilford's* no. 500 from Formosa with lobes 3½ in. long but only ¼ in. broad. These two are connected by numerous intermediate forms. As

the difference of leaf-form does not appear to be correlated with any other characters, it is impossible to define the various forms included rightly by Micheli under his variety *diversifolia*. The form known as *S. chinensis*, Sims (*S. macrophylla*, Bunge), is stated by Hance to be cultivated near Canton for the sake of its edible tubers.

Var. *oligocarpa*, Micheli, in DC. *Monogr. Phan.* iii. p. 68.

*Sagittaria pygmæa*, Miq. in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 138.

*Sagittaria sagittifolia*, L.?, Hook. et Arn. *Bot. Beechey's Voy.* p. 220.

CHINA (*Beechey*!); KIANGSI: Kiukiang (*Forbes*, 127, 188, 201, 1040!); HUPEH: Ichang (*A. Henry*, 4299!), Nanto (*A. Henry*, 1919!); SZECHUEN?, Suifu rice-fields (*Faber*, 975!); KWANGTUNG (*Ford*, 334!). Mus. Brit.; Herb. Kew.

Japan.

[*Sagittaria trifolia*, Linn. *Sp. Pl.* ed. 1, p. 993, said to be from China, is founded upon the figure in Petiver's 'Gazophylacium,' t. 19. fig. 5, and is referred by Micheli (in DC. *Monogr. Phan.* iii. p. 67) to *Sagittaria sagittifolia*, var. *diversifolia*, Micheli. It must, however, be regarded as a very doubtful plant. The leaves are represented with very narrow lobes.]

1. *Butomus umbellatus*, Linn. *Sp. Pl.* ed. 1, p. 372; *Kunth*, *Enum. Pl.* iii. p. 164; *Maxim. Ind. Fl. Pek. in Prim. Fl. Amur.* p. 477; *Reichb. Ic. Fl. Germ.* vii. t. 58; *Boiss. Fl. Orient.* v. p. 12; *Franch. Pl. David.* i. p. 315; *Micheli, in DC. Monogr. Phan.* iii. p. 85; *Hook. f. Fl. Brit. Ind.* vi. p. 562.

*Butomus junceus*, Turcz. in *Bull. Soc. Nat. Mosc.* (1837) no. vii. p. 157 (nomen), et xxvii. (1854) ii. p. 60.

*Butomus umbellatus*, var. *minor*, Ledeb. *Fl. Ross.* iv. p. 44.

*Butomus umbellatus*, var. *junceus*, Turcz. in *Bull. Soc. Nat. Mosc.* xxvii. (1854) ii. p. 60; *Micheli, in DC. Monogr. Phan.* iii. p. 86.

CHIHLI: Peking (*Wawra ex Micheli*), mountains west of Peking (*Bretschneider* in hb. *Forbes*, 1971!), Jehol (*David*, 1955!). Mus. Brit.; Herb. Kew.

Europe, Northern and Western Asia to the Punjab and Kashmir.

[Under *Damasonium indicum*, Willd., in *Bot. Mag.* t. 1201, it is stated that "Sir Joseph Banks has different species brought from China by Sir Geo. Staunton." I have not found the specimens referred to.]

## 156. NAIADACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Triglochin maritimum*, *Linn. Sp. Pl.* ed. 1, p. 339; *Act. Stockh.* (1742) p. 147, t. 6. figs. 4-5; *Lam. Illustr.* t. 270. fig. 2; *Fl. Dan.* t. 306; *Kunth, Enum. Pl.* iii. p. 145; *Maxim. Ind. Fl. Mongol. in Prim. Fl. Amur.* p. 485; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 17; *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 139; *Micheli, in DC. Monogr. Phan.* iii. p. 105; *Hook. f. Fl. Brit. Ind.* vi. p. 563.

*Ornithogalum japonicum*, *Buerger, ex Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 139.

CHIHLI: Peking (ex *Maximowicz*); KIANGSI: Nanshan near Sudshou (*Martin*!); KANSUH: Mudskik Mountains (*Przewalski*!). *Herb. Kew.*

Europe, Temperate Asia, North America, Patagonia.

2. *Triglochin palustre*, *Linn. Sp. Pl.* ed. 1, p. 338; *Kunth, Enum. Pl.* iii. p. 143; *Lam. Illustr.* t. 270. fig. 1; *Fl. Dan.* t. 490; *Maxim. Ind. Fl. Mongol. in Prim. Fl. Amur.* p. 485; *Micheli, in DC. Monogr. Phan.* iii. p. 98; *Hook. f. Fl. Brit. Ind.* vi. p. 563.

CHIHLI: near Peking (*Wawra*, 1018, ex *Micheli*); SHING-KING: Talienshan (*Swinhoe*!); SHANTUNG: Chefoo (*Forbes*, 406!); SZECHUEN: Tongolo (*Soulié*, 250!), near Sulphur springs, Haiaz (*Faber*, 1544!). *Mus. Brit.*; *Herb. Kew.*

Europe, Temperate Asia, North America, Temperate South America.

It is noteworthy in the distribution of this genus that five species are indigenous in Australia and four of them are endemic.

1. *Aponogeton monostachyon*, *Linn. fil. Suppl.* p. 214; *Roxb. Pl. Corom.* i. p. 58, t. 81; *Andr. Bot. Rep.* t. 406; *Thwaites, Enum. Pl. Zeyl.* p. 333; *Benth. Fl. Austral.* vii. p. 188; *Hance, in Journ. Bot.* xvi. (1878) p. 111; *Hook. f. Fl. Brit. Ind.* vi. p. 564.

*Aponogeton monostachys*, *Edgew. in Hook. Lond. Journ. Bot.* iii. (1844) p. 404, t. 17.

*Spathium monostachyon*, *Edgew. in Calc. Journ. Nat. Hist.* iii. (1843) p. 533, t. 16.

*Potamogeton indicus*, *Roth, ex Roem. et Schultes, Syst. Pl.* iii. p. 516.

*Saururus natans*, *Linn. Mant. alt.* p. 227; (*Rheede, Hort. Malab.* xi. p. 31, t. 15).

KWANGTUNG: Pools near Canton (*Sampson*, 1268, 15588!)  
Herb. Kew.

Tropical Asia and Australia.

2. *Aponogeton* sp.

COREA (*Faurie*!).

POTAMOGETON.

(By ARTHUR BENNETT, F.L.S.\*)

1. *Potamogeton angustifolius*, *Bercht. et Presl, Rostlin*, i. p. 19.

*Potamogeton Zizii*, *Roth, Enum. Pl. Germ.* i. sect. i. p. 531.

*Potamogeton heterophyllus*, var. *latifolius*, *Mert. et Koch, Deutsch. Fl.* i. p. 845.

YUNNAN: Plain of Tali (*Delavay*!).

Europe, Asia, Africa, and America.

2. *Potamogeton crispus*, *Linn. Sp. Pl.* ed. 1, p. 126; *Kunth, Enum. Pl.* iii. p. 133; *Reichb. Ic. Fl. Germ.* vii. t. 29. fig. 50; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 15; *Hook. f. Fl. Brit. Ind.* vi. p. 566.

CHIHLI (*Potanin*!), near Peking (*Bretschneider*, 893!);  
KIANGSU: Shanghai (*Maingay*, 422! *Faber*, 2!); YUNNAN  
(*Delavay*, 2500!); COREA (*Faurie*!); LUCHU ARCHIPELAGO  
(*Wright*, 323!). Mus. Brit.; Herb. Kew.

Var. *serratus*, *S. F. Gray, Nat. Arr. Brit. Pl.* ii. p. 34.

SHANGHAI (*Lamprey*!).

North and south temperate and subtropical regions.

The Corean plant is a narrow-leaved form.

3. *Potamogeton cristatus*, *Regel & Maack, Tent. Fl. Ussur.* p. 153, t. 10. figs. 3-6.

*Potamogeton hybridus*, *Makino, Ill. Fl. Jap.* i. no. 9, p. 2, t. 55, non *Michaux*; *Arth. Benn. in Journ. Bot.* xxx. (1892) p. 229.

---

\* Mr. Bennett had the Paris, St. Petersburg, and other collections of this genus on loan; hence this enumeration includes certain species and varieties not represented either at the British Museum or at Kew by Chinese specimens.—*W. B. H.*

KIUKIANG (*Maries*!); HUPEH: Ichang (*A. Henry*, 2377!).  
Mus. Brit.; Herb. Kew.

Ussuri, Mandshuria, and Japan.

4. *Potamogeton Delavayi*, *Arth. Benn. in Journ. Bot.* xxx. (1892) p. 228.

YUNNAN (*Delavay*, 4750!).

5. *Potamogeton Gaudichaudii*, *Cham. et Schlecht. in Linnæa*, ii. (1827) pp. 177 et 199.

*Potamogeton mucronatus*, *Presl, Epim. Bot.* p. 245; *Hook. f. Fl. Brit. Ind.* vi. p. 567.

*Potamogeton malaiana*, *Miq. Ill. Fl. Arch. Ind.* p. 46.

*Potamogeton japonicus*, *Franch. & Savat. Enum. Pl. Jap.* ii. p. 15.

*Potamogeton Wrightii*, *Morong, in Bull. Torrey Bot. Club*, xiii. (1886) p. 158.

*Potamogeton lucens*, *Vidal, Phan. Cuming. Philipp.* p. 155, non *Linn.*

CHIHLI: Peking (*Bretschneider*, 904!); FORMOSA: Takow (*A. Henry*, 1203!), Bankingsing (*A. Henry*, 1203 A!); YUNNAN: Lac de Koumpo (*Delavay*, 2727!); KWANGTUNG: Canton (*Graves* in herb. *Hance*, 22226!); COREA (*Faurie*!); LUCHU ARCHIPELAGO (*Wright*, 320!). Mus. Brit.; Herb. Kew.

India, Japan, Philippine Islands, Borneo, and Ladrone Islands.

This species was distributed by Maximowicz under the name of *Potamogeton tretocarpus*. Oldham's no. 635 from Formosa may be the young state of this species.

6. *Potamogeton Hillii*, *Morong, in Bot. Gaz.* vi. (1881) p. 290, *et in Mem. Torrey Bot. Club*, iii. (1893) pt. ii. p. 38, t. 46?

HUPEH: Ichang (*A. Henry*, 4375!). Herb. Kew.

North-eastern United States.

Dr. Henry's specimen is destitute of fruit, hence it cannot be positively asserted to belong to this species, although in all other characters it agrees.

7. *Potamogeton javanicus*, *Hassk. in Verh. Naturk. Ver. Nederl. Ind.* i. (1856) p. 26; *Hook. f. Fl. Brit. Ind.* vi. p. 566.

*Potamogeton tenuicaulis*, *F. Muell. Fragm.* i. pp. 90, 244.

*Potamogeton parvifolius*, *Buchenau, in Abhandl. Natur. Ver. Bremen*, vii. (1880) p. 32.

*Potamogeton hybridus*, *Hook. f. et Thoms. ex Hook. f. Fl. Brit. Ind.* vi. p. 566, non *Michaux*.

*Potamogeton heterophyllus*, *Buch.-Ham. in Wall. Cat.* no. 5181.

SHANGTUNG (*Maingay*, 170!); FORMOSA: Tamsui (*Oldham*, 381, 638!); COREA: Port Chusan (*Wilford*, 968!). Mus. Brit.; Herb. Kew.

Japan, Tropical Asia, Africa, Madagascar, and Australia.

8. *Potamogeton limosellifolius*, *Maxim. ex Korshinsky*, in *Act. Hort. Petrop.* xii. (1892) p. 393.

COREA (*Faurie*!); LUCHU ARCHIPELAGO (*Wright ex Korshinsky*).

Amur.

This may be a small, delicate variety or subspecies of *P. Miduhikimo*, Makino (*Ill. Fl. Jap.* t. 54).

9. *Potamogeton lucens*, *Linn. Sp. Pl.* ed. 1, p. 126; *Kunth, Enum.* iii. p. 132; *Hook. f. Fl. Brit. Ind.* vi. p. 567.

CHINA, without precise locality (*Korolkow and Krause*!). Mus. Brit.

Dispersed over all the continents, but rare in Australia.

10. *Potamogeton natans*, *Linn. Sp. Pl.* ed. 1, p. 126; *Kunth, Enum. Pl.* iii. p. 127; *Hook. f. Fl. Brit. Ind.* vi. p. 565.

SHINGKING (*James*!). Herb. Kew.

Very widely diffused, especially in temperate regions.

Abbé David's no. 1949 enumerated by Franchet (*Pl. David.* i. p. 315) as *P. natans*, Linn., is not that species, and may belong to either *P. Tepperi*, A. Benn., or *P. tricarinatus*, F. Muell. et A. Benn., but in the absence of fruit it cannot be determined with certainty.

11. *Potamogeton oxyphyllus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 161.

COREA (*Faurie*!).

Japan.

12. *Potamogeton pectinatus*, *Linn. Sp. Pl.* ed. 1, p. 127; *Kunth, Enum.* iii. p. 137; *Bunge, Enum. Pl. Chin. Bor.* p. 68; *Hook. f. Fl. Brit. Ind.* vi. p. 567.

NORTH CHINA: Tanschan (*Bunge*!); CHIHLI (*Potanin*!); KANSUH (*Potanin*!); COREA (*Faurie*!). Herb. Kew.

Common in the north and south temperate regions; less frequent in the tropics.

13. *Potamogeton perfoliatus*, *Linn. Sp. Pl.* ed. 1, p. 126; *Kunth, Enum.* iii. p. 133; *Hook. f. Fl. Brit. Ind.* vi. p. 566.

CHIHLI: near Peking (*Bretschneider*, 779!); CHINA, without precise locality (*Korolkow & Krause*!). *Herb. Kew.*

North temperate regions and Australia.

14. *Potamogeton polygonifolius*, *Pourr. in Mém. Acad. Toulouse*, iii. (1788) p. 325; *Franchet, in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 261.

CHIHLI: Peking (*Bretschneider*!); SHANTUNG: Chefoo (*Fauvel ex Franchet*); KANSUH (*Potanin*!).

Europe, including Britain.

15. *Potamogeton pusillus*, *Linn. Sp. Pl.* ed. i. p. 127; *Kunth, Enum. Pl.* iii. p. 136; *Franch. Pl. David.* i. p. 315; *Hook. f. Fl. Brit. Ind.* vi. p. 567.

CHIHLI: Jehol (*David*, 1916!); KANSUH (*Potanin*!), Tangut (*Przewalski*!); COREA (*Faurie*!). *Herb. Kew.*

Widely distributed, but absent from Australia.

16. *Potamogeton Robbinsii*, *Oakes*, var. *japonicus*, *Arth. Benn. in Bull. Herb. Boiss.* iv. (1896) p. 549.

*Potamogeton serrulatus*, *Regel et Maack, in Regel, Tent. Fl. Ussur.* p. 153.

COREA (*Faurie*!).

Japan.

17. *Potamogeton Tepperi*, *Arth. Benn. in Journ. Bot.* xxv. (1887) p. 178.

CHIHLI: Peking (*Bretschneider*, 778!); FORMOSA (*Oldham*, 636!); HUPEH: Ichang (*A. Henry*, 2366, 3603!); SZECHUEN: Hokiang (*Faber*, 314!); COREAN ARCHIPELAGO: Long Reach (*Oldham*, 824!). *Mus. Brit.*; *Herb. Kew.*

Siberia, India, Japan, and Australia.

Specimens from Yunnan, collected by Delavay, probably belong here.

18. *Potamogeton* sp.

YUNNAN (*Delavay*!).

This belongs to the group of *P. polygonifolius*, *Pourr.*, but the material is insufficient for determination.



1. *Ruppia maritima*, *Linn. Sp. Pl.* p. 127; *Willd. Sp. Pl.* i. p. 717; *Reichb. Pl. Crit.* ii. p. 66, t. 174. fig. 307; *Kunth, Enum. Pl.* iii. p. 123; *Benth. Fl. Austral.* vii. p. 174; *Mart. Fl. Bras.* iii. III. p. 699.

*Ruppia rostellata*, *Koch, in Reichb. Ic. Crit.* ii. p. 66, t. 174. fig. 306; *Hook. f. Fl. Brit. Ind.* vi. p. 568; *Syme, Engl. Bot.* ix. t. 928.

*Ruppia* sp., *Griff. Notul.* iii. p. 196; *Icon.* tt. 257-9.

FORMOSA: Takow (*A. Henry*, 1099!); COREAN ARCHIPELAGO (*Oldham*, 825!), Port Hamilton (*Oldham*, 1003!). *Herb. Kew.*

All temperate and subtropical regions.

1. *Zannichellia pedicellata*, *Buch.-Ham. in Wall. Cat.* n. 5185; *Fries, Mantissa*, iii. p. 133; *Syme, Engl. Bot.* ed. 3, ix. t. 926.

*Zannichellia palustris*, *Kunth, Enum. Pl.* iii. p. 124, *partim*.

*Zannichellia pedunculata*, *Reichb. Ic. Pl. Crit.* viii. p. 24, t. 760; *Ic. Fl. Germ.* vii. t. 16. fig. 21.

*Zannichellia palustris*, subsp. *pedicellata*, *Hook. f. Fl. Brit. Ind.* vi. p. 568.

KIANGSU (*Faber*, 1!); LUCHU ARCHIPELAGO (*Wright*, 326!). *Herb. Kew.*

Cosmopolitan, excluding Australia.

1. *Zostera marina*, *Linn. Sp. Pl.* ed. 1, p. 968; *Thunb. Fl. Jap.* p. 232; *Fl. Dan.* t. 15; *Kunth, Enum. Pl.* iii. p. 116; *Maxim. Prim. Fl. Amur.* p. 266; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 261; *Syme, Engl. Bot.* ed. 3, ix. t. 929; *Debeaux, Fl. Tchef.* p. 132.

SHANTUNG: Chefoo (ex *Debeaux & Franchet*).

Europe, Orient, Japan, and North America.

1. *Najas graminea*, *Delile, Fl. Egypte*, p. 282; *Kunth, Enum. Pl.* iii. p. 115; *A. Br. in Seem. Journ. Bot.* ii. (1864) p. 278; *Hook. f. Fl. Brit. Ind.* vi. p. 569; *K. Schum. in Mart. Fl. Bras.* iii. III. p. 730, t. 124. fig. 2; *Rendle, in Trans. Linn. Soc.* ser. 2, *Bot.* v. (1899) p. 424, t. 42. figg. 192-201.

*Najas minor*, *Benth. Fl. Hongk.* p. 345, non *Allioni*.

HONGKONG (*Wright*, 511!). *Mus. Brit.*; *Herb. Kew.*

Warmer parts of the Old World.

2. *Najas marina*, *Linn. Sp. Pl.* ed. 1, p. 1015.

*Najas major*, *Allioni, Fl. Pedem.* ii. p. 221.

Forma  $\alpha$ , *Rendle, in Trans. Linn. Soc.* ser. 2, *Bot.* v. (1899) p. 392?

CHIHLI: Peking (*Bretschneider*, 892!); COREA (*Faurie*!).  
Herb. Kew.

Forma  $\beta$ , *Rendle, in Trans. Linn. Soc.* ser. 2, *Bot.* v. (1899) p. 392.

KIANGSU: Kiangnang (*Staunton*!). Mus. Brit.  
Europe, India, Japan, North America, Australia.

### 157. ERIOCAULACEÆ.

(By C. H. WRIGHT, A.L.S.)

1. *Eriocaulon alpestre*, *Hook. f. et Thoms. ex Koern. in Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 163; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 99; *Maxim. Diagn. Pl. Asiat.* viii. p. 24; *Hook. f. Fl. Brit. Ind.* vi. p. 578; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 95.

CHEKIANG: Ningpo Mountains (*Faber ex Ruhland*).  
Himalayas.

2. *Eriocaulon australe*, *R. Br. Prodr.* p. 254; *Steud. Syn. Pl. Glum.* ii. p. 273; *Koern. in Linnæa*, xxvii. (1854) p. 686; *Kunth, Enum. Pl.* iii. p. 569; *Benth. Fl. Austral.* vii. p. 192; *Benth. Fl. Hongk.* p. 382; *Maxim. Diagn. Pl. Asiat.* viii. p. 27; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 66.

KWANGTUNG: Canton, White Cloud Hills (*Sampson*, 363!), Macao (*Vachell*!); HONGKONG (*Wright*, 549! *Hance*, 490!), Mount Victoria (*Lamont*, 818!). Mus. Brit.; Herb. Kew.

Australia: Queensland and New South Wales.

3. *Eriocaulon Buergerianum*, *Koern. in Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 163; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 100; *Maxim. Diagn. Pl. Asiat.* viii. p. 20; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 94.

KIANGSI: Kiukiang (*Shearer! Forbes*, 191!); HUPEH: Ichang (*A. Henry*, 37, 2767!), Nanto (*A. Henry*, 4424!); KWANGTUNG: Lofaushan (*Ford*, 135 pro parte!). Mus. Brit.; Herb. Kew.

Japan.

The apical cilia of the female petals appear to be caducous.

4. *Eriocaulon cinereum*, *R. Br. Prodr.* p. 254; *Kunth, Enum.* iii. p. 571; *Steud. Syn. Pl. Glum.* ii. p. 273; *Koern. in Linnæa*, xxvii. (1854) p. 618; *Benth. Fl. Austral.* vii. p. 193; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 111.

*Eriocaulon heteranthum*, *Benth. Fl. Hongk.* p. 382; *Maxim. Diagn. Pl. Asiat.* viii. p. 12.

HUPEH (*A. Henry*, 7786 partly!); KWANGTUNG: Canton (*Sampson*!), on way to Pakuan Mountains (*Hance*, 9666!); HONGKONG (*Wilford*, 293! *Wright*, 552! *Ford*, 74!). Mus. Brit.; Herb. Kew.

Australia.

5. *Eriocaulon cristatum*, *Mart. in Wall. Pl. As. Rar.* iii. p. 28, var. *brevicalyx*, *C. H. Wright*. ♀ petala sepalis duplo longiora.

*Eriocaulon cristatum*, *Benth. Fl. Hongk.* p. 382; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 84, partly.

HONGKONG (*Hance*, 795! *Forbes*, 465!), near Aberdeen (*Sampson*!). Mus. Brit.; Herb. Kew.

The type occurs in Northern India and Japan.

6. *Eriocaulon echinulatum*, *Mart. in Wall. Pl. As. Rar.* iii. p. 29; *Kunth, Enum. Pl.* iii. p. 568; *Steud. Syn. Pl. Glum.* ii. p. 272; *Koern. in Linnæa*, xxvii. (1854) p. 619; *Hance, in Journ. Bot.* xvi. (1878) p. 14; *Hook. f. Fl. Brit. Ind.* vi. p. 579; *Maxim. Diagn. Pl. Nov. Asiat.* viii. p. 11; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 106.

KWANGTUNG: Canton (*Sampson*, 958! [13453 in herb. *Hance*], *Ford*, 363!). Mus. Brit.; Herb. Kew.

Malay Peninsula.

7. *Eriocaulon Faberi*, *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 95.

CHEKIANG: Ningpo Mountains (*Faber ex Ruhland*).

8. *Eriocaulon Henryanum*, *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 86.

YUNNAN: Mengtze, 5000 ft. (*A. Henry*, 9443!). Herb. Kew.

9. *Eriocaulon luzulæfolium*, *Mart. in Wall. Pl. As. Rar.* iii. p. 28; *Cat. n.* 6071; *Kunth, Enum. Pl.* iii. p. 553; *Schnizl. Ic. t.* 46. fig. 2; *Steud. Syn. Pl. Glum.* ii. p. 270; *Koern. in*

*Linnæa*, xxvii. (1854) p. 636; *Hook. f. Fl. Brit. Ind.* vi. p. 582; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 88.

CHEKIANG: Ningpo Mountains (*Faber*, 90!); FORMOSA: Takow (*A. Henry*, 1801!); KWANGTUNG: Canton, swamp at the foot of the White Cloud Hills (*Sampson*, 127, 255, 7423 in herb. *Hance!*). Herb. Kew.

Throughout India and Ceylon.

10. *Eriocaulon Miquelianum*, *Koern. in Miq. Ann. Mus. Bot. Lugd.-Bat.* iii. p. 162; *Franch. & Savat. Enum. Pl. Jap.* ii. p. 99; *Maxim. Diagn. Pl. Asiat.* viii. p. 17; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 92.

CHEKIANG: Tientai Mt. (*Faber*, 88!), Ningpo Mountains (*Faber!*). Herb. Kew.

Japan.

11. *Eriocaulon nipponicum*, *Maxim. Diagn. Pl. Nov. Asiat.* viii. p. 9; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 38.

SHANTUNG: Chefoo Mts. (*Faber*, 89! 205, ex *Ruhland*). Herb. Kew.

12. *Eriocaulon officinale*, *Koern. in Mart. Fl. Bras.* iii. i. p. 480 in obs.; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 117.

CHINA (ex *Koernicke*).

13. *Eriocaulon setaceum*, *Linn. Sp. Pl.* ed. 1, p. 87; *Kunth, Enum. Pl.* iii. p. 550; *Steud. Syn. Pl. Glum.* ii. p. 270; *Benth. Fl. Hongk.* p. 383; *Thwaites, Enum. Pl. Zeyl.* p. 341; *Maxim. Diagn. Pl. Asiat.* viii. p. 13; *Hook. f. Fl. Brit. Ind.* vi. p. 572; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 89, excl. syn.

*Eriocaulon intermedium*, *Koern. in Linnæa*, xxvii. (1854) p. 601.

KWANGTUNG: Canton (*Sampson!*); HONGKONG (*Wright*, 553!). Mus. Brit.; Herb. Kew.

Malabar, Ceylon.

This species is stated by *Kurz* (*Journ. Bot.* xi. 1873, p. 193) to have been collected in Yunnan by *J. Anderson*, but I have seen no specimens to confirm this.

14. *Eriocaulon Sieboldianum*, *Sieb. & Zucc. ex Steud. Syn. Pl. Glum.* ii. p. 272; *Hook. f. Fl. Brit. Ind.* vi. p. 577; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 111, excl. syn.

*Eriocaulon sexangulare*, var.  $\beta$ , *Koern. in Linnæa*, xxvii. (1854) p. 613.

*Eriocaulon sexangulare*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 162, pro parte; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 99, pro parte; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1872) p. 130; *Maxim. Diagn. Pl. Asiat.* viii. p. 10; non *Linn.*

KWANGTUNG: Canton (*Sampson & Hance*, 9666!), Kowlung (*Sampson*, 7977! *Lamont*!). *Mus. Brit.*; *Herb. Kew.*

India, Tonkin, Philippines, Japan.

This species has the inner, or floral, bracts glabrous or nearly so, while those of *E. sexangulare*, *Linn.*, are bearded and villous. I have not seen the specimen collected by David in Kiukiang and placed by Franchet (*Pl. David.* i. p. 316) under *E. sexangulare*, *Linn.*

15. *Eriocaulon sinicum*, *Miq. Journ. Bot. Néerl.* i. (1861) p. 87; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 116.

KWANGTUNG: Fukwing (*Krone ex Miquel*).

16. *Eriocaulon truncatum*, *Buch.-Ham. in Wall. Pl. As. Rar.* iii. p. 29; *Wall. Cat.* n. 6076; *Kunth, Enum. Pl.* iii. p. 555; *Steud. Syn. Pl. Glum.* ii. p. 270; *Koern. in Linnæa*, xxvii. (1854) p. 631; *Benth. Fl. Hongk.* p. 382; *Maxim. Diagn. Pl. Asiat.* viii. p. 14; *Hook. f. Fl. Brit. Ind.* vi. p. 578; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 107.

CHEKIANG: Ningpo Mountains (*Faber*, 19!); KWANGTUNG: Lofaushan (*Ford*, 135 pro parte!), Canton (*Sampson*, in hb. *Hance*, 7423!); HONGKONG (*Wilford*, 102! *Wright*, 551!), race-course (*Lamont*, 819!), Wongneichung (*Hance*!). *Mus. Brit.*; *Herb. Kew.*

India, Ceylon, and Malaya.

17. *Eriocaulon Wallichianum*, *Mart. in Wall. Pl. As. Rar.* iii. p. 26, t. 249; *Kunth, Enum. Pl.* iii. p. 564; *Steud. Syn. Pl. Glum.* ii. p. 271; *Koern. in Linnæa*, xxvii. (1854) p. 687; *Benth. Fl. Hongk.* p. 381; *Bretschn. Early Res.* p. 181; *Maxim. Diagn. Pl. Asiat.* viii. p. 27.

*Eriocaulon longifolium*, *Koern. in Linnæa*, xxvii. (1854) p. 690, ex parte, non *Nees*; *Benth. Fl. Hongk.* p. 382, in adnot.; *Maxim. Diagn. Pl. Asiat.* viii. p. 28; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 41.

*Eriocaulon cantoniensis*, *Hook. et Arn. Bot. Beechey's Voyage*, p. 219.

*Eriocaulon sexangulare*, *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 110, pro parte.

CHINA (*Beechey*!); SOUTH CHINA (*Seemann*, 2400!); KIANGSI: Kiukiang (*Forbes*, 1055!); KWANGTUNG: Syngmun (*Meyen*!), Lofaushan (*Ford*!), Macao (*Gaudichaud*, 391! *Millett & Vachell ex Hooker & Arnott, Hance*, 10138!); HONGKONG (*Hinds*! *Fortune*, 120! *Wright*, 550! *Wilford*, 102 partly! *Hance*, 540, 541!), Happy Valley (*Lamont*, 817!); HAINAN (*A. Henry*, 8073!), Hoihau (*Bullock*!). Mus. Brit.; Herb. Kew.

India.

“*E. cantoniense*” is said to be used medicinally. See Kew Bulletin, 1898, p. 144.

18. *Eriocaulon Wightianum*, *Mart. in Wall. Pl. As. Rar.* iii. p. 28; *Wall. Cat.* n. 6067; *Kunth, Enum. Pl.* iii. p. 563; *Steud. Syn. Pl. Glum.* ii. p. 271; *Koern. in Linnæa*, xxvii. (1854) p. 658; *Hook. f. Fl. Brit. Ind.* vi. p. 576; *Ruhland, in Engl. Pflanzenr., Eriocaul.* p. 75.

KWANGTUNG: Pakhoi (*Playfair*, 113!). Herb. Kew.

India and Ceylon.

## 158. CYPERACEÆ.

(By C. B. CLARKE, F.R.S., F.L.S.)

1. *Pycneus angulatus*, *Nees, in Linnæa*, ix. (1834) p. 283; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 593.

*Cyperus unioloïdes*, *R. Br. Prodr.* p. 216; *Kunth, Enum. Pl.* ii. p. 112.

*Cyperus angulatus*, *Nees, in Wight, Contrib.* p. 73; *Boeck. in Linnæa*, xxxv. (1867-8) p. 465.

YUNNAN: Nienkiatze, prope Tapintze (*Delavay*, 1750!).

Scattered throughout the warmer regions of the whole world.

Var.  $\beta$ . *Wightii*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 593.

*Cyperus latispicatus*, *Boeck. MS. in Wight, Exsicc.* n. 2874, in herb. Berlin, nec *Boeck. in Flora*.

YUNNAN: Tapintze (*Delavay*, 1748!).

Madras.

2. ***Pycreus Baccha***, *Nees, in Linnæa*, ix. (1834) p. 283, *et in Mart. Fl. Bras.* ii. pars i. p. 10 in obs. [Bacha].

*Pycreus puncticulatus*, *Nees, in Mart. Fl. Bras.* ii. pars i. p. 10 in obs.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 593.

*Cyperus puncticulatus*, *Vahl, Enum.* ii. p. 348; *Kunth, Enum. Pl.* ii. p. 15; *Boeck. in Linnæa*, xxxv. (1867-8) p. 472, syn. *Roxburghii* deleto.

*Cyperus Baccha*, *Kunth, Enum. Pl.* ii. p. 115.

*Cyperus inundatus*, *Nees, in Wight, Contrib.* p. 76, non *Roxb.*

SHANTUNG: Chefoo (*Forbes!*).

Inhabits South India with Ceylon. This plant is, however, often passed by as the common *Cyperus pilosus*, *Vahl*, and not collected.

3. ***Pycreus Delavayi***, *C. B. Clarke*, n. sp.

Culmi cæspitiosi, 12-16 uncias longi, graciles, submonocephali. *Folia* 6-10 uncias longa, subulata. *Bracteæ* 2, ima usque ad 2 uncias longa, fere subulata. *Caput* 4-10-stachyum, laxum. *Spiculæ* subspicatae,  $\frac{1}{2}$  unciam longæ,  $\frac{1}{8}$  unciam latæ, 12-nucigeræ, lucide atro-castaneæ, compressæ. *Glumæ* naviculares, ovatae, vix acutæ, subdistantes, proventu paullo inflatæ. *Stylus* perbrevis; rami 2 longissimi. *Nux* lateraliter compressa, cum  $\frac{1}{2}$  parte glumæ vix æquilonga, ellipsoidea, nigrescens, levis; celluli extimi quadrato-hexagoni.

YUNNAN: Tali (*Delavay*, 413, in hb. *Paris.*!).

This species is near *P. globosus*, *Reichb.*, var.  $\beta$ . *nilagirica*, but the spikelets are larger, with loosely-packed glumes. The plant more resembles *P. Cooperi*, *C. B. Clarke* (in *Dyer, Fl. Cap.* vii. p. 160).

4. ***Pycreus globosus***, *Reichb. Fl. Excurs.* [1830-32], *Addend.* p. 140. n. 10.

*Pycreus capillaris*, *Nees, in Linnæa*, ix. (1834) p. 283; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 591; *Palibin, Fl. Koreæ*, iii. p. 20; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

*Cyperus globosus*, *Allioni, Fl. Pedem. Auctuarium*, p. 49; *Reichb. Ic. Pl. Crit.* iii. p. 26, t. 229, syn. *Beauv. excl.*, *et Ic. Fl. Germ.* viii. p. 32, t. 279. fig. 665; *Boeck. in Linnæa*, xxxv. (1867-8) p. 458; *Hance, in Journ. Bot.* xvii. (1879) p. 16; *Boeck. in Engl. Jahrb.* vi. (1885) p. 50.

*Cyperus capillaris*, *Roxb. Fl. Ind.* i. p. 198; *Hook. et Arn. Bot. Beechey's Voy.* p. 220.

*Cyperus vulgaris*, *Kunth, Enum. Pl.* ii. p. 4, syn. *C. conglomerato* aliisque exclusis; *Benth. Fl. Hongk.* p. 385; *Regel, Fl. Ussur.* p. 175; *Sieber, MS. ex Kunth.*

*Cyperus complanatus*, *Presl, in Oken, Isis*, xxi. (1882) p. 271; *Franch. Pl. David.* i. p. 316, et in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 261; cf. *Maxim. in Bull. Soc. Nat. Mosc.* liv. pars i. (1879) p. 63.

CHIHLI: Peking (*David!*); SHANTUNG: Chefoo (*Fauvel!*); FORMOSA (*Siebold!*), Takow (*A. Henry, 560!*); HUPEH (*A. Henry, 7520!*); KWANGTUNG: Macao (*Vachel!*); LUCHU ARCHIPELAGO (*Doederlein*). Mus. Brit.; Herb. Kew.

The North China and Japanese examples are nearly as the Mediterranean *C. vulgaris*, *Kunth* (which is widely spread in Africa and India), with rather broad spikelets, chestnut marked with green; the Southern Chinese examples have the spikelets slenderer and browner, and are the *C. complanatus*, *Presl*, which extends from the Mascarene Islands to New South Wales and Japan. I can draw no definite line between *Pycneus globosus* and *Cyperus complanatus*, nor between these and the three following varieties.

Var.  $\beta$ . *nilagirica*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 592; spiculis angustioribus, magis duris, non inflatis, sæpe castaneis interdum nigris.

*Cyperus nilagiricus*, *Hochst. ex Steud. Syn. Pl. Cyp.* p. 2; *Boeck. in Linnæa*, xxxv. (1867-8) p. 457.

NORTH CHINA (*Hance, 1387*); SHANTUNG (*Maingay, 152!*); KIANGSU: Shanghai (*Maingay, 637!*); CHEKIANG: Ningpo Mts. (*Faber, 223!*); HUPEH: Ichang (*A. Henry, 2574!*); NORTH CENTRAL CHINA: Kusan, Kisaoshian (*Hugh!*), Mt. Ngosan, Laoysan (*Hugh!*), Mt. Thaepeisan (*Hugh!*); SHENSI: Mt. Kifongsan (*Hugh!*); SZECHUEN: Tchenkeoutin (*Farges!*); YUNNAN: Tapintze (*Delavay, 415, 1753, 1754!*), Hokin, 5000 feet (*Delavay, 1751!*), Mosoyn (*Delavay, 1766!*), Mengtze, — 4500 feet (*A. Henry, 9119!*), Red River, 6000 feet (*A. Henry, 9640!*); KWANGTUNG: Canton (*Sampson, 1387!*). Mus. Brit.; Herb. Kew.

This var. extends from South and Equatorial Africa to Amurland, Malaya, Australia; it is especially a mountain form.



Var.  $\gamma$ . *stricta*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 592; spiculis abbreviatis, sæpius pallidis, non raro curvatis.

*Pycrus* Linneanus, *Nees*, in *Linnæa*, ix. (1834) p. 283.

*Cyperus strictus*, *Lam. Ill.* i. p. 146; *Kunth, Enum. Pl.* ii. p. 12.

*Cyperus tortuosus*, *Roxb. Fl. Ind.* i. p. 197 [ed. Wallich, p. 201]; *Kunth, Enum.* ii. p. 16.

*Cyperus* Linnæanus, *Kunth, Enum. Pl.* ii. p. 115.

CHIHLI: Peking (*Bretschneider*, 789!); SHANTUNG: Chefoo (*Faber*, 61! *Schottmueller*, 342!); HUPEH: Ichang (*A. Henry*, 83, 83 A!); YUNNAN: Tapintze (*Delavay*, 402!). *Mus. Brit.*; *Herb. Kew.*

This var., frequent in India, is scattered from South Africa and Persia to Japan and the Philippines.

Var.  $\delta$ . *erecta*, *C. B. Clarke*, var. nova; culmo foliisque robustioribus; umbella sæpe composita, umbellularum radiolis 3–5 usque ad 10–15 mm. longis; spiculis angustis, pallidis, sæpe erectis.

*Cyperus erectus* (sp.), *Roxburgh, MS.*

FORMOSA: Tamsui (*Oldham*, 581!); KWANGTUNG: Canton (*Sampson*, 933!). *Mus. Brit.*; *Herb. Kew.*

The same variety (or form) collected in India by Koenig and Roxburgh.

5. *Pycrus latespicatus*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 590.

*Cyperus latespicatus*, *Boeck. in Flora*, xlii. (1859) p. 433 bis, et in *Linnæa*, xxxv. (1867–8) p. 467.

YUNNAN: Tapintze (*Delavay*, 1752!), Mosoyn (*Delavay*, 1761, pro magna parte!), Milê (*A. Henry*, 9119!). *Herb. Kew.*

Frequent in Northern India.

6. *Pycrus polystachyus*, *Beauv. Fl. Owar.* ii. (1807) p. 48, t. 86. fig. 2; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* (1843) p. 55; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 592.

*Cyperus polystachyus*, *R. Br. Prodr.* p. 214; *Kunth, Enum. Pl.* ii. p. 13; *Hook. et Arn. Bot. Beechey's Voy.* p. 220; *Benth. Fl. Hongk.* p. 385; *Boeck. in Linnæa*, xxxv. (1867–8) p. 477; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, non *Rottb.*

*Cyperus odoratus*, *Linn. Sp. Pl.* ed. 1, p. 46, ed. 2, p. 68, et herb. propr., pro majore parte.

*Cyperus teretifructus*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62.

*Cyperus brunneus*, *Hook. et Arn. Bot. Beechey's Voy.* p. 99, non *Swartz.*

SHANTUNG (*Maingay*, 155 !); Chefoo (*Fauvel* !); KIANGSU : Shanghai (*Maingay*, 638 !); FORMOSA (*A. Henry*, 1047 !), Takow (*Playfair*, 135 ! *A. Henry*, 1060 !); KWANGTUNG (*Sampson*, 258 !), Lofaushan (*Ford* !), Macao (*Vachell*, 70 ! *da Silva* !), Whampoa (*Hance*, 1230 !); HONGKONG (*C. Wright*, 555 ! *Faurie*, 15800 !). *Mus. Brit.*; *Herb. Kew.*

In nearly all warm countries; especially common near the sea.

7. ***Pycrus pulvinatus***, *Nees, in Linnæa*, ix. (1834) p. 283.

*Pycrus nitens*, *Nees, in Linnæa*, ix. (1834) p. 283, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* (1843) p. 53; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 591.

*Cyperus pumilus*, *Linn. Amœn. Acad.* iv. (1759) p. 302, et *Sp. Pl.* ed. 2, p. 69; *Gaertn. Fruct.* i. p. 9, t. 2. fig. 2.

*Cyperus nitens*, *Retz. Obs.* v. (1789) p. 13 ?; *Kunth, Enum. Pl.* ii. p. 3; *Boeck. in Linnæa*, xxxv. (1867-8) p. 483.

*Cyperus pulvinatus*, *Nees, in Wight, Contrib.* p. 74; *Nees et Meyen, MS. e Nees.*

*Cyperus gymnoleptus*, *Steud. Syn. Pl. Cyp.* p. 3.

— *Pluk. Alm.* p. 179, t. 191. fig. 8.

KWANGTUNG : Canton, Lintin I. (*Meyen* !), North River (*Nevin* !), Pakhoi (*Playfair*, 204 !). *Mus. Brit.*; *Herb. Kew.*

In tropical and subtropical Africa and Asia, frequent; extending to Malaya and Australia. Also received from Florida and Martinique.

8. ***Pycrus sanguinolentus***, *Nees, in Linnæa*, ix. (1834) p. 283; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 590.

*Cyperus sanguinolentus*, *Vahl, Enum.* ii. p. 351; *Debeaux, Fl. Tchef.* p. 142.

*Cyperus Eragrostis*, *Vahl, Enum.* ii. p. 322, syn. *Retzii* excl. ?; *Roem. et Schult. Syst.* ii. p. 180; *Kunth, Enum. Pl.* ii. p. 7 (cf. p. 537); *Benth. Fl. Hongk.* p. 385; *Boeck. in Linnæa*, xxxv. (1867-8) p. 443, cum varr.; *Hance, in Journ. Bot.* xvii. (1879) p. 16; *Maxim. Fragm.* p. 64; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 26, et *Pl. David.* i. p. 316.

*Cyperus flavescens*, var. *rubro-marginata*, *Schrenk, Enum. Pl. Nov.* i. p. 3; *Regel, Fl. Ussur.* p. 174.

*Cyperus atratus*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62; *Boeck. in Linnæa*, xxxv. (1867-8) p. 446.

*Cyperus Teysmanni*, *Boeck. in Flora*, lviii. (1875) p. 259.

*Cyperus* sp., n. 36, *Rottb. Descr. et Ic.* p. 29, t. 9. fig. 4.

CHIHLI: Peking (*David!*); SHINGKING: Moukden (*James!*); SHANTUNG (*Maingay*, 156!), Chefoo (*Schottmueller*, 343!); KIANGSU: Kiangnang (*Staunton!*); FOKIEN: Amoy (*Hance*, 1387!), Tachiaotze (*Hance*, 3496!); FORMOSA (*Oldham*, 586!), Takow (*A. Henry*, 1120!); HUPEH: Ichang (*A. Henry*, 2907!), Patung (*A. Henry*, 5019 partim!); SZECHUEN: Tchenkeoutin, 4000 feet (*Farges*, 1131!); YUNNAN: Tapintze (*Delavay*, (1770!); KWANGTUNG (*Yvan! Sampson!*); Lofaushan (*Ford*, 122!), Macao (*da Silva*); HONGKONG (*Wilford*, 312!); HAINAN (*A. Henry*, 7949!). *Mus. Brit.*; *Herb. Kew.*

Common in Asia, alt. 0–9000 feet, extending from Amurland to Abyssinia and Australia.

1. *Juncellus inundatus*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 595.

*Cyperus inundatus*, *Roxb. Hort. Beng.* (1814) p. 6, et *Fl. Ind.* i. p. 201 (ed. Wallich, p. 205); *Hassk. in Flora*, xlv. (1862) p. 191, nec *R. Br.*, nec *Nees*.

*Cyperus Iria*, *Linn. Sp. Pl.* p. 67 partim; *Hassk. in Flora*, xlv. (1861) p. 743.

*Cyperus diluvialis*, *Schultes*, in *Roem. et Schult. Syst.* ii. *Mant.* p. 124.

— *Rheede, Hort. Mal.* xii. p. 105, t. 56.

SHINKING (*Ross*, 50!); KIANGSU: Shanghai (*Maingay*, 639!); FORMOSA (*A. Henry*, 702 A!), Takow (*A. Henry*, 1089, 1117, 1130, 1163!). *Herb. Kew.*

Bengal.

This plant, from its large size and flowering the first year, is often mixed with *C. digitatus*, *Roxb.*, and other large annual species of *Cyperus*. It appears to me closely allied to *Juncellus serotinus*, which differs by its long stolons, and the umbel usually smaller, often with the rhachilla of the spikes hairy.

2. *Juncellus pygmæus*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 596.

*Cyperus pygmæus*, *Rottb. Descr. et Ic.* p. 20, t. 14. figg. 4, 5; *Kunth, Enum. Pl.* ii. p. 18; *Boeck. in Linnæa*, xxxv. (1867–8) p. 493, var.  $\beta$  excl.; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 102.

*Isolepis Micheliana*, *Stev. in Mém. Soc. Mosc.* v. (1817) p. 356; *Regel, Fl. Ussur.* p. 178, non *Roem. et Schult.*

KIANGSU: Shanghai (*Maingay*, 634!).

Frequent throughout Boissier's Orient Region and in North India; thence scattered to Ussuri, the Philippines, and Queensland.

This plant closely resembles *Scirpus Michelianus*, Linn.; so closely, that Bentham and Boeckeler consider the two identical. Hance's Pekin *Cyperus pygmæus* is *Scirpus Michelianus*, Linn., while Maack's Ussuri *S. Michelianus* is *Juncellus pygmæus*.

3. *Juncellus serotinus*, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 594.

*Cyperus serotinus*, *Rottb. Descr. et Ic.* (1773) p. 31; *Kunth, Enum. Pl.* ii. p. 19; *Boeck. in Linnæa*, xxxv. (1867-8) p. 492; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 316; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

*Cyperus Monti*, *Linn. f. Suppl.* p. 102; *Host, Gram. Austr.* iv. p. 38, t. 67; *Regel, Fl. Ussur.* p. 175; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 88.

*Cyperus Japonicus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 140.

CHIHLI: Jehol (*David*), Peking (*Bretschneider*, 787! *Hance*, 6531!); SHINGKING: Moukden (*Webster*, 215!); SHANTUNG (*Maingay*, 157!), Chefoo (*Forbes! Fauvel!*); FORMOSA: Takow (*A. Henry*, 702, 1842!); SHENSI: Paokesan, Mt. Kifonsan (*Hugh!*); HUPEH: Ichang (*A. Henry*, 2249!), Hanyang (*Faber*, 1512!); YUNNAN: Mosoyu (*Delavay*, 1767, 2717!); KWANG-TUNG: Canton (*Macartney*). *Mus. Brit.*; *Herb. Kew.*

From South Europe to North India, Amurland, Japan.

1. *Cyperus albidus*, *Retz. Obs.* vi. p. 21; *Kunth, Enum. Pl.* ii. p. 101.

*Cyperus Retzii*, *Poir. in Lam. Encyc.* vii. p. 255.

CHINA (fide *Retz.*).

Not seen.

2. *Cyperus amuricus*, *Maxim. ! Prim. Fl. Amur.* (1859) p. 296; *Boeck. in Linnæa*, xxxv. (1867-8) p. 607; *Palibin, Fl. Koreæ*, iii. p. 19.

*Cyperus aristatus*, var. ?  $\beta$ . *Maingayi*, C. B. Clarke, in *Journ. Linn. Soc. Bot.* xxi. (1884) p. 94.

*Cyperus Iria*, var. (*amabilis*, *glumis submucronatis*), C. B. Clarke, in *Journ. Linn. Soc., Bot.* xxi. (1884) p. 138 in obs.

*Cyperus Iria*, *Linn.*, forma *japonica*, *Boeck. ! MS.*

CHIHLI : Peking (*Möllendorff!* *Hance*, 1147 pro parte !); SHINGKING (*Ross*, 86 !); SHANTUNG (*Maingay*, 84 !); SHENSI : Paokisan, Miaouansan (*Hugh!*); SZECHUEN : Tchenkeoutin (*Farges!*); COREA (*Bunge*). Mus. Brit.; Herb. Kew.

Amurland, Mandshuria, Japan.

3. *Cyperus aristatus*, *Rottb. Descr. et Ic.* (1773) p. 23, t. 6. fig. 1; *Kunth. Enum. Pl.* ii. p. 23; *Boeck. in Linnæa*, xxxv. (1867-8) p. 500; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 606.

*Cyperus squarrosus*, *Linn. Amœn. Acad.* iv. (1759) p. 303, et *Sp. Pl.* ed. 2, p. 66 partim.

*Scirpus intricatus*, *Linn. Mant.* p. 182 et herb. propr. !

YUNNAN : Mosoyn (*Delavay*, 1760 !).

A small weed, abundant in the warmer parts of Africa, India, Australia, and America, but apparently wanting in Malaya and China, except in the west.

4. *Cyperus castaneus*, *Willd.*; *Nees, in Wight, Contrib.* p. 79 (cum char. et syn.); *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 221.

CHINA (*Vachell*, n. 296).

Not seen. Hooker & Arnott add that the Chinese specimen of this which they had seen came from Prof. Henslow.

*Vachell* n. 296 may have been the true *C. castaneus*, *Willd.* (cf. *C. B. Clarke, in Hook. f. Fl. Ind.* vi. p. 598), but that is not a common plant, nor have I seen it from China. I think that *Vachell* n. 296 may have been *C. amuricus*, *Maxim.* = *Maingay* n. 84.

5. *Cyperus cephalotes*, *Vahl, Enum.* ii. p. 311; *Kunth, Enum. Pl.* ii. p. 48; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 597.

*Cyperus monogynus*, *Boeck. in Linnæa*, xxxv. (1867-8) p. 565.

*Scirpus glomeratus*, *Linn. Sp. Pl.* ed. 1, p. 52; *Hassk. in Flora*, xliv. (1861) p. 743.

*Schoenus coloratus*, var.  $\beta$ , *Linn. Sp. Pl.* ed. 2, p. 64.

*Anosporum monocephalum*, *Nees, in Wight, Contrib.* p. 92; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 411.

*Hydroschoenus kyllingioides*, *Moritzi, Syst. Verz. Zoll.* p. 95.

*Trentepohlia bifoliata*, *Boeck. in Bot. Zeit.* xvi. (1858) p. 249.

— *Rheede, Hort. Mal.* xii. p. 99, t. 53.

FOKIEN : Amoy (*Jenkins!*).

From Assam to Ceylon and North Australia.

6. *Cyperus compressus*, *Linn. Sp. Pl.* ed. 1, p. 46, ed. 2, p. 68; *Rottb. Descr. et Ic.* p. 27, t. 9. fig. 3; *Kunth, Enum. Pl.* ii. p. 23; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 56; *Boeck. in Linnæa*, xxxv. (1867-8) p. 517; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 605; *Henriq. in Bol. Soc. Brotero*, ii. p. 166.

*Cyperus pectiniformis*, *Nees, in Wight, Contrib.* p. 77 (*syn.* *C. pectinato*, *Roxb. excl.*); *Hook. et Arn. Bot. Beechey's Voy.* p. 221.

*Cyperus Meyenii*, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 57; *Nees et Arn. MS. fide Nees.*

FOKIEN: Amoy (*Sampson*, 257!); FORMOSA (*Campbell! A. Henry*, 1869!); SZECHUEN: Tcentosen, Mt. Uomisan (*Hugh!*); KWANGTUNG: Macao (*Tate!*), Whampoa (*C. Wright*, 556!); HAINAN (*A. Henry*, 8242!). *Mus. Brit.*; *Herb. Kew.*

In warm countries throughout the World, frequent.

7. *Cyperus dichroostachyus*, *Hochst. in Flora*, xxiv. (1841) pars i., *Intell.* p. 21; *Boeck. in Linnæa*, xxxv. (1867-8) p. 587; *C. B. Clarke, in Dyer, Fl. Trop. Afr.* viii. pp. 311, 331, with *syn.*

YUNNAN: south of Red River, from Maumei, 6000 feet (*A. Henry*, 9473!), Yunnanfu (*Ducloux*, 31!). *Herb. Kew.*

Africa and Madagascar.

8. *Cyperus difformis*, *Linn. Amœn. Acad.* iv. (1759) p. 302, *et Sp. Pl.* ed. 2, p. 67; *Rottb. Descr. et Ic.* p. 24, t. 9. fig. 2; *Benth. Fl. Hongk.* p. 385; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 89; *Debeaux, Fl. Tchef.* p. 144; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, *et Pl. David.* i. p. 317; *Kunth, Enum. Pl.* ii. p. 38; *Boeck. in Linnæa*, xxxv. (1867-8) p. 586; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 599.

*Cyperus fuscus*, *Maxim. Fragm.* p. 64? *non Linn.*

CHIHLI: Peking (*Hance!*); SHINKING (*Ross*, 85!), Neuchang (*Webster*, 314!); SHANTUNG (*Maingay*, 87!); KIANGSU: Shanghai (*Maingay*, 645! 646!); KIANGSI: Kiukiang (*David!*); FOKIEN: Amoy (*Hance*, 995!); FORMOSA: Tamsui (*Oldham*, 582!); SZECHUEN: Tcentosen, Mt. Uomisan (*Hugh!*); HUPEH: Ichang (*A. Henry*, 2370!); YUNNAN: Mengtze, 4600 feet (*A. Henry*, 11318!), Kiangyu (*Delavay*, 3078!), Hokin (*Delavay*, 1759!); KWANGTUNG: Canton (*Sampson*, 464!); HONGKONG (*C. Wright*, 558! *Harland*). *Mus. Brit.*; *Herb. Kew.*

The tropics and warm temperate zones of the Old World—a rice-field weed.

9. *Cyperus diffusus*, *Vahl*, *Enum.* ii. p. 321; *Kunth*, *Enum. Pl.* ii. p. 30; *Nees*, in *Nov. Act. Acad. Nat. Cur.* xix. *Suppl.* i. (1843) p. 58; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 603.

*Cyperus elegans*, *Swartz*, *Obs. Bot.* p. 30; *Kunth*, *Enum. Pl.* ii. p. 28; *Boeck.* in *Linnæa*, xxxv. (1867-8) p. 532, non *Linn.*

*Cyperus moestus*, *Kunth*, *Enum. Pl.* ii. p. 31.

*Cyperus nigro-viridis*, *Thwaites*, *Enum. Fl. Zeyl.* p. 344; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 131.

CHEKIANG: Chusan (*Yvan!*); KWANGTUNG: Whampoa (*Sampson*, *hb. Hance*, 2149!); HONGKONG (*Harland!*). *Mus. Brit.*; *Herb. Kew.*

The tropics in South-east Asia and in America; a sylvan species.

10. *Cyperus digitatus*, *Roxb.* *Hort. Beng.* p. 81, et *Fl. Ind.* (ed. *Wall. et Carey*) i. p. 209; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 618.

*Cyperus auricomus*, *Benth.* *Fl. Austral.* vii. p. 286.

*Cyperus venustus*, *Nees*, in *Flora*, xi. (1828) p. 333, et in *Wight Contrib.* p. 86; non *R. Br.*

*Cyperus Neesii*, *Kunth*, *Enum. Pl.* ii. p. 101, syn. *Linn.* excl.; *Boeck.* in *Linnæa*, xxxvi. (1869-70) p. 315.

*Cyperus verticillatus*, *Nees*, in *Wight, Contrib.* p. 87, pro parte.

FORMOSA: Takow (*A. Henry*, 1862, 1862 A!); YUNNAN: Mengtze, 5000 feet (*A. Henry*, 11320!); KWANGSI (*Wenyon!*), Lungchow (*A. Henry*, 685!). *Herb. Kew.*

Tropic and subtropic regions of the world; abundant in India, frequent in America, rare in Africa.

11. *Cyperus distans*, *Linn. f. Suppl.* p. 103; *Beauv.* *Fl. Owar.* i. p. 35, t. 20; *Benth.* *Fl. Hongk.* p. 387; *Kunth*, *Enum. Pl.* ii. p. 93; *Hook. et Arn.* *Bot. Beechey's Voy.* p. 223; *Boeck.* in *Linnæa*, xxxv. (1867-8) p. 612; *Franch.* in *Bull. Soc. Bot. France*, xxix. (1882) p. 12; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 607; *Linnæus*, *MS.*

FORMOSA (*A. Henry*, 1011, 1041!), Takow (*Playfair* 108!); KWANGTUNG: Lofaushan (*Ford*, 121, 124!), Macao (*d'Incarville!* *Fauvel!*). *Mus. Brit.*; *Herb. Kew.*

Throughout the warmer regions of the world, in rice-fields and ditches.

12. *Cyperus eleusinoides*, *Kunth, Enum. Pl.* ii. p. 39; *Boeck. in Linnæa*, xxxv. (1867-8) p. 596; *Hance, in Journ. Bot.* xvi. (1878) p. 112; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 608.

*Cyperus infra-apicalis*, *Nees, ex Aitch. Cat. Pl. Punjab*, p. 155; *Hance, l. c. in citat.*

FORMOSA: Takow (*Playfair* 56! *A. Henry*, 768, 1078!); YUNNAN: Tapintze (*Delavay*, 1769!), Mengtze 4600 feet (*A. Henry*, 10012!); KWANGTUNG (*Hance*, 19311!). *Mus. Brit.*; *Herb. Kew.*

Frequent in India; scattered west to Abyssinia and Syria, east to the Philippines and Queensland.

13. *Cyperus eriophorus*, *Steud. Syn. Pl. Cyp.* p. 50.

KWANGTUNG: Macao (ex *Steudel*).

Not seen. This has the leaf-sheaths fringed in their mouths and on their margins with woolly hairs, and must be a very remarkable *Cyperus*, if of that genus. Unfortunately, as in so many other cases, the author does not mention those points of structure which define the genus of the plant; so that there remains a doubt whether he looked at these points, which doubt cools the interest that otherwise would attach to such a species.

14. *Cyperus exaltatus*, *Retz. Obs.* v. p. 11; *Kunth, Enum. Pl.* ii. p. 70; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 319; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 617.

*Cyperus venustus*, *R. Br. Prodr.* p. 217.

*Cyperus alopecuroides*, *Roxb. Hort. Beng.* p. 6, et *Fl. Ind.* i. p. 208, nec *Rottb.*

*Cyperus racemosus*, *Boeck. in Flora*, lxii. (1879) p. 555 [neque in *Linnæa*, xxxvi. (1869-70) p. 310]; *Hance, in Journ. Bot.* xvi. (1878) p. 112, non *Retz.*

— *Rheede, Hort. Mal.* xii. p. 77, t. 42.

SHANTUNG (*Staunton*!); KWANGTUNG: Canton (*Sampson*! hb. *Hance*, 9308!), Macao (*Callery*!); HAINAN (*A. Henry*, 32, 8369!). *Mus. Brit.*; *Herb. Kew.*

In tropic and subtropic regions; common in Africa, Asia, and Australia; very rare in America.

15. *Cyperus fuscus*, *Linn. Sp. Pl.* ed. 1, p. 46, ed. 2, p. 69; *Kunth, Enum. Pl.* ii. p. 37; *Hook. in Sowerby, Engl. Bot., Suppl.* t. 2626; *Boeck. in Linnæa*, xxxv. (1867-8) p. 585; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 599.



NORTH CENTAL CHINA: Hukiascien (*Hugh!*); SHENSI: Mt. Kifongsan (*Hugh!*). Mus. Brit.

Throughout Europe and West Asia to the Himalayas.

16. *Cyperus glomeratus*, *Linn. Amœn. Acad.* iv. (1759) p. 301, *et Sp. Pl.* ed. 2, p. 68; *Kunth, Enum. Pl.* ii. p. 77; *Reichb. Ic. Fl. Germ.* t. 284. fig. 675; *Boeck. in Linnœa*, xxxv. (1867-8) p. 592; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 607; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, *et Pl. David.* i. p. 316.

CHIHLI: Jehol (*David!*), Peking (*Williams!*), Tachiaotze (*Möllendorff!*); SHINGKING (*Ross, 34!*); SHANTUNG (*Maingay, 155!*), Tsien Mts. (*Faber, 1514!*). Mus. Brit.; Herb. Kew.

In Europe frequent; scattered thence to Kashmir, North China, and Amurland.

17. *Cyperus Haspan*, *Linn. Sp. Pl.* ed. 2, p. 66 partim, *nec Linn. herb.*; *Kunth, Enum. Pl.* ii. p. 34; *Benth. Fl. Hongk.* p. 386, *et Fl. Austral.* vii. p. 270; *Boeck. in Linnœa*, xxxv. (1867-8) p. 574, var. a; *Bretschn. Early Research.* p. 111; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 600; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

*Cyperus pes-avium*, *Bertol. Misc. Bot.* viii. p. 30, t. 3. fig. 1.

*Cyperus*, n. 37, *Linn. Fl. Zeyl.* p. 15.

*Scirpus*, n. 78, *Rottb. Descr. et Ic.* p. 58, t. 17. fig. 3, syn. *Linn. excl.*

CHEKIANG: Ningpo Mts. (*Faber, 7, 1739!*); SZECHUEN: Tcentosen, Mt. Uomisau (*Hugh!*); \*HUPEH: Ichang (*A. Henry, 2250!*); KWANGTUNG: Canton (*Sampson, 260!*), Macao (*da Silva!*); Pakhoi (*Playfair, 108!*); HONGKONG (*Harland! Wilford, 48! C. Wright, 557! Schottmueller, 449!*). Mus. Brit.; Herb. Kew.

Throughout the world, in tropical and warm temperate regions.

18. *Cyperus Iria*, *Linn. Sp. Pl.* ed. 1, p. 45, ed. 2, p. 67, tab. Rheedei cit. excl.; *Thunb. Fl. Japon.* p. 36; *Kunth, Enum. Pl.* ii. p. 38; *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 223, *et in Nov. Act. Nat. Cur.* xix. *Suppl.* (1843) p. 70; *Benth. Fl. Hongk.* p. 386; *Boeck. in Linnœa*, xxxv. (1867-8) p. 595, *et in Engl. Jahrb.* vi. (1885) p. 51; *Bretschn. Early Research.* p. 111; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 606; *Debeaux, Fl. Tchef.* p. 154, *et in Act. Soc. Linn. Bordeaux*, xxxiii. (1879)

p. 45; *Franch. in Bull. Soc. Bot. de France*, xxix. (1882) p. 12, in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 317; *Henriq. in Bol. Soc. Brotero*, ii. p. 166; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227; *Palibin, Fl. Koreae*, iii. p. 19.

*Cyperus Santonici*, *Rottb. Descr. et Ic.* p. 41, t. 9, fig. 1.

CHIHLI: Peking (*Hance*, 1147 pro parte!), Tientsin (*Debeaux*!); SHANTUNG: Chefoo (*Forbes! Debeaux! Fauvel*!); KIANGSU: Shanghai (*Maingay*, 635!), Kiangnan (*Staunton*!); CHEKIANG: Ningpo Mts. (*Faber*!), Chusan (*Yvan*!); KIANGSI (*David*!); FORMOSA (*A. Henry*, 731!); HUPEH (*E. H. Wilson*, 1657 a!), Ichang (*A. Henry*, 2161, 2770!); SZECHUEN: Tchenkeoutin (*Farges*!); YUNNAN: Tapintze (*Delavay*, 1764!); KWANGTUNG: Canton (*Sampson*, 259!), Macao (*Vachell! Meyen*!), Lienchau (*Lamont*!); HONGKONG (*Wilford*, 266! *Harland*!); COREA (*Kalinowsky*); LUCHU ARCHIPELAGO (*Döderlein*). *Mus. Brit.*; *Herb. Kew.*

Through the warmer regions of the world—a weed in rice-fields.

Var. *paniciformis*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 607.

*Cyperus paniciformis*, *Franch. et Savat. Pl. Jap.* ii. pp. 103, 357.

*Cyperus parviflorus*, *Nees! in Wight, Contrib.* p. 87 (syn. omnibus excl.); *Hook. et Arn. Bot. Beechey's Voy.* p. 223; non *Vahl*.

*Cyperus Iria*, *Thwaites! Enum. Pl. Zeyl.* p. 344.

FORMOSA (*Campbell*), Tamsui (*Oldham*, 585!); HUPEH: Ichang (*A. Henry*, 302, 680!), Suinlinglen (*Faber*!); HONGKONG (*C. Wright*, 359! *Didrichsen*, 3310!). *Mus. Brit.*; *Herb. Kew.*

Scattered from the Himalaya and Japan to Ceylon and Timor.

19. *Cyperus malaccensis*, *Lam. Ill.* i. (1791) p. 146; *Kunth, Enum. Pl.* ii. p. 74; *Boeck. in Linnæa*, xxxv. (1867–8) p. 603, var.  $\beta$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 608.

*Cyperus odoratus*, *Linn. Sp. Pl.* ed. 1, p. 46, ed. 2, p. 68, et herb. propr. partim.

*Cyperus spaniophyllus*, *Steud. Syn. Pl. Cyp.* p. 21.

*Cyperus Fortunei*, *Steud. Syn. Pl. Cyp.* p. 21.

— *Rheede, Hort. Mal.* xii. p. 93, t. 50.

CHINA (*Fortune*, 7, 1004!); KWANGTUNG: Macao (herb. *Berol.*!), Whampoa (*Hance*, 6193!); LUCHU ARCHIPELAGO (*Dickins*!). *Mus. Brit.*; *Herb. Kew.*

Frequent in India, extending west to Persia, east to the Philippines and North Australia.

20. *Cyperus niveus*, *Retz. Obs.* v. p. 12; *Kunth, Enum. Pl.* ii. p. 45; *Boeck. in Linnæa*, xxxv. (1867-8) p. 530; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 601.

SZECHUEN: Mount Omei (*Faber*, 1122!); YUNNAN: Hokin (*Delavay*, 1758!), Tienfang, 900 feet (*A. Henry*, 9511!). Herb. Kew. - 7

Throughout India and Burma, in turf.

21. *Cyperus nutans*, *Vahl, Enum.* ii. p. 363; *Kunth, Enum. Pl.* ii. p. 94; *Boeck. in Linnæa*, xxxv. (1867-8) p. 597; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 607.

*Cyperus Jacquini*, *Schrad. in Linnæa*, xi. (1837) *Litt.-Bericht.* p. 87; *Fenzl, in Denkschr. Akad. Wiss. Wien*, viii. (1854) p. 45, t. 1.

YUNNAN: Mengtze, 4600 feet (*A. Henry*, 9737!). Herb. Kew. -  
In India common, extending east to Khasia and Assam.

22. *Cyperus pilosus*, *Vahl, Enum.* ii. p. 354; *Kunth, Enum. Pl.* ii. p. 80; *Benth. Fl. Hongk.* p. 387; *Boeck. in Linnæa*, xxxv. (1867-8) p. 598; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 609; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

*Cyperus marginellus*, *Nees, in Wight, Contrib.* p. 83, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 222; *Kunth, Enum. Pl.* ii. p. 75.

*Cyperus subalatus*, *Steud. Syn. Pl. Cyp.* p. 31.

*Cyperus pauciflorus*, *Steud. Syn. Pl. Cyp.* p. 34.

*Cyperus piptolepis*, *Steud. Syn. Pl. Cyp.* p. 40.

CHINA (*Millett!*); HUPEH: Ichang (*Henry*, 2358, 5019 partim!); YUNNAN: Szemao, 4000 feet (*A. Henry*, 13140!); KWANGTUNG: Macao (*Vachell*, 681!), Canton (*Sampson*, 8193!), Lofaushan (*Ford!*), Lienchau (*Lamont!*), Whampoa (*Hance*, 13138!); HONGKONG (*Urquhart! C. Wright*, 562!). Mus. Brit.; Herb. Kew.

Rare in Tropical Africa; abundant in India, Burma, and Malaya, extending to Japan, the Philippines, and Queensland.

23. *Cyperus procerus*, *Rottb. Descr. et Ic.* p. 29, t. 5. fig. 3; *Kunth, Enum. Pl.* ii. p. 72; *Hance, in Journ. Bot.* xvi. (1878) p. 113; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 610.

*Cyperus amoyensis*, *Hance, in Ann. Sc. Nat.* 5<sup>me</sup> sér. v. (1866) p. 249.

*Cyperus Heynei*, *Boeck. in Linnæa*, xxxv. (1867-8) p. 600.

*Cyperus punctulatus*, *Trimen, Cat. Pl. Ceylon*, p. 99, non *Vahl*.

FOKIEN: Amoy (*Hance*, 1392 partim!); KWANGTUNG (*Sampson!*). Mus. Brit.; Herb. Kew.

Frequent in India ; extending to Malaya and Queensland.

This plant, from its red colour, is much mixed in the old herbaria with *C. puncticulatus*, Vahl, which never has a trifid style, and is a true *Pycneus*.

Hance described his *C. amoyensis* from his Amoy plant, Hance 1392. Subsequently Hance put his n. 1392 on a plant which he received from India, which is not at all the same. Hance's numbers refer to his numbered list—not to his collections.

24. *Cyperus radicans*, *Nees et Meyen, ex Nees, in Linnæa*, ix. (1834) p. 285, *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 63 ; *Benth. Fl. Hongk.* p. 386 ; *Boeck. in Linnæa*, xxxv. (1867–8) p. 515 ; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 261 ; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 605 ; *Henriq. in Bol. Soc. Brotero*, ii. (1883) p. 166.

*Cyperus radicans*, *Kunth, Enum. Pl.* ii. p. 95 (errore typogr.).

*Cyperus Griffithii*, *Steud. Syn. Pl. Cyp.* p. 316 ; *C. B. Clarke, Journ. Linn. Soc., Bot.* xxi. (1884) p. 101 ; *Ridl. in Journ. Straits Branch Asiat. Soc.* xxiii. (1891) p. 5.

*Cyperus macropus*, *Miq. Fl. Ind. Bat., Suppl.* pp. 260, 599, non *Boeck.*

*Cyperus sinensis*, *Debeaux, in Act. Soc. Linn. Bordeaux*, xxxi. (1877) p. 14, t. 2 ; xxxviii. (1878) p. 30.

SHANTUNG : Chefoo (*Debeaux!*) ; KWANGTUNG : Cape Syngmun (*Meyen!*), Macao (*Hance, 10135!*), Pakhoi (*Playfair, 84!*) ; HONGKONG (*C. Wright, 565!*). Mus. Brit. ; Herb. Kew.

Malay Peninsula and Islands.

25. *Cyperus radiatus*, *Vahl, Enum.* ii. p. 369 ; *Kunth, Enum. Pl.* ii. p. 71 ; *Boeck. in Linnæa*, xxxvi. (1869–70) p. 317 ; *Hance, in Journ. Bot.* xvi. (1878) p. 113 ; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 617.

*Cyperus verticillatus*, *Roxb. Hort. Beng.* p. 6, *et Fl. Ind.* i. p. 206 ; *Hook. et Arn. Bot. Beechey's Voy.* p. 222.

*Cyperus macrosciadion*, *Steud. Syn. Pl. Cyp.* p. 37.

FORMOSA : Takow (*A. Henry, 1819!*) ; KWANGTUNG (*Sampson, 1460, 1617! Naumann!*). Mus. Brit. ; Herb. Kew.

In tropic and subtropic regions of the world ; common in Africa and India, less abundant in America.

26. *Cyperus rotundus*, *Linn. Sp. Pl.* ed. 2, p. 67, *nec Linn. herb. propr.* ; *Thunb. Fl. Japon.* p. 36 ; *Lour. Fl. Cochinch.* ed.

## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

Part I. for 1901, consisting of 378 pages, price 21 shillings, was issued on July 10, 1902; Part II., completing the volume, is now in the press.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.

---

VOL. XXXV. is still in course of issue, the Parts already published being Nos. 242-245. Those still unpublished are Parts 246-248 inclusive.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."

**THE JOURNAL**  
OF  
**THE LINNEAN SOCIETY.**

VOL. XXXVI.

BOTANY.

No. 252 ✓

## CONTENTS.

	Page
<p>An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.—</p> <p>PART XVII. ....</p>	217

See Notice on last page of Wrapper.

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1903.

712

# LINNEAN SOCIETY OF LONDON.

---

## LIST OF THE OFFICERS AND COUNCIL.

Elected 25th May, 1903.

---

### PRESIDENT.

Prof. Sydney Howard Vines, M.A., D.Sc., F.R.S.

### VICE-PRESIDENTS.

C. B. Clarke, M.A., F.R.S.  
Frank Crisp, LL.B., B.A.

Prof. J. B. Farmer, M.A., D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

Prof. G. B. Howes, D.Sc., LL.D., F.R.S. | D. H. Scott, M.A., Ph.D., F.R.S.

### COUNCIL.

Rev. R. A. Bullen, F.G.S.  
Charles Baron Clarke, M.A., F.R.S.  
Frank Crisp, LL.B., B.A.  
Prof. J. B. Farmer, M.A., D.Sc., F.R.S.  
Prof. J. Reynolds Green, M.A., D.Sc.,  
F.R.S.  
Prof. G. B. Howes, D.Sc., LL.D., F.R.S.  
B. Daydon Jackson, Esq.  
George Massee, Esq.

W. G. Ridewood, D.Sc., F.Z.S.  
G. S. Saunders, F.E.S.  
Dukinfield H. Scott, M.A., Ph.D.,  
F.R.S.  
A. C. Seward, M.A., F.R.S.  
Col. C. Swinhoe, F.Z.S., F.E.S.  
Prof. Sydney H. Vines, D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

---

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

---

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*; the former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meets at 4 P.M., as required during the Session. The Members for 1903-1904, in addition to the officers, are:—

Herbert Druce, F.Z.S.  
Antony Gepp, M.A.  
Herbert Goss, F.G.S.  
Albert C. L. G. Günther, M.D., F.R.S.  
W. Botting Hemsley, F.R.S.

P. Chalmers Mitchell, M.A., F.Z.S.  
G. R. M. Murray, F.R.S.  
A. B. Rendle, M.A., D.Sc.  
A. G. Tansley, M.A.



*Willd.* p. 53; *Kunth, Enum. Pl.* ii. p. 58 (*C. bulboso*, Vahl, sp. aliisque excl.); *Boeck. in Linnæa*, xxxvi. (1869-70) p. 283; *Benth. Fl. Hongk.* p. 387; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 615; *Debeaux, Fl. Tchefou*, p. 145, et in *Act. Soc. Linn. Bordeaux*, xxx. (1875) p. 65, et xxxiii. (1879) p. 45; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 317; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 166; *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

*Cyperus hexastachyus*, *Rottb. Descr. et Ic.* p. 28, t. 14. fig. 2; *Hook. et Arn. Bot. Beechey's Voy.* p. 272 [neque *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 60].

*Cyperus odoratus*, *Osbeck, Voy. ed. Forst.* ii. p. 342; *Bretschn. Early Research.* p. 111 (*oderatus*), non *Linn.*

*Cyperus lævissimus*, *Steud. Cyp.* p. 32.

CHINA (*Staunton! Hance, 1247!*); CHIHLI: Tientsin (*Debeaux!*); SHANTUNG: Chefoo (*Debeaux! Forbes, 1247!*); NORTH CENTRAL CHINA: Hukiascien (*Hugh!*); KIANGSU: Shanghai (*Maingay, 640, 763!*); CHEKIANG: Ningpo (*Faber, 223!*); KIANGSI (*David!*); FORMOSA (*Swinhoe! A. Henry, 1014, 1035, 1165!*), Tamsui (*Oldham, 583!*), Takow (*A. Henry, 1774!*); HUPEH: Ichang (*A. Henry, 214!*); YUNNAN: Tapintze (*Delavay, 405!*); KWANGTUNG: Macao (*Gaudichaud, 140!*), Canton (*Sampson, 262!*), Pakhoi (*Playfair, 208!*); HONGKONG (*Harland! C. Wright, 560 partim! 561 in herb. Kew! Lamont, 833!*); HAINAN (*A. Henry, 8240!*); LUCHU ARCHIPELAGO (*Beechey!*). *Mus. Brit.*; *Herb. Kew.*

In all tropical and warm temperate countries, most abundant; of sedges the worst pest in rice-fields.

27. *Cyperus stolonifer*, *Retz. Obs.* iv. p. 10; *Boeck. in Linnæa*, xxxv. (1867-8) p. 489 (syn. *C. tuberoso* et var.  $\beta$  excl.); *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 615.

*Cyperus littoralis*, *R. Br. Prodr.* p. 216.

*Cyperus tuberosus*, *Bojer, Hort. Maurit.* p. 379; *Kunth, Enum. Pl.* ii. p. 50, non *Rottb.*

*Cyperus rotundus*, "partim," *Kunth, Enum. Pl.* ii. p. 59; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 284, non *Linn.*

*Cyperus bulboso-stoloniferus*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62, et *Cyp.* p. 18.

*Cyperus conjunctus*, *Steud. Cyp.* p. 18.

*Cyperus arenarius*, *Hance, in Ann. Sc. Nat.* 4<sup>me</sup> sér. xviii. (1862) p. 233, in *Obs.*, non *Retz.*

FOKIEN : Amoy (*Hance* !); FORMOSA : Takow (*Playfair*, 115 !; *A. Henry*, 1957 !); KWANGTUNG : Macao (*Hance*, 1386 !). Mus. Brit.; Herb. Kew.

Mauritius, India, Malaya, Tropical Australia.

28. *Cyperus subfuscus*, *Debeaux*, in *Act. Soc. Linn. Bordeaux*, xxxi. (1877) t. 3, et xxxii. (1878) p. 30; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262.

*Cyperus fuscus*, var. *chinensis*, *Debeaux*, l. c. xxxii. (1878) p. 30.

SHANTUNG (*Staunton* !); Chefoo (*Debeaux*). Mus. Brit.

No type of this critical species has been seen; Staunton's example is referred here by comparison with Debeaux's figure.

29. *Cyperus tegetiformis*, *Roxb. Hort. Beng.* p. 6; *Kunth, Enum. Pl.* ii. p. 56; *Hance, in Journ. Bot.* xvii. (1879) p. 102; *Dyer, in Journ. Bot.* xviii. (1880) p. 224; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 612.

*Cyperus malaccensis*, var.  $\beta$ . *brevifolia*, *Boeck. in Linnæa*, xxxv. (1867-8) p. 604, syn. excl.

FORMOSA (*Hosie*, 3! *A. Henry*, 790, 1129 !); SZECHUEN (*Swinhoe*); KWANGTUNG : Macao (*Hornemann*, 103 !), Whampoa (*Hance*, 6193, 1392 partim ! nec 1392 in Amoy lecta); KWANGSI (*Wenyon* !). Mus. Brit.; Herb. Kew.

Extends from India to Japan.

Is used extensively for making mats; in Formosa, *vide* Hosie, cultivated for that purpose.

30. *Cyperus truncatus*, *Turcz. ! in Bull. Soc. Mosc.* (1838) p. 103 (nomen nudum), et *Fl. Baical.-Dahur.* ii. pars I. p. 245; *Maxim. Prim. Fl. Amur.* p. 297; *Boeck. in Linnæa*, xxxv. (1867-8) p. 604, non *A. Rich.*

SHINKING : Moukden (*James*, 6, 7 !); CHEKIANG : Ningpo Mountains (*Faber*, 84 !). Herb. Kew.

Dahuria, Amurland, Japan.

Var.  $\beta$ . *orthostachya*, *C. B. Clarke.*

*Cyperus orthostachyus*, *Franch. et Sav. Pl. Jap.* ii. pp. 106, 539.

*Cyperus fimbriatus*, *Franch. et Sav. Pl. Jap.* ii. p. 107; *Miquel ! MS.*, non *Nees* (an *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 142 ?).

HUPEH : Ichang (*A. Henry*, 5019 !). Herb. Kew.

Japan.

31. *Cyperus tuberosus*, *Rottb. Descr. et Ic.* p. 28, t. 7. fig. 1; *Kunth, Enum. Pl.* ii. p. 50, pro parvâ parte; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 285 partim; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 616.

*Cyperus rotundus*, *Benth. Fl. Austral.* vii. p. 279, pro magnâ parte.

*Cyperus tenuiflorus*, *Roxb. Hort. Beng.* p. 5, et *Fl. Ind.* i. p. 199, nec *Rottb.*

*Cyperus Retzii*, *Nees, in Wight, Contrib.* p. 82; *Kunth, Enum. Pl.* ii. p. 73.

FORMOSA: Takow (*A. Henry*, 778! 1037!). Mus. Brit.; Herb. Kew.

Tropical Africa, India, and Australia.

This species is so hardly separable from tall forms of *C. rotundus* that the very copious synonymy which has been accumulated is not here cited. The Formosa examples match very closely the Indian *C. tenuiflorus*, Roxb.

32. *Cyperus uncinatus*, *Poir. in Lam. Encyc.* vii. (1806) p. 247; *Kunth, Enum. Pl.* ii. p. 21; *Boeck. in Linnæa*, xxxv. (1867-8) p. 502.

*Cyperus cuspidatus*, *H. B. K. Nov. Gen. et Sp.* i. p. 204; *Kunth, Enum. Pl.* ii. p. 22; *Boeck. in Linnæa*, xxxv. (1867-8) p. 496; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 598.

*Cyperus castaneus*, *Miq. Fl. Ind. Bat.* iii. p. 261; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 130, vix *Willd.*

*Cyperus solutus*, *Steud. Cyp.* p. 14.

YUNNAN: Szemao, 5000 feet (*A. Henry*, 13390!); KWANG-TUNG: Macao (*Lay!*), Canton (*Dods!*), Whampoa (*Hance*, 7461!). Mus. Brit.; Herb. Kew.

In the tropics and warm temperate regions of Asia, Africa, and America. A common rice-field weed and on moist banks. To this is closely allied the true *C. castaneus*, Willd., a very much rarer species.

33. *Cyperus Zollingeri*, *Steud. in Zoll. Verz. Ind. Archip.* ii. (1854) p. 62; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 352; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 613.

*Cyperus bracteatus*, *Steud. Cyp.* p. 19.

*Cyperus hexastachyus*, *Nees, in Nov. Act. Acad. Nat. Cur.* xix. *Suppl.* i. (1843) p. 60.

*Cyperus tenuiculmis*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 286, nec *Boeck. in Flora*, lxii. (1879) p. 554.

*Cyperus Wightii*, *Hance*, in *Journ. Bot.* xvi. (1878) p. 14, non *Nees*.

*Cyperus lucidulus*, *C. B. Clarke*, in *Journ. Linn. Soc., Bot.* xxi. (1884) p. 99, et xxv. (1889) p. 80; nec *Klein*, nec *Kunth*.

— YUNNAN: Szemao, 5000 feet (*A. Henry*, 12186!); KWANGTUNG: Macao (*Abel*!), Whampoa (*Hance*, 19480!); HONGKONG (*C. Wright*, 560 partim!). *Mus. Brit.*; *Herb. Kew*.

In Tropical Africa and in India plentiful; extending to Luzon and to Queensland.

1. *Mariscus albescens*, *Gaud.* in *Freycinet, Voy.* (1826) p. 415; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 623.

*Cyperus stuppeus*, *Forst. f. Ins. Austral. Prodr.* p. 89.

*Cyperus pennatus*, *Lam. Ill.* i. (1791) p. 144; *Kunth, Enum. Pl.* ii. p. 80; *Benth. Fl. Hongk.* p. 387; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 131; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 166.

*Cyperus canescens*, *Vahl, Enum.* ii. p. 355, syn. *Rheede* excl.; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 340; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 61; *Hook. et Arn. Bot. Beechey's Voy.* p. 222.

*Cyperus bromoides*, *Vachell*, ex *Nees*, l. c.

CHINA (*Gaudichaud*, 42!); FORMOSA (*A. Henry*, 763, 783, 1032!), Takow (*Playfair*, 107! *A. Henry*, 1007!); KWANGTUNG: Macao (*Vachell*, 67!); HONGKONG (*Harland*! *C. Wright*, 563! *Seemann*, 556). *Mus. Brit.*; *Herb. Kew*.

South-east Asia, Tropical Australia, Polynesia, Sandwich Isles.

2. *Mariscus cyperinus*, *Vahl, Enum.* ii. p. 377; *Nees*, in *Hook. et Arn. Bot. Beechey's Voy.* p. 223; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 621.

*Kyllinga cyperina*, *Retz. Obs.* vi. (1791) p. 21.

*Cyperus umbellatus*, *Benth. Fl. Hongk.* p. 386, non *Roxb*.

FORMOSA (*Oldham*, 584! *A. Henry*, 1118!), Takow (*A. Henry*, 1700!); YUNNAN: Hsenkai (*A. Henry*, 9563!); KWANGTUNG: Macao (*Vachell*, 73 a! *Calléry*!); HONGKONG (*C. Wright*, 566! *Hance*, 1244! *Lamont*, 831!). *Mus. Brit.*; *Herb. Kew*.

A common Malay and Polynesian plant, extending to Japan and Kamtschatka.

This "species" (as *M. philippensis* and numerous others) is united with *M. Sieberianus* by Bentham under the name *Cyperus umbellatus*, for which an older name is *Mariscus biglumis*, *Gaertn. Fruct.* ii. (1788) p. 12, t. 2, fig. 8.

3. *Mariscus microcephalus*, *Presl, Rel. Haenk.* i. p. 182; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 624.

*Mariscus dilutus*, *Nees, in Wight, Contrib.* p. 90.

*Cyperus giganteus*, *Boeck. in Flora*, xlii. (1859) pp. 443, 496.

*Cyperus compactus*, *Retz. Obs.* v. p. 10, fide auctorum.

*Cyperus dilutus*, *Vahl, Enum.* ii. p. 357; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 65; *Kunth, Enum. Pl.* ii. p. 92.

*Cyperus septatus*, *Steud. Cyp.* p. 46.

KWANGTUNG: Canton (*Sampson, 1612!* hb. *Hance, 1367!*); Lofaushan (*Ford!*). *Mus. Brit.*; *Herb. Kew.*

Common in India and Malaya, extending to Mascarenia.

4. *Mariscus philippensis*, *Steud. Cyp.* p. 66, char. penitus mutato.

*Mariscus umbellatus*, *Presl, Rel. Haenk.* i. p. 181.

*Cyperus cylindrostachys* (partim), *Boeck. in Linnæa*, xxxvi. (1869-70) p. 383.

HAINAN (*A. Henry, 8!*). *Herb. Kew.*

Philippine Islands.

See observation above under *M. cyperinus*. The Hainan plant of A. Henry closely matches at all events the Philippine type of Steudel.

5. *Mariscus Sieberianus*, *Nees, in Linnæa*, ix. (1834) p. 286; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 622.

*Mariscus umbellatus*, *Vahl, Enum.* ii. p. 376, pro parte; *Kunth, Enum. Pl.* ii. p. 118, pro majore parte.

*Mariscus cyperinus*, *Presl, in Oken, Isis*, xxi. (1828) p. 270; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 71, var.  $\alpha$  et partim var.  $\beta$ : *Hook. et Arn. Bot. Beechey's Voy.* p. 272.

*Scirpus cyperoides*, *Linn. Mant.* p. 181, et herb. propr.!

*Kyllinga umbellata*, *Roxb. Fl. Ind.* i. p. 182.

*Cyperus umbellatus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 142.

*Cyperus ovularis*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 383, var.  $\alpha$  partim, *in Engl. Jahrb.* vi. (1885) p. 51, nec *Torrey*.

*Cyperus cylindrostachyus*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 383, pro magnâ parte.

*Cyperus Sieberianus*, *Diels, in Engl. Jahrb.* xxix. (1901) p. 227.

CHINA (*Meyen!*); CHEKIANG: Ningpo Mountains (*Faber, 225!*); FORMOSA: Tamsui (*A. Henry, 1456!*); HUPEH (*A. Henry, 1330!*), Patung (*E. H. Wilson, 1657!*); SZECHUEN: Tchenkeoutin (*Farges!*); YUNNAN: Tapintze (*Delavay, 1773!*), Mengtze

(*A. Henry*, 13726 !), Ichatong (*Delavay*, 414 bis !); HONGKONG (*Lamont* !); LUCHU ARCHIPELAGO (*Beechey* ! *Döderlein* !). Mus. Brit.; Herb. Kew.

Throughout all tropical and subtropical regions ; both general and abundant in South-east Asia.

1. *Torulinium confertum*, *Ham. Prodr. Ind. Occ.* p. 15.

*Torulinium ferox*, *Kunth, Enum. Pl.* ii. p. 90 in citat.

*Cyperus odoratus*, *Linn. Sp. Pl.* ed. 1, p. 46, quoad tab. Sloan. citatam neque ultra, nec *Sp. Pl.* ed. 2, p. 68, neque herb. propr.; *Vahl, Enum.* ii. p. 356 partim; *Kunth, Enum. Pl.* ii. p. 67; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 407.

*Cyperus ferax*, *Rich. in Act. Soc. Hist. Nat. Paris*, i. (1792) p. 106 *Kunth, Enum. Pl.* ii. p. 89; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 399.

*Cyperus ferox*, *Vahl, Enum.* ii. p. 357, pro parte.

*Cyperus flexuosus*, *Vahl, Enum.* ii. p. 359.

*Cyperus densiflorus*, *G. F. W. Meyer, Fl. Esseq.* p. 34; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 67, quoad syn. citat.; *Kunth, Enum. Pl.* ii. p. 104.

*Cyperus auriculatus*, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 67; *Kunth, Enum. Pl.* ii. p. 83; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 354.

*Cyperus lomentaceus*, *Nees et Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 60; *Kunth, Enum. Pl.* ii. p. 83.

*Cyperus multiceps*, *Hook. et Arn. Bot. Beechey's Voy.* p. 100; *Kunth, Enum. Pl.* ii. p. 102.

*Cyperus Prescottianus*, *Hook. et Arn. Bot. Beechey's Voy.* p. 100; *Kunth, Enum. Pl.* ii. p. 102.

*Cyperus strigosus*, *Hook. et Arn. Bot. Beechey's Voy.* p. 99; *Kunth, Enum. Pl.* ii. p. 88 partim; non *Linn.*

*Papyrus odoratus*, *Willd. in Abhandl. Akad. Berlin* (1812), p. 73, partim.

*Mariscus Pohlianus*, *Nees, in Mart. Fl. Bras.* ii. pars I. p. 50.

*Mariscus ferax*, *C. B. Clarke, in Hook. Fl. Brit. Ind.* vi. p. 624.

*Diclidium ferox*, *Schrader, ex Nees, in Mart. Fl. Bras.* ii. pars I. p. 54.

FORMOSA: Takow (*A. Henry*, 716, 729 !). Mus. Brit.; Herb. Kew.

Throughout the Tropics. Abundant in the New World, where it extends to Buenos Ayres. Besides the synonyms above cited, I have sixty-three others, several of which have been accepted as species by Kunth, many by Boeckeler. The confusion arose with Linnæus; his *C. odoratus* was founded on two

plants—*Pycneus polystachyus*, Beauv., mixed with *Cyperus malaccensis*, Lam. The specific name of L. C. Rich, "*ferax*," was early improved (?) by authors into "*ferox*."

1. *Kyllinga brevifolia*, *Rottb. Descr. et Ic.* p. 13, t. 4. fig. 3; *Boeck. in Linnæa*, xxxv. (1867–8) p. 424; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 588; *Ridl. in H. O. Forbes, Eastern Archip.* p. 521; *Diels, in Engl. Jahrb.* xxix. (1901) p. 228; *Palibin, Fl. Koreæ*, iii. p. 20.

*Kyllinga triceps*, *Thunb. Fl. Japon.* p. 35, non *Rottb.*

*Kyllinga sororia*, *Kunth, Enum. Pl.* ii. p. 131.

*Kyllinga gracilis*, *Kunth, Enum. Pl.* ii. p. 134.

*Schoenus coloratus*, *Linn. Sp. Pl.* ed. 1, p. 64, et herb. propr.!

*Scirpus glomeratus*, *Linn. Sp. Pl.* ed. 1, p. 52; *Osbeck, ex Bretschn. Early Research.* p. 111?

NORTH CENTRAL CHINA: Tcentosen, Mt. Momisan (*Hugh*!); Hukiascien (*Hugh*!); FORMOSA (*Oldham*, 249! 588 in hb. Kew! 587 in hb. Mus. Brit.), Apes' Hill (*Playfair*, 281!); HUPEH: Ichang (*A. Henry*, 2908, 3644, 3985, 4268, 7375!); SZECHUEN: Tchenkeoutin (*Farges*!); YUNNAN: Kichau (*Delavay*, 4035!), Tchenfongchang (*Delavay*!), Mosoyu (*Delavay*, 1772!), Mengtze, 5000 feet (*A. Henry*, 10728!); KWANGTUNG: Canton (*Yvan*!), Macao (*Calléry*!); HONGKONG (*Lamont*, 835! *C. Wright*, 567 partim!); COREA (*Oldham*, 906!); LUCHU ARCHIPELAGO (*C. Wright*, 359!). Mus. Brit.; Herb. Kew.

Throughout the world, in tropical and warm temperate regions.

2. *Kyllinga cylindrica*, *Nees, in Wight, Contrib.* p. 91; *Boeck. in Linnæa*, xxxv. (1867–8) p. 415; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 588.

YUNNAN: Mengtze, 4600 feet (*A. Henry*, 10966!). Herb. Kew. — 2

Scattered in the tropical and subtropical regions of the Old World, from West Africa to New South Wales. Represented in the New World by *K. odorata*, Vahl, hardly separable as a species.

3. *Kyllinga intermedia*, *R. Br. Prodr.* p. 219; *Kunth, Enum. Pl.* ii. p. 138; *Benth. Fl. Austral.* vii. p. 251.

*Kyllinga brevifolia*, *Boeck. MS. in Amalia Dietrich, Exsicc.* n. 717.

Australia.

Var.  $\beta$ . *oligostachya*, C. B. Clarke.

*Kyllinga oligostachya*, Boeck. in *Linnæa*, xxxv. (1867-8) p. 407.

*Kyllinga monocephala*, Seem. *Fl. Vit.* p. 318, pro majore parte.

FORMOSA: Takow, Apes' Hill (A. Henry, 1012, A, B!).  
Herb. Kew.

Fiji.

This species is perhaps, as Boeckeler sometimes thought, only a small-headed var. of *K. brevifolia*, Rottb.

4. *Kyllinga melanosperma*, Nees, in *Wight, Contrib.* p. 91; *Kunth, Enum. Pl.* ii. p. 131; Boeck. in *Linnæa*, xxxv. (1867-8) p. 419; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 588.

*Kyllinga fuscescens*, Boeck. in *Linnæa*, xxxv. (1867-8) p. 421 partim.

*Kyllinga vaginata*, Steud. in *Zoll. Verz. Ind. Archip.* ii. p. 63, var.  $\beta$  incl.

YUNNAN: Mengtze, 4600 feet (A. Henry, 10164!), Tapintze (Delavay, 1747!). Herb. Kew.

Scattered in the tropic and subtropic regions of the Old World, from Central and South Africa to Java.

5. *Kyllinga monocephala*, Rottb. *Descr. et Ic.* p. 13, t. 4. fig. 4; *Kunth, Enum. Pl.* ii. p. 129; Benth. *Fl. Hongk.* p. 388 partim; Debeaux, *Fl. Tchefou*, p. 146?; Franch. in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262?; Boeck. in *Linnæa*, xxxv. (1867-8) p. 427; in *Engl. Jahrb.* vi. (1885) p. 51; Ridl. in *H. O. Forbes, Eastern Archip.* p. 521; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 588.

*Kyllinga triceps*, *Linn. f. Suppl.* p. 104 partim; non Rottb.

*Kyllinga brevifolia*, Moritzi, *Verz. Zoll. Pfl.* p. 95.

*Kyllinga sororia*, Miq. *Fl. Ind. Bat.* iii. p. 293 partim; non *Kunth*.

*Kyllinga gracilis*, Steud. in *Zoll. Verz. Ind. Archip.* ii. p. 63, non *Kunth*.

*Kyllinga mindorensis*, Steud. *Cyp.* p. 67.

*Kyllinga intermedia*, Seem. in *Bonplandia*, ix. (1861) p. 261, non R. Br.

*Kyllinga monocephala*, var. *subtriceps*, *Kunth, Enum. Pl.* ii. p. 129.

*Schoenus niveus*, *Linn. Syst. Veget.* ed. Murray, xiii. p. 81.

*Scirpus cephalotes*, Jacq. *Hort. Vindob.* i. p. 42, t. 97.

*Thryocephalon nemorale*, J. R. et G. Forst. *Gen. Pl.* p. 130, t. 65.

*Cyperus leucocephalus*, Hassk. *Pl. Jav. Rar.* p. 87, non Retz.

FORMOSA: South Cape (A. Henry, 1329!) KWANGTUNG: Macao (Calléry, 377! Tate!), Pakhoi (Playfair, 202!); HONGKONG (C. Wright, 567 partim!); KWANGSI (Wenyon!); HAINAN



(*A. Henry*, 30!); LUCHU ARCHIPELAGO (*Döderlein*!). Mus. Brit.; Herb. Kew.

Throughout the Old World, in tropic and subtropic regions, especially near the sea.

6. *Kyllinga triceps*, *Rottb. Descr. et Ic.* p. 14, t. 4. fig. 6, tab. Rheedei excl.; *Linn. f. Suppl.* p. 104 partim; *Kunth, Enum. Pl.* ii. p. 133, saltem pro majore parte; *Boeck. in Linnæa*, xxxv. (1867-8) p. 413, var.  $\beta$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 587.

*Kyllinga nana*, *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 224; *Kunth, Enum. Pl.* ii. p. 138.

*Kyllinga monocephala*, *Nees, in Wight, Contrib.* p. 191, in *Hook. et Arn. Bot. Beechey's Voy.* p. 224; non *Rottb.*

KWANGTUNG: Macao (*Vachell*, 72, a, b!). Herb. Kew.

In tropical and subtropical regions of the Old World, from Senegambia to China and Australia.

1. *Eleocharis acicularis*, *R. Br. Prodr.* p. 224 in notâ; *Kunth, Enum. Pl.* ii. p. 141; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 628.

*Eleocharis Chætaria*, *Hance, in Journ. Bot.* xvii. (1878) p. 112, non *Roem. et Schult.*

*Scirpus acicularis*, *Linn. Sp. Pl.* ed. 1, p. 48, ed. 2, p. 71, partim neque in hb. propr.; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262.

*Heleocharis acicularis*, *Koch, Syn. Fl. Germ.* ed. 2, p. 853; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 431; *Diels, in Engl. Jahrb.* xxix. (1901) p. 228.

*Elæocharis acicularis*, *Ledeb. Fl. Ross.* iv. p. 243; *Maxim. Prim. Fl. Amur.* p. 298.

*Chætocyperus costulatus*, *Nees et Meyen, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 96.

*Chætocyperus Limnocharis*, *Nees, in Hook. Journ. Bot.* ii. (1840) p. 397; *Hook. et Arn. Bot. Beechey's Voy.* p. 272.

SHANTUNG: Chefoo (*Fauvel*!); KIANGSU: Shanghai (*Main-gay*, 633!); CHEKIANG: Ningpo Mountains (*Faber*, 71!); FORMOSA: Takow (*A. Henry*, 1839!); HUPEH (*A. Henry*, 7367!); Ichang (*A. Henry*, 2368!); YUNNAN: Mosoyn (*Delavay*, 811!); Kanhaytze (*Delavay*, 4803!); KWANGTUNG: Canton (*Hance*, 7883!); COREA: Chusan (*Wilford*, 443!); LUCHU ARCHIPELAGO (*C. Wright*, 357!). Mus. Brit.; Herb. Kew.

Throughout Europe and North Asia, extending southward only in China. Extends in America throughout the continent.

2. *Eleocharis afflata*, *Steud. in Zoll. Verz. Ind. Archip.* ii. (1854) p. 62, *et Cyp.* p. 76; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 629.

*Eleocharis pellucida*, *Presl, Rel. Haenk.* i. p. 196; *Kunth, Enum. Pl.* ii. p. 152, e *descript.*

*Eleocharis subprolifera*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62, *et Cyp.* p. 80.

*Scirpus afflatus*, *Benth. Fl. Hongk.* p. 394.

*Scirpus attenuatus*, *Franch. et Savat. Pl. Jap.* ii. pp. 110, 543.

*Heleocharis ochrostachys*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 452, *non Eleocharis ochrostachys, Steud.*

*Heleocharis chlorocarpa*, *Boeck. in Flora*, lxi. (1878) p. 34.

*Heleocharis Kuntzei*, *Boeck. in Cyp. Novæ*, i. (1888) p. 14.

*Heleocharis afflata*, *Naves, in Blanco, Fl. Philipp. Append.* 1880, p. 307; *Diels, in Engl. Jahrb.* xxix. (1901) p. 228.

*Heleocharis acicularis*, var. *japonica*, *Diels, in Engl. Jahrb.* xxix. (1901) p. 228?

CHEKIANG: Ningpo Mountains (*Faber*, 1518!); SZECHUEN: Tcentosen, Mt. Uomisan (*Hugh*!); HUPEH: Ichang (*A. Henry*, 3644! 4295! 7786!); YUNNAN: Tchenfongchang (*Delavay*!), Ontohay (*Delavay*, 408!); KWANGTUNG: Canton (*Sampson*!); HONGKONG (*Wilford*, 273! *Harland*, 543! *C. Wright*, 594! *O. Kuntze*, 3561!). Mus. Brit.; Herb. Kew.

Abundant in East Bengal and in Japan. The specific name "*pellucida*" is older, but less certain, than *afflata*.

3. *Eleocharis atropurpurea*, *Kunth, Enum. Pl.* ii. p. 151; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 627.

*Scirpus atropurpureus*, *Retz. Obs.* v. p. 14.

*Isolepis atropurpurea*, *Roem. et Schult. Syst.* ii. p. 106, *et Mant.* p. 61; *Kunth, Enum. Pl.* ii. p. 190.

*Eleogenus atropurpureus*, *Nees, in Wight, Contrib.* p. 113.

*Heleocharis atropurpurea*, *Koch, Syn. Fl. Germ.* ed. ii. p. 853; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 458, var.  $\gamma$  *excl.*

SZECHUEN: Tcentosen, Mt. Uomisan (*Hugh*!); KWANGTUNG: Macao (*Lay*!). Mus. Brit.

Throughout the world, in tropical and warm temperate climates.

4. *Eleocharis capitata*, *R. Br. Prodr.* p. 225; *Kunth, Enum. Pl.* ii. p. 150, syn. quibusdam excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 627.

*Eleocharis setacea*, *R. Br. Prodr.* p. 225, nec p. 224.

*Scirpus capitatus*, *Linn. hb. propr.*, partim; *Benth. Fl. Hongk.* p. 394.

*Eleogenus capitatus*, *Nees, in Wight, Contrib.* p. 112.

*Heleocharis capitata*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 461.

FORMOSA (*A. Henry*, 1042!), Takow (*Playfair*, 134! *A. Henry*, 704!); HONGKONG (*C. Wright*, 593!). *Mus. Brit.*; *Herb. Kew.*

Throughout the world in tropical and warm regions, more abundant in America; not yet seen from Europe.

5. *Eleocharis fistulosa*, *Link, Jahrb.* iii. (1820) p. 78; *Kunth, Enum. Pl.* ii. p. 155; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 626.

*Eleocharis acutangula et E. media*, *Schultes, in Roem. et Schult. Syst.* ii. *Mant.* p. 91.

*Scirpus fistulosus*, *Poir. in Lam. Encycl.* vi. p. 749.

*Limnochloa acutangula et L. media*, *Nees in Wight, Contrib.* p. 114.

*Heleocharis fistulosa*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 472 (syn. et exemplis Amer. excl. ?); *Diels, in Engl. Jahrb.* xxix. (1901) p. 228.

HUPEH: Ichang (*A. Henry*, 4102, 4102 A!). *Mus. Brit.*; *Herb. Kew.*

In the tropic and subtropic regions of the Old World; abundant in India. The representative common American species is *E. mutata*, *R. Br.*, which is united with *E. fistulosa* by Boeckeler, not without reason.

6. *Eleocharis palustris*, *R. Br. Prodr.* p. 224 in notâ; *Bunge, Enum. Pl. China Bor.* p. 68; *Kunth, Enum. Pl.* ii. p. 147; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 96; *Regel, Fl. Ussur.* p. 244; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 628.

*Eleocharis uniglumis*, *Schultes, in Roem. et Schult. Syst.* ii. *Mant.* pp. 88, 539.

*Eleocharis pileata*, *A. Gray! in Mem. Amer. Acad.* n. s. vi. (1858-9) p. 417 in notâ.

*Scirpus palustris*, *Linn. Sp. Pl.* ed. 1, p. 47, ed. 2, p. 70 partim; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 317.

*Scirpus uniglumis*, *Link, Jahrb.* iii. p. 77; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 318.

*Scirpus pileatus*, *Franch. et Sav. Enum. Pl. Jap.* ii. p. 110.

*Scirpus kamtschaticus*, *C. A. Meyer, in Mém. Sav. Etr. Pétersb.* i. (1831) p. 198, t. 1.

*Heleocharis palustris*, *Lindl. Syn. Brit. Fl.* p. 280; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 466, exemplis 3-stigmatis excl.

CHIHLI: Jehol (*David*, 1910, 2712!), Peking (*Bretschneider*, 865!); SHANTUNG: Chefoo (*Fauvel*!); HUPEH: Patung (*E. H. Wilson*, 1645!), 6700 feet (*E. H. Wilson*, 1645 a!); SZECHUEN: Tchenkeoutin, 4500 feet (*Farges*, 4185!); YUNNAN: 5000 feet (*J. Anderson*!), Mengtze, 4500 feet (*A. Henry*, 10862!), Kanhaytze, 9500 feet (*Delavay*, 2182!), Tapintze (*Delavay*, 244, 416, 912!). *Mus. Brit.*; *Herb. Kew.*

Throughout the world, except South-east Asia and Australia. The Australian plants referred to *E. palustris* by Boeckeler have the style 3-fid; *cf.* *Benth. Fl. Austral.* vii. pp. 294-296. The high-level Yunnan *E. palustris* is a frontier of the species.

7. *Eleocharis plantaginea*, *R. Br. Prodr.* p. 224 in notâ; *Bunge, Pl. Chin. Bor.* p. 68; *Kunth, Enum. Pl.* ii. p. 153; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 625.

*Scirpus plantaginoides*, *Rottb. Descr. et Ic.* p. 45, t. 15, fig. 2.

*Scirpus plantagineus*, *Retz. Obs.* v. p. 14; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 131.

*Limnochloa plantaginea et L. tumida*, *Nees, in Wight, Contrib.* p. 114.

*Heleocharis plantaginea*, *Boeck. in Linnæa*, xxxvi. (1869-70) p. 474, descr. et exemplis Americanis excl.; *Diels, in Engl. Jahrb.* xxix. (1901) p. 228.

FOKIEN: Amoy (*Hance*!); FORMOSA: Takow (*A. Henry*, 734!); HUPEH: Ichang (*A. Henry*, 4247!); YUNNAN: Mosoyñ (*Delavay*, 4075!); HONGKONG (*Hance*, 1391! *Lamont*, 860!). *Mus. Brit.*; *Herb. Kew.*

In the Old World, from Angola to Japan and Fiji; common in India. The representative species in America is *E. interstincta*, *R. Br.*

8. *Eleocharis tetraquetra*, *Nees, in Wight, Contrib.* p. 113; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 630; *Palibin, Fl. Koreæ*, iii. p. 19.

*Eleocharis erythrochlamys*, *Miq. Fl. Ind. Bat.* iii. p. 300.

*Scirpus Wichurai*, *Franch. et Sav. Pl. Jap.* ii. p. 544, non *Boeck.*

*Scirpus hakonensis et S. Onaei*, *Franch. et Sav. Pl. Jap.* ii. pp. 110, 111, 544.

*Scirpus petasatus*, Maxim. in *Bull. Soc. Mosc.* liv. (1879) p. 64.

*Heleocharis tetraquetra*, Boeck. in *Linnaea*, xxxvi. (1869-70) p. 447; Diels, in *Engl. Jahrb.* xxix. (1901) p. 229.

*Heleocharis Wichurai*, Boeck. in *Linnaea*, xxxvi. (1869-70) p. 448.

*Heleocharis alta*, Boeck. *Cyp. Novæ*, i. (1888) p. 17.

CHINA (*A. Henry*, 2298!); CHEKIANG: Ningpo Mts. (*Faber*, 238!); HUPEH: Ichang (*A. Henry*, 4232!). Mus. Brit.; Herb. Kew.

Frequent in the Himalayas, from 1500 to 11,000 feet altitude, extending thence to Japan and New South Wales. The species is strongly defined by the long-fimbriate, not merely bristly, hypogynous setæ; but, in its great range, it varies much in size and in the size of its heads; whence its copious synonymy.

9. *Eleocharis tuberosa*, Schultes, in *Roem. et Schult. Syst.* ii. *Mant.* p. 86; *Miq. Fl. Ind. Bat.* iii. p. 302; *Debeaux*, in *Act. Soc. Linn. Bordeaux*, xxx. (1875) p. 66 (*cf.* Thozet, 'Roots and Tubers of Queensland,' 1866).

*Eleocharis plantaginea*, var., Nees, in *Wight, Contrib.* p. 114.

*Scirpus tuberosus*, Roxb. *Pl. Corom.* iii. p. 25, t. 231, et *Fl. Ind.* i. p. 210; *Hance*, in *Ann. Sc. Nat.* 5<sup>m</sup>e sér. v. (1866) p. 249.

*Scirpus plantagineus*, Franch. *Pl. David.* i. p. 318.

*Cyperus dulcis*?, Rumph. *Amb.* v. 6, p. 7, t. 3. fig. 1.

*Hippuris indica*, Lour. *Fl. Cochinch.* p. 16, fide syn. citati.

CHIHLI: Peking, culta (*Bretschneider*, 797! *David*, 109!); KIANGSU: Shanghai, culta (*Debeaux*); CHEKIANG: Ningpo Mts. (*Faber*, 1688); KWANGTUNG: Whampoa (*Hance*, 10142!). Mus. Brit.; Herb. Kew.

Both Nees and Franchet regard this important Chinese food-plant as only a cultivated form of *E. plantaginea*. The authentic example of *Bretschneider* has neither flower-heads nor rhizome. The plant, from China, cultivated at Buitenzorg (Miquel's *tuberosa*?) is gigantic, the culms more than 3 feet high, more than  $\frac{1}{2}$  in. in diam., the spike  $\frac{1}{2}$ - $\frac{1}{3}$  in. thick; but nothing is known about its tubers.

10. *Eleocharis variegata*, Kunth, *Enum. Pl.* ii. p. 153.

In Mascarenia.

Var.  $\beta$ . *laxiflora*, Ridley, in *Journ. Straits Br. Asiat. Soc.* n. 23 (1891) p. 14; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 626.

*Eleocharis* ? *spiralis*, *hb. Hance*.

*Scirpus laxiflorus*, *Thwaites, Enum. Pl. Zeyl.* p. 435; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 131.

*Heleocharis Græffeana*, *Boeck. in Flora*, lviii. (1875) p. 109.

HONGKONG (*Sampson* in *hb. Hance*, 13333!). *Mus. Brit.*; *Herb. Kew.*

In Asia, common, from Bengal and Ceylon to Hongkong and Samoa in W. Polynesia; nor can I distinguish the Central-American *Scirpus plicarrhachis*, *Griseb.*

1. ***Fimbristylis Actinoschoenus***, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 650.

*Arthrostylis filiformis*, *Thw. Enum. Pl. Zeyl.* p. 352; *Boeck. in Linnæa*, xxxvii. (1873) p. 524.

*Actinoschoenus filiformis*, *Benth. in Hook. Ic. Pl.* xiv. p. 33, t. 1346. *Ceylon.*

**Var. *chinensis***, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 651.

*Arthrostylis chinensis*, *Benth. Fl. Hongk.* p. 397; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 133; *cf. Hance, in Journ. Bot.* vii. (1869) p. 63.

*Actinoschoeni* sp., *Benth. in Hook. Ic. Pl.* xiv. p. 33.

HONGKONG (*C. Wright*, 599! *Hance*, 7493! *Lamont*, 867!). *Mus. Brit.*; *Herb. Kew.*

This variety occurs also in Perak.

2. ***Fimbristylis acuminata***, *Vahl, Enum.* ii. p. 285; *Kunth, Enum. Pl.* ii. p. 221; *Benth. Fl. Hongk.* p. 391; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 167; *Boeck. in Linnæa*, xxxvii. (1871) p. 3, var.  $\beta$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 631.

*Fimbristylis acicularis*, *R. Br. Prodr.* p. 226.

*Scirpus acuminatus*, *Poir. Encyc. Suppl.* v. p. 100, non *Muhl.*

HONGKONG (*Hance*!). *Mus. Brit.*; *Herb. Kew.*

From the North-west Himalaya to Singapore, Malaya, and New South Wales.

3. ***Fimbristylis æstivalis***, *Vahl, Enum.* ii. p. 288; *Kunth, Enum. Pl.* ii. p. 225; *Benth. Fl. Hongk.* p. 392; *Boeck. in Linnæa*, xxxvii. (1871) p. 11; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 637.

*Fimbristylis laxa*, *Regel, Fl. Ussur.* p. 245, non *Vahl*.

*Fimbristylis leiocarpa*, *Maxim. Prim. Fl. Amur.* p. 301, non *Miq.*

*Scirpus æstivalis*, *Retz. Obs.* iv. p. 12.

FORMOSA (*Oldham*, 591!); KWANGTUNG: Canton (*Sampson*, 942! *C. Wright*, 582! *Ford*, 32!); HONGKONG (*Wilford!* *Hance*, 1090!); HAINAN (*Swinhoe*). *Mus. Brit.*; *Herb. Kew.*

Eastern Asia and Australia; from Nepaul and Amurland to Victoria and Viti. The representative species in Brazil, *F. limosa*, *Kunth*, may be esteemed a geographic form only of *F. æstivalis*.

4. *Fimbristylis complanata*, *Link, Hort. Berol.* i. (1827) p. 292; *Kunth, Enum. Pl.* ii. p. 228; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 132; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 167; *Franch. in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 263; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 646; *Diels, in Engl. Jahrb.* xxix. (1901) p. 229.

*Fimbristylis autumnalis*, var. *a*, *Boeck. in Linnæa*, xxxvii. (1871) p. 38.

*Fimbristylis amblyphylla*, *Steud. Cyp.* p. 116, pl. in *Obs. excl.*

*Fimbristylis quinquangularis*, *Munro, in Seem. Bot. Voy. 'Herald,'* p. 422, non *Kunth*.

*Scirpus complanatus*, *Retz. Obs.* v. p. 14.

*Scirpus anceps*, *Willd. Ges. Naturf. Fr. Berl. Mag.* ii. (1808) p. 287, t. 8. fig. 2.

*Trichelostylis complanata*, *Nees, in Wight, Contrib.* p. 103, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 84.

SHANTUNG: Chefoo (*Fauvel!* *Schottmueller*, 331!), Chefoo Mts. (*Faber*, 69!); KIANGSU: Shanghai (*Faber*, 235!); CHEKIANG: Ningpo (*Faber*, 376!); FORMOSA: Takow (*A. Henry*, 772!); HUPEH: Ichang (*A. Henry*, 4311!); YUNNAN: Mosoyu (*Dela-  
vay*, 1779, 4552!); KWANGTUNG: Macao (*da Silva!*), Canton (*Hance*, 7483!). *Mus. Brit.*; *Herb. Kew.*

Common in all warm regions.

Var. *Kraussiana*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 646.

*Fimbristylis Kraussiana*, *Krauss, in Flora*, xxviii. (1845) p. 757; *Hochst. MS.*

*Fimbristylis connectens*, *Thw. Enum. Pl. Zeyl.* p. 349.

SHANTUNG (*Maingay*, 175!); KIANGSU: Shanghai (*Maingay*, 693!); KWANGTUNG: Canton (*Sampson*, 462 partim!). *Mus. Brit.*; *Herb. Kew.*

Both in Africa and South-east Asia.

Var. *microcarpa*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 646.

*Fimbristylis microcarpa*, *F. Muell. Fragm. Phyt. Austral.* i. p. 200.

*Fimbristylis tenera*, *Boeck. in Flora*, lviii. (1875) p. 111, var. excl.; non *Roem. et Schult.*

*Fimbristylis* sp., a *F. autumnali* vix distinguenda, *F. Muell. Fragm. Phyt. Austral.* ix. p. 11.

KWANGTUNG: Canton (*Sampson*, 462 partim); HONGKONG (*Hance*, 7488!). *Mus. Brit.*; *Herb. Kew.*

In South-east Asia and Tropical Australia.

*F. complanata*, as here understood, is common nearly throughout the warmer regions of the World, not, however, extending to Europe or the Mediterranean Region. The two varieties pass into the main species, so that no separate areas can be defined for them. In China, the series graduates from the stout form (*A. Henry*, 4311) down to the delicate Australian *F. microcarpa*, admitted as a species by Bentham, which, again, Boeckeler regards as inseparable from the *F. autumnalis*, Linn., abundant in America.

5. *Fimbristylis cymosa*, *R. Br. Prodr.* p. 228; *Kunth, Enum. Pl.* ii. p. 244; *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 98; *Benth. Fl. Austral.* vii. p. 318.

*Fimbristylis glomerata*, *Boeck. in Linnæa*, xxxvii. (1871) p. 47. partim.

*Fimbristylis multifolia*, *Boeck. in Linnæa*, xxxviii. (1874) p. 397.

*Scirpus cymosus*, *Lam. Ill.* i. p. 141.

LUCHU ARCHIPELAGO (*C. Wright*, 358!). *Herb. Kew.*

Common in Polynesia, extending from the Sandwich Isles to Tropical Australia and Java.

Boeckeler first placed this plant as identical with the subcapitate form of *F. spathacea*, Roth; and three years later, forgetting this, reconstituted it as a species under a new name.

6. *Fimbristylis cyperoides*, *R. Br. Prodr.* p. 228; *Kunth, Enum. Pl.* ii. p. 244; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 650.

*Gussonea cyperoides*, *Presl, Rel. Haenk.* i. p. 183, t. 33.

In warm Australia and the Carolines.



Var. *cinnamometorum*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 650.

*Fimbristylis cinnamometorum*, *Kunth*, *Enum. Pl.* ii. p. 229; *Boeck.* in *Linnæa*, xxxvii. (1871) p. 35.

*Scirpus cinnamometorum*, *Vahl*, *Enum.* ii. p. 278.

KWANGTUNG: Canton (*Sampson*, 462!). Mus. Brit.; Herb. Kew.

This variety occurs in Ceylon, the mountains of South India, and in Pegu.

7. *Fimbristylis dichotoma*, *Vahl*, *Enum.* ii. p. 287; *Kunth*, *Enum. Pl.* ii. p. 225; *Boeck.* in *Linnæa*, xxxvii. (1871) p. 12, varr.  $\beta$ ,  $\gamma$ , syn. *Royenianâ* et exemplis Americanis excl.; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 635; *Hance*, in *Journ. Bot.* xvii. (1878) p. 112.

*Fimbristylis pallescens*, *Nees*, in *Wight, Contrib.* p. 101.

*Scirpus dichotomus*, *Linn. Sp. Pl.* ed. 1, p. 50; ed. 2, p. 74.

SHANTUNG (*Maingay*, 176!); NORTH CENTRAL CHINA: *Hukiasien* (*Hugh*!); SZECHUEN: *Icentosen*, Mt. *Uomisan* (*Hugh*!); YUNNAN: *Tali* (*Delavay*, 4753!); KWANGTUNG: *Macao* (*Calléry*, 379!); HONGKONG (*Lamont*!); KWANGSI (*Wenyon*!). Mus. Brit.; Herb. Kew.

An annual weed throughout the warmer parts of the Old World, from South Europe and the Cape of Good Hope to New South Wales. Abundant in India; little collected in China.

8. *Fimbristylis diphylla*, *Vahl*, *Enum.* ii. p. 289; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 81; *Benth. Fl. Hongk.* p. 392; *Henriq.* in *Bolet. Soc. Broter.* ii. (1883) p. 166; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 636; *Diels*, in *Engl. Jahrb.* xxix. (1901) p. 229.

*Fimbristylis laxa*, *Vahl*, *Enum.* ii. (1806) p. 292; *Kunth*, *Enum. Fi.* ii. p. 232.

*Fimbristylis pilosa*, *Presl, Rel. Haenk.* i. p. 191, et auctorum plur., non *Vahl*.

*Fimbristylis tomentosa*, *Vahl*, *Enum.* ii. p. 290; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 81, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 225; *Debeaux, Fl. Tchefou*, p. 150; *Franch.* in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 264.

*Fimbristylis depauperata*, *R. Br. Prodr.* p. 227.

*Fimbristylis ovalis*, *Nees*, in *Wight, Contrib.* p. 98; *Hance*, in *Journ. Bot.* xvii. (1879) p. 16.

*Fimbristylis podocarpa*, *Nees*!, in *Wight, Contrib.* p. 98 partim, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 77, var.  $\beta$  et a partim nec  $\gamma$ , in *Hook. et Arn. Bot. Beechey's Voy.* p. 225 partim; *Benth. Fl. Hongk.* p. 391; non *F. podocarpa*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 638.

*Fimbristylis communis*, *Kunth, Enum. Pl.* ii. p. 234, syn. quibusdam excl.; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 264.

*Fimbristylis polymorpha*, *Boeck.*! in *Vidensk. Meddel. Kjoeb.* (1869) pp. 141, 158, et in *Linnaea*, xxxvii. (1871) p. 14, syn. paucis excl.

*Scirpus diphyllus*, *Retz. Obs.* v. p. 15.

*Scirpus dichotomus*, *Linn. Sp. Pl.* ed. 1, p. 50, quoad exempl. Zeylanicum "typicum," e *Trimen*, in *Journ. Linn. Soc., Bot.* xxiv. (1888) p. 135.

CHIHLI: Peking (*Bretschneider*, 862!); SHANTUNG (*Maingay*, 176!), Chefoo (*Schottmueller*, 352!), Chefoo Mts. (*Faber*, 285!, *Forbes*, 289, 390!); KIANGSU: Shanghai (*Maingay*, 641, 642!, *Faber*, 234!); KIANGSI: Kiukiang (*Miss Reid*, 57!); FORMOSA: (*A. Henry*, 786 A, 788, 789, 1040, 1101!), Takow (*A. Henry*, 1148!); HUPEH: (*A. Henry*, 4248!), Ichang (*A. Henry*, 2248, 2296, 4251!); YUNNAN: Tchenfongchan (*Delavay*!), Tapintze (*Delavay*, 401, 1765, 4821!); KWANGTUNG (*Forbes*, 182, 186, 187, 189!), Canton (*Sampson*, 269!), Macao (*Abel*!); SOUTH CHINA (*Hance*, 1260! *Didrichsen*, 3197!); HONGKONG (*Wilford*, 278! *Hance*, 7482! *C. Wright*, 585, 587, 588!). *Mus. Brit.*; *Herb. Kew.*

Common in all warm countries throughout the world; the most widespread and one of the most abundant of Cyperaceæ.

The range of forms here supposed to be included under *F. diphylla* is rather narrower than that included by *Kunth* or by *Boeckeler*, who gave the species a new name (*communis*, *Kunth*, or *polymorpha*, *Boeck.*) to indicate that their species had wider limits than *F. diphylla*, *Vahl*. In the narrower sense here understood, the species has 140 names besides those above cited. The variation in development and in hairiness is very great. The stems and leaves are sometimes stout and rigid, the umbel with 50–100 spikelets, sometimes almost capillary with only one spikelet or very few (= var. *leptophylla*, *Benth. Fl. Hongk.* p. 392). The colour of the spikelets is usually a ferruginous brown, sometimes nearly black. The common form is glabrous or nearly so; but a hairy form (= var. *tomentosa*, *Benth. Fl.*

Hongk. p. 392) is common in Malay, frequent in China. Finally the spikelets are sometimes clustered subcapitate, as in Delavay 1765 !

9. **Fimbristylis dipsacea**, *Benth. et Hook. f. Gen. Pl.* iii. (1883) p. 1049; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 635.

*Scirpus dipsaceus*, *Rottb. Descr. et Ic.* p. 56, t. 12. fig. 1; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 736.

*Echinolytrum dipsaceum*, *Desv. in Journ. Bot.* i. (1808) p. 21, t. 1.

*Isolepis dipsacea*, *Roem. et Schult. Syst.* ii. p. 119; *Kunth, Enum. Pl.* ii. p. 205.

*Isolepis verrucifera*, *Maxim. Prim. Fl. Amur.* p. 300.

YUNNAN (*J. Anderson* !); KWANGTUNG: Canton (*Lamont* !). *Mus. Brit.*

A common Indian weed, extending to the Philippines and Amurland; received also lately from the Congo.

10. **Fimbristylis Eragrostis**, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 132.

*Fimbristylis quinquangularis*, *Munro, in Seem. Voy. 'Herald,'* p. 422.

*Abildgaardia Eragrostis*, *Nees, in Wight, Contrib.* p. 95, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 224; *Kunth, Enum. Pl.* ii. p. 249; *Benth. Fl. Hongk.* p. 390; *Boeck. in Linnæa*, xxxvii. (1871) p. 55 partim.

CHEKIANG: Mt. Pakwan (*Hance, 7482* !); FORMOSA (*Campbell*); KWANGTUNG (*Ford, 190, 191, 192* !), Lofaushan (*Ford, 126, 131* !), Cape Syngmun (*Meyen* !), Macao (*Gaudichaud* !), Whampoa (*Didrichsen* !), Canton (*Wichura, 1708* !); HONGKONG (*C. Wright, 573, 586, 588* ! *Seemann, 224* partim !). *Mus. Brit.*; *Herb. Kew.*

Also received from Tonkin.

11. **Fimbristylis ferruginea**, *Vahl, Enum.* ii. (1806) p. 291; *Hook. et Arn. Bot. Beechey's Voy.* p. 312; *Benth. Fl. Hongk.* p. 391; *Boeck. in Linnæa*, xxxvii. (1871) p. 16; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 638.

*Fimbristylis arvensis*, *Vahl, Enum.* ii. p. 291; *Kunth, Enum. Pl.* ii p. 237.

*Fimbristylis tristachya*, *R. Br. Prodr.* p. 226; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i (1843) p. 76.

*Scirpus ferrugineus*, *Linn. Sp. Pl.* ed. 1, p. 50; ed. 2, p. 74.

*Scirpus arvensis*, *Retz. Obs.* iv. p. 11.

FORMOSA (*A. Henry*, 712, 733, 773!), Takow (*A. Henry*, 1066!); KWANGTUNG: Pakhoi (*Playfair*, 206!); HONGKONG (*Harland*, 679! *Hance*, 1260 partim! 7481!). Mus. Brit.; Herb. Kew.

Common in all warm countries, especially near the sea.

The numbers of Hance are not the field-numbers of the collectors, but the list number of his own herbarium (as are the numbers cited mostly by European and by "British" botanists); they are therefore sometimes made up of several distinct collections, and misleading.

12. *Fimbristylis Fordii*, *C. B. Clarke*, n. sp.

Annua, debilis, fere glabra. Culmi cæspitiosi, 4-8 uncias longi, 2-4-spiculigeri. Folia pauca, subsetacea, culmum superantia. Umbella simplex; radii usque ad  $1\frac{3}{4}$  unc. longi; bracteæ usque ad 1 unc. longæ. Spiculæ  $\frac{1}{4}$ - $\frac{1}{3}$  unc. longæ, lineari-lanceolatæ. Glumæ acuminatæ, mucronatæ. Stylus 2-fidus. Nux obovoidea, straminea, in utraque facie longitudinaliter 9-striata, et (præsertim in lateribus apiceque) valide tuberculata.

KWANGTUNG: Lofaushan (*Ford*, 132!). Herb. Kew.

Next *F. gracilentata*, Hance; but the nut is different, resembling somewhat that of *F. polytrichoides*.

13. *Fimbristylis fusca*, *Benth. et Hook. f. Gen. Pl.* iii. (1883) p. 1048; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 649.

*Fimbristylis cinnamometorum*, *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 132, non *Kunth*; cf. *Benth. Fl. Austral.* vii. p. 318, in notâ.

*Fimbristylis Kamphoveneri*, *Boeck. in Engl. Jahrb.* v. (1884) p. 505.

*Gussonea pauciflora*, *Brongn. in Duperrey, Voy. 'Coquille,'* pars 2 (1829) p. 171, t. 34 B.

*Schoenus puberulus*, *C. A. Meyer*, in *Mém. Sav. Etr. Pétersb.* i. (1831) p. 196, t. 1; *Kunth, Enum. Pl.* ii. p. 337.

*Abildgaardia fusca*, *Nees*, in *Wight, Contrib.* p. 95; *Benth. Fl. Hongk.* p. 390; *Boeck. in Linnæa*, xxxvii. (1871) p. 54.

*Abildgaardia pauciflora*, *Kunth, Enum. Pl.* ii. p. 249.

*Rhynchospora* ? anomala, *Steud. in Zoll. Verz. Pfl. Ind. Archip.* ii. p. 61, et *Cyp.* p. 149.

*Isolepis longispica*, *Steud. Cyp.* p. 104.

KWANGTUNG: Canton, West River (*Hance*, 13111 in herb. Paris! foliis perbrevibus); HONGKONG (*C. Wright*, 574! *Hance*, 356!). Mus. Brit.; Herb. Kew.

A Malay species extending to Bengal and Canton.

Var. **cantoniensis**, n. var., *C. B. Clarke*; foliis culmo fere æquilongis; bracteis 2, inferiore inflorescentiâ æquilongâ, planâ, suberectâ, apice subito angustatâ.

*Fimbristylis fulvescens*, *Hance*, in *Journ. Bot.* xvi. (1878) p. 112, non *Thwaites*.

KWANGTUNG: Canton, Pakwan (*Sampson* in hb. *Hance*, 13111!). *Mus. Brit.*; *Herb. Kew.*

The bracts in this plant of *Sampson's* are very like those of *F. fulvescens*, *Thwaites*, and I can find no similar ones in *F. fusca*, which has very short setaceous bracts.

14. **Fimbristylis globulosa**, *Kunth*, *Enum. Pl.* ii. p. 231; *Boeck. in Linnæa*, xxxvii. (1871) p. 45; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 644.

*Scirpus globulosus*, *Retz. Obs.* vi. p. 19.

*Trichelostylis globulosa*, *Nees*, in *Wight, Contrib.* p. 105.

KIANGSU: Shanghai (*Maingay*, 692! in hb. *Kew*, 693! in hb. *Calcutta*); KIANGSI: Kiukiang (*Shearer*!); SZECHUEN: Mt. Omei, alt. 3000 feet (*Faber*, 1134!). *Herb. Kew.*

Common in India and Malaya.

Var.  $\beta$ . **Torresiana**, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 645.

*Fimbristylis Torresiana*, *Gaudich. in Freycinet, Voy.* p. 413; *Kunth, Enum. Pl.* ii. p. 231.

LUCHU ARCHIPELAGO (*Maximowicz*!). *Herb. Kew.*

This variety is also known from Bengal, Japan, Marianne Isles.

15. **Fimbristylis gracilentata**, *Hance*, in *Journ. Bot.* vi. (1868) p. 89.

KWANGTUNG: Whampoa (*Hance*, 13227!), Canton (*Sampson*, 919). *Mus. Brit.*

Also from Tonkin (*Balansa*, 2786!).

The stem often carries 10–12 spikelets in a loose compound umbel; the spikelets are very slender and the nut smooth; in all these points differing from *F. Fordii*.

16. **Fimbristylis Henryi**, *C. B. Clarke*, n. sp.

Fere glabra. *Culmi* annui, dense cæspitosi, 6 uncias longi. *Folia* plura culmos cum inflorescentia superantia,  $\frac{1}{10}$  unc. lata. *Umbellæ* bis ter compositæ, c. 20-stachyæ, rigidæ; bracteæ

inflorescentiam superantes, foliis similes. *Spiculæ*  $\frac{1}{5}$  unc. longæ, anguste oblongæ, teretes, lucide albo-virides. *Glumæ* apice admodum acuminatæ, mucronatæ. *Stylus* persistens, basi bulbosus, nuce longior, glaber; rami 2, stylo multo breviores. *Nux* angustius obovoidea, levis, luteo-brunea; striæ longitudinales numerosæ, pertenues, subobsoletæ.

HUPEH: Ichang (*A. Henry*, 2774!); NORTH CENTRAL CHINA: Hansunfu, Mt. Leansan (*Hugh*!). Mus. Brit.; Herb. Kew.

This species resembles bright examples of *F. dichotoma*; the nut wants the 5-7 strong longitudinal furrows on each face of *F. dichotoma*.

17. **Fimbristylis insignis**, *Thwaites*, *Enum. Pl. Zeyl.* p. 349; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 132 (syn. *Trichelostylis* n. 20 herb. Hook. f. et T. Thoms. excl.); *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 645.

*Fimbristylis Thwaitesii*, *Boeck. in Linnæa*, xxxvii. (1871) p. 34.

*Fimbristylis amblyphylla*, *Steud. Cyp.* p. 116, quoad plantam in obs. descriptam.

KWANGTUNG: Canton (*Hance*, 9858!), Pakwan (*Sampson*, 9858!). Mus. Brit.; Herb. Kew.

Also in Ceylon, Madras, Tonkin, Borneo.

18. **Fimbristylis kwantungensis**, *C. B. Clarke*, n. sp.

Fere glabra. *Culmi* c. 8 uncias longi, graciles, dense cæspitosi. *Folia* plura,  $\frac{1}{12}$  unc. lata, cum  $\frac{1}{3}$  parte culmi subæquilonga. *Umbella* semel composita, laxa, c. 12-stachya; radii 4-7; bracteæ  $\frac{1}{4}$ - $\frac{3}{4}$  unc. longæ, subsetaceæ. *Spiculæ*  $\frac{1}{4}$  unc. longæ, anguste lanceolatæ, læte brunæ. *Glumæ* elliptico-lanceolatæ, caudato-acuminatæ. *Stylus* tenuis, glaber, basi pyramidalis; rami 3. *Nux* parvula, cum  $\frac{1}{3}$  parte glumæ æquilonga, oblongo-ellipsoidea, rotundo-trigona, lutea brunescens, levis, apicem versus plus minus tuberculosa, leviter striata et transversim lineata.

KWANGTUNG (*Ford*, 193!). Herb. Kew.

This species is allied to *F. quinquangularis*, and more closely to *F. debilis*, *F. Mueller*, from North Australia.

19. **Fimbristylis leptoclada**, *Benth. Fl. Hongk.* p. 393, nec *Benth. Fl. Austral.* vii. p. 314; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 647.

*Fimbristylis retusa*, *Thwaites, Enum. Pl. Zeyl.* p. 349; *Boeck. in Linnæa*, xxxvii. (1871) p. 46; *Hance, in Journ. Bot.* xvi. (1878) p. 112.

*Fimbristylis micrantha*, *Boeck. l. c.*

KWANGTUNG: Macao (*Calléry!*), Whampoa (*Hance, 9925!*); HONGKONG (*C. Wright, 579!*). Mus. Brit.; Herb. Kew.

Received also from Ceylon, Malacca, Borneo.

Bentham made a new sp. *F. leptoclada* in the 'Flora Australiensis,' forgetting that he had already used the name for a Chinese plant.

20. *Fimbristylis longispica*, *Steud. Cyp.* p. 118; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 639.

*Fimbristylis Buergeri*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 144; *Debeaux, Fl. Tchefou*, p. 149.

*Fimbristylis spadicea*, var. *major*, *Boeck. in Linnæa*, xxxvii. (1871) p. 20.

*Fimbristylis Didrichseni*, *Boeck. in Engl. Jahrb.* v. (1884) p. 505, var.  $\beta$ . excl.

CHINA (*Horne!*); SHANTUNG: Chefoo (*Debeaux*); KIANGSI: Woosung (*Didrichsen, 3000!*); YUNNAN: Tapintze (*Delavay, 1757!*); HONGKONG (*Faber, 221!*). Herb. Kew.

From Malacca to Japan.

This species differs little from *F. subbispicata*, Nees, but by having more spikelets on the culm; while strong examples hardly differ from the common American *F. spadicea*, Vahl.

21. *Fimbristylis miliacea*, *Vahl, Enum.* ii. (1806) p. 287; *Kunth, Enum. Pl.* ii. p. 230; *Benth. Fl. Hongk.* p. 393; *Boeck. in Linnæa*, xxxvii. (1871) p. 43, et in *Engl. Bot. Jahrb.* vi. (1885) p. 51; *Debeaux, Fl. Tchefou*, p. 151; *Bretschn. Early Research.* p. 181; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 167; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263, et *Pl. David.* i. p. 318; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 644; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 229.

*Scirpus miliaceus*, *Thunb. Fl. Jap.* p. 37; *Burm. f. Fl. Ind.* p. 22, t. 9. fig. 2, non *Linn. herb. propr.*

*Trichelostylis miliacea*, *Nees, in Wight, Contrib.* p. 103, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 226, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 84, et in *Seem. Bot. Voy. 'Herald,'* p. 222.

NORTH CENTRAL CHINA: Hansunfu, Mt. Leansan (*Hugh!*); KIANGSU: Shanghai (*Maingay, 423 partim! 435*); CHEKIANG: Ningpo Mts. (*Faber, 232!*), Chusan (*Yvan!*); KIANGSI: Kiu-

kiang (*Forbes*, 193 !); FOKIEN: Amoy (*Hance*, 1393 !); FORMOSA (*Oldham* !), South Cape (*A. Henry*, 249 !), Takow (*A. Henry*, 786, 786 B, 1113 !); HUPEH: Ichang (*A. Henry*, 4147, 4297, 7785 !); SZECHUEN (*Faber*, 1171 !), Tcentosen, Mt. Uomisan (*Hugh* !); YUNNAN: Tchenfongchan (*Delavay*, 5114 !); KWANGTUNG (*Tate* !), Cape Syngmun (*Meyen* !), Whampoa (*Hance*, 1393 !), Canton (*Sampson*, 272 ! *Hance*, 1393 !), Pakhoi (*Playfair*, 122 !); HONGKONG (*Wilford*, 368 ! *C. Wright*, 584 ! *Hance*, 1393 !); LUCHU ARCHIPELAGO (*Döderlein* !). Mus. Brit.; Herb. Kew.

In rice-fields, in all warm parts of the world; abundant in India, Malaya, China.

22. *Fimbristylis monostachya*, *Hassk. Pl. Jav. Rar.* p. 61; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 132; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 649.

*Cyperus monostachyus*, *Linn. Mant.* p. 180, et in herb. propr.!

*Abildgaardia monostachya*, *Vahl, Enum.* ii. p. 296; *Kunth, Enum. Pl.* ii. p. 247; *Benth. Fl. Hongk.* p. 389; *Boeck. in Linnæa*, xxxvii. (1871) p. 53; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 166.

*Abildgaardia compressa*, *Presl, Rel. Haenk.* i. p. 179; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) pp. 74, 75.

*Abildgaardia Rottboelliana*, *Nees, in Wight, Contrib.* p. 95, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 272.

FORMOSA (*Oldham*, 593 ! *Playfair*, 41 ! *A. Henry*, 732, 759 !), Takow (*A. Henry*, 1029 !); YUNNAN: Tapintze (*Delavay*, 403 !); KWANGTUNG: Macao (*da Silva* ! *Calléry* !); LUCHU ARCHIPELAGO (*Beechey* ! *Döderlein* !). Mus. Brit.; Herb. Kew.

Common through the tropical and warmer parts of the world.

23. *Fimbristylis nigro-brunnea*, *Thwaites, Enum. Pl. Zeyl.* p. 434; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 648.

*Fimbristylis subtetrestachya*, *Boeck. in Linnæa*, xxxvii. (1871) p. 50.

*Fimbristylis pycnostachya*, *Hance, in Journ. Bot.* xv. (1877) p. 338.

*Abildgaardia Eragrostis*, *Boeck. in Linnæa*, xxxvii. (1871) p. 55, pro parte, vix *Nees*.

*Trichelostylis* sp. n. 20, *herb. Ind. Or. Hook. f. et T. Thoms.*

YUNNAN: Tali (*Delavay*, 414 !).

Originally described from Ceylon and South Madras, this species is abundant in North-east Bengal, extending to Muney-poor, Cambodia, Yunnan and the Nicobars. The Yunnan example is the Khasi form *F. subtetrestachya*, *Boeck.* This species



is difficult to separate from *F. Eragrostis*, Hance, affording one more instance of the uncertain foothold afforded by the phyllotaxis of leaves or glumes as a character to stand upon

24. **Fimbristylis nutans**, *Vahl, Enum.* ii. p. 285; *Kunth, Enum. Pl.* ii. p. 221; *Benth. Fl. Hongk.* p. 391; *Boeck. in Linnæa*, xxxvii. (1871) p. 5; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 632.

*Scirpus nutans*, *Retz. Obs.* iv. p. 12.

KWANGTUNG: Cantou (*Sampson*, 1161); HONGKONG (*Harland! C. Wright*, 575, 576! *Hance*, 1259! *Didrichsen*, 3272!). *Mus. Brit.*; *Herb. Kew.*

Southern India, Malaya, Australia to New South Wales.

25. **Fimbristylis podocarpa**, *Hook. f. ex C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 638; *Nees, in Wight, Contrib.* p. 98 partim (char. emend.), *et in Hook. et Arn. Bot. Beechey's Voy.* p. 225 partim, *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 77, var.  $\gamma$  et  $\alpha$  partim nec  $\beta$ .

*Fimbristylis communis* (forma), *Kunth, Enum. Pl.* ii. p. 234.

*Fimbristylis polymorpha* (forma), *Boeck. in Linnæa*, xxxvii. (1871) p. 14.

HONGKONG (*Hance*, 1091! *Forbes!*). *Mus. Brit.*; *Herb. Kew.*

Frequent in India, extending to Java and the Marianne Isles.

The species, as here understood, is destitute of the 5-7 longitudinal furrows so prominent on each face of the nut in *F. diphylla*, *Vahl* (*F. communis*, *Kunth*, or *F. polymorpha*, *Boeck.*).

In my old Chinese list, this Hance 1091 is noted as in herb. Kew, but I do not now find it. Possibly it is on a mixed sheet somewhere.

26. **Fimbristylis polytrichoides**, *Vahl, Enum.* ii. p. 248 (polytrichoides); *R. Br. Prodr.* p. 226; *Hance, in Journ. Bot.* xvi. (1878) p. 112; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 632.

*Fimbristylis juncea*, *Boeck. in Linnæa*, xxxvii. (1871) p. 4, non *Roem. et Schult.*

*Scirpus polytrichoides*, *Retz. Obs.* iv. p. 11.

*Abildgaardia javanica*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 63, *et Cyp.* p. 72.

FOKIEN: Amoy (*Hance*, 10153!); FORMOSA: Takow (*A. Henry*, 1100!). Mus. Brit.; Herb. Kew.

Tropical Africa, South India, Philippines.

Var.  $\beta$ . *halophila*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 632.

*Fimbristylis Didrichseni*, var.  $\beta$ . *minor*, *Boeck. in Engl. Jahrb.* vi. (1884) p. 505.

CHUSAN ARCHIPELAGO (*Didrichsen*, 3569!). Herb. Kew.  
Calcutta; Madras.

27. *Fimbristylis quinquangularis*, *Kunth, Enum. Pl.* ii. (1837) p. 229; *Boeck. in Linnæa*, xxxvii. (1871) p. 42 (syn. *F. salbundia*, *Thwaites*, excl.); *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 644.

*Fimbristylis salbundia*, *Boeck. in Linnæa*, xxxvii. (1871) p. 44.

*Scirpus quinquangularis*, *Vahl, Enum.* ii. p. 279.

*Scirpus miliaceus*, *Linn. herb. propr.* (an *Sp. Pl.* ed. ii. p. 75?).

CHEKIANG: Ningpo Mts. (*Faber*, 237!); YUNNAN: Tapintze (*Delavay*, 1763!). Mus. Brit.; Herb. Kew.

Abundant in rice-fields and on banks throughout India and Malaya, extending to the Philippines, Tropical Australia, and Mauritius.

28. *Fimbristylis rigidula*, *Nees, in Wight, Contrib.* p. 99; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 640; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 131, syn. et *Thwaitesii* exemplo excl.; *Diels, in Engl. Jahrb.* xxix. (1901) p. 229.

*Fimbristylis Hanceana*, *Boeck. in Linnæa*, xxxviii. (1874) p. 394.

*Fimbristylis communis*, *Kunth, Enum. Pl.* ii. p. 235 partim.

9 — KIANGSI: Kiukiang (*Shearer*!); HUPEH: Hankow (*Faber*, 1133!), Ichang (*A. Henry*, 545, 1234 partim!); YUNNAN: Mengtze (*A. Henry*, 13728!), Tapintze (*Delavay*, 404, 1755!), Mosoyñ (*Delavay*, 4464!); HONGKONG (*Hance*, 1188!). Mus. Brit.; Herb. Kew.

Common in North India, at 1000–5000 feet; also in the Philippines.

The examples named "*rigidula*, *Nees*," which were in the hands of *Kunth* and of *Boeckeler*, were correctly referred by them to our *F. diphylla*, *Vahl*. The true *F. rigidula*, *Nees*, is well marked by the woody rhizome and by the many paired spikelets.

29. *Fimbristylis schœnoides*, *Vahl, Enum.* ii. (1806) p. 286; *Kunth, Enum. Pl.* ii. p. 222; *Benth. Fl. Hongk.* p. 391; *Boeck. in Linnæa*, xxxvii. (1871) p. 5, var.  $\beta$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 634.

*Fimbristylis bispicata*, *Nees, in Wight, Contrib.* p. 97, pro majore parte; *Boeck. in Linnæa*, xxxvii. (1871) p. 6 partim.

*Fimbristylis polymorpha*, var. *depauperata*, *Boeck. in Flora*, lviii. (1875) p. 111.

*Scirpus schœnoides*, *Retz. Obs.* v. p. 14.

*Scirpus bispicatus*, *Roxb. Hort. Beng.* p. 6, et *Fl. Ind.* i. p. 220.

*Abildgaardia nervosa*, *Presl, Rel. Haenk.* i. p. 180.

SHANTUNG: Chefoo (*Forbes!*); FORMOSA: Takow (*A. Henry, 1912!*); KWANGTUNG: Canton (*Sampson, 267!*); HONGKONG (*C. Wright, 578, 580! Hance, 7479! Lamont, 846!*). Mus. Brit.; Herb. Kew.

Common throughout India; extending thence to the Philippines and to Queensland.

30. *Fimbristylis sericea*, *R. Br. Prodr.* p. 228, char. emend.; *Kunth, Enum. Pl.* ii. p. 244; *Boeck. in Linnæa*, xxxvii. (1871) p. 22; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 641.

*Fimbristylis decora*, *Nees, in Wight, Contrib.* p. 101, in *Hook. et Arn. Bot. Beechey's Voy.* p. 225, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 83; *Kunth, Enum. Pl.* ii. p. 240; *Henriq. in Bol. Soc. Broter.* ii. (1883) p. 167.

*Fimbristylis velutina*, *Franch. in Bull. Soc. Bot. France*, xxvi. (1879) p. 88.

*Scirpus sericeus*, *Poir. Encycl. Suppl.* v. p. 99.

CHINA (*Meyen! Millett! Parkes!*); FOKIEN: Amoy (*Didrichsen, 3044!*); FORMOSA: Takow (*A. Henry, 1007, 1008!*); KWANGTUNG: Macao (*da Silva! Hance, 10131! Tate!*), Canton (*Yvan!*), Pakhoi (*Playfair, 86!*); HAINAN (*A. Henry, 8195!*). Mus. Brit.; Herb. Kew.

On sandy sea-shores in South-east Asia and Tropical Australia, extending from Coromandel to Japan.

31. *Fimbristylis Sieboldii*, *Miq. Cat. Mus. Bot. Lugd.-Bat.* (1870) p. 118, name only; *Franch. et Sav. Pl. Jap.* ii. p. 118; *Debeaux, Fl. Tchefou*, p. 149; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263.

*Fimbristylis leiocarpa*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 144, non *Maxim.*

SHANTUNG: Chefoo (*Debeaux!*).

Also in Japan.

32. *Fimbristylis spathacea*, *Roth, Nov. Pl. Sp.* p. 24; *Kunth, Enum. Pl.* ii. p. 246; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 640.

*Fimbristylis Wightiana*, *Nees, in Wight, Contrib.* p. 99; *Kunth, Enum. Pl.* ii. p. 241; *Benth. Fl. Hongk.* p. 392.

*Fimbristylis glomerata*, *Nees, in Linnæa*, ix. (1834) p. 290, nomen; *Kunth, Enum. Pl.* ii. p. 246; *Boeck. in Linnæa*, xxxvii. (1871) p. 47 partim, *i. e.* plantis 3-stigmatis excl.

*Fimbristylis rigida*, *Kunth, Enum. Pl.* ii. p. 246; *Henriq. in Bol. Soc. Broter.* ii. (1883) p. 167.

*Fimbristylis junciformis*, *Munro, in Seem. Bot. Voy. 'Herald,'* p. 422, non *Kunth*.

*Fimbristylis biumbellata*, *Boeck. in Flora*, xli. (1858) p. 603; cf. *Boeck. in Flora*, xliii. (1860) p. 34.

*Scirpus glomeratus*, *Retz. Obs.* iv. p. 11.

*Isolepis rigescens*, *Presl, in Oken, Isis*, xxi. (1828) p. 268.

*Isolepis Haenkei*, *Presl, Rel. Haenk.* i. p. 187 saltem partim.

FORMOSA: Takow (*A. Henry*, 1071, 1071 A, 1834, 1860! *Playfair*, 136!); KWANGTUNG: Macao (*da Silva*!); HONGKONG (*Seemann*, 542! *Munro*! *C. Wright*, 589! *Lamont*! *Hance*!). *Mus. Brit.*; *Herb. Kew.*

A common tropical plant near the sea, from Arabia and the Mascarene Isles to the Philippines; also in Trop. America.

Some examples of this have the umbel contracted nearly into a head, and the species has thus been merged into *F. cymosa*, R. Br., by Boeckeler. In the fine specimens of A. Henry (1071, 1834) the umbel is open, compound.

33. *Fimbristylis squarrosa*, *Vahl, Enum.* ii. p. 289; *Kunth, Enum. Pl.* ii. p. 224; *Benth. Fl. Hongk.* p. 392; *Boeck. in Linnæa*, xxxvii. (1871) p. 10; *Debeaux, Fl. Tchefou*, p. 150; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 635; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263.

*Scirpus squarrosus*, *Poir. Encyc. Suppl.* v. p. 100, non *Linn.*

*Pogonostylis squarrosa*, *Bertol. Fl. Ital.* i. p. 312.

SHANTUNG: Chefoo (*Debeaux*, 42! *Fauvel*!); FORMOSA: Tam-sui (*Oldham*, 591!); YUNNAN: Tali (*Delavay*, 2629!); HONGKONG (*Harland*! *C. Wright*, 581! *Wilford*, 267!); HAINAN (*Swinhoe*!). *Mus. Brit.*; *Herb. Kew.*

A rice-field annual, extending from South Europe to the Cape of Good Hope, from Amurland to New Zealand, from Cuba to Buenos Ayres; but not known in North America.

34. *Fimbristylis Stauntoni*, *Debeaux*, *Fl. Tchefou* (1877), p. 151, t. 3. fig. 1, char. emend.; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263.

SHANTUNG (*Debeaux*, 144!), Chefoo (*Fauvel*); KIANGSU: Shanghai (*Maingay*, 423 in hb. Kew!); CHEKIANG (*Staunton*!); HUPEH: Ichang (*A. Henry*, 2774!). Mus. Brit.; Herb. Kew.

This is a critical species, characterized by the very parallel-sided oblong nut. The style varies from 2-fid to 3-fid (a rare variation in the genus *Fimbristylis*); I noted at Calcutta that Maingay n. 423 contains *F. Stauntoni* and *F. miliacea* mixed.

35. *Fimbristylis stolonifera*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. (1893) p. 637.

YUNNAN: Tapintze (*Delavay*, 1756, 1757, 1768!).

North-east India; plentiful in the hills.

Stolons are very rare in the genus *Fimbristylis*.

36. *Fimbristylis subbispicata*, *Nees et Meyen*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 75; *Benth. Fl. Hongk.* p. 391; *Hance*, in *Journ. Bot.* xvii. (1879) p. 16; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 634.

*Fimbristylis japonica*, *Sieb. et Zucc. ex Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62, et *Cyp.* p. 107.

*Fimbristylis bispicata*, *Nees*, in *Hook. et Arn. Bot. Beechey's Voy.* p. 224, neque in *Wight, Contrib.* p. 97; *Boeck. in Linnæa*, xxxvii. (1871) p. 6 partim.

CHIHLI: Peking (*Hance*), Tachiaotze (*Möllendorff*); SHINGKING: Kinchow (*Ross*, 195!); SHANTUNG (*Debeaux*!), Chefoo (*Forbes*, 391, 392! *Fauvel*! *Schottmueller*, 344!); KIANGSU: Shanghai (*Maingay*, 641, 643, 691!); FOKIEN: Amoy (*Hance*, 998! *Lamont*, 845!); KWANGTUNG: Macao (*Gaudichaud*! *Vachell*! *Calléry*, 380!); HONGKONG (*Harland*, 681! *C. Wright*, 577! *Faurie*, 15775!). Mus. Brit.; Herb. Kew.

Common in Japan; as in China mainly by the sea-side, extending throughout Malaya and to Orissa.

The limits of this species are not settled; Boeckeler places here the mountain form of *F. schænoides*. The smaller form with the head usually solitary and only  $\frac{1}{2}$ - $\frac{1}{4}$  in. long is *F. cæsia*, *Miq.* (*Fl. Ind. Bat.* iii. p. 315); and to this subspecies or var.

belongs Maingay n. 643. Nees's original *F. bispicata*, founded in Wight, Contrib. 1834, p. 97, was, as Arnott saw, typical *F. schænoides*; his *F. bispicata*, in Hook. et Arn. Beechey's Voy. p. 224 [published in 1836], was Vachell's plant, *F. subbispicata*. It is preferred, in such confusion by the author himself, to follow Bentham in taking up the later name, *F. subbispicata*.

37. **Fimbristylis tetragona**, *R. Br. Prodr.* p. 226; *Kunth, Enum. Pl.* ii. p. 242; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 631.

*Fimbristylis cylindrocarpa*, *Kunth, Enum. Pl.* ii. p. 222; *Boeck. in Linnæa*, xxxvii. (1871) p. 7; *Hance, in Journ. Linn. Soc.* xiii. (1873) p. 132.

*Fimbristylis abjiciens*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 62, et *Cyp.* p. 107.

*Fimbristylis Arnottii*, *Thwaites, Enum. Pl. Zeyl.* p. 348.

*Scirpus tetragonus*, *Poir. Encycl. Suppl.* v. p. 98, neque in *Lam. Encycl.* vi. p. 767.

*Mischospora efoliata*, *Boeck. in Flora*, xliii. (1860) p. 113.

KWANGTUNG: Canton (*Sampson*, 475!), Wongneichung (*Hance*, 7484!); HONGKONG (*Lamont*, 846, 854 A!), Lofaushan (*Ford*!). Mus. Brit.; Herb. Kew.

Common throughout India, extending thence to New South Wales and South China.

The example of this in herb. Hance is marked by Hance "*F. schænoides*, Vahl," to which it bears strong superficial resemblance.

38. **Fimbristylis Thomsonii**, *Boeck. in Linnæa*, xxxvii. (1871) p. 37; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 646.

*Fimbristylis complanata*, *Benth. Fl. Hongk.* p. 393, vix *Link.*

*Trichelostylis complanata*, *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 226, non *Nees, in Wight, Contrib.* p. 103.

*Trichelostylis* sp. n. 12, *herb. Ind. Or. Hook. f. et T. Thoms.*

CHINA (*Millett*!); YUNNAN: Mengtze, 5000 feet (*A. Henry*, 13729!); KWANGTUNG: Macao (*Vachell*, 61! *Lamont*, 52!); HONGKONG (*Harland*! *Seemann*, 539! *C. Wright*, 583! *Hance*, 1188! 1255! *Furet*, 377!). Mus. Brit.; Herb. Kew.

Bengal and Assam, frequent; extending thence through Burma to China and Tonkin.

This species is a split-off by Boeckeler from *F. complanata*,

Link. Munro had Seemann n. 539, but did not name it; nor do I know how he catalogued it in Bot. Voy. 'Herald.'

39. *Fimbristylis yunnanensis*, C. B. Clarke, n. sp.

Undique sparsim tenuiter pilosa. Culmi 8-15 unc. longi, graciles, præter basin nudi, monocephali, tenuiter pilosi. Folia cum  $\frac{1}{3}$ - $\frac{1}{2}$  parte culmi æquilonga,  $\frac{1}{8}$  unc. lata. Caput globosum,  $\frac{1}{2}$  unc. in diam., stellatim 7-12-stachyum; bracteæ c. 3, usque ad  $\frac{1}{2}$ -1 unc. longæ, subsetaceæ. Spiculæ  $\frac{1}{4}$  unc. longæ, bruneæ, minute pilosæ, 7-10-glumæ, 2-4-nuciferæ. Glumæ subdistichæ, naviculares, mucronatæ. Stylus linearis, glaber, basi pyramidalis, persistens, rami 3. Nux parva, cum  $\frac{1}{3}$ - $\frac{1}{2}$  parte glumæ æquilonga, obovoidea, rotundo-trigona, obscure reticulata, alba.

YUNNAN: Mengtze, 5000 feet (A. Henry, 11150!). Herb. Kew. -2

This species is very closely allied to *F. Actinoschænus*, the structure of the spikelet being identical.

1. *Bulbostylis barbata*, Kunth, ex C. B. Clarke, in Hook. f. *Fl. Brit. Ind.* vi. p. 651; Palibin, *Fl. Koreæ*, iii. (1901) p. 18.

*Scirpus barbatus*, Rottb. *Descr. et Ic.* p. 52, t. 17. fig. 4; Hance, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 131; Boeck. in *Linnæa*, xxxvi. (1869-70) p. 751, var.  $\beta$  excl.; Franch. in *Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263.

*Scirpus capillaris*, Linn. *Sp. Pl.* ed. 2, p. 73 partim, nec Linn. *Mant.*; cf. R. Br. *Prodr.* p. 122, in Obs.

*Isolepis barbata*, R. Br. *Prodr.* p. 222; Nees, in Hook. et Arn. *Bot. Beechey's Voy.* p. 226; Benth. *Fl. Hongk.* p. 393; Henriq. in Bolet. Soc. Broter. ii. (1883) p. 67.

*Isolepis capillaris*, Debeaux, *Fl. Tchefou*, p. 148, fide Franchet, non Rœm. et Schult.

*Fimbristylis barbata*, Benth. *Fl. Austral.* vii. p. 321.

SHANTUNG (Maingay, 142, 149!), Chefoo (Fauvel! Debeaux! Schottmueller, 300!), Chefoo Mts. (Faber, 72!); KIANGSI: Kiu-kiang (Shearer!); FORMOSA: Tamsui (Oldham, 590!); YUNNAN: Mosoyñ (Delavay, 3971!); KWANGTUNG: Macao (Vachell!), Whampoa (Hance, 996!), Canton (Sampson, 644!), Pakhoi (Playfair, 210!); HONGKONG (Harland! C. Wright, 591!); HAINAN (A. Henry, 8241!). Mus. Brit.; Herb. Kew.

In tropical and warm regions of the Old World, very common. In South U.S. rare, perhaps only introduced.

2. **Bulbostylis capillaris**, *Kunth*, ex *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 652.

*Scirpus capillaris*, *Linn. Sp. Pl.* ed. 2, p. 73, pro magnâ parte.

From Canada to Patagonia, abundant.

Var. **trifida**, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 652.

*Bulbostylis trifida*, *Kunth*, *Enum. Pl.* ii. p. 213.

*Scirpus capillaris*, *Linn. Mant.* p. 321, et herb. propr.; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263.

*Scirpus trifidus*, *Hance*, in *Journ. Bot.* xvi. (1878) p. 112.

*Isolepis trifida*, *Nees*, in *Wight, Contrib.* p. 108.

*Fimbristylis capillaris*, *F. Muell. Fragm. Phyt. Austral.* ix. p. 7.

SHANTUNG (*Maingay*, 108!), Chefoo (*Fauvel*!); CHEKIANG: Ningpo Mts. (*Faber*, 1738!); SZECHUEN: Tchenkeoutin (*Farges*, 1099 ter!); YUNNAN: Tali (*Delavay*, 3639!), Tchenfongchan (*Delavay*, 5116!); KWANGTUNG: Mt. Pakwan (*Hance*!). *Mus. Brit.*; *Herb. Kew.*

In tropical and warm countries, very common.

1. **Scirpus Caricis**, *Retz. Fl. Scand.* p. 11; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 660.

*Scirpus compressus*, *Pers. Syn.* i. p. 66; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 707.

*Schoenus compressus*, *Linn. Sp. Pl.* ed. 1, p. 43; ed. 2, p. 65.

*Blysmus compressus*, *Link, Hort. Berol.* i. p. 278, ii. p. 310; *Kunth, Enum. Pl.* ii. p. 328.

YUNNAN: Mosoyn (*Delavay*, 4463!), Kanhaytze (*Delavay*).

Common in the Palæarctic Region, extending south to the Himalaya.

2. **Scirpus erectus**, *Poir. in Lam. Encycl.* vi. p. 761; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 656; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 228.

*Scirpus debilis*, *Pursh, Fl. Amer. Sept.* i. p. 55; *Kunth, Enum. Pl.* ii. p. 159; *Benth. Fl. Austral.* vii. p. 332.

*Scirpus juncoides*, *Roxb. Hort. Beng.* p. 81, et *Fl. Ind.* i. p. 216; *Kunth, Enum. Pl.* ii. p. 160; *Benth. Fl. Hongk.* p. 395.

*Scirpus luzonensis*, *Presl, Rel. Haenk.* i. p. 193; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 93.

*Scirpus timorensis*, *Kunth, Enum. Pl.* ii. p. 162; *Hance, in Journ. Bot.* xvi. (1878) p. 112.

*Scirpus supinus*, varr.  $\beta$ ,  $\gamma$ , *Boeck. in Linnæa*, xxxvi. (1869-70) p. 600.

CHIHLI: Peking (*Bretschneider*! *Möllendorff*!); SHANTUNG



(*Maingay*, 159!), Chefoo Mts. (*Faber*, 1517!); Kiangsi: Kiu-kiang (*Forbes*, 192!); FORMOSA: Takow (*Playfair*, 171! *A. Henry*, 1088!); HUPEH: Ichang (*A. Henry*, 2740, 4212, 4296!), Patung (*A. Henry*, 78!), Yangtzekiang (*Faber*, 1123!); SZECHUEN (*Farges*!); YUNNAN: Tapintze (*Delavay*, 1771, 1778!), Mosoyu (*Delavay*, 4075! rhizomate longius repente); KWANGTUNG: Macao (*Calléry*, 426!), Canton (*Nevin*!); HONGKONG (*Wilford* 272! *Harland*!). Mus. Brit.; Herb. Kew.

Extends from Madagascar to Amurland, the Philippines, and Tropical Australia. The American form extends from Georgia to Labrador.

3. *Scirpus Eriophorum*, *Michx. Fl. Bor.-Amer.* i. p. 33 *Boeck. in Linnæa*, xxxvi. (1869-70) p. 731; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 661; *Boeck. in Engl. Bot. Jahrb.* vi. (1885) p. 51?

*Eriophorum cyperinum*, *Linn. Sp. Pl.* ed. 2, p. 77; *Benth. et Hook. f. Gen. Pl.* iii. p. 1052.

SHANTUNG: Chefoo Mts. (*Faber*, 68!); CHEKIANG: Tientai Mts. (*Faber*, 70!); HUPEH (*E. H. Wilson*, 1632!), alt. 6700 feet (*E. H. Wilson*, 1637 a!); SZECHUEN (*A. Henry*, 8836, 8912!), Tachienlu, alt. 9000-13,500 feet (*Pratt*, 420!); LUCHU ARCHIPELAGO (*Döderlein*). Herb. Kew.

Very common in cooler North America and in Japan, extending to the East Himalaya and Khasi Mountains, where it is frequent.

Bentham and Hooker put this species back into *Eriophorum*, on the stated ground that the setæ are altogether as in *Eriophorum alpinum* and not as in *Scirpus*. I find the setæ in *Scirpus Eriophorum* very slender, setaceous, undivided, as of *S. sylvaticus*, Linn., while in *E. alpinum* the setæ are ligulate, often, sometimes deeply, laciniate. But it may well be held that the whole section of *Scirpus* called *Sylvatici* (= *Seidlia*, genus Opiz) is more nearly allied to *Eriophorum* than to *Scirpus* by the leafy stem and elongate inflorescence.

4. *Scirpus filipes*, *C. B. Clarke*, n. sp.

Glabra. *Rhizoma* robustius, repens. *Culmi* 12-20 unc. longi, graciliores, per totam longitudinem remote nodosi, apice triquetri. *Folia* 8-15 unc. longa, 2 lin. lata. *Umbellæ* radii 1-6, pertenues, inæquales, 1-6-stachyi; *bractea* ima longa (usque ad

1 pedem), suberecta. *Spiculæ* 4 lin. longæ, digitato-fasciculatæ. *Glumæ* elliptico-oblongæ, integræ, in lateribus rubro-brunæ. *Setæ* 6, nucem superantes, rubro-ferrugineæ, subretrorsim scabræ. *Stylus* 3-fidus. *Nux* parva, anguste obovoidea, apice acuminata, luteo-brunæ.

FOKIEN (*De Grijs*, herb. *Hance*, 6695 !). Herb. Kew.

Also, without habitat, herb. Link, 1028 ! herb. Berlin 889 !

From the nodes on the stem, from the acuminate top to the nut, and from the setæ, I should think this species allied to the Japanese *S. fuirenoides*, Maximowicz, but Boeckeler has marked three sheets of it in herb. Berlin "*S. maritimus*, Linn."

5. *Scirpus fuirenoides*, *Maxim. MS. in hb. Kew.*

Glabra, elata. *Culmus* per totam longitudinem nodosus, foliiger. *Folia* saltem 18 unc. longa,  $2\frac{1}{2}$  lin. lata, lenta. *Corymbi* longepedunculati, in paniculam longam 18 unc. dispositi; terminalis bis compositus radiis radiolisque longis; *bracteæ* cum foliis consimiles, paniculam superantes. *Spiculæ* 5-9nim fasciculatæ,  $2\frac{1}{2}$  lin. longæ, 2 lin. latæ (quam in sp. affinibus majores), virides, rubescentes, densæ. *Glumæ* 1 lin. longæ, ellipticæ obtusæ. *Setæ, nux, stylus* omnino ut *S. sylvatici*.

SHANTUNG: Chefoo Mts. (*Faber*, 67 !). Herb. Kew.

Many examples received from Japan.

This is a typical *Seidlia*, i. e. one of the *Sylvatici*, and close allied to *S. polystachyus*, F. Muell.; but the spikelets are twice as broad.

6. *Scirpus grossus*, *Linn. f. Suppl.* p. 104; *Presl, Rel. Haenk.* i. p. 195; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 723; *Hook. f. Fl. Brit. Ind.* vi. p. 659.

Hymenochæte grossa, *Nees, in Wight, Contrib.* p. 110.

Isolepis maxima, *A. Dietr. Sp. Pl.* ii. p. 130.

KWANGSI: Lungchow (*A. Henry*, 684 !). Herb. Kew.

Frequent in India, Malaya, extending through Tonkin to the Philippines.

7. *Scirpus lacustris*, *Linn. Sp. Pl.* ed. 1, p. 48, ed. 2, p. 72, et herb. propr.; *Bunge, Enum. Pl. Chin. Bor.* p. 68; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 712; *Regel, Fl. Ussur.* p. 176; *Hook. et Arn. Bot. Beechey's Voy.* p. 98; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 658.

*Scirpus Tabernæmontani*, *C. C. Gmel. Fl. Baden.* i. p. 101; *Regel, Fl. Ussur.* p. 176; *Maxim. Prim. Fl. Amur.* p. 298.

*Scirpus Meyenii*, *Nees, in Linnæa*, ix. (1834) p. 293 nomen, *et in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 293.

CHIHLI: Peking (*Bretschneider*, 796!), Kantai et Tanschan (*Bunge*!); FORMOSA (*A. Henry*, 754, 1028, 1057!), Takow (*A. Henry*, 1059!); YUNNAN: Mengtze 5500 feet (*A. Henry*, 11135!), Szemao, "cultivated for making mats" (*A. Henry*, 12194!); LUCHU ARCHIPELAGO (*C. Wright*, 356!). *Mus. Brit.*; *Herb. Kew.*

Frequent in fresh water nearly throughout the world, except South America and Malaya.

8. *Scirpus maritimus*, *Linn. Sp. Pl.* ed. 1, p. 51, ed. 2, p. 74, *et herb. propr.*; *Bunge, Enum. Pl. Chin. Bor.* p. 68; *Maxim. Prim. Fl. Amur.* p. 298; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 722; *Hook. et Arn. Bot. Beechey's Voy.* p. 98; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, *et Pl. David.* i. p. 317; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 658.

CHIHLI: Peking (*Bretschneider*, 784!), Kantai et Tanschan (*Bunge*), Jehol (*David*, 1909!); SHINGKING: Moukden (*Webster*, 104!); NORTH CENTRAL CHINA (*Hugh*!); FORMOSA: Takow (*A. Henry*, 1818!); HUPEH: Yangtzekiang (*Faber*, 1126!). *Mus. Brit.*; *Herb. Kew.*

Throughout the Old World, inland as well as near the sea; less common in the tropics. Varieties extend throughout America.

Var.  $\beta$ . *affinis*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 659.

*Scirpus affinis*, *Roth, Nov. Pl. Sp.* p. 30.

*Scirpus strobelinus*, *Roxb. Hort. Beng.* p. 6, *et Fl. Ind.* i. p. 219.

*Scirpus macrostachys*, *Boeck. Cyp. Novæ*, i. (1888) p. 19.

*Scirpus maritimus*, var. (verosimiliter sp.), *Maxim. Prim. Fl. Amur.* p. 299.

SHANTUNG: Chefoo (*Perry*!). *Herb. Kew.*

This well-marked variety is plentiful in India; extends to Turcomania, Songaria, Amurland, and Burma.

9. *Scirpus Michelianus*, *Linn. Sp. Pl.* ed. 1, p. 52, ed. 2, p. 76; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 662.

*Isolepis Micheliana*, *Roem. et Schult. Syst.* ii. p. 114; *Maxim. Prim. Fl. Amur.* p. 300; *Debeaux, Fl. Tchefou*, p. 147.

*Cyperus pygmæus*, var.  $\beta$ , *Boeck. in Linnæa*, xxxv. (1867-8) p. 494, et in *Flora*, liv. (1871) p. 159; *Benth. et Hook. f. Gen. Pl.* iii. p. 1044 pro parte.

*Fimbristylis Micheliana*, *Reichb. Fl. Germ. Excurs.* p. 75.

CHIHLI: Peking (*Williams*, 12187!); SHINGKING: Moukden (*James*); SHANTUNG (*Staunton! Debeaux*, 199!); KIANGSU: Shanghai (*Maingay*, 634! *Faber*, 230!), Tsien Mts. (*Faber*, 1513!); NORTH CENTRAL CHINA: Hukiascien (*Hugh!*); YUNNAN: Tali (*Delavay*, 2628!); KWANGTUNG: Canton (*Debeaux*). *Mus. Brit.*; *Herb. Kew.*

Frequent throughout the Palæarctic area, extending in East Asia to Tonkin, Java, and Japan.

10. *Scirpus mucronatus*, *Linn. Sp. Pl.* ed. 1, p. 50, ed. 2, p. 73, pro majore parte et herb. propr.; *Kunth, Enum. Pl.* ii. p. 161; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 703; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 657; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 228.

*Scirpus triangulatus*, *Roxb. Hort. Beng.* p. 81, et *Fl. Ind.* i. p. 217; *Hance, MS. in herb. propr.*

*Scirpus cognatus*, *Hance, in Ann. Sc. Nat. sér. 4*, xv. (1861) p. 228.

FORMOSA: Takow (*A. Henry*, 1190!); HUPEH (*A. Henry*, 4213, 4255!), Ichang (*A. Henry*, 2295, 2930!); KWANGTUNG: Whampoa (*Hance*, 6682!), Canton (*Sampson*, 266!). *Mus. Brit.*; *Herb. Kew.*

Old World; extending from Portugal and Madagascar to Japan and New South Wales; abundant in India.

11. *Scirpus radicans*, *Schkuhr, in Usteri, Ann.* iv. (1793) p. 49, t. 1; *Maxim. Prim. Fl. Amur.* p. 299; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 727; *Franch. Pl. David.* i. p. 317.

*Scirpus sylvaticus*, var. *radicans*, *Willd. Sp. Pl.* i. p. 308; *Regel, Fl. Ussur.* p. 177.

*Seidlia radicans*, *Opiz, Natur. Tausch.* ix. p. 349.

CHIHLI: Jehol (*David*).

Common in North Asia, extending west to Germany, east to Japan. It is, in analytical tables, differentiated from *S. sylvaticus* by the pedicellate, not clustered, spikelets. As pedicellate as well as clustered spikelets occur frequently on one culm, there is some confusion in the synonymy, and such examples constitute

the *S. sylvatico-radicans*, Baenitz, assumed a hybrid, which it is not. There exist many paper hybrids, made out of analytical tables.

12. *Scirpus Rosthornii*, Diels, in *Engl. Bot. Jahrb.* xxix. (1901) p. 228.

HUPEH: Nan Chuan (*Rosthorn*).

Not seen. One of the *Sylvatici* (*Siedlia*, Opiz), and on account of the 2-fid style placed by Diels next *S. ternatensis*, Miq. (*S. chinensis*, Monro). The general description of Diels, and the habitat, might do for *S. fuirenoides*, Maxim.; but in this the style is always 3-fid.

13. *Scirpus setaceus*, Linn. *Sp. Pl.* ed. 2, p. 73 partim, neque herb. propr.; *Sm. in Sowerby, Engl. Bot.* t. 1693; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 500, varr.  $\beta$ ,  $\gamma$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 654.

*Isolepis setacea*, *R. Br. Prodr.* p. 222.

NORTH CENTRAL CHINA: Thaepeisan (*Hugh!*). Mus. Brit.

Throughout Europe and West Asia; extends to the Cape, India, and Australia.

14. *Scirpus squarrosus*, Linn. *Mant.* p. 181, et herb. propr.; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 734; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 263; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 663.

*Scirpus chinensis*, *Osbeck, Voy. ed. Forster*, i. p. 354; non *Munro*; cf. *Bretschn. Early Research.* p. 111 [*S. sinensis*].

*Scirpus* sp. n. 65, *Rottb. Descr. et Ic.* p. 49, t. 17. fig. 5.

*Ascolepis tenuior*, *Steud. Cyp.* p. 105.

*Lipocarpa microcephala*, *Hance, in Ann. Sc. Nat. sér. 5*, v. (1866) p. 249, non *R. Br.*

CHEKIANG (*Staunton!*); KWANGTUNG: Whampoa (*Hance, 10152!*). Mus. Brit.; Herb. Kew.

In the tropical and warm regions of Africa and Asia; widely spread, common in India.

15. *Scirpus Stauntoni*, *C. B. Clarke*, n. sp.

Annua, cæspitosa. Culmi 3-4 unc. longi, monocephali. Folia cum culmo fere æquilonga, fere setacea, omnia fere ad basin culmi sita. Caput 4 lin. in diam., globosum, polystachyum; bracteæ 7, setacæ, usque ad 2½ unc. longæ. Spiculæ iis

*S. Micheliani* similes. *Glumæ* ovatæ, acuminatæ, apice obtusæ. *Setæ* hypogynæ 0. *Stylus* trifidus. *Nux* parvula, obovoidea, acutius, subæqualiter trigona, cellulis extimis quadratis astrictis.

CHEKIANG (*Staunton!*). Mus. Brit.

This may be a variety of *S. Michelianus*; but it does not match it satisfactorily, and the nut is obovoid, instead of oblong with parallel sides and hyaline cells.

16. *Scirpus subcapitatus*, *Thwaites, Enum. Pl. Zeyl.* [1864] p. 351; *Boeck. in Linnæa*, xxxvi. (1869–70) p. 704; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 661.

*Scirpus subaphyllus*, *Boeck. MS.*

KIANGSI: Kiukiang (*Shearer! Faber, 1135! Maries!*). Herb. Kew.

Ceylon and mountains of South India. Also from Malaya, but Dr. Stapf considers the Bornean plant specifically distinct.

17. *Scirpus supinus*, *Linn. Sp. Pl.* ed. 1, p. 49, ed. 2, p. 73, neque herb. propr.; *Boeck. in Linnæa*, xxxvi. (1869–70) p. 699, varr.  $\beta$ ,  $\gamma$  excl.; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 655; *Bretschn. Early Research.* p. 181.

*Isolepis supina*, *R. Br. Prodr.* p. 221; *Kunth, Enum. Pl.* ii. p. 196; *Benth. Fl. Hongk.* p. 394.

HONGKONG (*Wilford, 271! C. Wright, 590! Hance!*). Mus. Brit.; Herb. Kew.

Extends from France to South Australia, in all the warmer parts of the Old World; abundant in India. Also a variety in North America.

18. *Scirpus sylvaticus*, *Linn. Sp. Pl.* ed. 1, p. 51, ed. 2, p. 75, et herb. propr.; *Boeck. in Linnæa*, xxxvi. (1869–70) p. 726; *Maxim. Prim. Fl. Amur.* p. 299; *Regel, Fl. Ussur.* p. 177, var.  $\gamma$  excl.

*Scirpus microcarpus*, *Presl, Rel. Haenk.* i. p. 195.

*Seidlia sylvatica*, *Opiz, Natur. Tausch.* ix. p. 349.

SHINGKING: Moukden (*James!*).

Through the sub-arctic and cool north temperate zone, extending in America southwards to California.

19. *Scirpus ternatensis*, *Reinw. ex Miq. Fl. Ind. Bat.* iii. p. 307, et in *De Vriese, Pl. Reinw.* p. 140; *C. B. Clarke, in Journ. Linn. Soc., Bot.* xxxiv. (1898) p. 83.

*Scirpus chinensis*, *Munro*, in *Seem. Bot. Voy. 'Herald'*, p. 423; *Benth. Fl. Hongk.* p. 395; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 662; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 228.

*Scirpus Eriophorum*, var. *conglomeratus*, *Boeck. MS. in herb. Berol.*

FORMOSA (*A. Henry*, 593!), South Cape (*A. Henry*, 244!); Takow (*A. Henry*, 609!), Tamsui (*Oldham*, 589); HUPEH (*Faber*, 1120!), Ichang (*A. Henry*, 1400, 1854, 3983!); SZECHUEN: Tchenkeoutin (*Farges*, 1099 bis); YUNNAN: Mengtze 6000 feet (*A. Henry*, 10557!), Tchenfongchang (*Delavay*, 5168!), Tapintze (*Delavay*, 1774!), Tali (*Delavay*, 2646, 3525!); HONGKONG (*Seemann*, 545! *C. Wright*, 592! *Ford*, 76!). Mus. Brit.; Herb. Kew.

From the North-west Himalaya to Celebes, the Philippines, and the Bonin Isles; plentiful in the East Himalaya and North Burma.

20. *Scirpus triqueter*, *Linn. Mant.* p. 29; *Kunth, Enum. Pl.* ii. p. 163; *Hance*, in *Journ. Bot.* xii. (1874) p. 329, et xvii. (1879) p. 16; *Benth. Fl. Austral.* vii. p. 334 in notâ; *Regel, Fl. Ussur.* p. 176; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 658.

*Scirpus Pollichii*, *Gren. et Godr. Fl. France*, iii. p. 374; *Boeck. in Linnæa*, xxxvi. (1869-70) p. 711; *Franch. in Mém. Soc. Sc. Nat. Cherb.* xxiv. (1884) p. 262, et *Pl. David.* i. p. 317.

CHIHLI: Peking (*Bretschneider*!), Tachiaotze (*Möllendorff*), Jehol (*David*!); SHINGKING (*Ross*, 47!); SHANTUNG, Chefoo (*Fauvel*); NORTH CENTRAL CHINA: Kusar, Kisaoshian (*Hugh*!); CHEKIANG: Ningpo (*Faber*, 231!); FORMOSA: Tamsui (*A. Henry*, 1761!); SHENSI: Paokiscen, Mt. Kifonsan (*Hugh*!); SZECHUEN: Tcentosen, Mt. Uomisan (*Hugh*!); KWANGTUNG: Canton (*Hance*, 13752!). Mus. Brit.; Herb. Kew.

From Europe to North India, China, Japan. Also has been received from South Africa.

1. *Eriophorum angustifolium*, *Roth, Neue Beitr.* p. 94; *Boeck. in Linnæa*, xxxvii. (1871) p. 95.

*Eriophorum polystachion*, *Linn. Sp. Pl.* ed. 1, p. 52; ed. 2, p. 56 partim.

*Eriophorum polystachion*, var. *angustifolia*, *Regel, Fl. Ussur.* p. 178.

SHINGKING: Moukden (*Webster*!); YUNNAN: Outchay (*Delavay*, 407!).

In the arctic and cool temperate zone throughout the world. Also received from the Transvaal.

2. *Eriophorum comosum*, *Nees, in Wight, Contrib.* p. 110, var.  $\beta$  excl.; *Boeck. in Linnæa*, xxxvii. (1871) p. 98; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 664; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 228.

HUPEH (*A. Henry*, 7734! *E. H. Wilson*, 538!), Ichang (*A. Henry*, 575, 4000!), Nanto (*A. Henry*, 2623!), Chung city (*Faber*, 1124!); SZECHUEN (*Faber*, 1130, 1132!), Tcentosen, Mt. Uomisan (*Hugh*!); YUNNAN, Tapintze (*Delavay*, 2554, 4568!), Yunnanfu (*Duchoux*, 319!). Mus. Brit.; Herb. Kew.

Abundant in North India, especially at 1000–5000 feet alt.; extending to Burma, Penang, Tonkin.

3. *Eriophorum latifolium*, *Hoppe, Taschenb.* (1800) p. 108; *Maxim. Prim. Fl. Amur.* p. 299; *Boeck. in Linnæa*, xxxvii. (1871) p. 96.

*Eriophorum polystachion*, *Linn. Sp. Pl.* ed. 1, p. 52, et ed. 2, p. 76 partim.

SHINGKING: Moukden (*James*).

In the arctic and cool temperate zone, throughout the world.

1. *Fuirena glomerata*, *Lam. Ill.* i. p. 150; *Benth. Fl. Hongk.* p. 396; *Boeck. in Linnæa*, xxxvii. (1871) p. 107; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 666; *Henriq. in Bolet. Soc. Broter.* ii. (1883) p. 167.

*Fuirena Rottboellii*, *Nees, in Wight, Contrib.* p. 94, et in *Hook. et Arn. Bot. Beechey's Voyage*, p. 224.

*Scirpus ciliaris*, *Linn. Mant.* p. 182.

CHINA (*Yvan*!); FORMOSA: Takow (*A. Henry*, 1084!); KWANGTUNG: Macao (*da Silva*!), Pakhoi (*Playfair*, 121!); HONGKONG (*Wilford*, 274! *C. Wright*, 570!). Mus. Brit.; Herb. Kew.

Hot regions of the Old World, from the Niger to Queensland; abundant throughout India in rice-swamps.

2. *Fuirena umbellata*, *Rottb. Descr. et Ic.* p. 70, t. 19 (*i. e.* t. 18 altera), fig. 3; *Benth. Fl. Hongk.* p. 395; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 74; *Boeck. in Linnæa*,



xxxvii. (1871) p. 110; *C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi. p. 666.*

*Fuirena pentagona*, *Nees, in Wight, Contrib. p. 93.*

CHINA (*Meyen, 244! Millett!*); KWANGTUNG (*Ford, 183!*), Macao (*fide Nees*); HONGKONG (*C. Wright, 571! Lamont, 861!*). *Mus. Brit.; Herb. Kew.*

In tropical and warm-temperate regions throughout the globe, widespread and in quantity, up to 3000 feet alt.

3. *Fuirena Wallichiana*, *Kunth, Enum. Pl. ii. (1837) p. 182; C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi. p. 665.*

*Fuirena pubescens*, *Boeck. in Linnæa, xxxvii. (1871) p. 104 partim.*

*Scirpus cuspidatus*, *Roth, Nov. Pl. Sp. p. 31.*

YUNNAN: Mengtze, 6000 feet (*A. Henry, 9327!*). *Herb. Kew.*

In North India; but not known from the eastern side.

1. *Lipocarpa argentea*, *R. Br. in Append. Tuckey, Congo, p. 459; Benth. Fl. Hongk. p. 388; Boeck. in Linnæa, xxxvii. (1871) p. 114 (exempl. Amer. excl.); C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi. p. 667.*

*Hypaelyptum argenteum*, *Vahl, Enum. ii. p. 283.*

*Kyllinga albescens*, *Steud. Cyp. p. 68.*

FOKIEN: Amoy (*Hance, 1404!*); HONGKONG (*Wilford, 270! C. Wright, 568!*); YUNNAN: Kichau, 6000 feet (*Delavay, 4034!*). *Mus. Brit.; Herb. Kew.*

Throughout the tropical and subtropical regions of the Old World, from the Niger to the Philippines and Queensland; common in India up to 5000 feet alt.

2. *Lipocarpa microcephala*, *Kunth, Enum. Pl. ii. p. 268; Benth. Fl. Hongk. p. 388; Boeck. in Linnæa, xxxvii. (1871) p. 118; Debeaux, Fl. Tchefou, p. 146, t. 3. fig. 2; Franch. in Mém. Soc. Sc. Nat. Cherb. xxiv. (1884) p. 264; C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi. p. 668; Diels, in Engl. Bot. Jahrb. xxix. (1901) p. 227.*

*Hypaelyptum microcephalum*, *R. Br. Prodr. p. 220.*

*Ascolepis kyllingioides*, *Steud. in Zoll. Verz. Pfl. Ind. Archip. ii. p. 63, et Cyp. p. 105 pro magnâ parte.*

*Scirpus leptocarpus*, *F. Muell. in Trans. Phil. Soc. Vict. i. (1855) p. 109.*

*Isolepis squarrosa*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 211, non *Roem. et Schult.*

SHANTUNG: Chefoo (*Debeaux, Fauvel*); HUPEH: Ichang (*A. Henry, 4542!*); HONGKONG (*Wilford!*). Mus. Brit.; Herb. Kew.

Eastern Asia; from Japan to Singapore, Macassar, Queensland.

1. *Hypolytrum latifolium*, *Rich. in Pers. Syn.* i. p. 70; *Benth. Fl. Hongk.* p. 389; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 678, vix *Boeck.*

*Tunga diandra*, *Roxb. Fl. Ind.* i. p. 184.

*Aibikkia scirpoides et A. schœnoides*, *Presl, Rel. Haenk.* i. p. 185, tt. 34, 35.

FORMOSA (*Oldham, 592!*); KWANGTUNG: Canton (*Hance, 1368!*); HONGKONG (*Harland, 1031! C. Wright, 569! Lamont, 838!*), Lofaushan (*Ford!*); HAINAN (*A. Henry!*). Mus. Brit.; Herb. Kew.

Plentiful in India, extending to Queensland and Polynesia. The representative Mascarene species is hardly separate; and Boeckeler unites also the Tropical African *A. nemorum*, Beauv.

1. *Lepironia mucronata*, *Rich. in Pers. Syn.* i. p. 70; *Boeck. in Linnæa*, xxxvii. (1871) p. 140; *Hance, in Journ. Bot.* xvii. (1879) p. 100; *Goebel, in Ann. Jard. Bot. Buitenz.* vii. (1888) p. 126, t. 14. figg. 12, 13; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 684.

*Restio articulatus*, *Retz. Obs.* iv. p. 14.

*Scirpus coniferus*, *Poir. in Lam. Encycl.* vi. p. 756, et *Suppl.* v. p. 90.

*Chondrache articulata*, *R. Br. Prodr.* p. 220.

*Choricarpa aphylla*, *Boeck. in Flora*, xli. (1858) p. 20.

HONGKONG (*Luetjens!*); KWANGTUNG (*Hance!*). Mus. Brit.

Spread from Mascarenia to New South Wales and Viti, a maritime plant. Hance says that, in Kwangtung, it was only cultivated for the making of mats.

1. *Remirea maritima*, *Aubl. Pl. Guian.* i. p. 45, t. 16; *Kunth, Enum. Pl.* ii. p. 139; *Boeck. in Linnæa*, xxxv. (1867-8) p. 435; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 677.

*Miegia maritima*, *Willd. Sp. Pl.* i. p. 311.

*Mariscus capitatus*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 63, et *Cyp.* p. 60.

*Mariscus pungens*, *Miq. Fl. Ind. Bat.* iii. p. 288.

*Mariscus maritimus*, *Miq. Fl. Ind. Bat., Suppl.* p. 600.

*Cyperus Kegelianus*, *Steud. Cyp.* p. 60.

*Lipocarpa foliosa*, *Miq. Fl. Ind. Bat.* iii. p. 332, *et Suppl.* p. 262.

CHINA (*Millett!*); KWANGTUNG: Canton (*Yvan!*). Herb. Kew.

On tropical sea-coasts throughout the world.

There are many other synonyms.

1. *Rynchospora aurea*, *Vahl, Enum.* ii. p. 229; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 670.

*Rynchospora aurea*, *R. Br. Prodr.* p. 230; *Benth. Fl. Hongk.* p. 396; *Boeck. in Linnæa*, xxxvii. (1873) p. 626.

*Rynchospora articulata*, *Spreng. Syst.* i. p. 197.

*Scirpus corymbosus*, *Linn. Amœn. Acad.* iv. p. 303.

*Cephaloschoenus articulatus*, *Nees, in Wight, Contrib.* p. 115.

FORMOSA: Takow (*A. Henry, 1843!*); HONGKONG (*Harland!*); HAINAN (*A. Henry, 8551!*). Mus. Brit.; Herb. Kew.

Throughout the world in and near the tropics, in swamps.

2. *Rynchospora Faberi*, *C. B. Clarke*, n. sp.

Annua. Culmi 4–12 unc. longi, graciles. Folia setacea aut fere capillacea. Corymbi  $2\frac{1}{2}$ –4 lin. in diam., distantes, 3–5-stachyi. Spiculæ  $1\frac{1}{2}$  lin. longæ, brunæ, 1-floræ. Setæ 6, cum nuce subæquilongæ, rigidæ, pallide brunæ, dense retrorsim scabræ. Nux  $\frac{3}{4}$  lin. longa, obovoidea, luteo-brunæ, obscure transversim undulata; rostrum cum  $\frac{1}{2}$  parte nucis æquilongum.

CHEKIANG: Lu Mts., 3300 feet (*Faber, 247!*). Herb. Kew.

This species is allied to *R. capillacea* and *R. Knieskernii*, but has the nut much broader.

3. *Rynchospora glauca*, *Vahl, Enum.* ii. p. 233; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 671.

*Rynchospora glauca*, *Roem. et Schult. Syst.* i. p. 85; *Boeck. in Linnæa*, xxxvii. (1873) p. 585, forma "majore" excl.

*Rynchospora laxa*, *R. Br. Prodr.* p. 230.

SHANTUNG: Chefoo Mts. (*Faber, 286!*). Herb. Kew.

Common throughout the world in the tropics and warm temperate zone.

Var.  $\beta$ . *chinensis*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 672.

*Rhynchospora chinensis*, Boeck. in *Linnæa*, xxxvii. (1873) p. 586.

*Rhynchospora triflora*, Nees, in *Linnæa*, ix. (1834) p. 297, e Nees, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 124 b, nec Vahl.

*Rhynchospora glauca*, Boeck. in *Linnæa*, xxxvii. (1873) p. 585 partim.

*Rhynchospora lavarum*, Hook. et Arn. *Bot. Beechey's Voy.* p. 98, non Gaudich.

*Rhynchospora laxa*, Benth. *Fl. Hongk.* p. 397.

CHINA (Meyen!); CHEKIANG: Ningpo Mts. (Faber, 228, 229!); HONGKONG (C. Wright, 595! Wilford, 277! Hance, 1071! Faber, 1886! Didrichsen, 3271!). Mus. Brit.; Herb. Kew.

Scattered from Madagascar to the Sandwich Isles.

[*R. chinensis*, Nees (in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 108), is founded on a plant collected at Cape Syngmun by Meyen, and is characterized by having 8–9 retrorsely scabrid setæ. It could therefore have been no known species except *R. alba*, Vahl, which though unknown from China is very common in Japan. But from the way this species is united by Nees with Indian known plants, and from Boeckeler's way of dealing, it seems probable that the description by Nees was radically erroneous.]

4. ***Rynchospora gracillima***, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 671.

*Rhynchospora gracillima*, Thwaites, *Enum. Pl. Zeyl.* p. 435; Boeck. in *Linnæa*, xxxvii. (1873) p. 597.

*Rhynchospora Kamphoeveneri*, Boeck. in *Engl. Bot. Jahrb.* v. (1884) p. 508.

HONGKONG (Lamont!). Mus. Brit.

Ceylon, Tinnevely, Khasia, Nicobars.

5. ***Rynchospora Wallichiana***, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. (1893) p. 668.

*Rhynchospora Haenkei*, Presl, *Rel. Haenk.* i. (1828) p. 199; Kunth, *Enum. Pl.* ii. p. 290.

*Rhynchospora Wallichiana*, Kunth, *Enum. Pl.* ii. p. 289; Benth. *Fl. Hongk.* p. 396; Boeck. in *Linnæa*, xxxvii. (1873) p. 542 (exempl. Amer. excl.), in *Engl. Bot. Jahrb.* vi. (1885) p. 51; Henriq. in *Bolet. Soc. Broter.* ii. (1883) p. 167.

*Morisia Wallichii*, Nees, in *Edinb. New Phil. Journ.* (1834) n. 34, p. 265.

*Haplostylis Meyenii*, Nees, in *Edinb. New Phil. Journ.* (1834) n. 34, p. 265, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 227.

*Sphæroschœnus Wallichii*, *Arn. et Nees, in Nov. Act. Nat. Cur. xix. Suppl. i.* (1843) p. 97.

*Cephaloschœnus parvus*, *Nees, in Linnæa, ix.* (1834) p. 295, *et in Nov. Act. Nat. Cur. xix. Suppl. i.* (1843) p. 100.

CHINA (*Millett!*); YUNNAN: Mengtze, 5000 feet (*A. Henry, 11150 partim!*); KWANGTUNG (*Meyen! Ford, 181!*), Pakhoi (*Playfair, 207!*); HONGKONG (*C. Wright, 596! Harland! Hance, 1237! Didrichsen, 3208! Champion, 539! Faurie, 15831!*); LUCHU ARCHIPELAGO (*Döderlein!*). *Mus. Brit.; Herb. Kew.*

Old World, in the tropics and warm regions, from the Niger to Queensland and Japan.

1. *Schœnus apogon*, *Roem. et Schult. Syst. ii.* p. 77.

*Schœnus imberbis*, *Poir. Encyc. Suppl. ii.* p. 251, nec p. 250, nec *R. Br.*

*Schœnus Brownii*, *Hook. f. Handb. Fl. New Zeal. p.* 298; *Benth. Fl. Austral. vii.* p. 373.

*Chætospora imberbis*, *R. Br. Prodr. p.* 233; *Boeck. in Linnæa, xxxviii.* (1874) p. 299.

*Chætospora japonica*, *Franch. et Sav. Enum. Pl. Jap. ii.* pp. 122, 548.

*Isolepis margaritifera*, *Nees, in Ann. Nat. Hist. ser. 1, vi.* (1849) p. 46.

*Scirpus margaritiferus*, *Boeck. in Linnæa, xxxvi.* (1869-70) p. 697.

LUCHU ARCHIPELAGO (*C. Wright, 360!*). *Herb. Kew.*

Frequent in Australia and New Zealand; also on Kinabalu, in Borneo.

2. *Schœnus falcatus*, *R. Br. Prodr.* (1810) p. 232; *Kunth, Enum. Pl. ii.* p. 336; *Benth. Fl. Austral. vii.* p. 372.

*Schœnus elatus*, *Boeck. in Flora, lviii.* (1875) p. 117.

FORMOSA: Takow (*A. Henry, 1943!*), Ape's Hill (*Playfair, 395!*). *Herb. Kew.*

Tropical Australia and Queensland; Borneo.

1. *Lepidosperma chinense*, *Nees, in Linnæa, ix.* (1834) p. 302, *et in Hook. et Arn. Bot. Beechey's Voy. p.* 228, *et in Nov. Act. Nat. Cur. xix. Suppl. i.* (1843) p. 117; *Kunth, Enum. Pl. ii.* p. 320; *Benth. Fl. Hongk. p.* 398; *Seem. Bot. Voy. 'Herald,' p.* 422; *Boeck. in Linnæa, xxxviii.* (1874) p. 329 (*L. confini* excl.); *C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi.* p. 676.

CHINA (*Millett! Schottmueller, 414! O. Kuntze, 3163!*); KWANGTUNG (*Seemann, 2399!*), Macao (*Vachell, 39 b!*), Cape

Syngmun (*Meyen*!); HONGKONG (*Hinds*! *Seemann*! *Fortune*! *Wilford*, 275! *Harland*, 682! *Hance*, 999!). Mus. Brit.; Herb. Kew.

Also on Mt. Ophir, near Malacca.

1. *Cladium ensigerum*, *Hance*, in *Journ. Bot.* xxiii. (1885) p. 80.

HONGKONG: Pokfulum (*Ford*!). Herb. Kew.

Only one sheet of excellent specimens known.

2. *Cladium glomeratum*, *R. Br. Prodr.* p. 237; *Benth. Fl. Austral.* vii. p. 404; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 575, non *Baumea glomerata*, *Gaudich.*

*Chapelliera glomerata*, *Nees*, in *Lehm. Pl. Preiss.* ii. p. 76; *Franch. et Sav. Pl. Jap.* ii. p. 121.

*Baumea rubiginosa* et *B. Brownei*, *Boeck. in Linnæa*, xxxviii. (1874) pp. 241, 242.

HONGKONG (*Faber*, 226! *Faurie*, 15792!). Herb. Kew.

Australia and New Zealand, extending to Bengal and Japan.

3. *Cladium jamaicense*, *Crantz, Inst.* i. p. 362.

*Cladium germanicum*, *Schrad. Fl. Germ.* i. t. 5. fig. 7.

*Cladium Mariscus*, *R. Br. Prodr.* p. 236; *Benth. Fl. Hongk.* p. 397; *Boeck. in Linnæa*, xxxviii. (1874) p. 232; *Hook. f. Fl. Brit. Ind.* vi. p. 673.

*Cladium leptostachyum*, *Nees*, in *Linnæa*, ix. (1834) p. 301, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 115.

*Cladium chinense*, *Nees*, in *Linnæa*, ix. (1834) p. 301, et in *Hook. et Arn. Bot. Beechey's Voy.* p. 227, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 116; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 132.

*Schoenus Mariscus*, *Linn. Sp. Pl.* ed. 1, p. 42; ed. 2, p. 62.

*Gahnia Mariscus*, *F. Muell. Key Victoria Pl.* i. p. 456.

KWANGTUNG: Macao (*Vachell*, 59!); HONGKONG (*Hance*, *Ford*, 75!). Mus. Brit.; Herb. Kew.

Extends from Sweden to the Cape of Good Hope, from Japan to Victoria, from New Orleans to Rio Janeiro; also throughout Polynesia; but absent in India and West Malaya.

1. *Gahnia javanica*, *Moritzi, Verz. Zoll. Pfl.* (1845-6) p. 98; *Boeck. in Linnæa*, xxxviii. (1874) p. 339; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 676.

*Phakellanthus multiflorus*, *Steud. in Zoll. Verz. Ind. Archip.* ii. p. 61.

YUNNAN: Mengtze, 7000 feet (*A. Henry*, 9168!). *Herb. Kew.*

In Malaya frequent; extending to New Guinea and Viti. Also a variety from Penang and Singapore.

2. *Gahnia tristis*, *Nees, in Linnæa*, ix. (1834) p. 301 (nomen), *in Hook. et Arn. Bot. Beechey's Voy.* p. 228; *Benth. Fl. Hongk.* p. 398.

*Gahnia Wichurai*, *Boeck. in Linnæa*, xxxviii. (1874) p. 348.

KWANGTUNG: Macao (*Vachell*, 39 c!); HONGKONG (*Wilford*, 276! *Harland*, 742! *C. Wright*, 397! *Faurie*, 15858!). *Mus. Brit.*; *Herb. Kew.*

Received also from Singapore, Anam, Borneo.

1. *Scleria annularis*, *Kunth, Enum. Pl.* ii. p. 359; *Boeck. in Linnæa*, xxxviii. (1874) p. 456; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 687.

*Hypoporum annulare*, *Nees, in Linnæa*, ix. (1834) p. 303 nomen.

HUPEH: Ichang (*A. Henry*, 4253!). *Herb. Kew.*

Scattered throughout India. Also in the Philippines.

2. *Scleria biflora*, *Roxb. Hort. Beng.* p. 66, *et Fl. Ind.* iii. p. 573; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 687.

*Scleria Steudeliana*, *Miq. Fl. Ind. Bat.* iii. p. 344, *et Suppl.* p. 262; *Boeck. in Linnæa*, xxxviii. (1874) p. 475.

*Scleria tessellata*, *Benth. Fl. Hongk.* p. 399, *non Willd.*

HONGKONG (*Hance*, 5232!). *Mus. Brit.*; *Herb. Kew.*

Frequent in India and Malaya. But hardly differs from *S. tessellata* except in the minute character taken from the lobes of the disc.

3. *Scleria chinensis*, *Kunth, Enum. Pl.* ii. (1837) p. 357; *Benth. Fl. Hongk.* p. 400; *Boeck. in Linnæa*, xxxviii. (1874) p. 486; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 690 (*Ridley*, n. 1556 excl.).

*Scleria ciliaris*, *Nees, in Wight, Contrib.* (1834) p. 117, *et in Hook. et Arn. Bot. Beechey's Voy.* p. 229, *non Mich.*

*Scleria scrobiculata*, *Moritzi, Verz. Zoll. Pfl.* p. 98 partim.

CHINA (*Beechey! Hinds!*); KWANGTUNG: Macao (*Vachell*, 40! *Calléry!*); HONGKONG (*Champion*, 291! *Wilford*, 47! *C. Wright*,

601! *Harland*, 736! *Hance*, 840, 1054! *Faurie*, 15808!). Mus. Brit.; Herb. Kew.

Also in Malaya, to Singapore and Queensland. This species differs but little from *S. scrobiculata*, Nees.

4. *Scleria elata*, *Thwaites*, *Enum. Pl. Zeyl.* p. 353 (Thwaites, n. 825 excl.); *Boeck. in Linnæa*, xxxviii. (1874) p. 487; *Hance*, in *Journ. Bot.* xvii. (1879) p. 16; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 690.

*Scleria Rinkiana*, *Boeck. Cyp. Novæ*, ii. p. 30.

*Diaphora cochinchinensis*, *Lour. Fl. Cochinch.* p. 709.

FORMOSA: South Cape (*A. Henry*, 608!); KWANGTUNG: Mts. Pakwan (*Sampson!* hb. *Hance*, 12962!). Mus. Brit.; Herb. Kew.

This species, abundant in India, extends through Malaya to New Guinea. It differs little from *S. scrobiculata* but by the conspicuously 3-winged sheaths to the leaves. Henry's n. 608 may be called var. *decolorans*, *C. B. Clarke* (in *Hook. f. Fl. Brit. Ind.* vi. p. 690); while *Hance* n. 12962 has the middle stem-leaves approximated in twos or threes, and may therefore be named *S. scrobiculata* by some.

5. *Scleria Harlandi*, *Hance*, in *Ann. Sc. Nat.* sér. 5, v. (1866) p. 248, et in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 133.

*Scleria purpurascens*, *Benth. Fl. Hongk.* p. 400, non *Steud.*

HONGKONG (*Harland*, 735! *Lamont*, 875!). Mus. Brit.; Herb. Kew.

6. *Scleria hebecarpa*, *Nees*, in *Wight, Contrib.* p. 117; *Kunth*, *Enum. Pl.* ii. p. 357; *Hance*, in *Ann. Sc. Nat.* sér. 5, v. (1866) p. 249; *Boeck. in Linnæa*, xxxviii. (1874) p. 478; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 689.

*Scleria japonica*, *Steud. Cyp.* p. 169.

*Scleria aspera*, *Boeck. in Linnæa*, xxxviii. (1874) p. 483 partim.

*Scleria Wichurai*, *Boeck. in Engl. Bot. Jahrb.* v. (1884) p. 510.

CHEKIANG: Ningpo Mts. (*Faber*, 288 bis!); YUNNAN: Syemoo, 5000 feet (*A. Henry*, 12230!); KWANGTUNG: Macao (*Abel*), Whampoa (*Hance*, 10112!), Lofaushan (*Ford*, 188!), Pakhoi (*Playfair*, 146!); KWANGSI: Lungchow (*A. Henry*, 690!); HAINAN (*A. Henry*, 8009!). Mus. Brit.; Herb. Kew.

Common in India, extending thence to Japan, Viti, and New South Wales.



Var.  $\beta$ . *pubescens*, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 689.

*Scleria pubescens*, Steud. in *Zoll. Verz. Ind. Archip.* ii. p. 61, et *Cyp.* p. 168; *Benth. Fl. Hongk.* p. 400.

*Scleria hebecarpa*, Munro, in *Seem. Bot. Voy. 'Herald,'* p. 423.

*Scleria vestita*, Boeck. in *Linnæa*, xxxviii. (1874) p. 482.

HUPEH: Ichang (*A. Henry*, 4095!); KWANGTUNG: Lofaushan (*Ford*, 123!); HONGKONG (*C. Wright*, 600! *Harland!*). Mus. Brit.; Herb. Kew.

Frequent in North-east India; also in Java.

Boeckeler founded his *S. vestita* on Hance, n. 1157, which number is in herb. Kew *S. Radula*. But, as before noted, the Hance numbers are not of collections, but of names in a list.

7. *Scleria Hookeriana*, Boeck. in *Linnæa*, xxxviii. (1874) p. 498; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 691.

HUPEH (*E. H. Wilson*, 1633!); YUNNAN: Tchenfongchan (*Delavay!*), Szemao, 5000 feet (*A. Henry*, 12230 A!). Herb. Kew.

Plentiful in Khasia and Manipur.

8. *Scleria lithosperma*, Swartz, *Prodr.* p. 18, et *Fl. Ind. Occ.* i. p. 92 in notâ; *Kunth, Enum. Pl.* ii. p. 349; *Benth. Fl. Hongk.* p. 399; Boeck. in *Linnæa*, xxxviii. (1874) p. 451; *Henriq. in Bol. Soc. Broter.* ii. (1883) p. 167; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 685.

*Scleria filiformis*, Swartz, *Prodr.* p. 19, et *Fl. Ind. Occ.* i. p. 91; *Kunth, Enum. Pl.* ii. p. 348.

*Scirpus lithospermus*, *Linn. Sp. Pl.* ed. 1, p. 51.

*Schoenus lithospermus*, *Linn. Sp. Pl.* ed. 2, p. 65 pro parte.

*Olyra orientalis*, *Lour. Fl. Cochinch.* ii. p. 674.

KWANGTUNG: Macao (*da Silva*); HONGKONG (*Harland!* *C. Wright*, 598!). Mus. Brit.; Herb. Kew.

In the tropics of both hemispheres. Boeckeler places the Hongkong plant as a variety of the American *S. filiformis*, Swartz; but I consider the Hongkong plant identical with the Indian.

9. *Scleria margaritifera*, *Willd. Sp. Pl.* iv. p. 312; *Kunth, Enum. Pl.* ii. p. 341; Boeck. in *Linnæa*, xxxviii. (1874) p. 511.

*Scleria polycarpa*, Boeck. in *Linnæa*, xxxviii. (1874) p. 509.

*Scleria Græffiana*, Boeck. in *Flora*, lviii. (1875) p. 121.

*Scleria Doederleiniana*, *Boeck. in Engl. Bot. Jahrb.* vi. (1885) p. 51.

*Carex lithosperma*, *Forst. f. Prodr. Fl. Ins. Austral.* p. 65.

LUCHU ARCHIPELAGO (*Döderlein*!).

Also in Ladrones, Viti, Samoa, Solomon Isles, Tanna, North-east Australia.

10. *Scleria pergracilis*, *Kunth, Enum. Pl.* ii. (1837) p. 354; *Boeck. in Linnæa*, xxxviii. (1874) p. 438; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 685.

*Hypoporum pergracile*, *Nees, in Edinb. Phil. Journ.* xvii. (1834), p. 267, et in *Wight, Contrib.* p. 118.

YUNNAN (*Delavay, 1557*!), Tapintze (*Delavay, 1657*!), Mosoyn (*Delavay, 4039*!), Hokin (*Delavay, 2490*!).

In tropical and warm-temperate Africa and India.

11. *Scleria Radula*, *Hance, in Ann. Sc. Nat. sér. 4*, xviii. (1862) p. 232, et in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 133; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 691.

*Scleria lævis*, var. *scaberrima*, *Benth. Fl. Hongk.* p. 400, non *S. levis*, *Retz.*

*Scleria aspera*, *Boeck. in Linnæa*, xxxviii. (1874) p. 483 partim.

CHINA (*Meyen*); HONGKONG (*Harland, 730*!). Mus. Brit.; Herb. Kew.

Also in Perak.

12. *Scleria Ridleyi*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. (1893) p. 686.

*Scleria lævis*, *Hance, in Ann. Sc. Nat. sér. 4*, xviii. (1862) p. 232; *Benth. Fl. Hongk.* p. 400, not of *Retz.*

HONGKONG (*Hance, 2985*! *Naumann*). Mus. Brit.; Herb. Kew.

Singapore.

13. *Scleria scrobiculata*, *Nees, in Wight, Contrib.* p. 117, et in *Nov. Act. Nat. Cur. xix. Suppl. i.* (1843) p. 119; *Kunth, Enum. Pl.* ii. p. 342; *Boeck. in Linnæa*, xxxviii. (1874) p. 508.

*Scleria tessellata*, *Decne. in Nouv. Ann. Mus.* iii. (1834) p. 362, non *Willd.*

*Scleria Neesiana*, *Hook. et Arn. Bot. Beechey's Voy.* p. 229, nec *S. Neesii*, *Kunth.*

*Scleria waigiouensis*, *Steud. Cyp.* p. 173.

*Scleria keyensis*, *K. Schum. in Engl. Bot. Jahrb.* xiii. (1891) p. 267.

CHINA (*Beechey!* *Meyen!*); FORMOSA (*Oldham*, 594! *A. Henry*, 478!); YUNNAN: *Mengtze*, 6000 feet (*A. Henry*, 10484!).—  
Mus. Brit.; Herb. Kew.

*A. Henry* notes on his n. 478, "Used for making brooms": on his n. 10484 "Was 9 ft. 1 in. high, leaves mostly in the upper half"; the specimen is very young, and may or may not be *S. scrobiculata*.

Frequent in Malaya, from Java to the Philippines and New Guinea.

14. *Scleria tessellata*, *Willd. Sp. Pl.* iv. p. 315, tab. *Rumphii* citatâ excl.; *Kunth, Enum. Pl.* ii. p. 343; *Boeck. in Linnæa*, xxxviii. (1874) p. 470, nec *Boeck. in Flora*, lxii. (1879) p. 571; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 686; non *Benth.*

CHEKIANG: *Ningpo Mts.* (*Faber*, 288!); YUNNAN: *Tchen-fongchan* (*Delavay!*). Herb. Kew.

Plentiful in India; frequent in Japan; occurs scattered southward to Queensland.

1. *Diplacrum caricinum*, *R. Br. Prodr.* p. 241; *Kunth, Enum. Pl.* ii. p. 360; *Boeck. in Linnæa*, xxxviii. (1874) p. 434; *Goebel, in Ann. Jard. Buitenz.* vii. (1888) p. 132, t. 15. figg. 21-29.

*Scleria caricina*, *Benth. Fl. Austral.* vii. p. 426; *Benth. et Hook. f. Gen. Pl.* iii. p. 1071; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 688.

HONGKONG (*Lamont*, 869 A!, *Hance*, 9932!). Mus. Brit.; Herb. Kew.

Frequent in India, extending thence to the Philippines and Queensland.

1. *Kobresia cercostachys*, *C. B. Clarke, MS.*

*Carex cercostachys*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1894) p. 27, in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 191, 209.

SZECHUEN: *Tachienlu*, 9000-13,500 feet (*Pratt*, 741!); YUNNAN: *Yengtzehay*, 12,250 feet (*Delavay*, 3403!). Herb. Kew.

2. *Kobresia fragilis*, *C. B. Clarke*, n. sp.

*Culmi* 12-16 unc. longi, graciliores, teretes. *Folia* omnia basalia, cum  $\frac{1}{3}$  parte culmi vix æquilonga,  $\frac{1}{10}$  unc. lata. *Panicula*  $1\frac{1}{2}$  unc. longa, 3 lin. lata; bractea ima 2-3 unc. longa, 1 lin.

lata; spicæ 12–30,  $1\frac{1}{2}$ –2 lin. longæ, fere omnes 2-floræ, in sicco caducæ. *Gluma* (prophyllon) compressa, alte fissa, levis, apice truncata dentata. *Rhachilla* cum glumâ æquilonga, lata, 2-costata, apice florem masculum cum rudimento (floris alteri masculi rhachillâ) gerens. *Stylus* brevis; rami 3 longi rubro-brunei. *Nux* prophyllon parum brevior, exacte ellipsoidea, trigona, multum compressa, fere biconvexa, levis, nigro-brunea, rostro vix ullo.

SZECHUEN: Tongolo in Kiala (*Soulié*, 731!). Herb. Kew.

3. **Kobresia graminifolia**, *C. B. Clarke*, n. sp.

*Culmi*  $15\frac{1}{2}$  unc. longi, apice acutius trigoni. *Folia* 6–8 unc. longa,  $1\frac{1}{2}$  lin. lata, glabra, in marginibus viscoso-rubro-scabrida; vaginæ fissæ, apice scarioso-ligulatæ. *Spica* subbracteata, 14–19 lin. longa,  $3\frac{1}{2}$  lin. lata, densiuscula, matura intense brunea. *Spiculæ* omnes 1-nucigeræ; axeos rudimentum sæpe, floris rudimentum rarius additur. *Gluma* 2 lin. longa, elliptico-oblonga, obtusiuscula, in carinâ et in apice minute scabro-pilosa. *Nux* 2 lin. longa, anguste oblonga, trigona, lineari-conice rostrata.—Species *K. Prattii* affinis.

SHENSI: Mt. Miao Uansan (*Hugh*!). Mus. Brit.

4. **Kobresia Prattii**, *C. B. Clarke*, n. sp.

*Culmi* visi omnes juveniles, usque ad 3 unc. longi, masculi, basi arcte cæspitosi a vaginis fusce bruneis fibrillosis incrassati. *Folia* juvenilia cum culmis æquilonga, vix  $1\frac{1}{4}$  lin. lata. *Spica* fere 1 unc. longa,  $2\frac{1}{2}$  lin. lata, viridi-ferruginea. *Bracteæ* (si mavis glumæ masculæ) 5 lin. longæ, lineares, apice perobtusæ.—Species *K. cercostachydi* affinis et similis, ob bracteas obtusas muticas diversa.

SZECHUEN: Tachienlu, 9000–13,500 feet (*Pratt*, 744!). Herb. Kew.

5. **Kobresia Royleana**, *Boeck. in Linnæa*, xxxix. (1875) p. 8: *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 698.

*Trilepis Royleana*, *Nees*, in *Edinb. New Phil. Journ.* xvii. (1834) p. 267, in *Wight, Contrib.* p. 119; *Kunth, Enum. Pl.* ii. p. 535.

*Elyna stenocarpa*, *Kar. et Kir. in Bull. Soc. Nat. Mosc.* xv. (1842) p. 526.

SZECHUEN: Tongolo, in Kiala (*Soulié*, 771!).

Central Asia, Himalaya tota.

6. **Kobresia trinervis**, *Boeck. in Linnæa*, xxxix. (1875) p. 548 in notâ; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 695, cum syn.

*Carex mutans*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1894) p. 27, in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 191, 209, syn. excl.; non *Boott.*

YUNNAN: Hokin, Mt. Koualapo, 12,250 feet (*Delavay*, 1049!). Himalaya.

7. **Kobresia uncinoides**, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. (1894) p. 698.

*Carex uncinoides*, *Boott, Carex*, p. 8, t. 23; *Boeck. in Linnæa*, xxxix. (1875) p. 109; *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1894) p. 28, in *Nouv. Arch. Mus.* sér. 3, ix. (1896) pp. 192, 209.

YUNNAN: Tapintze (*Delavay*), Tsangchan, 12,250 feet (*Delavay*, 2653!), Yanginchan, 9750 feet (*Delavay*, 2339 bis).

Sikkim, 10,000–16,000 feet.

1. **Carex acutiformis**, *Ehrh. Beitr.* iv. (1789) p. 43; *Boeck. in Linnæa*, xli. (1877) p. 289; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 740; *Franch. in Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 83, 89; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 233.

*Carex paludosa*, *Gooden. in Trans. Linn. Soc.* ii. (1794) p. 202; *Sowerby, Engl. Bot.* t. 807; *Meinsh. in Act. Hort. Petrop.* xviii. (1901) pp. 298, 377.

HUPEH (*A. Henry*, fide *Franchet* et fide *Diels*).

From England to Tunis and to Central Asia. Also at the Cape of Good Hope. I have seen no plant of Henry's at Kew that could be *C. acutiformis*.

2. **Carex agglomerata**, *C. B. Clarke*, n. sp.

*Culmi* pedales, longi, cæspitiosi, apice trigoni leves. *Folia* culmo parum breviora,  $\frac{1}{8}$ – $\frac{1}{6}$  in. lata, in facie inferiore densius puberula. *Spicæ* 4 approximatae, fere agglomeratae, ellipsoideae, virides, densae; terminalis basi mascula, apice feminea. *Bractea* ima cum inflorescentiâ subæquilonga, vel (raro paullo distans) inflorescentiam superans. *Glumæ* femineae (mucrone incluso) utriculis breviores, ovatae, virides aut pallidae. *Styli* rami 3 cum  $\frac{1}{2}$  parte utriculi subæquilongi. *Utriculus* rostro incluso 2 lin. longus, ellipsoideus, viridis, vix nervatus, levis, glaber, sessilis,

apice subito angustatus. *Rostrum* cum  $\frac{1}{2}$ - $\frac{2}{3}$  parte utriculi æquilongum, breviter 2-dentatum.

SHENSI: Paokisan, Mt. Kifonsan (*Hugh*!). Mus. Brit.

3. *Carex albomas*, *C. B. Clarke*, n. sp.

Hemiscaposa, præter utriculos glabra. *Culmi* 5-7 unc. longi; vaginæ imæ fimbriatæ, lentæ, fusco-bruneæ. *Folia* culmum sæpe superantia,  $1\frac{1}{2}$ -2 lin. lata. *Spicæ* 1 unc. longæ, distantes, terminalis mascula alba, femineæ 2-3 albo-virides; *bractea* ima inflorescentiâ brevior,  $\frac{3}{4}$ -2 unc. longa; *pedunculus* imus e vaginâ brevi breviter exsertus. *Glumæ* femineæ parvæ, albæ, longius viridi-aristatæ. *Styli* rami 3, longi, atro-rubri. *Utriculus* immaturus ovoideus, apice attenuatus, pilosus; rostri dentes 2, longiusculi, scabri.

SHINGKING: Korean Gate (*Ross*, 57!), Fungshan (*Ross*, 47!), Moukden to Yaloo River (*Webster*!). Herb. Kew.

These examples are all young, and are marked in hb. Kew as allied to *C. præcox*, Jacquin, which is, I think, the true affinity of *C. albomas*; it differs from *C. præcox*, Jacq., in the distant, green, narrow female spikes and the large white male spike.

In *Nouv. Arch. Mus. sér. 3, ix.* (1897) p. 186, Franchet has reduced three of his own species to *C. præcox*, Jacq., which he makes a variable species universally distributed in Japan. None of this Japanese *C. præcox* is the same as *C. albomas*; I believe it all quite distinct from *C. præcox*, Jacq., as it all has the bulbous style-base characteristic of so many Japan species of small *Carex*. But of this *C. præcox*, Franchet, so universal in Japan, I find no example from China.

4. *Carex alliiformis*, *C. B. Clarke*, n. sp.

*Culmus* 5-17 unc. *Folia* basalia plura, 7-20 unc. longa, 3-6 lin. lata, glabra. *Spicæ* 5-6, distantes, terminalis mascula  $\frac{1}{2}$ -1 unc. longa,  $1\frac{1}{4}$  lin. lata, brunea; femineæ 1-2 unc. longæ, 3 lin. latæ, juniores viridi-purpureæ, maturæ albæ, imus in pedunculo  $1\frac{1}{2}$ -unciali suberectus; *bracteæ* foliiformes, 4-6 lin. latæ, inflorescentiam longe superantes. *Glumæ* femineæ lanceolatæ acutæ, utriculos paullo superantes, in dorso virides in lateribus rubro-purpureæ. *Styli* rami 3, longi. *Utriculus*  $1\frac{1}{4}$ - $1\frac{1}{2}$  lin. in diam., globosus, glaber, enervis, inflatus, maturus albi-papyraceus, in rostrum sublineare integrum subito angustatus.

*Nux ellipsoidea*, trigona, apice depresso-pyramidalis, erostrata, cum  $\frac{1}{2}$  parte utriculi vix æquilata.

HUPEH: Ichang (*A. Henry*, 3409, 3483!), 6700 feet (*E. H. Wilson*, 1660!). Herb. Kew.

The spikes of fruit suggested Lilliputian ropes of onions.

5. *Carex alopecuroides*, *D. Don*, in *Trans. Linn. Soc.* xiv. (1825) p. 332, *Prodr. Fl. Nepal.* p. 43; *Boott*, *Carex*, p. 88, t. 258; *Boeck. in Linnæa*, xli. (1877) p. 172.

*Carex Emodorum*, *Spreng. Syst.* iii. p. 818.

*Carex patens*, *Franch. in Bull. Soc. Philom.* sér. 8, vii. (1894-5) p. 90.

*Carex consocialis*, *Steud. Cyp.* p. 222.

*Carex japonica*, var. *alopecuroides*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 737.

HUPEH: Patung (*A. Henry*, 3744!). Herb. Kew.

Frequent in North-east India and in Japan.

The female spikes are 1-2 $\frac{1}{4}$  in. long, but not more than  $\frac{1}{4}$  in. broad, except in the var.  $\beta$ . In ripe fruit the beaks of the utricles are recurved.

Var.  $\beta$ . *chlorostachya*, *C. B. Clarke*; robustior, foliis 4-5 lin. latis; spicis fructiferis usque ad 4 lin. latis.

*Carex japonica* (typ.), *Boott*, *Carex*, t. 257; *Boeck. in Linnæa*, xli. (1877) p. 283.

*Carex chlorostachya*, *D. Don*, in *Trans. Linn. Soc.* xiv. (1825) p. 330 (partim).

*Carex Doniana*, *Spreng. Syst.* iii. p. 825; *Kunth*, *Enum. Pl.* ii. p. 503; *Drejer*, *Symb. Caricol.* p. 26, t. 13.

*Carex Zollingeri*, *Steud. Cyp.* p. 221.

*Carex japonica*, var. *alopecuroides*, *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 233.

HUPEH: Patung (*A. Henry*, 5655! *E. H. Wilson*, 1638, 1653!), Chieneht (*E. H. Wilson*, 1644!); SZECHUEN: Wushan (*A. Henry*, 5500!), Mount Omei alt. 3500 feet (*Faber*, 1144!). Mus. Brit.; Herb. Kew.

I have taken *Boott's Carex* t. 257 as the "type" of this variety, *i. e.* it is the most strongly marked form, and it was drawn from *Boott's* specimen of "*C. Doniana*, *Spreng.*" (*i. e.* *C. chlorostachya*, *D. Don*); but most of *D. Don's* own *C. chlorostachya* appears to be nearer *C. alopecuroides*, *D. Don*, between which and my variety  $\beta$ . *chlorostachya* I see no good line.

6. *Carex alta*, Boott, in *Proc. Linn. Soc.* i. (1845) p. 254, in *Trans. Linn. Soc.* xx. (1848) p. 130, et *Carex*, p. 59, t. 153; Boeck. in *Linnæa*, xxxix. (1875) p. 126; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 707; Franch. in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 236; Diels, in *Engl. Bot. Jahrb.* xxix. (1900) p. 230.

*Carex Brizopyrum*, Kunze, *Suppl. Schkuhr*, p. 169, t. 43.

CHEKIANG (*Faber*, 1024!); SZECHUEN: Tchenkeoutin, 7800 feet (*Farges*, 1244!); YUNNAN: Longki (*Delavay*). Herb. Kew. North-east India, Java.

Var.  $\beta$ . *Rochebruni*, Franch. in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 236; Diels, in *Engl. Bot. Jahrb.* xxix. (1900) p. 230.

*Carex Rochebruni*, Franch. et Sav. *Enum. Pl. Jap.* ii. pp. 126, 555; Franch. *Pl. David.* ii. p. 139.

SZECHUEN: Moupin (*David*!).

Japan.

7. *Carex arenicola*, F. Schmidt, *Reise Amurl.* (1868) p. 191; Franch. in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 221.

*Carex chætorrhiza*, Franch. et Sav. *Enum. Pl. Jap.* ii. p. 552.

*Carex curaica*, Franch. et Sav. *Enum. Pl. Jap.* ii. p. 124.

*Carex muricata*, Boott, in *Mem. Amer. Acad.* new ser. vi. (1859) p. 417 in Obs.

NORTH CENTRAL CHINA: Kaolinsien (*Hugh*!). Mus. Brit. In Sachalin and Japan, frequent.

8. *Carex aristata*, R. Br. in *Franklin, Polar Sea* (1823), p. 751; Kunth; *Enum. Pl.* ii. p. 519.

*Carex trichocarpa*, Muhl., var. *aristata*, L. H. Bailey, in *Bot. Gaz.* x. (Indianapolis, 1885) p. 294.

Cold America and Europe.

Var. *orthostachys*, C. B. Clarke; culmis basi 1-1½ lin. in diam., a vaginis parum incrassatis, quam in *C. aristatâ*, R. Br., typ. gracilioribus.

*Carex orthostachys*, C. A. Meyer, in *Ledeb. Ic. Fl. Ross.* iv. (1833) p. 10, t. 324; Kunth, *Enum. Pl.* ii. p. 491; Maxim. *Prim. Fl. Amur.* p. 316; Franch. *Pl. David.* i. p. 321.

*Carex aristata* (pars), Boott, *Carex*, p. 22; Boeck. in *Linnæa*, xli. (1877) p. 315; Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 82, 99.

*Carex drymophila*, Turcz. in *Bull. Soc. Nat. Mosc.* (1838) p. 104.

*Carex trichocarpa*, var. *orthostachys*, Kuek. in *Bot. Centralbl.* lxxvii. (1899) p. 92.



*Carex amurensis*, *Kuek. in Bot. Centralbl.* lxxvii. (1899) p. 94; *Meinsh. in Act. Hort. Petrop.* xviii. (1900) p. 407.

CHIHLI: Jehol (*David*, 1743!), Peking Mountains (*Bodinier*, 98!); SHINGKING: Liao River (*Ross*, 16!); SHANTUNG: Chefoo (*Fauvel*!); KIANGSU: Shanghai (*Wykeham Perry*, *Helol et d'Argy*, 39, 42!); ANWHEI: Nanking (*Faber*, 1525, 1526!). Mus. Brit; Herb. Kew.

Western Mongolia, Japan, Sachalin.

In this series of *Carex*, united by the American botanists under *C. aristata*, R. Br., the lower leaf-sheaths are of loose texture, with transverse lines and scattered needle-like white hairs; these hairs are continued up the under surface of the leaf, more or less. The utricle is glabrous in the lower half; in the upper portion it has sometimes white fine hairs on the angles or is slightly scabrous-pilose on the beak. In the 'Bot. Centralbl.' lxxvii. (1899) Kuekenthal has elaborately reviewed the material, and arranged it in four species and numerous varieties. It is true that the stems are much narrower in the Chinese plant than in the subarctic *C. aristata* type of R. Brown. The plant in Kuekenthal's *C. amurensis* sometimes is 30 in. high. The sheath of the lower bracts is  $\frac{1}{4}$  in. or  $2\frac{1}{2}$  in. in the North-American type *aristata*. In *C. trichocarpa*, Muhl., the utricles have numerous minute small hairs between the nerves, different in character from the white needle-like hairs sometimes present on the beak of *C. aristata*. I have not seen any Asiatic plant that is = the American *C. trichocarpa*. At all events, I cannot accept Kuekenthal's proposal to divide *C. orthostachys*, C. A. Meyer, into two species, and to put one of these into the North-American *C. trichocarpa*. The species are very close hereabout; and I think Bailey's proposal to collect the whole series under *C. aristata*, R. Br., quite worthy consideration.

9. *Carex Arnellii*, *Christ, in Kongl. Svensk. Vet.-Akad. Handl. Ny Folg.* xxii. no. 10 (1888) p. 177; *Meinsh. in Act. Hort. Petrop.* xviii. (1900) p. 363 in notâ, p. 411.

*Carex sylvatica*, *Maxim. Prim. Fl. Amur.* p. 312.

*Carex sylvatica*  $\beta$ , *Turcz. Fl. Baical.-Dahur.* ii. pars I. p. 279.

*Carex Maximowiczii*, *Boeck. in Linnæa*, xli. (1877) p. 237; *Franch. Pl. David.* i. p. 320, non *F. Schmidt*.

*Carex Turczaninowiana*, *Meinsh. in Act. Hort. Petrop.* xii. (1892) p. 411, xviii. (1900) pp. 296, 363; *Franch. in Nouv. Arch. Mus. sér. 3*, x. (1898) pp. 59, 97.

CHIHLI: Jehol (*David*).

North and Central Asia.

The utricle is pale, loose; the style-branches short; the nut obovoid—quite distinct from *C. sylvatica*, Huds.

10. *Carex atrata*, *Linn. Sp. Pl.* ed. 1, p. 976, ed. 2, p. 1387; *Schkuhr, Riedgr.* i. p. 52, ii. p. 42, t. x. fig. 77; *Kunth, Enum. Pl.* ii. p. 433; *Boott, Carex*, p. 114, t. 362; *Boeck. in Linnæa*, xl. (1876) p. 398; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 731; *Franch. Pl. David.* ii. p. 140, in *Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 36, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 146, 161; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 231.

SHINGKING: Changpeishan, 1000–7000 feet (*James!*); HUPEH, 6700 feet (*E. H. Wilson, 1019!*); SZECHUEN: Tachienlu (*Soulié*), Moupin (*David!*); YUNNAN: Yentzehay, alt. 11,500 feet (*Delavay ex Franchet*), Tsangchan, alt. 13,250 feet (*Delavay, 867, 2807!*). Herb. Kew.

Cold parts of the North Hemisphere.

The plants of James were probably collected on the Manchurian side of the frontier.

11. *Carex baccans*, *Nees, in Wight, Contrib.* (1834) p. 122; *Kunth, Enum. Pl.* ii. p. 513; *Boott, Carex*, p. 83 (syn. *C. curvirostri*, *Kunze, excl.*), tt. 234–236 et 238, 239; *Boeck. in Linnæa*, xl. (1876) p. 339; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 722; *Franch. in Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 251, 260; *Hook. f. Bot. Mag.* t. 7288.

CHINA (*Staunton*); FORMOSA: Tamsui (*A. Henry, 1402!*). Herb. Kew.

Tonkin, India, Malaya.

12. *Carex basiflora*, *C. B. Clarke, n. sp.*

*Folia* basalia 12–16 unc. longa, usque ad 4 lin. lata. *Culmi* frutescentes 5, usque ad 4–5 unc. longi. *Spicæ* 3–4 distantes, terminalis mascula, fusce olivaceæ. *Bracteæ* inflorescentiâ breviores. *Utriculus* rostro incluso 2 lin. longus, in humeris sparsim aciculari-pilosus, globosus; rostrum lineare cum  $\frac{3}{4}$  parte utriculi æquilongum, breviter 2-dentatum. *Nux* ellipsoidea, prava, trigona, apice pyramidata erostrata; styli basis linearis torta.

HUPEH: Patung (*A. Henry*, 3748!); NORTH CENTRAL CHINA: Laosan (*Hugh*!). Mus. Brit.; Herb. Kew.

*C. brevicuspis*, huic peraffinis, ob nucem longe rostratam præcipue differt.

13. *Carex bidentula*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1894-5) p. 41, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 132, 159 (excluding the examples with 2-fid style).

*Carex rigidula*, *Franchet, MS.*

SHINGKING: Moukden to Yalos R. (*Webster*, 181!). Herb. Kew.

Frequent in Japan.

Franchet says the affinity of this species is with *C. vulgaris*; this can only refer to the examples with bifid style that Franchet included here.

14. *Carex biwensis*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 28, et in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 197, t. 2. fig. 2.

HUPEH: Ichang (*A. Henry*, 7865!). Herb. Kew.

Japan.

This species differs very little from *C. capillacea*, Boott.

15. *Carex Boottiana*, *Hook. et Arn. Bot. Beech. Voy.* (1841) p. 273; *Kunth, Enum. Pl.* ii. p. 515, nec Boott, nec Bentham, MS.

*Carex Bongardi*, Boott, in *Trans. Linn. Soc.* xx. (1846) p. 144, et *Carex*, p. 160, t. 531.

*Carex wahuensis*, Boott, *Carex*, t. 351, quoad utriculum dextrorsum nucem superiorem (neque aliter).

BONIN ISLES (*Beechey*, n. 70; *C. Wright*, n. 370).

Beechey's plant has (as Hook. et Arn. describe) an oblong-lanceolate utricle, which is  $\frac{1}{4}$  inch long, stramineous, with 20-24 nerves. Boott's tab. 351 (except the utricle and nut above cited) is taken from the Sandwich plant which Boott subsequently published as a distinct species, viz. *C. nuptialis*, Boott (*Carex*, p. 175, t. 591). The nut has no beak, which character shuts it off from *C. tenebrosa*, Boott.

Var.  $\beta$ . *bracteosa*, *C. B. Clarke*. *Bracteæ* longissimæ, 2 inferiores usque ad 8-12 unc. longæ, inflorescentiam longe superantes. *Spica* mascula sessilis. *Glumæ* femineæ longe aristatæ, inferiores subbracteiformes. *Bractea* ima longius vaginata.

*Carex wahuensis*, *Boott*, in *Mem. Amer. Acad.* new ser. vi. (1859) p. 417; *Carex*, pp. 111, 112 partim, t. 354; *Palibin*, *Fl. Koreæ*, iii. p. 26 partim.

LUCHU ARCHIPELAGO (*C. Wright*, 363!). Herb. Kew.

The utricles agree closely with those of *C. Boottiana*, Hook. et Arn.; but it might be made a distinct species. The bracts are sometimes more than twice the length depicted in Boott's tab. 354. Though Boott included this Luchu plant in *C. wahuensis* (*Carex*, pp. 111, 112), yet he subsequently (p. 160) joined it with his *C. Bongardi* (i. e. *C. Boottiana*, Hook. et Arn.); for his locality for *C. Bongardi*, viz. "Bonin Island, Loochoo, Bongard (herb. Hooker), Wright, N. Pacific Exped. U.S. (herb. Gray)," must mean that he put together the Bonin and Luchu plants.

16. *Carex breviculmis*, *R. Br. Prodr.* p. 242; *Kunth*, *Enum. Pl.* ii. p. 522; *Boott*, *Carex*, p. 181,  *Ic. Ined.* 614-616; *Benth. Fl. Austral.* vii. pp. 436, 445; *Hook. f. Fl. New Zeal.* i. p. 283, t. 63; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 746; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1880) p. 388; *Franch. in Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 192, x. (1898) p. 95; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 232; *Palibin*, *Fl. Koreæ*, iii. p. 22.

*Carex Royleana*, *Nees*, in *Wight, Contrib.* p. 127; *Kunth*, *Enum. Pl.* ii. p. 441; *Boott*, *Carex*, p. 6, t. 19.

*Carex leucochlora*, *Bunge*, in *Mém. Sav. Étrang. St. Pétersb.* ii. (1835) p. 142; *Meinsh. in Act. Hort. Petrop.* xviii. (1900) pp. 295, 360.

*Carex Langsdorffii*, *Boott*, in *Trans. Linn. Soc.* xx. (1847) p. 144.

*Carex puberula*, *Boott*, in *Perry, Exped.* ii. (1857) p. 324.

*Carex discoidea*, *Boott*, in *Mem. Amer. Acad.* vi. (1859) p. 419.

CHINA BOR. (*Bunge*!); CHIHLI: Peking (*Bushell*!); SHING-KING (*Ross*, 66!); KIANGSU: Shanghai (*Maingay*, 644! *Faber*, 227!); CHEKIANG: Ningpo Mountains (*Faber*, 1523, 1524!), Puntó Isle (*Faber*, 24!); KIANGSI: Kiukiang (*Faber*, 1138, 1141! *Maries*!), Nankangfoo (*Miss Reid*!); FOKIEN: Amoy (*Grys*, 9964!); FORMOSA (*Oldham*, 597!); HUPEH: Patung (*A. Henry*, 3732! *E. H. Wilson*, 1659!); NORTH CENTRAL CHINA: Scekintsuen (*Hugh*!); COREA: Port Hamilton (*Wilford*, 706, 709! *Oldham*, 916!); LUCHU ARCHIPELAGO (*C. Wright*, 364!). Mus. Brit.; Herb. Kew.

India, Australia, New Guinea, New Zealand; abundant in Japan.

This species varies much in the points dear to Cyperologists:

the female glume has usually a long green arista, but sometimes the glume is acute or hardly mucronate; the spikes are usually approximate, the bract to the lowest spike much overtopping the inflorescence, but sometimes the lowest spike is somewhat distant and the bract short. From these variations, Booth and Franchet each constructed several species. But the material having now increased, Franchet has reduced the whole to *C. breviculmis*, R. Br.

The species, in its characteristic state, with long bracts, is not very like *C. pisiformis*, Boott; but the examples with short bracts, and glumes but little aristate, can hardly be distinguished from *C. pisiformis* but by their small, slender, sessile male spike.

17. *Carex brevicuspis*, C. B. Clarke, n. sp.

Fusco-viridis. *Folia* 2-pedalia, 3-4 lin. lata, fere glabra. *Culmi* 12-16 unc. longi. *Bracteæ* inflorescentiam non superantes. *Spicæ* 4, distantes, terminalis omnino mascula pedunculata 1 unc. longa,  $\frac{1}{5}$  unc. lata, rufo-brunea. *Spica* feminea 2 unc. longa, 3-4 lin. lata, fusco-viridis. *Glumæ* femineæ utriculis subbreviores, anguste lanceolatæ 8-nerviæ, apice acuminatæ lineares, vix mucronatæ. *Stylus* 3-fidus. *Utriculus* globosus, subinflatus, olivaceus, multistriatus, fere glaber, in rostrum subito contractus; rostrum lineare, cum utriculi corpore subæquilongum, apice brevi-2-cuspidatum. *Nux* trigona; rostro cylindrico, pravo (nuci *C. tenebrose* simillima).

KIANGSI: Kiukiang (*Maries!* *Shearer!*); CHEKIANG: Ningpo Mountains (*Faber*, 62!). Herb. Kew.

18. *Carex Brownii*, Tuckerm. *Enum. Caric.* (1849) pp. 15, 21; *Boott, Carex*, p. 161, t. 532; *Franch. et Sav. Enum. Pl. Jap.* ii. p. 578; *Boeck. in Linnæa*, xli. (1877) p. 151; *Franch. in Nouv. Arch. Mus. sér. 3, x.* (1898) pp. 49, 96; *Palibin, Fl. Koreæ*, iii. p. 22, syn. *C. lacistomate* excl.

*Carex striata*, R. Br. *Prodr.* p. 243; *Drejer, Symb. Caricol.* p. 28, t. 15; *Kunth, Enum. Pl.* ii. p. 458; non *Mich.*

*Carex ringens*, *Boott, in Mem. Amer. Acad. n. s. vi.* (1859) p. 419; *Franch. et Sav. Enum. Pl. Jap.* ii. p. 149.

*Carex bulbosa*, *Boott, MS.*

COREA: Chusan (*Wilford*, 926!). Herb. Kew.

Japan, Australia.

19. *Carex brunnea*, *Thunb. Fl. Jap.* (1784) p. 38; *Schkuhr, Riedgr.* ii. p. 16, t. Xx. fig. 111; *Kunth, Enum.* ii. p. 392; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 125; *Boeck. in Linnæa*, xxxix. (1875) p. 145; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 705; *Franch. in Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 241, 259; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 230.

*Carex gracilis*, *R. Br. Prodr.* p. 242; *Boott, Carex*, p. 59, tt. 154, 155, 156.

*Carex Commersoniana*, *Kunth, Enum. Pl.* ii. p. 391.

*Carex Fabri*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 90; *Franch. in Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 138.

CHINA (*Sir E. Home!*); KIANGSU: Shanghai (*Hance*, 10238!); CHEKIANG: Ningpo (*Swinhoe! Hance*, 9945!); FORMOSA: Kelung (*Oldham*, 598!), Tamsui (*A. Henry*, 1401!), Takow (*A. Henry*, 2009!); HUPEH: Patung (*A. Henry*, 5020, 5156!), Ichang (*A. Henry*, 3341!); SZECHUEN: Tchenkeoutin, 5500 feet (*Farges*, 225!); YUNNAN: Mosoyu (*Delavay*, 4090!), Mengtze, 5000 feet (*A. Henry*, 1401!); LUCHU ARCHIPELAGO (*C. Wright*, 368!). *Mus. Brit.; Herb. Kew.*

From Madagascar and India to Japan, the Sandwich Isles, and Australia; a frequent plant.

20. *Carex cæspitosa*, *Linn. Sp. Pl.* ed. 1, p. 978, ed. 2, p. 1388, pro parvâ parte; *Palmstr. Svensk Bot.* iv. (1805) t. 222; *Boeck. in Linnæa*, xl. (1876) p. 427; *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 87, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 127, 159.

CHIHLI: Peking (*Bodinier*).

Cool parts of the North Hemisphere; not rare in Japan, but much less common than *C. Goodenowii*, *J. Gay*. No Chinese specimen has been seen by me.

21. *Carex capillacea*, *Boott, Carex*, p. 44 (1858), t. 110; *Boeck. in Linnæa*, xxxix. (1875) p. 37; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 713; *Franch. in Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 197, 209 (var. *yunnanensi* excl.; syn. *C. nana*, *C. ontakensis* excl.).

*Carex rara*, *Palibin, Fl. Koreæ*, iii. p. 25, quoad exemplum in Seoul lecta?

*Carex capitellata*, *Boiss. Fl. Orient.* v. p. 399.

*Carex simplicissima*, *F. Muell. Fragm. Phyt. Austral.* ix. p. 191.

CHEKIANG : Ningpo Mountains, 2000 feet (*Faber*, 59!); YUNNAN : Outchay (*Delavay*, 406), Langkong (*Delavay*, 2078). Mus. Brit.; Herb. Kew.

Scattered from Laristan to Sachalin to New South Wales.

22. *Carex capilliformis*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 89, et in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 66, 97, t. 6. fig. 1; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

NORTH CENTRAL CHINA : Mount Thaepersan (*Hugh*); SZECHUEN : Tchenkeoutin (*Farges*). Mus. Brit.

23. *Carex cardiolepis*, *Nees, in Wight, Contrib.* (1834) p. 127; *Kunth, Enum. Pl.* ii. p. 516; *Boott, Carex*, p. 6, t. 17; *Boeck. in Linnæa*, xli. (1877) p. 319; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 744; *Franch. in Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 172, x. (1898) p. 93; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

SZECHUEN : Tchenkeoutin (*Farges*!).

Cabul and North-west Himalaya.

24. *Carex cercidascus*, *C. B. Clarke*, n. sp.

Viridis, 10-uncialis, debilior. *Folia* 10 unc. longa, vix 2 lin. lata, glabra. *Spicæ* c. 4, terminalis mascula,  $1\frac{1}{4}$  unc. longæ, graciles, virides, albidæ, paullo distantes (ima 1-2 unc. remota pedunculata); bracteæ foliiformes, inflorescentiam non superantes. *Glumæ* parvæ, in dorso fusco-virides, in lateribus albo-scariosæ, mucrone lanceolato incluso cum utriculis æquilongæ. *Stylus* 3-fidus. *Utriculus* (immaturus)  $1\frac{1}{4}$  lin. longus, anguste ellipsoideus, utrinque angustatus, levis subnervis, glaber, ore integro.

CHEKIANG : Tientai Mountains, 3000 (? feet) (*Faber*, 60!). Herb. Kew.

The young utricle is like a shuttle. The long, slender, green-white spikes resemble no Chinese species of this subgenus except *C. ichangensis*.

25. *Carex cernua*, *Boott, Carex*, iv. (1867) p. 171, t. 578; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 708, non *Franch.*

*Carex phacota*, *Franch. in Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 117, 158, non *Spreng.*

KIANGSU: Shanghai (*Carles*, 540 !); CHEKIANG (*Faber*, 248 b !).  
Herb. Kew.

Assam, Tonkin, Japan.

[*C. cernua*, Franchet, in *Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 116, is partly = *C. shimidzensis*, Franch., partly = *C. Prescottiana*, Boott—two species hardly separable; the terminal spike in both male.]

26. *Carex chinensis*, Retz. *Obs.* iii. (1783) p. 42; *Kunth*, *Enum. Pl.* ii. p. 517; *Boott*, *Carex*, p. 13, t. 36; *Benth. Fl. Hongk.* p. 402; *Boeck. in Linnæa*, xli. (1877) p. 231; *Franch. in Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 182, x. (1898) p. 94; *Finet et Franch. in Bull. Soc. Bot. France*, sér. 3, vi. (1899) p. 214.

*Carex Retzii*, *Nees, in Wight, Contrib.* p. 128 (*Retziana*, p. 132).

FOKIEN: Kuatun Mountains (*De la Touche*); HUPEH: Ichang (*A. Henry*, 3530 !); KWANGTUNG: Macao (*Gaudichaud! Cal-léry*, 42, 397 !); HONGKONG (*Harland*, 712 ! *C. Wright*, 80, 603 ! *Faurie*, 15866 ! *Bon*, 131 !), Pic Victoria (*Bodinier*, 541 !).  
Mus. Brit.; Herb. Kew.

In this species the nut is "apiculate," *i. e.* contracted to a very narrow neck below the style-base. The female glumes have usually long awns, but in *A. Henry's* 3530 they are short.

27. *Carex condensata*, *Nees, in Wight, Contrib.* (1834) p. 123 partim (*i. e.* *Royle*, n. 83); *Boott, Carex*, p. 86, tt. 247, 248; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 716; *Franch. in Nouv. Arch. Mus. sér. 3*, viii. (1896) pp. 252, 260 partim; non *Kunth*.

*Carex bengalensis*, *Boeck. in Linnæa*, xl. (1876) p. 347 pro magnâ parte.

*Carex indica*, *Nees, in Wight, Contrib.* p. 123 (*i. e.* *Royle*, n. 85); non *Linn.*

*Carex cruciata*, *Franch. in Nouv. Arch. Mus. sér. 3*, viii. (1896) p. 253 partim.

YUNNAN (*A. Henry*, 86 !), Mengtze 6000 feet (*A. Henry*, 9576, 9576 A, 9576 B !), Tsangchan (*Delavay*, 4174 !).  
Herb. Kew.

India; common.

28. *Carex conica*, *Boott, in Perry, Exped.* ii. (1857) p. 325; *Franch. et Sav. Enum. Pl. Jap.* ii. p. 143; *Boeck. in Linnæa*, xl.



(1876) p. 451; *Franch. in Nouv. Arch. Mus. sér. 3, ix.* (1897) p. 169.

COREA: Tsusima (*Wilford*, 778!). Herb. Kew.  
Japan.

29. *Carex continua*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi.* p. 717.

YUNNAN (*A. Henry*, 10161!). Herb. Kew.  
North India, Philippines.

30. *Carex cremostachys*, *Franch. in Bull. Soc. Philom. Paris, sér. 8, xi.* (1895) p. 34, *et in Nouv. Arch. Mus. sér. 3, ix.* (1897) pp. 119, 159.

YUNNAN: Yentzsehay 10,000 feet (*Delavay*, 4691).

31. *Carex cruciata*, *Wahlenb. in Kongl. Vet.-Akad. Stockh. Handl. xxiv.* (1803) p. 149; *C. B. Clarke, in Hook. f. Fl. Brit. Ind. vi.* p. 715; *Franch. in Nouv. Arch. Mus. sér. 3, viii.* (1896) pp. 253, 260; *Diels, in Engl. Bot. Jahrb. xxix.* (1900) p. 230.

*Carex benghalensis*, *Roxb. Fl. Ind. iii.* p. 572; *Benth. Fl. Hongk. p. 401*; *Boott, Carex*, p. 85 partly, *i. e. t.* 243.

*Carex valida*, *Nees, in Wight, Contrib. p. 123*; *Kunth, Enum. Pl. ii.* p. 513.

*Carex indica*, *Munro, in Seem. Voy. 'Herald,' p. 423*, non *Linn.*

*Carex stramentitia*, *Franch. in Bull. Soc. Philom. sér. 8, vii.* (1895) p. 33, *fide Franchet*; non *Boott.*

CHINA (*Fortune*, 115!); FORMOSA: Kelung (*Oldham*!), Bankinsing (*A. Henry*, 840!); HUPEH: Changyang (*A. Henry*, 7753!), Nanto Mts. (*A. Henry*, 4628!), Patung (*E. H. Wilson*, 1572!); SZECHUEN: Mt. Omei (*Faber*, 1121!); YUNNAN: Mengtze, 4000–5000 feet (*A. Henry*, 9971, 11151, 11440!), Szemao, 4500 feet (*A. Henry*, 12404!), Tchenfongchan (*Delavay*, 5078!); KWANGTUNG (*Ford*, 180! *Tate*! *Seemann*, 2398!); HONGKONG (*Furet*, 13! *Harland*, 693! *Bodinier*, 985! *C. Wright*, 606! *Hance*, 2248!). Mus. Brit.; Herb. Kew.

Malaya, Tonkin, India, Madagascar.

Part of *Franchet's C. cruciata* is here referred to *C. condensata*, *Nees*; and part of his *C. condensata* to *C. cruciata*, *Wahl.*

32. *Carex cryptostachys*, *Brongn. in Duperrey, Voy. 'Coquille'* (1829), p. 152, t. 25; *Kunth, Enum. Pl. ii.* p. 513; *Benth.*

*Fl. Hongk.* p. 403; *Boott, Carex*, p. 103, t. 310; *Boeck. in Linnæa*, xl. (1876) p. 327; *Franch. in Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 193, x. (1898) p. 95.

*Carex cyrtostachys*, *C. B. Clarke* (sphalm.), in *Hook. f. Fl. Brit. Ind.* vi. p. 714.

HONGKONG (*C. Wright*, 604! *Wilford*, 256! *Harland*, 791! *Hance*, 1158! *Faurie*, 15777! *Bodinier*, 103, 107!). Mus. Brit.; Herb. Kew.

Tonkin. Malaya Peninsularis et Insularis.

33. *Carex cylindrostachys*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 32, et in *Nouv. Arch. Mus. sér. 3*, ix. (1897) pp. 114, 158.

YUNNAN: Tapintze (*Delavay*, 1742!).

34. *Carex Davidi*, *Franch. Pl. David.* i. (1884) p. 319, et in *Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 165, x. (1898) p. 93; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 231.

*Carex ascotreta*, *Franch. in Nouv. Arch. Mus. sér. 3*, ix. (1897) p. 182, x. (1898) p. 94; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 232.

SHANSI (*David*!); HUPEH: Ichang (*Henry*, 7899!), Patung (*Henry*, 5320!); NORTH CENTRAL CHINA: Scekintsuen (*Hugh*!). Mus. Brit.; Herb. Kew.

35. *Carex Delavayi*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 29, et in *Nouv. Arch. Mus. sér. 3*, viii. (1896) pp. 208, 210, t. 2. fig. 1.

YUNNAN: Tapintze, 6000 feet (*Delavay*, 1739!).

There is, in two of the plants, a second remote peduncled axillary spike. The species is remote from the sect. *Raræ*, where Franchet has arranged it; but it is not easy to find a good place for it.

36. *Carex Dickinsii*, *Franch. et Sav. Enum. Pl. Jap.* ii. (1879) pp. 153, 581; *Franch. in Nouv. Arch. Mus. sér. 3*, x. (1898) pp. 80, 99.

*Carex retrorsa*, *Palibin, Fl. Koreæ*, iii. p. 25; *Maximowicz, MS.*, non *Schweinitz*.

*Carex retrorsa*, var. *minor*, *Boott, MS.*

COREA: Chusan (*Wilford*, 927!). Herb. Kew.

Frequent in Japan.

Franchet separated this plant as a species from the North-American *C. retrorsa*, Schwein., which has longer spikes with the beaks of the utricles prominently deflexed. *C. retrorsa*, Schwein., has also spongiose transversely lineolate leaves, which will not match.

37. *Carex digitata*, Linn. *Sp. Pl.* ed. 1, p. 975, ed. 2, p. 1384; *Schkuhr, Riedgr.* i. p. 76, ii. p. 57, t. H. fig. 38; *Kunth, Enum. Pl.* ii. p. 473; *Boeck. in Linnæa*, xli. (1877) p. 193; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1880) p. 389.

SHINGKING: E. of Fungwhangshan (*Ross*, 118!), Korean Gate (*Ross*, 424!). Herb. Kew.

A common European species, extending to North Asia.

38. *Carex dineuros*, C. B. Clarke, n. sp.

*Culmus* 2-3-pedalis, imâ basi foliis squamiformibus intectus. *Folia* supra culmi basin usque ad 12 unc. longa, 4 lin. lata, glabra. *Spicæ* 4-5, distantes, terminalis mascula  $1\frac{3}{4}$ - $2\frac{1}{2}$  unc. longa, pallide purpureo-brunea; femineæ  $1\frac{3}{4}$ - $2\frac{1}{2}$  unc. longæ, 2 lin. latæ, fusce luteo-virides; bractea ima pedalis, inflorescentiam multo superans. *Glumæ* ovatæ, obtusæ vel vix acutæ, femineæ cum utriculis æquilongæ. *Stylus* 3-fidus. *Utriculus*  $\frac{3}{4}$  lin. longus, obovoideo-ellipsoideus, glaber, trigonus, fusce viridis obscure purpureo-maculatus, apice contractus; in 2 angulis conspicue nervatus; rostrum perminutum, lineare, ore integro. *Nux* ellipsoidea, trigona, apice breviter pyramidata acuminata.

SHINGKING: Chienshan (*Ross*, 533!); CHEKIANG: Tsien Mts. (*Faber*, 1527!). Herb. Kew.

39. *Carex dispalata*, Boott, in *Perry, Exped.* ii. (1859) p. 325; *Boott, Carex* [1], p. 205; *F. Schmidt, Reise Amurl.* p. 196, t. 7. figg. 6-10; *Franch. et Sav. Enum. Pl. Jap.* ii. pp. 151, 580; *Franch. in Nouv. Arch. Mus. sér. 3, x.* (1898) pp. 74, 98.

*Carex marginaria*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1894-5) p. 40; *Franch. in Nouv. Arch. Mus. sér. 3, ix.* (1897) pp. 133, 160.

SHINGKING: Moukden (*Webster*, 287!). Herb. Kew.

Sachalin. Frequent in Japan.

The style is always 3-fid; whereas Franchet says, in both his descriptions, that the style in his *C. marginaria* (Toya, n. 1084) is bifid. But in *C. marginaria*, herb. Faurie (Toya, n. 10184),

the style is always 3-fid; and I believe Faurie has rightly inscribed this *C. marginaria*, Franch.

40. *Carex drepanorrhyncha*, Franch. *Pl. David*. ii. p. 141, in *Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 178, t. 4. fig. 2, x. (1898) p. 94; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 232.

SZECHUEN: Moupin (*David*).

Not seen. Franchet says that this species was very near *C. thibetica*, as doubtless it was; for in his description he says that the stems were clothed at base with short leaves *only*, and that the leaves of the innovations were longer than the stems; but his picture shows the stem with long leaves, and, I assume, requires correction.

41. *Carex Duthiei*, C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. (1894) p. 731, in *Journ. Linn. Soc., Bot.* xxxiv. (1898) p. 127.

*Carex atrata*, var. *pullata*, *Boott, Carex*, p. 114, t. 364 (spicis nucigeris nimis latis).

SZECHUEN: Tachienlu, 9000–13,500 feet (*Pratt*, 556!). Herb. Kew.

Himalaya.

*Pratt* n. 556 in herb. Mus. Brit. is *C. ustulata*, *Wahlenb.*; and Franchet has noted in MS. that the *Pratt* n. 556 in herb. Paris is "altogether different" from n. 556 in hb. Kew.

42. *Carex fastigiata*, Franch. in *Bull. Soc. Philom. Paris*, sér. 8, vii. (1894) p. 35, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 113, 158, viii. (1896) t. 12. fig. 2.

YUNNAN: Mosoyn (*Delavay*).

Not seen.

43. *Carex ferruginea*, Scop. *Fl. Carniol.* ed. 2 (1772), p. 225; *Boeck. in Linnæa*, xli. (1877) p. 228.

*Carex Scopoliana*, *Willd. Sp. Pl.* iv. p. 292; *Schkuhr, Riedgr.* ii. p. 77, t. Nnnn. fig. 199.

*Carex Mielichhoferi*, *Schkuhr, Riedgr.* ii. p. 66, t. Mmmm. fig. 198, non *J. E. Smith*.

A common species of Central Europe, extending to Turkestan.

Var. *tatsiensis*, Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 55, 97.

SZECHUEN: Tachienlu (*Soulié*, 834).

44. *Carex filicina*, Nees, in *Wight, Contrib.* (1834) p. 123; *Kunth, Enum. Pl.* ii. p. 510; *Boott, Carex*, p. 105 (varr.  $\alpha$  et  $\gamma$ ), tt. 311, 312; *Boeck. in Linnæa*, xl. (1876) p. 352; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 717; *Franch. in Nouv. Arch. Mus. sér. 3, viii.* (1896) pp. 254, 260; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 231.

HUPEH: Ichang, Nanto Mts. (*A. Henry*, 2669!); YUNNAN: Mengtze, 7000–8000 feet (*Henry*, 10210, 10210 A!), Tapintze, 7000 feet (*Delavay*, 2400!), Tchenfongchan (*Delavay*), foot of Kichau (*Delavay*, 4001). Herb. Kew.

Malaya, Tonkin, India.

45. *Carex filipes*, *Franch. et Sav. Enum. Pl. Jap.* ii. (1879) pp. 148, 576; *Franch. in Nouv. Arch. Mus. sér. 3, x.* (1898) pp. 68, 98.

*Carex sparsinux*, *Franch. in Nouv. Arch. Mus. sér. 3, x.* (1898) pp. 66, 97; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 233.

HUPEH: Patung (*Henry*, 3745!), Changlo (*Henry*, 6271!). Herb. Kew.

Also in Japan.

46. *Carex finitima*, *Boott, Carex*, p. 44 (1857), t. 112; *Boeck. in Linnæa*, xli. (1877) p. 247; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 736; *Franch. in Nouv. Arch. Mus. sér. 3, x.* (1898) pp. 65, 97; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

SZETCHUEN: Mt. Omei (*Faber*, 1149!). Herb. Kew.

Central Himalaya.

47. *Carex foraminata*, *C. B. Clarke*, n. sp.

Hemiscaposa. *Folia* basalia 18 unc. longa, 3 lin. (et ultra) lata, fere glabra. *Culmi* florentes 12–16 unc. longi, efoliati; bracteæ vix unciales, setaceæ. *Spicæ* 5, remotæ, terminalis omnino mascula pedunculata 3 unc. longa, 2–2½ lin. lata, intensius brunea. *Spica* feminea 2 unc. longa, 2 lin. lata; glumæ ellipticæ, acutæ, utriculos paullo superantes, in dorso 3-nerviæ viridi-luteæ, in lateribus brunæ. *Stylus* 3-fidus. *Utriculus* obovoideus, ¾ lin. longus, apice erostris, trigonus, multi-striatus, undique tenuiter microscopice pilosus. Nux anguste obovoidea, trigona, erostris in medio impressa (prava).

CHEKIANG: Ningpo Mts., 1800 feet (*Faber*, 63!); KIANGSI: Kiukiang Mts. (*Faber*, 1139!), Chinkiang et Kiukiang (*Maries*!). Herb. Kew.

This species is next *C. Davidi*, Franchet, of which it has the utricle. The male spike is much longer; the female glume is exaristate.

48. *Carex Forbesii*, *C. B. Clarke*, n. sp.

*Culmi* 14 unc. longi, recti. *Folia* 9 unc. longa,  $2\frac{1}{2}$  lin. lata, glabra. *Spicæ* 4, paullo distantes, rufo-bruneæ, 2 superiores masculæ 1 unc. longæ,  $2\frac{1}{2}$  lin. latæ, 2 inferiores femineæ parum pedunculatæ  $1\frac{1}{2}$  unc. longæ, 4 lin. latæ; bracteæ filiformes, inflorescentiam paullo superantes. *Glumæ* femineæ ovatæ, mucronatæ, in lateribus bruneæ, utriculis multo breviores. *Stylus* 3-fidus. *Utriculus*  $2\frac{1}{2}$ –3 lin. longus, ellipsoidei-conicus, suberostratus, coriaceus, glaber, obscure glanduloso-puberulus, a nervis numerosis striatus, ore breviter 2-dentato. *Nux* obovoidea, trigona, apice apiculata, basi substipitata.

SHANTUNG: Chefoo (*Forbes*!). Herb. Kew.

This has been arranged as a possible variety of *C. pumila* in hb. Kew, to which species it is closely allied. The long straight stems, the leaves being shorter than the stems, give it a different aspect, while the utricles are beakless with very numerous longitudinal striations.

49. *Carex Forficula*, *Franch. et Sav. Enum. Pl. Jap.* ii. (1879) pp. 131, 557; *Franch. in Nouv. Arch. Mus. sér. 3, ix.* (1897) pp. 135, 160, viii. t. 8. fig. 2; *Palibin, Fl. Koreæ*, iii. p. 22.

*Carex notha*, *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1880) p. 389, non *Kunth*.

*Carex notha*, var., *Boott, MS.*

SHINGKING: Kwandien Mts. (*Ross, 75*!); SHENSI: Mt. Miao-waungsan (*Hugh*!); COREA (*Kalinowski*). Mus. Brit.; Herb. Kew.

Japan, abundant.

*C. notha*, *Kunth*, is very like *C. Forficula*, but has the utricle nerved, its wall dotted over by transparent ferruginous elliptic glands sunk in the utricle-wall, its beak shorter.

50. *Carex gentilis*, *Franch. in Bull. Soc. Philom. Paris, sér. 8, vii.* (1895) p. 84, et *Nouv. Arch. Mus. sér. 3, viii.* (1896) pp. 246, 260, t. 4. fig. 2; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 230.

KIANGSI: Kiukiang Mts. (*Faber, 1188*!); YUNNAN: Tcheng-fongchan Mts. (*Delavay*!). Herb. Kew.

This species is very near *C. longicruris*, Nees; it has shorter style-branches and a smaller utricle.

51. *Carex gibba*, *Wahlenb. in Kongl. Vet.-Akad. Stockh. Handl.* xxiv. (1803) p. 148, char. emend.; *Schkuhr, Riedgr.* ii. p. 22; *Kunth, Enum. Pl.* ii. p. 420; *Boott, Carex*, p. 187; *Boeck. in Linnæa*, xxxix. (1875) p. 127; *Franch. et Sav. Pl. Jap.* ii. p. 132; *Franch. in Nouv. Arch. Mus. sér. 3, viii.* (1896) pp. 236, 241; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 230.

*Carex remota*, *Thunb. Fl. Jap.* p. 37, non *Linn.*

*Carex anomala*, *Boott, in Perry, Exped. Jap.* p. 327; *Palibin, Fl. Koreæ*, iii. p. 21, non *Steud.*

*Carex pteroloma*, *Kunze ex Steud. Cyp.* p. 242.

KIANGSU: Shanghai (*Maingay*, 360, 362!, *Perry!*); CHEKIANG (*Faber*, 246, 1025!), Ningpo Mts. (*Faber*, 222!); NORTH CENTRAL CHINA: Hykiascien (*Hugh!*); HUPEH: Patung (*A. Henry*, 1746, 3862), Ichang (*A. Henry*, 3479!); HUNAN: Shihmen (*A. Henry*, 7544!); SZECHUEN: Wushan (*A. Henry*); COREA: Port Hamilton (*Wilford*, 710!), Chusan (*Wilford*, 933 partim!). *Mus. Brit.*; *Herb. Kew.*

Japan.

It would be perhaps preferable to call this *C. gibba*, Boott; for Wahlenberg (also Willdenow and Schkuhr) say their *C. gibba* had the style 2-fid; so that their plant was not our *C. gibba*. On this ground, Boott started the new name *C. anomala*, but he afterwards accepted "*gibba*."

52. *Carex Goodenovii*, *J. Gay, in Ann. Sc. Nat. sér. 2, xi.* (1839) p. 191 (*Godenowii err. typogr.*); *Hook. f. Student's Fl. Brit.* ed. 3, p. 455.

*Carex cæspitosa*, *Linn. Sp. Pl.* ed. 1, p. 978, ed. 2, p. 1388, pro magnâ parte; *Gooden. in Trans. Linn. Soc.* ii. (1798) p. 195, t. 21; *Schkuhr, Riedgr.* i. p. 57, ii. p. 27, t. Aa. figg. 85 a, 85 b; *Quensel, in Svensk Bot.* iv. t. 222.

*Carex vulgaris*, *Fries, Novit. Mant.* iii. (1842) pp. 153-6; *Boott, Carex*, p. 166 (varieties all excl.), tt. 557, 558, 562, 563, 565; *Boeck. in Linnæa*, xl. (1876) p. 416; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 711; *Franch. in Nouv. Arch. Mus. sér. 3, ix.* (1897) pp. 124, 159.

*Carex decidua*, *Boott, Carex*, p. 63 (*i. e. n. 1 planta Douglasi*) et auctt. Amer.

CHIHLI: Peking (*David*); SHANTUNG: Chefoo (*Fauvel*); KIANGSI (*Miss Reid*), Kiukiang (*Shearer!*). *Herb. Kew.*

Common throughout the cool North Hemisphere; very common in Japan.

On referring to J. Gay's classical paper on *C. Goodenovii*, it will be seen that it only differs from *C. cæspitosa*, Palmstr., by the stronger nervation of the utricle. Linnæus included both (and other plants) under his "*cæspitosa*"; he probably had but a grain of the comparatively rare *C. cæspitosa*, Palmstr., to several ounces of the very common *C. Goodenovii*; but the name *C. cæspitosa* (Linn. partim) is reserved by botanists for the plant of Palmstruch.

53. *Carex Hancei*, C. B. Clarke, n. sp.

Undique pilosa. *Rhizoma* repens. *Culmi* 20 unc. longi, hemiscaposi. *Folia* usque ad 20 unc. longa, 3 lin. lata, in utrâque facie pilosa. *Spicæ* 2, terminalis mascula in pedunculo robusto piloso 3½-4 unc. longo, altera feminea in pedunculo subincluso; *bracteæ* 9 lin. longæ, patulæ, lanceolatæ, vagina 1 unc. longa, inflata, fissa. *Spica* mascula 9 lin. longa, 2-2½ lin. lata, ferrugineo-brunea; *glumæ* lanceolatæ, vix mucronatæ, in carinâ lutescentes. *Spica* feminea 9 lin. longa, 6 lin. lata, 20-nucigera, fusce viridis. *Glumæ* ovatæ, obtusæ, breviter aristatæ, utriculis (aristâ inclusâ) breviores, in lateribus scarioso-albæ. *Stylorum* rami 3, longi. *Utriculus* 2 lin. longus, ovoideus, subinflatus, multi-nervatus, dense pilosus; *rostrum* 1½-2 lin. longum, lineare, dense pilosum, dentibus 2 longis linearibus. *Nux* prava, stylobasi tenui, tortâ.

KIANGSU: Hills of Fengwangshan (*Forbes!*); CHEKIANG: Hills near Huchau (*Forbes!*); KIANGSI: Kiukiang (*Shearer!*). Mus. Brit.; Herb. Kew.

This was pasted down in herb. Hance with *C. Sampsoni*, Hance, but was called *C. longerostrata*, C. A. Meyer, in hb. Kew, which the male spike resembles.

54. *Carex Hancockiana*, Maxim. in *Bull. Soc. Mosc.* liv. (1879) pp. 66-68; *Franch. in Journ. de Bot.* iv. (1890) p. 320, in *Nouv. Arch. Mus. sér. 3, ix.* (1897) pp. 149, 161.

*Carex Buxbaumii*, *Franch. Pl. David.* i. p. 130, non *Wahlenb.*

*Carex Peiktusani*, *Komarov, in Act. Hort. Petrop.* xviii. (1901) p. 445.

CHIHLI: Peking, Mts. Pohuashan (*Bretschneider!*), Mts. Siao-



wutaishan, 3500–5000 feet (*Möllendorff*, 89!), Trappist Monastery (*Bodinier*, 100!), Jehol (*David*, 1821!). Herb. Kew.

Mandshuria.

55. *Carex Harlandi*, *Boott*, *Carex* (1860), p. 87, t. 255; *Benth. Fl. Hongk.* p. 403; *Franch. in Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 176 (*Boottii* citat. p. 132 excl.), x. (1898) p. 94.

HUPEH: Changyang, 6000 feet (*E. H. Wilson*, 435!); HONGKONG (*Harland*, 690! *Lamont*, 883! *Faurie*, 15778!), West Point (*Wilford*, 313! *C. Wright*), Mt. Gough (*Bodinier*, 101, 102!). Mus. Brit.; Herb. Kew.

*Franchet l. c.* cites also *Boott*, *Carex*, iv. p. 132, where *Boott* writes of “*C. manca*, *Boott*, var. *a. Harlandi*,” which *Franchet* supposes to mean a reduction of the sp. *C. Harlandi*, *Boott*, to a var. of *C. manca*.

56. *Carex heterolepis*, *Bunge*, in *Mém. Soc. Étrang. St. Pétersb.* ii. (1834) p. 137, et *Enum. Pl. China Bor.* p. 69.

NORTH-EAST CHINA: Panschan Mts. (*Bunge*!); CHIHLI: Peking (*Williams*, 13829!); SHINGKING: Jehol (*David*!). Mus. Brit.; Herb. Kew.

57. *Carex heterostachya*, *Bunge*, in *Mém. Soc. Étrang. St. Pétersb.* ii. (1834) p. 142; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1880) p. 389.

*Carex Bungeana*, *Debeaux*, *Fl. Tientsin*, p. 45.

*Carex songarica*, *Franch. in Nouv. Arch. Mus.* sér. 3, x. (1898) p. 92 in Obs.

CHINA BOREALIS (*Bunge*!); CHIHLI: Peking (*Williams*, hb. *Hance*, 11446!); SHINGKING: Liao R. (*Ross*, 14!), Jaolin (*Ross*, 391!); SHANTUNG: Tchefou (*Wykeham Perry*!); NORTH CENTRAL CHINA: Mt. Laoyan (*Hugh*!). Mus. Brit.; Herb. Kew.

This species has been very generally united with *C. nutans*, *Host*; but *Franchet* says correctly that it is nearer *C. songarica*, *Kar. et Kir.*, of which it has nearly the utricles. It differs a good deal from *C. nutans*, *Host*; it is slenderer, the inflorescence erect with 3–4 approximate spikes (the spikes are nearly always remote in *C. nutans*); the leaves are much narrower and glaucous beneath by reason of their being coated with grey, capitate glands. The utricles have fewer nerves; when young they have

in the lower half many large immersed yellow oil-glands, when ripe they are brown, not shining, nearly nerveless. The utricles are narrowed into a longer narrower beak (than in *C. nutans*) and the two teeth are smaller. The stolons are long, slender, hardening into long, slender wiry roots. *C. songorica*, Kar. et Kir., is a larger plant with remote spikes; the ripe fruits are chestnut-red, almost black, shining, and subbaccate.

58. *Carex humilis*, *Leysser, Fl. Hal.* ed. 1 (1761) p. 175, ed. 2 (1783) p. 235; *Boeck. in Linnæa*, xli. (1877) p. 198.

*Carex clandestina*, *Gooden. in Trans. Linn. Soc.* ii. (1794) p. 167; *Schkuhr, Riedgr.* i. p. 81, ii. p. 58, t. K. fig. 43; *Kunth, Enum. Pl.* ii. p. 476; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1880) p. 389.

SHINGKING: S. side Fungwhangshan (*Ross, 25!*). Herb. Kew.  
A European species, extending to the Altai.

59. *Carex ichangensis*, *C. B. Clarke*, n. sp.

*Culmi* 6–9 unc. longi, rigide cæspitiosi. *Folia* cum culmo æquilonga,  $1\frac{1}{2}$  lin. lata, glabra, supra parce scabrida, subtus obscure glanduloso-impressa. *Spicæ* 5, distantes; terminalis mascula, 1 unc. longa,  $\frac{1}{10}$  unc. lata, albo-viridis aut alba; spicæ femineæ 1 unc. longæ (juniores) perangustæ; *bracteæ* cum inflorescentiâ fere æquilongæ, foliiformes, virides. *Glumæ* femineæ oblongo-ellipticæ, albo-scariosæ, 3-nerviæ, nervo centrali viridi in aristam excurrente. *Stylus* 3-fidus. *Utriculus* 1 lin. longus, oblongo-ellipsoideus, stipitatus, trigonus, multi-striatus, viridis, undique tenuius pilosus, apice angustatus vix rostratus, ore subintegro.

HUPEH: Ichang (*Henry, 7860, 7894!*). Herb. Kew.

Henry n. 7860 is *C. lanceolata* var. *macrosandra*, Franchet, in *Nouv. Arch. Mus. sér. 3, ix.* (1897) p. 169; but the description is so unlike *C. ichangensis* that I suppose two plants have been mixed under the number 7860.

60. *Carex immanis*, *C. B. Clarke*, n. sp.

*Culmus* 2–3-pedalis. *Folia* 2 pedes longa, 6 lin. lata. *Spicæ* 6–8,  $4\frac{1}{2}$  unc. longæ,  $2\frac{1}{2}$  lin. latæ, terminalis basi mascula apice feminea. *Glumæ* femineæ utriculis breviores, truncatæ, mucrone brevi. *Styli* rami 2. *Utriculus*  $1\frac{1}{4}$ –2 lin. longus, ovoideus, levis, glaber, nervatus, non (vel minute) rostratus.

This species is perhaps nearest to *C. teres*, Boott (*Carex*,

t. 167), a Sikkim sp. In the fine specimens of Henry, many of the utricles contain two nuts.

HUPEH: Patung (*Henry*, 6113!). Herb. Kew.

61. *Carex incurva*, *Lightf. Fl. Scot.* (1777) p. 544, t. 24. fig. sinistr.; *Schkuhr, Riedgr.* i. p. 27, t. Hh. fig. 95; *Kunth, Enum. Pl.* ii. p. 375; *Boott, Carex*, p. 210; *Boeck. in Linnæa*, xxxix. (1875) p. 48; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 700; *Franch. in Nouv. Arch. Mus. sér. 3*, viii. (1896) pp. 223, 240.

SZECHUEN: Tongolo, in Kiala, on Tibet frontier (*Soulié*, 579!).

Cold places in the north hemisphere.

62. *Carex intermedia*, *Good. in Trans. Linn. Soc.* ii. (1794) p. 154; *Schkuhr, Riedgr.* i. p. 16, ii. p. 9, t. B. fig. 7; *Kunth, Enum. Pl.* ii. p. 376; *Regel, Fl. Ussur.* p. 180; *Meinsh. in Act. Hort. Petrop.* xviii. (1901) pp. 288, 320.

*Carex disticha*, *DC. Fl. Franç.* iii. p. 104; *Boott, Carex*, p. 125, t. 410; *Boeck. in Linnæa*, xxxix. (1875) p. 139, varr.  $\beta$ ,  $\delta$ , excl.; *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 29, in *Nouv. Arch. Mus. sér. 3*, viii. (1896) pp. 210, 239; non (aut pro parvâ parte) *Hudson, Fl. Angl.* (1762) p. 347.

*Carex siccata*, *Franch. Pl. David.* i. p. 319, non *Dewey*; *vide Franch. in Nouv. Arch. Mus. sér. 3*, viii. (1896) p. 210.

SHINGKING: Moukden (*Webster*, 188!), Jaoling (*Ross*, 381!); SHANTUNG: Chefoo (*Fauvel*); MONGOLIA CHINENSIS (*David*); NORTH CENTRAL CHINA (*Hugh*!). Mus. Brit.; Herb. Kew.

Cool places in the North Hemisphere.

For his *C. disticha*, type, Hudson cites in his first edition *Pluk. Phyt.* t. 34. fig. 7, drawn from a plant now in the Mus. Brit. which is *C. arenaria*, Linn. In his second edition, Hudson repeats this citation, and adds *Micheli, Gen.* t. 33. fig. 3, an excellent drawing, which is, beyond doubt of mine, *C. arenaria*, Linn., also.

As to Hudson's *C. disticha*  $\beta$ , he cites, in both first and second editions, only *Pluk. Phyt.* t. 34. fig. 4, which is totally unlike *C. intermedia*, Gooden.

It is possible that Hudson, among the material for his *C. disticha*, had plants of *C. intermedia*, and merely cited the pictures to give a general idea of his species. If that had been so, I prefer the name *C. intermedia*, Good.

63. *Carex ischnostachya*, Steud. *Cyp.* 1855, p. 222; Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) p. 47 cum syn. p. 96.

*Carex Ringoldiana*, Boott, in *Mem. Amer. Acad.* n.s. vi. (1859) p. 419, *Carex*, p. 199,  *Ic. Ined. in hb. Kew.* 671.

LUCHU ARCHIPELAGO (*C. Wright*, 365, 367!). *Herb. Kew.*  
Frequent in Japan.

64. *Carex Jackiana*, Boott, in *Proc. Linn. Soc.* i. (1846) p. 260, in *Trans. Linn. Soc.* xx. (1847) p. 132, *Carex*, p. 9, t. 25; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 735; Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 43, 46.

*Carex oxyphylla*, Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 57, 97.

*Carex instabilis*, Boott, *MS.*

YUNNAN (*Delavay*, 2669!), Mosoyn (*Delavay*, 2524!).

Khasia, Java.

Franchet, by inadvertence, founded his *C. oxyphylla* on the same two collections of Delavay which he had cited as the sole ground for the occurrence of *C. Jackiana* in China. The area of this species depends on its diagnosis; a var.  $\beta$ . *minor* is found in Ceylon and the Nilghiris; see Hook. f. in Trimen *Fl. Ceylon*, vol. v. p. 113. Several closely allied species occur in Japan.

65. *Carex japonica*, Thunb. *Fl. Jap.* (1784) p. 38; Schkuhr, *Riedgr.* i. p. 115, t. Ww. fig. 110; Baker et S. Moore, in *Journ. Linn. Soc., Bot.* xvii. (1880) p. 389; Franch. in *Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 77, 78, var.  $\alpha$  et  $\beta$  partim, non Boott.

*Carex Motoskei*, Miq. in *Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-6) p. 148; Franch. et Sav. *Enum. Pl. Jap.* ii. pp. 140, 584.

*Carex aphanolepis*, Franch. et Sav. *Enum. Pl. Jap.* ii. pp. 152, 580.

*Carex trichostyles*, Franch. et Sav. *Enum. Pl. Jap.* ii. pp. 152, 581.

*Carex Doniana*, var. *minor*, Boott, *MS.*

CHIHLI: Peking (*Bretschneider*!); SHINGKING: Chiensan (*Ross*, 432, 579!); NORTH CENTRAL CHINA: Mt. Ngosan (*Hugh*!), Kusan (*Hugh*, 166!). *Mus. Brit.*; *Herb. Kew.*

Common in Japan.

This is a slender plant, with scattered spikes; the largest female spike, figured excellently by Schkuhr, is 7 lin. long, ellipsoid; Thunberg described it as "ovate." Boott united it with the Indian *C. alopecuroides*, D. Don, which view is by no means impossible; but then, taking for his "type" the finest form of his broad species, fig. 257 (Boott's *C. japonica*, type) has spikes

2¼ in. long, and is the plant recorded in the present list as *C. alopecuroides*, D. Don, var. *chlorostachya*, C. B. Clarke. *C. japonica*, Thunb., as understood here, differs from *C. alopecuroides*, D. Don, by its smaller size and shorter spikes; moreover the under surface of the leaves is glabrate, not glaucous by reason of a dense covering of capitate white glands; the leaf-sheaths are not fimbriate; there is very often a leaf, or empty bract, added about the middle of the stem.

66. *Carex lamprosandra*, Franch. in *Bull. Soc. Philom. Paris*, sér. 8, vii. (1894) p. 45, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 179, x. (1898) p. 94.

YUNNAN: Mosoyn, 8000 feet (*Delavay*, 1743!).

Not examined.

67. *Carex lancifolia*, C. B. Clarke, n. sp.

Hemiscaposa. *Folia* basalia 6–8 unc. longa, 4 lin. lata, glabra. *Culmi* florentes 4–8 unc. longi, efoliati. *Bracteæ* vix 6 lin. longæ, vaginantes, limbo setaceo. *Spicæ* 4, distantes, terminalis omnino mascula 6 lin. longa, ¾ lin. lata, purpurea. *Spica* feminea 8 lin. longa, 2 lin. lata, lanceolata, 4–6-nucigera. *Glumæ* 2 lin. longæ, oblongæ, apice truncatæ vix mucronatæ, carinâ angustâ viridi lateribus læte rubro-bruneis. *Stylus* 3-fidus. *Utriculus* 2–3 lin. longus, anguste oblanceolatus, in dimidiâ parte inferiore contractus, sublinearis, trigonus, subnervius, minute pilosus; rostrum minimum, lineare, apice integrum. *Nux* ellipsoidea, trigona, basi in pseudo-stipitem linearem contracta; stylobasis angusta.

HUPEH (*Henry*, 5467!), Patung (*Henry*, 5368!), 6500 feet (*E. H. Wilson*, 1650!). Mus. Brit.; Herb. Kew.

68. *Carex laticeps*, Franch. in *Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 178, x. (1898) p. 94; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 232.

KIANGSU: Shanghai (*Faber*, 241 f!); ANWHEI: Nanking (*Faber*, 1181!); Kiangsi: Chinkiang et Kiukiang (*Maries*!); HUPEH: Ichang (*A. Henry*, 3332!). Herb. Kew.

69. *Carex Lehmanni*, Drej. *Symb. Caricol.* (1844) p. 13, t. 2; *Boott*, *Carex*, p. 113, t. 361; *Boeck*, in *Linnæa*, xl. (1876) p. 395; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 730; Franch. in

*Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 35, in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 148, 161.

HUPEH: Fang, 9000 feet (*Henry*, 6837!); SZECHUEN: Mt. Omei summit (*Faber*, 1125!), Tachienlu, alt. 9000–13,500 feet (*Pratt*, 646! *Soulié*, 637), Tongolo (*Soulié*, 331!), Tchenkeoutin (*Farges*!); YUNNAN: Tsangchan (*Delavay*, 1738!), Maeulchan 11,500 feet (*Delavay*, 4677!). *Mus. Brit.*; *Herb. Kew.*

Himalaya.

70. *Carex leiorrhyncha*, *C. A. Meyer*, in *Mém. Sav. Etrang. St. Pétersb.* i. (1831) p. 217, t. 9; *Kunth*, *Enum. Pl.* ii. p. 388; *Kunze*, *Suppl. Schkuhr*, p. 9, t. 2. fig. 1; *Boott*, *Carex*, p. 189; *Boeck.* in *Linnæa*, xxxix. (1875) p. 96; *Franch. Pl. David.* i. p. 319, et in *Journ. de Bot.* iv. (1890) p. 320, et in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 213; *Palibin*, *Fl. Koreæ*, iii. p. 23; cf. *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 702.

*Carex leiocarpa*, *Franch.* in *Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 239 err. typogr., non *C. A. Meyer*.

CHIHLI: Jehol (*David*, 1868), Peking (*Bodinier*, 86!); SHING-KING: Chienshan (*Ross*, 392, 495, 528!), Jaoling (*Ross*, 382!), Moukden (*James*); COREA: Soul (*Kalinowski*). *Herb. Kew.*

Central Asia, Mandshuria.

Bongard printed, in the list of his collections at Sitka, *C. leiocarpa*, *C. A. Meyer*; by an early slip this was copied as *C. leiorrhyncha*, *C. A. Meyer*, which species is still being vainly looked for in Alaska.

71. *Carex ligata*, *Boott*, in *Benth. Fl. Hongk.* (1861) p. 402, et *Carex* (1867), p. 211; *Franch.* in *Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 164, t. 7. fig. 1, x. (1898) p. 93.

*Carex hongkongensis*, *Franch.* in *Nouv. Arch. Mus.* sér. 3, x. (1898) p. 93 in notâ.

HONGKONG (*Harland*, 790 ex parte! *C. Wright*, 8, 78, 79, 602! *Faurie*, 15774! *Bon*, 138!), Mt. Kellet (*Bodinier*, 1432!), Mt. Gough (*Wilford*, 263! *Bodinier*, 501!), Victoria Peak (*Bodinier*, 108, 109!). *Herb. Kew.*

72. *Carex ligulata*, *Nees*, in *Wight, Contrib.* (1834) p. 127; *Kunth*, *Enum. Pl.* ii. p. 516; *Boott*, *Carex*, p. 45, t. 113; *Boeck.* in *Linnæa*, xli. (1877) p. 305; *C. B. Clarke*, in *Hook. f. Fl. Brit.*

*Ind.* vi. p. 747; *Franch. in Nouv. Arch. Mus.* sér. 3, x. (1898) pp. 69, 98; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 233.

CHEKIANG: Lu Mts., 3000 feet (*Faber*, 249), Konit Cheon (*Simon*); HUPEH: 6700 feet (*E. H. Wilson*, 1009!), Patung (*A. Henry*, 1864!), Nanto (*A. Henry*, 3948!); SZECHUEN (*Farges*, 910 a!); COREA: Tsusima (*Wilford*, 902!). *Mus. Brit.*; *Herb. Kew.*

India, Japan.

73. *Carex litorrhyncha*, *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 84, et in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 196, 209; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 229.

YUNNAN: Tchengfongchan (*Delavay*, 4907!).

This species is allied to *C. capillacea*, Boott; its utricle has a much longer beak.

74. *Carex longicruris*, *Nees, in Wight, Contrib.* (1834) p. 124; *Kunth, Enum. Pl.* ii. p. 417; *Boott, Carex*, p. 59, t. 157; *Boeck. in Linnæa*, xxxix. (1875) p. 148; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 705; *Franch. in Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 85, et in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 245, 260.

Western India; from the Himalayas to Ceylon.

Var.  $\beta$ . *Henryi*; paniculâ magis effusâ, ramo imo sæpe 6–10 unc. longo; utriculo minore.

*Carex Henryi*, *C. B. Clarke, ex Franch. in Nouv. Arch. Mus.* sér. 3, viii. (1896) p. 243 in Obs.

*Carex longicruris*, *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 230.

CHEKIANG: Ningpo Mts. (*Faber*, 10!); HUPEH: Ichang (*A. Henry*, 4266!), Nanto (*A. Henry*, 4495!), Changyang (*A. Henry*, 7502!), Patung (*A. Henry*, 5185, 5186!); SZECHUEN: Mt. Omei, 4000 feet (*Faber*, 1153!); YUNNAN: Tchenfongchan (*Delavay*!). *Mus. Brit.*; *Herb. Kew.*

Neither *C. longicruris*, type, nor this var. *Henryi* has been seen from North-east India.

75. *Carex longipes*, *D. Don, in Trans. Linn. Soc.* xiv. (1825) p. 329; *Kunth, Enum. Pl.* ii. p. 418; *Boott, Carex*, p. 190; *Boeck. in Linnæa*, xl. (1876) p. 376; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 704; *Franch. in Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 244, 260; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 230.

*Carex Bodinieri*, Franch. in *Bull. Soc. Philom.* sér. 8, vii. (1895) p. 85, in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 243, 260.

HUPEH: Patung (*A. Henry*, 4713!); HONGKONG: Happy Valley (*Bodinier*, 334!). Herb. Kew.

India.

76. *Carex longistolon*, Franch. in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 243, 260; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 230.

HUPEH: Ichang (*A. Henry*, 86, 2776!); SHENSI: Paokisan, Mt. Misouansan (*Hugh*!). Mus. Brit.; Herb. Kew.

Endemic.

77. *Carex longkiensis*, Franch. in *Bull. Soc. Philom. Paris*, sér. 8, vii. (1895) p. 86, et in *Nouv. Arch. Mus.* sér. 3, ix. (1897) p. 181, x. (1898) p. 94; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 232.

YUNNAN: Longki (*Delavay*).

Not seen. Allied to *C. fragilis*, Boott, *fide* Franchet.

78. *Carex luctuosa*, Franch. in *Nouv. Arch. Mus.* sér. 3, ix. (1897) pp. 130, 160; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 231.

SZECHUEN: Tchenkeoutin (*Farges*!).

“Style 2-fid; uppermost spike male; utricule ovate, attenuate into a short, rough beak; next *C. rigida*.”—*Franchet*.

79. *Carex macrocephala*, Spreng. *Syst.* iii. (1826) p. 808; *Kunth*, *Enum. Pl.* ii. p. 428; *Hook. Fl. Bor.-Amer.* ii. p. 215, t. 216; *Regel*, *Fl. Ussur.* p. 180, t. 12. figg. 8–12; *Boott*, *Carex*, p. 27, t. 69; *Boeck.* in *Linnæa*, xxxix. (1875) p. 66; *Debeaux*, *Fl. Tchefou*, p. 153; *Franch.* in *Nouv. Arch. Mus.* sér. 3, viii. (1896) pp. 237, 241; *Palibin*, *Fl. Koreae*, iii. p. 23; *hb. Willd.* MS. *fide* *Kunth*.

*Carex anthericoides*, Presl, *Rel. Haenk.* i. (1828) p. 204.

SHANGTUNG (*Favel*); CHEKIANG (*Faber*, 18!), Ningpo (*Hance*, 13411!); COREA (*Oldham*, 918, 1076!). Mus. Brit.; Herb. Kew.

Japan to Alaska and Oregon.

Culms unisexual.



## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

Part I. for 1901, consisting of 378 pages, price 21 shillings, was issued on July 10, 1902; Part II., completing the volume, is now in the press.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.

---

VOL. XXXV. is still in course of issue, the Parts already published being Nos. 242-247. Part 248 will complete the volume.

VOL. XXXVI. Parts 249-252 are published.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."

THE JOURNAL  
OF  
THE LINNEAN SOCIETY.

---

VOL. XXXVI. BOTANY. No. 253.

---

CONTENTS.

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.— PART XVIII. ....	297

---

See Notice on last page of Wrapper.

---

LONDON:  
SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;  
AND BY  
LONGMANS, GREEN, AND CO.,  
AND  
WILLIAMS AND NORWATE.  
1904.

F12

# LINNEAN SOCIETY OF LONDON.

---

## LIST OF THE OFFICERS AND COUNCIL.

Elected 25th May, 1903.

---

### PRESIDENT.

Prof. Sydney Howard Vines, M.A., D.Sc., F.R.S.

### VICE-PRESIDENTS.

C. B. Clarke, M.A., F.R.S.  
Frank Crisp, LL.B., B.A.

Prof. J. B. Farmer, M.A., D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

D. H. Scott, M.A., Ph.D., F.R.S.

Rev. T. R. R. Stebbing, M.A., F.R.S.

### COUNCIL.

Rev. R. A. Bullen, F.G.S.  
Charles Baron Clarke, M.A., F.R.S.  
Frank Crisp, LL.B., B.A.  
Prof. J. B. Farmer, M.A., D.Sc., F.R.S.  
Prof. J. Reynolds Green, M.A., D.Sc.,  
F.R.S.  
B. Daydon Jackson, Esq.  
George Masee, Esq.  
W. G. Ridewood, D.Sc., F.Z.S.

G. S. Saunders, F.E.S.  
Dukinfield H. Scott, M.A., Ph.D.,  
F.R.S.  
A. C. Seward, M.A., F.R.S.  
Rev. T. R. R. Stebbing, M.A., F.R.S.  
Col. O. Swinhoe, F.Z.S., F.E.S.  
Prof. Sydney H. Vines, D.Sc., F.R.S.  
Alfred O. Walker, Esq.

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

---

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

---

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*; the former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meets at 4 P.M., as required during the Session. The Members for 1903-1904, in addition to the officers, are:—

Herbert Druce, F.Z.S.  
Antony Gepp, M.A.  
Herbert Goss, F.G.S.  
Albert C. L. G. Günther, M.D., F.R.S.  
W. Botting Hemsley, F.R.S.

P. Chalmers Mitchell, M.A., F.Z.S.  
G. R. M. Murray, F.R.S.  
A. B. Rendle, M.A., D.Sc.  
A. G. Tansley, M.A.

80. *Carex maculata*, *Boott*, in *Trans. Linn. Soc.* xx. (1848) p. 128, et *Carex*, i. p. 9, t. 26; *Boeck*. in *Linnæa*, xli. (1877) p. 191; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 572; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 735; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1893) pp. 44, 96; *Palibin, Fl. Koreæ*, iii. p. 23.

*Carex micans*, *Boott*, in *Mem. Amer. Acad. n. s.* vi. (1859) p. 419.

CHEKIANG: Lu Mts. (*Faber*, 244, 248!); HUPEH: Ichang (*A. Henry*, 3650!); KWANGTUNG: Canton (*Sampson*, 368, 1879!); COREA: Chusan (*Wilford*, 933!). Mus. Brit.; Herb. Kew.

Japan, Australia, India.

81. *Carex manca*, *Boott*, in *Benth. Fl. Hongk.* (1861) p. 402, et *Carex*, iv. (1867) p. 131, t. 425; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 175, x. (1898) p. 93.

HUPEH: Patung (*E. H. Wilson*, 1658!); HONGKONG (*Harland*, 790 ex parte! *Hance*, 6648!). Mus. Brit.; Herb. Kew.

82. *Carex manciformis*, *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 63, 97; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

HUPEH (*E. H. Wilson*, 544!), Ichang (*A. Henry*, 3390, 7839!); Changyang (*A. Henry*, 5222!); SZECHUEN: Chunking (*Faber*, 1143!). Mus. Brit.; Herb. Kew.

The leaves are glaucous beneath by reason of the dense clothing of stalked grey glands. This is one of the Chinese group of *Carex* in which the nut has a cylindric neck dilated at top into a disk (*Franchet*).

83. *Carex Maubertiana*, *Boott, Carex*, i. (1858) p. 45, t. 114.

*Carex hebecarpa*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 747, vix *C. A. Meyer*.

*Carex hebecarpa*, var.  $\beta$ . *Maubertiana*, *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) p. 70; cf. *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 233.

*Carex ligulata*, varr. *minor et angustifolia*, *Boott!* MS.

HUPEH: Nanto, R. Yangtze (*E. H. Wilson*, 775!); Koni-tcheou (*Perny*). Herb. Kew.

North-east India; Assam.

84. *Carex melinacra*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 135, 160.

YUNNAN: foot of Tsangchan, near Tali (*Delavay*, ex *Franchet*).

Not seen. "Style 2-fid; uppermost spike female at top; utricle orbicular, resinous-punctate, suddenly contracted into a minute beak"—*Franchet*, who says it is allied to *C. sadoensis*, one of the Sect. *Vulgares*; but, from the description, it should be very near *C. cernua*, Boott, or even identic with it.

*C. notha*, Kunth [*Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 87], from Tali at the foot of Tsangchan, does not appear taken up in *Nouv. Arch. Mus.* (1897), unless *C. melinacra* is it.

85. *Carex Michauxiana*, *Boeck. in Linnæa*, xli. (1877) p. 336; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 81, 99.

*Carex rostrata*, *Mich. Fl. Bor.-Amer.* ii. (1803) p. 173; *Boott, Carex*, ii. p. 91, t. 267; non *Stokes*.

*Carex folliculata*, *Rudge, in Trans. Linn. Soc.* vii. (1804) p. 98, t. 9. fig. 4; non *Linn.*

*Carex xanthophysa*, varr. *nana et minor*, *Dewey, in Sillim. Amer. Journ.* ser. I. xiv. (1828) p. 353, t. R. figg. 57, 58.

*Carex abacta*, *L. H. Bailey, in Bull. Torrey Bot. Club*, xx. (1893) p. 427.

*Carex multinervia*, *Kuekenthal! MS. in Faber n. 243 a.*

KIANGSU: Shanghai (*Faber, 243 a!*). *Herb. Kew.*

Frequent in Japan; extending to Kamschatka, and in North America to North Labrador.

86. *Carex microglochin*, *Wahl. in Vet. Akad. Stockh.* xxiv. (1803) p. 140; *Schkuhr, Riedgr.* ii. p. 32, t. Ssss. fig. 110; *Kunth, Enum. Pl.* ii. p. 424; *Boott, Carex*, iv. p. 174, t. 589; *Boeck. in Linnæa*, xxxix. (1875) p. 32; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 712; *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 29, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 206, 210.

*Uncinia microglochin*, *Spreng. Syst.* iii. p. 830.

SZECHUEN: Tongolo, in Kiala (*Soulié!*).

Cool parts of the North Hemisphere; also South Andes.

87. *Carex microstoma*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1894) p. 48.

*Carex pallescens*, Linn., forma, Franch. in *Bull. Soc. Philom. Paris*, 3me sér., vii. (1895) p. 89.

YUNNAN: Longki (*Delavay*).

Not seen. I do not find either *C. pallescens* or *C. microstoma* in the Index to Franchet's *Carex* in *Nouv. Arch. Mus.* 3me sér., x. (1898) p. 114.

88. *Carex mollicula*, Boott, *Carex*, iv. (1867) p. 192, et  *Ic. ined. in hb. Kew.* t. 650; Franch. in *Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 73, 98.

COREA: Tsusima (*Wilford*, 842!); NORTH CENTRAL CHINA: Tonsianfu (*Hugh*!). *Mus. Brit.*; *Herb. Kew.*

Japan, frequent.

This is very near *C. japonica*, Thunb., *i. e.* its small form. In short, Boott's first specimen of *C. japonica* in his own herbarium, named by his hand, is *C. Wright* n. 15, which I call *C. mollicula*, Boott. Boott lays stress on the red glands immersed in the utricle of *C. mollicula*, which are prominent in *Wilford* n. 842, but wanting in most of the Japan material. *C. mollicula* must be diagnosed as having a very small sessile male spike, and the style-branches shorter than the utricle.

89. *Carex mosoynensis*, Franch. in *Bull. Soc. Philom. Paris*, 3me sér., vii. (1895) p. 31, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 242, 259.

YUNNAN: Mosoyn (*Delavay*, 4458, 4831!).

This species, again, may be called *C. brunnea*, Thunb., in a broad sense. It differs from both *C. Tonnerrei* and *C. gentilis* in its larger utricle; also from *C. Tonnerrei* in its strong creeping rhizome, and from *C. gentilis* by its nearly glabrous utricle.

90. *Carex moupinensis*, Franch. in *Nouv. Arch. Mus.* 2me sér., x. (1888) p. 102, 3me sér., viii. (1896) pp. 257, 260, t. 7. fig. 2, et *Pl. David.* ii. p. 140; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 231.

*Homolostachys sinensis*, Boeck. *Cyp. Novæ*, Heft 1 (1888) p. 38.

HUPEH: Ichang (*Henry*, 1234!); SZECHUEN: Moupin (*David*!); YUNNAN: Keoupatan (*Delavay*!). *Mus. Brit.*; *Herb. Kew.*

91. *Carex nemostachys*, *Steud. in Flora*, xxix. (1846) p. 23, et *Cyp.* p. 218; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 746; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 233.

*Carex excurva*, *Boott, Carex*, i. p. 57, t. 150; *Boeck. in Linnæa*, xli. (1877) p. 245.

*Carex Zollingeri*, *Kunze, ex Boeck. in Linnæa*, xli. (1877) p. 242.

KIANGSI: Kiukiang (*Shearer!*); HUPEH: Ichang (*Henry, 2931, 3531!*). *Mus. Brit.*; *Herb. Kew.*

India; Japan.

“Le n. 3531 (Hupeh, Ichang) du Dr. Henri s'éloigne” (from *C. gracilirostris*, *Franch.*) “par ses utricles pubescents,” *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 139, l. 17.

92. *Carex neurocarpa*, *Maxim. Prim. Fl. Amur.* (1859) p. 306; *Regel, Fl. Ussur.* p. 181; *Boott, Carex*, iv. p. 193; *Boeck. in Linnæa*, xxxix. (1875) p. 60; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 125; *Franch. in Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 215, 239; *Palibin, Fl. Koreæ*, iii. p. 24.

CHIHLI: Peking (*Bretschneider, 790!*), Mountains north of Peking (*Tatarinow*); NORTH CENTRAL CHINA: Mt. Laoysan (*Hugh!*), Kusan (*Hugh, 175!*); SHENSI: Paokisan, Mt. Kifonsan (*Hugh!*); COREA: Soul (*Gottsche*). *Mus. Brit.*; *Herb. Kew.*

Mandshuria, Amurland, Nippon.

93. *Carex nexa*, *Boott, in Benth. Fl. Hongk.* (1861) p. 402, et *Carex*, iv. (1867) p. 194; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 174, x. (1898) p. 93.

KIANGSI: Kiukiang (*Miss Reid!* young?); HUPEH: Patung (*E. H. Wilson, 1654!*); HONGKONG (*C. Wright, 100, 607! Hance, 1058! Harland, 790 ex parte! Lamont, 879! Wilford! Faurie, 15867, 15870! Faber*), Mt. Gough (*Wilford, 349! Bodinier, 502*), Pic Victoria (*Bodinier, 105, 106!*), Happy Valley (*Bodinier, 104*). *Mus. Brit.*; *Herb. Kew.*

This species differs from *C. ligata*, *Boott*, by having the male spikes 3 in. long,  $\frac{1}{8}$  in. broad, with bright straw-coloured glumes  $\frac{1}{4}$  in. long.

94. *Carex nubigena*, *D. Don, in Trans. Linn. Soc.* xiv. (1825) p. 326; *Kunth, Enum. Pl.* ii. p. 385; *Boott, Carex*, i. p. 1, t. 2; *Boeck. in Linnæa*, xxxix. (1875) p. 90; *Franch. Pl. David.* ii.



p. 139, *et in Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 214, 239; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 702; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 229.

*Carex albata*, *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* iii. p. 123; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 553; *Franch.* in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 216, 239; *Palibin*, *Fl. Koreæ*, iii. p. 21.

*Carex argyrolepis*, *Maxim. ex Franch. et Savat. Enum. Pl. Jap.* ii. p. 126.

*Carex yedoensis*, *Boeck.* in *Engl. Bot. Jahrb.* v. (1884) p. 515.

*Carex fallax*, *Steud.* in *Zoll. Verz. Ind. Archip.* ii. p. 60, *et Syn. Pl. Cyp.* p. 189.

HUPEH: Patung, 6000 feet (*A. Henry*, 6103 !), Hsingshan, 9000 feet (*A. Henry*, 6928 !); SZECHUEN: Moupin (*David*); YUNNAN: Mosoyn (*Delavay*, 3831, 4830 !), Koualapo (*Delavay*, 2203); COREA: Soul (*Kalinowsky*). *Mus. Brit.*; *Herb. Kew.*

Throughout India; Cabul, Malaya, Japan.

The type of *C. albata* has been reduced by Boott's hand to *C. nubigena*; and Franchet has reduced *C. argyrolepis* to *C. albata*. The three species are for me identic.

Var.  $\beta$ . *fallax*, *C. B. Clarke*.

*Carex fallax*, *Steud.* in *Zoll. Verz. Ind. Archip.* ii. p. 60, *et Syn. Pl. Cyp.* p. 189; *Boeck.* in *Linnæa*, xxxix. (1875) p. 57.

*Carex pterolepta*, *Franch.* in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 215, 239.

YUNNAN: Mosoyn (*Delavay*, 4459, 4829 !); Tsangchan, 8000 feet (*Delavay*, 1745 !).

Java.

95. *Carex nutans*, *Host*, *Gram. Austr.* i. (1801) p. 61, t. 83; *Schkuhr*, *Riedgr.* ii. p. 82, t. Qqqq. fig. 205; *Kunth*, *Enum. Pl.* ii. p. 418; *Boott*, *Carex*, iv. p. 194; *Franch. et Savat. Enum. Pl. Jap.* ii. (1879) pp. 154, 582 (?); *Ledeb. Fl. Ross.* iv. p. 315; *Boeck.* in *Linnæa*, xli. (1877) p. 296, forma  $\alpha$ ; *Franch. Pl. David.* i. p. 321.

CHIHLI: Peking (*Bretschneider*, 793 !), Jehol (*David*); NORTH CENTRAL CHINA: Kusan (*Hugh*, 77 !). *Mus. Brit.*; *Herb. Kew.*

This species extends from East Europe through Central Asia.

Bretschneider's plant matches the type of Host; all the rest of the Chinese "*nutans*" is *C. heterostachya*, Bunge, which only slightly differs.

96. *Carex obscura*, Nees, in *Wight, Contrib.* (1834) p. 126; *Kunth, Enum. Pl.* ii. p. 515; *Boott, Carex*, i. p. 70, t. 192; *Boeck. in Linnæa*, xl. (1876) p. 395; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 731, var.  $\beta$  incl.

*Carex Souliei*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 35, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 147, 161.

SZECHUEN: Tachienlu (*Soulié*, 412!); YUNNAN: Tongolo (*Soulié*, 958!).

Himalaya.

97. *Carex pachyrrhiza*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1894) p. 45, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 179, x. (1898) p. 94.

YUNNAN: Tsangchan, near Tapintze, 12,500 feet (*Delavay*, 2508).

Not examined.

98. *Carex panicea*, *Linn. Sp. Pl.* p. 1387; *Schkuhr, Riedgr.* i. p. 110, ii. p. 68, t. II. fig. 100; *Kunth, Enum. Pl.* ii. p. 460; *Ledeb. Fl. Ross.* iv. p. 291; *Bunge, in Mém. Sav. Etrang. St. Pétersb.* ii. (1835) p. 143; *Boott, Carex*, iv. p. 147, t. 472; *Boeck. in Linnæa*, xli. (1877) p. 154; *L. H. Bailey, in Proc. Amer. Acad.* xxii. (1886) p. 117.

CHINA (fide *Boott*); NORTH CHINA: Lunzuanssy (*Bunge*).

Cool parts of Europe and Asia.

*Boott* attributes this plant to China, but it does not appear on what authority, and there is no Chinese example in his herbarium. The plant is only introduced in America, fide *Bailey, l. c.*; and is not admitted by *Franchet* in his *Carex* of East Asia. The Chinese habitat, therefore, hangs almost entirely on the correctness of *Bunge's* identification.

99. *Carex parva*, Nees, in *Wight, Contrib.* (1834) p. 120; *Kunth, Enum. Pl.* ii. p. 419; *Boott, Carex*, i. p. 56, t. 418; *Boeck. in Linnæa*, xxxix. (1875) p. 38; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 712; *Franch. in Bull. Soc. Philom. Paris*, 8me

sér., vii. (1895) p. 29, *et in Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 207, 210.

*Carex macrorrhyncha*, *Kar. et Kir. in Bull. Soc. Nat. Mosc.* iii. (1842) p. 521.

YUNNAN: Tapintze Mts. (*Delavay*!), Mosoyn (*Delavay*, 797, 1746!).

Himalaya, Central Asia.

100. *Carex pediformis*, *C. A. Meyer*, in *Mém. Sav. Etrang. St. Pétersb.* i. (1831) p. 219, t. 10, fig. infer.; *Kunze*, *Suppl. Schkuhr*, p. 65, t. 16; *Kunth*, *Enum. Pl.* ii. p. 475; *Bunge*, *Enum. Pl. Chin. Bor.* p. 69; *Maxim. Prim. Fl. Amur.* pp. 309-311 (var. *pedunculatâ* incl.); *Boott*, *Carex*, iv. p. 196; *Regel*, *Fl. Ussur.* p. 181; *Boeck. in Linnæa*, xli. (1877) p. 197; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1880) p. 389; non *Franchet*.

*Carex lanceolata*, *Boott*, in *Perry, Exped.* ii. (1857) p. 326; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 134; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 167; *Palibin, Fl. Koreae*, iii. p. 23.

CHIHLI (*Bunge*); SHINGKING: south of Korean Gate (*Ross*, 65!), south of Fungwhangshan (*Ross*, 24!), Kwandien Mts. (*Ross*, 106!); SHANTUNG: Chefoo (*Fauvel*!); NORTH CENTRAL CHINA: Paoki, Chifengsan (*Hugh*, 66!); ANWHEI: Nanking (*Faber*, 242, 1142!); HUPEH (*E. H. Wilson*, 1666!), Patung (*A. Henry*, 3747, 5321!), Changlo (*A. Henry*, 6326!), Ichang (*A. Henry*, 7890, 7897!). *Mus. Brit.*; *Herb. Kew.*

Cool parts of Europe and Asia.

Var. *rostrata*, *Maxim. Prim. Fl. Amur.* p. 310; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 166.

SZECHUEN: Tachienlu (*Soulié*, 833!).

Amurland.

Var. *macrosandra*, *C. B. Clarke*.

*Carex lanceolata*, var. *macrosandra*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 169; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

HUPEH: Ichang (*Henry*, 1288!), Changyang (*Henry*, 7846!). *Herb. Kew.*

101. *Carex phacodes*, *Spreng.* [err. typogr. "phacota"] *Syst.* iii. (1826) p. 826; *Boott, Carex*, i. p. 63, t. 168; *C. B. Clarke, in*

*Hook. f. Fl. Brit. Ind.* vi. p. 708; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 117 pro parvâ parte.

*Carex lenticularis*, *D. Don, in Trans. Linn. Soc.* xiv. (1825) p. 331, non *Mich.*

*Carex platycarpa*, *Steud. Syn. Pl. Cyp.* p. 214.

*Carex rubro-brunea*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 123, 159; non *C. B. Clarke.*

— YUNNAN: Szemao Mts., 5000 feet (*A. Henry, 11759!*); KWANGTUNG: Canton (*Sampson, 1879 partim!*). *Mus. Brit.*; *Herb. Kew.*

Frequent in North India, extending to Japan and New Caledonia; also a form in South Africa.

102. *Carex phænocarpa*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 90, et *in Nouv. Arch. Mus.* 3me sér., x. (1898) p. 50.

YUNNAN: Outchay (*Delavay, 411*).

"Habit of *C. japonica*, Thunb." (*Franchet*). Not seen. *Franchet* says the male spike is solitary and 3-4 cm. long, the utricles glabrous, olivaceous, attenuated into a 2-toothed beak,—on which characters the species may have been near *C. pollens*, q. v.

103. *Carex Pierotii*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. (1865-66) pp. 148, 211; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 154; *Boeck. in Linnæa*, xli. (1877) p. 157; *Debeaux, Fl. Shanghai*, p. 66, et *Fl. Tchefou*, p. 189; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 90, 99; *Palibin, Fl. Koreae*, iii. p. 24.

*Carex suberea*, *Boott, in Hodgson, Nagasaki* (1861), p. 325 (nomen nudum), et *Carex*, iv. p. 165 (1867), t. 546; *Boeck. in Linnæa*, xli. (1877) p. 186.

KIANGSU: Shanghai (*Maingay, 363! Faber, 290!*), Shanghai and Woosung (*Debeaux*); CHEKIANG: Ningpo Mts. (*Faber, 25!*); COREA: Tsusima (*Wilford, 768!*), Port Hamilton (*Oldham, 1001!*). *Herb. Kew.*

Frequent in Japan.

104. *Carex pisiformis*, *Boott, in Perry, Exped.* ii. (1857) p. 324; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 142; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1880) p. 389; *Franch. in Nouv.*

*Arch. Mus.* 3me sér., ix. (1897) p. 185, t. 7. fig. 2, x. (1898) p. 94.

SHINGKING: Chienshan (*Ross*, 583!), south of Hingying (*Ross*, 304!); KIANGSI: Kiukiang (*Shearer*!). *Herb. Kew.*

Japan.

All the above-cited examples from China are very young. I believe they have been correctly referred here by Baker and Moore; but I could hardly have ventured to name them myself.

105. *Carex pollens*, *C. B. Clarke*, n. sp.

Robusta. *Culmi* apice crassi, leves. *Bractea* ima inflorescentiam superans, usque ad  $\frac{1}{2}$  unc. lata. *Spica* mascula 1, usque ad 2-3 unc. longa,  $\frac{1}{4}$ - $\frac{1}{3}$  unc. lata. *Spicæ* femineæ 5, distantes, usque ad 3 unc. longæ, densæ, luride virides, utriculis rectangulatim patulis. *Utriculi* (rostro lineari incl.)  $\frac{1}{8}$  unc. longi, globosi, inter se compressi, leves, glabri; rostri dentes 2 lineari-lanceolati.—Species *C. rhynchophysæ*, *C. A. Meyer*, peraffinis, ob spicam masculam unicam maximam diversa.

SHINGKING: Moukden, Yalu R. (*James! Webster*, 180!), Laoyeh Mts. (*James!*), Sungari R. (*James!*), Changdien (*Ross*, 161!). *Herb. Kew.*

Also collected at Tokio by Bisset.

Var.  $\beta$ . *angustior*, *C. B. Clarke*. *Culmi* apice triquetri, sæpe scabri. *Folia*  $\frac{1}{4}$  unc. lata. *Spica* mascula usque ad 4 unc. longa.

HUPEH (*A. Henry*, 7858!), Ichang (*A. Henry*, 3356!), Changyang (*A. Henry*, 7101!). *Herb. Kew.*

James's Moukden collections show the utricles nearly ripe; the species is described solely from these. The remainder of the material is young, and is put with the "type" merely on account of the very long, thick, solitary male spike.

The var. *angustior* may indicate a new species, all the examples being young; the male spike is similar in all.

106. *Carex Prainii*, *C. B. Clarke*, n. sp.

*Culmus* 2-pedalis, 40-60-stachyus. *Folia* cum culmo æqui-longa,  $\frac{1}{2}$  unc. lata. *Inflorescentia* 14 unc. longa; *bractea* ima foliiformis, vaginans; *paniculæ* partiales 4-5 unc. longæ. *Spicæ* 2 unc. longæ, lineares, fusce stramineæ, plures basi femineæ. *Glumæ* femineæ, aristellâ inclusâ, utriculis breviores. *Styli* rami 3. *Utriculi*  $\frac{1}{4}$  unc. longi, ellipsoidei trigoni, utrinque

angustati, stipitati, in parte superiore pilosuli; rostrum conico-lineare cum  $\frac{1}{2}$ – $\frac{2}{3}$  parte utriculi æquilongum.—Species *C. arridenti*, *C. B. Clarke* affinis.

— YUNNAN: Fengcheulin, 9500 feet (*Henry*, 10839!). Herb. Kew.

Naga Hills of N.E. India.

107. *Carex Prescottiana*, *Boott*, in *Trans. Linn. Soc.* xx. (1846) p. 135, et *Carex*, i. p. 45 partim (*i. e.* excluding tab. 115 and the Hawaiian plants); *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 710; *Franch.* in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 122, 159.

*Carex cernua*, *Franch.* in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 117, 159 pro magnâ parte; non *Boott*.

*Carex Fargesii*, *Franch.* in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 134, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 119, 159, t. 2. fig. 1; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 231.

SZETCHUEN: Tchenkeoutin, 5000 feet (*Farges*, 761!).

North-east India; Japan.

108. *Carex pruinosa*, *Boott*, in *Proc. Linn. Soc.* i. (1845) p. 255, in *Ann. Nat. Hist.* xvii. (1846) p. 52, in *Trans. Linn. Soc.* xx. (1846) p. 131, et *Carex*, i. p. 65, t. 174; *Miq. Fl. Ind. Bat.* iii. p. 352; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 709; *Palibin*, *Fl. Koreæ*, iii. p. 24.

*Carex picta*, *Boott*, in *Mem. Amer. Acad.* vi. (1859) p. 418; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 131; non *Steud.*

*Carex pruinosa*, var.  $\beta$ . *picta*, *Boott*, *Carex*, iv. p. 198; *Franch.* in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 155, 162; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 231.

*Carex dimorpholepis*, *Steud. Syn. Pl. Cyp.* p. 214.

*Carex Maximowiczii*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. (1866) p. 150.

SHANTUNG: Chefoo (*Fauvel*); CHEKIANG: Ningpo Mts. (*Faber*, 64, 1519!); HUPEH: Ichang (*A. Henry*, 3501!), Patung (*E. H. Wilson*, 1652!); YUNNAN: Longki (*Delavay*, 4999, 5162!), Tchenfongchan (*Delavay*, 417!); HONGKONG: Mt. Kellet (*Bodinier*, 1431!); COREA (*Kalinowsky*), Port Chusan (*Wilford*, 937!). Herb. Kew.

North-east Bengal; Java, Japan.

109. *Carex psycrophila*, *Nees*, in *Wight, Contrib.* (1834) p. 127; *Kunth*, *Enum. Pl.* ii. p. 463; *Boott*, *Carex*, i. p. 70, t. 191; *Boeck.*

in *Linnæa*, xl. (1876) p. 402; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 732; *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 88, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 146, 161; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 231.

*Carex celsa*, *Boott, Carex*, iii. p. 108, t. 330.

SZECHUEN: Tchenkeoutin (*Farges*!).

Himalaya.

110. *Carex pumila*, *Thunb. Fl. Jap.* (1784) p. 39; *Schkuhr, Riedgr.* ii. p. 82, t. Yy. fig. 112; *Kunth, Enum. Pl.* ii. p. 517; *Boott, Carex*, iv. p. 217; *Benth. Fl. Hongk.* p. 402; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 89, 99; *Debeaux, Fl. Shanghai*, p. 67; *Palibin, Fl. Koreæ*, iii. p. 21.

*Carex littorea*, *Labill. Pl. Nov. Holl.* ii. p. 69, t. 219.

*Carex Urvillei*, *Brongn. in Duperrey, Voy.* p. 157; *Kunth, Enum. Pl.* ii. p. 517.

*Carex nutans*, var.  $\delta$ . *pumila*, *Boeck. in Linnæa*, xli. (1877) p. 298.

SHANTUNG: Chefoo (*Perry*), Taliensuan (*Birnie*!); KIANGSU: Shanghai et Woosung (*Debeaux*); HONGKONG (*C. Wright*, 6, 605!); COREA: Chusan (*Wilford*, 930!); LUCHU ARCHIPELAGO (*C. Wright*, 362!). *Herb. Kew.*

Creeping in sea-sand, in Japan, Australia, New Zealand, Chile.

111. *Carex remota*, *Linn. Amœn. Acad.* iv. (1788) p. 293, et *Sp. Pl.* ed. 2, p. 1383 partim; *Schkuhr, Riedgr.* i. p. 46, ii. p. 22, t. E. fig. 23; *Kunth, Enum. Pl.* ii. p. 404; *Boott, Carex*, iv. p. 198; *Boeck. in Linnæa*, xxxix. (1875) p. 129; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 706; *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 84, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 235, 241; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 230.

SZECHUEN: S. Wushan (*Henry*, 5722!). *Mus. Brit.*; *Herb. Kew.*

Cool parts of the North Hemisphere.

Henry n. 5722 is young, and very possibly should be referred to one of the following varieties.

Var.  $\beta$ . *Rochebruni*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. (1894) p. 706, syn. *Franch. et Savat. excl.* Utriculi nervi minus conspicui, faciei planæ nulli; rostrum basi valide marginatum aut anguste alatum.

KIANGSI (*Miss Reid*).

This variety is frequent in the Himalaya, from Kashmir to Khasia. Also occurs in Java.

Var.  $\gamma$ . *reptans*, *Franch. in Nouv. Arch. Mus.* 3me sér., viii. (1896) p. 235; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 230.

SZECHUEN (*Farges*, 910), Mt. Omei (*Faber*, 1148!). *Herb. Kew.*

In the examples seen, the rhizome is woody, obliquely descending, about 1 in. preserved; in *Farges* 910, not seen, *Franchet* describes the rhizome as "reptans."

112. *Carex repanda*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. (1894) p. 720.

*Carex Wightiana*, *Boott, Carex*, i. p. 11 partim.

YUNNAN; Szemao Mts., 5000 feet (*A. Henry*, 9971 A, 11719!). *Herb. Kew.*

Khasia, Perak.

Both the specimens of *Henry* are very young, and possibly represent one of the species nearly allied to *C. repanda*.

113. *Carex rhynchophora*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 90, et in *Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 53, 97; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

SZECHUEN: Tchenkeoutin (*Farges*).

Not examined. Next *C. longerostrata*, *C. A. Meyer*, ex *Franchet*.

114. *Carex rigida*, *Gooden.!* in *Trans. Linn. Soc.* ii. (1792) pp. 193-195, t. 22. fig. 10; *Schkuhr, Riedgr.* i. p. 56, ii. p. 25, t. U. fig. 71; *Boeck. in Linnæa*, xl. (1876) p. 414 (excluding var. *trigyna*); *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 711 (excluding *C. orbicularis*, *Boott*); *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 40, in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 129, 159 (excluding the plants with three style-branches).

*Carex saxatilis*, *Oeder, Fl. Dan.* t. 159; *Schkuhr, Riedgr.* i. p. 54, ii. p. 25, t. I. fig. 40, t. Tt. fig. 40; *J. Gay, in Ann. Sc. Nat.* 2me sér., xi. (1839) pp. 187, 191.

*Carex vulgaris*, var. *alpina*, *Boott, Carex*, iv. p. 167, tt. 568, 570, 572.

SZECHUEN: Tachienlu, 9000-13,500 feet (*Pratt*, 738!), Tongolo (*Soulié*, 842)! *Mus. Brit.*; *Herb. Kew.*



Very common in the cool part of the North Hemisphere, extending to the Himalaya; but no example from Japan or from North-west America has been seen.

115. *Carex Rouyana*, *Franch. in Bull. Soc. Philom. Paris*, 3me sér., vii. (1894-5) p. 51, *et in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 67, 98, t. 7. fig. 1.

HUPEH: Patung (*E. H. Wilson*, 1651!).

In Japan, *e Franchet*.

The species is, as Franchet says, near *C. filipes*, Franchet, differing by the short leaves and large peduncled deeply-purple male spike. I identify Wilson's plant with *C. Rouyana* only by comparison with Franchet's picture.

116. *Carex rubrobrunea*, *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. (1894) p. 710, excluding citations.

YUNNAN: Szemao, 4500 feet (*Henry*, 11722!). *Herb. Kew.*

Frequent in Khasia, North-east India.

117. *Carex rupestris*, *All. Fl. Pedem.* ii. (1785) p. 264, t. 92. fig. 1; *Schkuhr, Riedgr.* ii. p. 36, t. Nnnn. fig. 280; *Kunth, Enum. Pl.* ii. p. 426; *Boott, Carex*, iv. p. 158, t. 516. figg. 1-5; *Boeck. in Linnæa*, xxxix. (1875) p. 34; *Franch. in Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 205, 209.

*Carex petraea*, *Schkuhr, Riedgr.* ii. p. 35, t. Kkk. fig. 139.

SHINGKING: Changpeishan (*James*!). *Herb. Kew.*

Cool parts of the North Hemisphere.

118. *Carex Sampsoni*, *Hance, in Journ. Bot.* vi. (1863) p. 89.

KWANGTUNG?, Kulangsu Island (*Sampson*, hb. *Hance*, 13059!). *Mus. Brit.*

Next to *C. manca*, Boott, fide Hance. Male spike  $\frac{3}{4}$  by  $\frac{1}{8}$  in. Female spike rather lax, with 8-12 utricles; peduncle up to  $1\frac{1}{4}$  in. Utricles ellipsoid, narrowed into a linear beak, sparingly pubescent.

Appears to have been overlooked by Franchet.

119. *Carex scaposa*, *C. B. Clarke, in Bot. Mag.* cxiii. (1887) t. 6940; *Franch. in Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 255, 260; *Finet et Franch. in Bull. Soc. Bot. France*, 3me sér., vi. (1899) p. 214; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 231.

FOKIEN: Amoy Interior (*Swinhoe!*), Kualun Mts. (*De la Touche*); YUNNAN: Tchenfongchang (*Delavay*, 410), Longki (*Delavay*, 5160); KWANGTUNG: Lofaushan, 3200 feet (*Ford*, 49!). Mus. Brit.; Herb. Kew.

Tonkin.

120. *Carex sclerocarpa*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 91, *et in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 71, 98; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

YUNNAN: Mts. of Tchenkeoutin (*Farges*).

"Allied to *C. setosa*, Boott."—*Franchet*.

Not examined.

121. *Carex siderosticta*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 89; *Franch. Pl. David.* i. p. 320, *et in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 195, t. 5. fig. 2, x. (1898) p. 95; cf. *Meinsh. in Act. Hort. Petrop.* xviii. (1900) pp. 300, 308.

Pseudo-*Carex plantaginea*, *Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 146.

CHIHLI: Peking, Mountain Valleys (*Bodinier*, 97), Jehol (*David*, 1812!); SHINGKING: Korean Gate (*Ross*, 433!); SHANTUNG: Chefoo (*Fauvel!*); HUPEH: Ichang (*A. Henry*, 3758!); NORTH CENTRAL CHINA: Paoki, Chifongsan (*Hugh*, 45!). Mus. Brit.; Herb. Kew.

Japan.

122. *Carex simulans*, *C. B. Clarke*, n. sp.

Hemiscaposa. *Innovationum folia* 18 unc. longa,  $\frac{1}{8}$ – $\frac{1}{6}$  in. lata, in paginâ inferiore a glandulis capitatis dense inspersa. *Spicæ* 1–1 $\frac{1}{4}$  unc. longæ, remotæ, terminalis mascula; glumæ ferrugineæ, carinâ viridi in aristellam excurrente. *Bractea* ima 2 unc. longa, basi vaginans. *Spicæ* femineæ laxæ. *Utriculi* rostro incluso fere  $\frac{1}{5}$  unc. longi, late ellipsoidei, trigoni, fusce olivacei, in humeris obscure pilosuli; rostrum cum  $\frac{1}{2}$  parte utriculi æquilongum, dentibus 2 lanceolatis. *Nux* æqualis in collum late cylindricum breve apice angustata.

CHEKIANG: Ningpo Mts. (*Faber*, 1522!); SZECHUEN: Hohkiang (*Faber*, 66); HUPEH: Ichang (*A. Henry*, 7892!), Patung (*A. Henry*, 3712!). Herb. Kew.

Resembles generally, and in the utricles, *C. manca*, Boott,

which species differs by having an unequal nut (*prava*, Boott) without a beak, the leaves not densely glandular beneath. *C. simulans* is a dusky olive colour, while *C. manca* is yellowish.

123. *Carex sociata*, Boott, in *Mem. Amer. Acad.* n. s. vi. (1859) p. 420, et *Carex*, iv. p. 200 ( *Ic. ined. in hb. Kew.* t. 674); Franch. in *Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 59, 97.

FORMOSA (*Swinhoe!* Oldham, 595!); LUCHU ARCHIPELAGO (*C. Wright*, 57 ex parte, 77, 361, 366!).

The utricle is distinctly hairy. Franchet says (and he appears to be right) that he had never seen this species, which is near *C. ligata*, Boott, but has a brown much thicker male spike.

124. *Carex Spachiana*, Boott, *Carex*, i. (1858) p. 43, t. 108; Franch. in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 137, 160.

KWANGTUNG: Macao (*Calléry*, 40, 398!). Herb. Kew.

In this species the style is 3-fid, the terminal spike female at top; Franchet, to whom (as to several modern caricologists) such characters are of the very lowest import, compares *C. Spachiana* with the Brunæ, or suggests it may be nearer his *C. Forficula*. *C. Spachiana* appears to me as remote from *C. brunea* and from *C. Forficula* as is any species in the genus.

125. *Carex stellulata*, Gooden. in *Trans. Linn. Soc.* ii. (1798) p. 144; Schkuhr, *Riedgr.* i. p. 45, ii. p. 19, t. C. fig. 14; Kunth, *Enum. Pl.* ii. p. 399; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 126.

*Carex echinata*, Boeck. in *Linnæa*, xxxix. (1875) p. 124; Franch. in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 232, 240; non Murray.

YUNNAN: Outchay (*Delavay*, 409!).

Cool parts of the North Hemisphere; plentiful in the Himalaya and Japan. Also in Australia, New Zealand.

Murray's *C. echinata*, the "type" described by him, is in Mus. Brit.; and is *C. divulsa*, Gooden.

126. *Carex stenantha*, Franch. et Savat. *Enum. Pl. Jap.* ii. (1879) pp. 146, 573; Franch. in *Nouv. Arch. Mus.* 3me sér., x. (1898) p. 50, t. 3. fig. 2.

*Carex stenolepis*, Franch. in *Nouv. Arch. Mus.* 3me sér., x. (1898) p. 96; non Torrey.

SHINGKING: Chieusan (*Ross*, 493, 535!), Moukden (*James!*). Herb. Kew.

Frequent in Japan.

127. *Carex stenophylla*, *Wahlenb. in Vet.-Akad. Stockh. Handl.* xxiv. (1803) p. 142; *Bunge, Pl. Chin. Bor.* p. 68; *Kunth, Enum. Pl.* ii. p. 374; *Maxim. Prim. Fl. Amur.* p. 304, et in *Bull. Soc. Mosc.* liv. (1879) p. 66; *Boott, Carex*, iv. p. 200; *Boeck. in Linnæa*, xxxix. (1875) p. 50; *Baker et S. Moore, in Journ. Linn. Soc., Bot.* xvii. (1880) p. 388; *Franch. Pl. David.* i. pp. 318, 319, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 220, 240; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 700.

CHIHLI: Peking (*Bushell, Bretschneider*, 866! *Williams*, 11447! *David*, 1786, 2373, *Bodinier*, 95, 96); SHINGKING: Ngaiyang (*Ross*, 230!); SHANTUNG: Chefoo (*Hancock*, 31! *Fauvel*); NORTH CENTRAL CHINA: Kisansien (*Hugh*!). *Mus. Brit.*; *Herb. Kew.*

Cool parts of the North Hemisphere.

128. *Carex stipitinux*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 31, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 246, 260, char. emend.; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 230.

HUPEH (*A. Henry*, 5971, 6610!), Patung (*A. Henry*, 4852!), Chiensih (*A. Henry*, 5838!); SZECHUEN: Tchenkeoutin (*Farges*!). *Herb. Kew.*

This species is closely allied to *C. brunnea*, Thunb., as stated by Franchet in *Bull. Soc. Philom.*, where it is placed at the head of the section "spiculæ apice masculæ." The terminal spike is male at top, usually for the greater part of its length, and frequently male altogether.

But, in *Henry 5971*, which Franchet worked from in *Nouv. Arch. Mus.*, there are about  $\frac{1}{8}$  of the female spikes in which the lower female flowers have not perfected utricles (as not infrequently in this section); Franchet has taken these as male flowers apparently (*i. e.* I find no males) and has created a new section for this species only, having "spiculæ inferne masculæ, vel interrupte masculæ et femineæ," the character of the section *Remotæ*, from which *C. stipitinux* is alien.

129. *Carex sutchuensis*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1894) p. 45.

*Carex setchuensis*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 180, x. (1898) p. 94; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 232.

SZECHUEN: Mt. Omei (*Faber*, 1146!), Tchenkeoutin (*Farges*).  
Not examined.

130. *Carex taliensis*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 34, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 139, 160, t. 2. fig. 2; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 231.

*Carex stachydesma*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 85, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 118, 159; *Diels, in Engl. Bot. Jahrb.* xxix. (1900) p. 231.

*Carex gracilirostris*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 138; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 231.

*Carex angusticurvis*, *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 160.

HUPEH (*A. Henry*, 5431!), Patung (*A. Henry*, 1725!);  
YUNNAN: Tali (*Delavay*), Longki (*Delavay*), Tchenfongchan (*Delavay*, 412!), Szemao Mts. (*Henry*, 11722!); NORTH CENTRAL CHINA: Laoyan (*Hugh!*). *Mus. Brit.*; *Herb. Kew.*

131. *Carex tapintzensis*, *Franch. in Bull. Soc. Philom. Paris*, 8me sér., vii. (1894) p. 44, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 164, x. (1898) p. 93.

YUNNAN: Tapintze (*Delavay*, 1740, 4532), Yengtzehay (*Delavay*, 3973).

Not examined.

132. *Carex tenebrosa*, *Boott, Carex*, ii. (1860) p. 88, t. 256; *Benth. Fl. Hongk.* p. 402; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 51, 96.

HONGKONG (*Harland*, 691! *hb. Kew.* 547!), Happy Valley woods (*Wilford*, 264!), Pic Victoria (*Bodinier*, 515). *Herb. Kew.*

The whole plant is olive-black, at least in the dried state.

133. *Carex tenuissima*, *Boott, in Proc. Linn. Soc.* i. (1845) p. 288.

*Carex panicea*, *hb. Bunge.*

NORTH CHINA: in *herb. Fielding, e Boott.*

This species is not in *herb. Boott*, nor is the name in the *Index to Boott's Carex*; but *Boott* says it was "*C. paniceæ*, L., affinis."

This could hardly have been the *C. tenuissima*, *Franch. et*  
LINN. JOURN.—BOTANY, VOL. XXXVI. Y

Savat. Enum. Pl. Jap. ii. p. 147, Franch. in *Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 63, 97, which resembled (except as to the glabrous utricle) *C. capilliformis*, Franch. Of this species Franchet has named an example in herb. Kew, which has minute lax spikes of 2-5 utricles. It is hardly conceivable that Bunge should have marked this "*C. panicea*, L." and that Boott should have described it as "*C. paniceæ*, L., affinis."

134. *Carex thibetica*, Franch. *Pl. David*. ii. (1888) p. 141, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 177, x. (1898) p. 94; Diels, in *Engl. Bot. Jahrb.* xxix. (1901) p. 232.

*Carex dichroa*, Franch. in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 86.

*Carex bella*, C. B. Clarke, ex Franchet, in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 177.

HUPEH (*A. Henry*, 6332!), Nanto Mts. (*A. Henry*, 3833!), Changjang (*A. Henry*, 5245!), Patung (*E. H. Wilson*, 1636!); SZECHUEN: Moupin (*David*), Tchenkeoutin (*Farges*, 910 c!); YUNNAN: Outchay (*Delavay*!). *Mus. Brit.*; *Herb. Kew.*

135. *Carex Thomsoni*, Boott, *Carex*, i. (1858) p. 1, t. 1; Boeck. in *Linnæa*, xxxix. (1875) p. 92; C. B. Clarke, in *Hook. f. Fl. Brit. Ind.* vi. p. 703.

*Carex Thomsoni*, Franch. in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 29, et in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 212, 239; Diels, in *Engl. Bot. Jahrb.* xxix. (1901) p. 229.

HUPEH: Ichang (*A. Henry*, 3392!); SZECHUEN: Souifou (*Delavay*, 141). *Herb. Kew.*

Bengal to Tonkin.

136. *Carex Tonnerrei*, C. B. Clarke, n. sp.

*Culmi* numerosi, dense cæspitosi, pedales. *Folia* cum culmis æquilonga,  $\frac{1}{2}$  unc. lata. *Panicula* 4 unc. longa,  $\frac{1}{2}$  unc. lata. *Gluma* feminea utriculo brevior. *Styli* rami 2, cum utriculo vix æquilongi. *Utriculi*  $\frac{1}{2}$  unc. longi, conspicue stipitati, glabri, leves, vel in rostri apice minute sparsim scabri.

SZECHUEN: Mount Omei (*Faber*, 1140!). *Herb. Kew.*

This species is close to *C. gentilis*, Franchet; from which it differs in its densely tufted stems, narrow leaves, and its stalked smooth glabrous utricle.

137. *Carex transversa*, Boott, in *Perry, Exped.* ii. (1857) p. 324, et *Carex*, iv. p. 202, *lc. ined. in hb. Kew.* t. 681; Franch.

*et Savat. Enum. Pl. Jap.* ii. p. 149; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 48, 96, t. 3. fig. 1.

KIANGSU: Shanghai (*Forbes, Faber, 441 b!*); CHEKIANG: Ningpo Mts. (*Faber, 1520!*), Punto Isle (*A. Henry, 24!*). Herb. Kew.

Japan.

*C. transversa* resembles *C. rigens*, i. e. *C. Brownii*, Tuckerman, in the leaden colour of the utricles contrasted against the scarious-white glumes, but has the beak of the utricle much longer.

138. *Carex Trappistarum*, *Franch. in Journ. de Bot.* iv. (1890) p. 320, *et in Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 133, 160.

CHIHLI: Peking (*Bodinier, 94*); KIANGSI: Kewkiang (*Shearer!*). Herb. Kew.

Japan.

Shearer's plant is young, and was named *C. heterolepis*, Bunge, in herb. Kew. Franchet has omitted altogether *C. heterolepis*, Bunge. The two, *C. heterolepis*, Bunge, and *C. Trappistarum*, Franch., are very close together, and may perhaps be united in one when more material has come to hand.

139. *Carex tristachya*, *Thunb. Fl. Jap.* (1784) p. 38; *Schkuhr, Riedgr.* ii. p. 48, t. Ww. fig. 109; *Kunth, Enum. Pl.* ii. p. 474; *Boeck. in Linnæa*, xli. (1877) p. 196; *Boott, Carex*, iv. p. 131, t. 424; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 135; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 162, x. (1898) p. 93; *Diels, in Engl. Bot. Jahrb.* xxix. (1901) p. 231.

*Carex monadelpha*, *Boott, in Perry, Exped.* ii. (1857) p. 323.

*Carex pocilliformis*, *Boott, Carex*, iv. p. 175, t. 593; *Boeck. in Linnæa*, xli. (1877) p. 195; *Franch. in Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 163, x. (1898) p. 93.

ANWHEI: Nanking (*Faber, 240!*); CHEKIANG: Ningpo Mts. (*Faber, 65!*); HUPEH: Ichang (*A. Henry, 7898!*), Antelope Glen (*A. Henry, 7901!*); KWANGTUNG: Whampoa (*Hance, 7198!*), Canton (*Sampson, 1150!*); COREA: Tsusima (*Wilford, 763!*). Mus. Brit.; Herb. Kew.

Frequent in Japan.

140. *Carex truncatigluma*, *C. B. Clarke*, n. sp.

Hemiscaposa. *Folia* basalia 18 unc. longa,  $\frac{1}{3}$  unc. et ultra lata, glabra. *Culmi* florentes 6-9 unc. longi, efoliati; bracteæ vix unciales, setaceæ. *Spicæ* 6, parum distantes, terminalis

sessilis omnino mascula,  $\frac{1}{2}$  unc. longa,  $\frac{1}{16}$  unc. lata, straminea. *Spica* feminea  $1\frac{1}{2}$  unc. longa,  $\frac{1}{8}$  unc. lata, fusce lutescens; glumæ femineæ utriculis multo breviores, late obovatæ, obtusæ, stramineæ, sublucidæ. *Stylus* 3-fidus. *Utriculus*  $\frac{1}{8}$  unc. longus, lanceolatus, trigonus, utrinque angustatus vix rostratus, multistriatus undique tenuiter microscopice pilosus, apice vix 2-dentatus. *Nux* anguste lanceolata, trigona, apice late cylindrica, utriculo parum angustior.

CHEKIANG: Ningpo Mts. (*Faber*, 1541!). Herb. Kew.

141. *Carex tsangensis*, *Franch.* in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1894) p. 44, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 181, x. (1898) p. 94.

YUNNAN: Mt. Tsangchan, alt. 7600 feet (*Delavay*, 2615).

Not examined.

142. *Carex tumida*, *Boott*, *Carex*, i. (1858) p. 66, t. 181; *Boeck.* in *Linnæa*, xli. (1877) p. 243; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 741.

YUNNAN: Mengtze, 5500 feet (*A. Henry*, 10833!). Herb. Kew.

Himalaya and Khasia.

143. *Carex unisexualis*, *C. B. Clarke*, n. sp.

*Culmi* omnino unisexuales. *Inflorescentia* feminea 3 unc. longa,  $\frac{1}{2}$  unc. lata, 30-stachya, densa; inflorescentia mascula multo angustior; bractea ima brevis. *Utriculi*  $\frac{1}{10}$  unc. longi, a glandulis nigro-rubris inspersi, in margine superiore hispidi; rostrum breve lineari-oblongum; ceteroquin ut *C. fluviatilis*, *Boott*, *Carex*, iv. t. 582.

*Carex fluviatilis*, *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 703, quoad plantas chinenses citatas; *Franch.* in *Nouv. Arch. Mus.* 3me sér., viii. (1896) pp. 213, 239; *Diels*, in *Engl. Bot. Jahrb.* xxix. (1901) p. 229.

KIANGSU: Shanghai (*Maingay*, 361!); KIANGSI: Kiukiang (*Shearer*!), Kiukiang Mts. (*Faber*, 1186!); HUPEH: Ichang (*Henry*, 594!). Herb. Kew.

Endemic.

144. *Carex ustulata*, *Wahlenb.* in *Vet.-Akad. Stockh. Handl.* xxiv. (1803) p. 156; *Kunth*, *Enum. Pl.* ii. p. 462; *Boott*, *Carex*, i. p. 70, t. 193. figg. 1, 2, et t. 194. fig. 1; *Boeck.* in *Linnæa*, xli.



(1877) p. 260; *C. B. Clarke*, in *Hook. f. Fl. Brit. Ind.* vi. p. 734; *Franch.* in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 38, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 155, 162.

*Carex atrofusca*, *Schkuhr*, *Riedgr.* i. (1801) p. 106, t. Y. fig. 82.

*Carex Prattii*, *Franch.* in *Bull. Soc. Philom. Paris*, 8me sér., vii. (1895) p. 32.

*Carex Moorcroftii*, *Franch.* in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 150, 161; non *Boott*.

SZECHUEN: Tachienlu (*Pratt*, 556! *Soulié*); YUNNAN: Mts. Likiang, 11,000 feet (*Delavay*, 2472!). Mus. Brit.

Cold parts of the North Hemisphere, down to the Himalaya. The Kew example of *Pratt* 556 differs from *Pratt* 556 at the Mus. Brit.

145. *Carex vaginata*, *Tausch.* in *Flora*, iv. (1821) p. 557; *Kunth*, *Enum. Pl.* ii. p. 451; *Kunze*, *Suppl. Schkuhr*, *Riedgr.* pp. 60-64, t. 15; *Boott*, *Carex*, iv. p. 148, t. 478; *Boeck.* in *Linnæa*, xli. (1877) p. 156; *Franch.* in *Nouv. Arch. Mus.* 3me sér., ix. (1897) p. 199, x. (1898) p. 95.

*Carex Mielichhoferi*, *Smith*, *Engl. Fl.* iv. p. 98, et in *Sowerby*, *Engl. Bot.* t. 2293; non *Schkuhr*.

*Carex phæostachya*, *Smith*, *Engl. Fl.* iv. p. 99, et in *Sowerby*, *Engl. Bot.* t. 2731.

*Carex scotica*, *Spreng. Syst.* iii. p. 819.

*Carex falcata*, *Turcz.* in *Bull. Soc. Mosc.* (1838) p. 104, xxviii. (1855) pars I. p. 341; *Maxim. Prim. Fl. Amur.* p. 311; *Baker et S. Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1880) p. 389; *Korshinsky*, in *Act. Hort. Petrop.* xii. (1892) p. 409.

SHINGKING: Kwandien Mts. (*Ross*!). Herb. Kew.

A European species extending to Kamschatka.

146. *Carex vesicaria*, *Linn. Sp. Pl.* ed. 1, p. 979, ed. 2, p. 1388 partim; *Schkuhr*, *Riedgr.* i. p. 124, ii. p. 85, t. Ss. fig. 106; *Kunth*, *Enum. Pl.* ii. p. 494; *Bunge*, *Enum. Pl. Chin. Bor.* p. 69; *Ledeb. Fl. Ross.* iv. p. 317; *Boott*, in *Hook. London Journ.* v. (1846) p. 69 (exemplis Americanis excl.), et *Carex*, iv. p. 162 pro minore parte, t. 536; *Boeck.* in *Linnæa*, xli. (1877) p. 319 partim.

NORTH CHINA: Ssijuissy (*Bunge*).

Europe, scattered also in North Asia.

The plant of *Bunge*, not seen, was doubtless *C. vesicaria* as understood by *Boott* and *Boeckeler*; but all the "*C. vesicaria*"

of Japan and East Siberia (also of the Himalaya) is now referred to subspecies of "*vesicaria*" identic with (or allied to) American forms.

147. *Carex villosa*, Boott, *Carex*, iv. (1867) p. 203,  *Ic. ined. in hb. Kew.* 683; *Franch. et Savat. Enum. Pl. Jap.* ii. pp. 142, 567 partim; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 44, 96; *Palibin, Fl. Koreæ*, iii. p. 25.

COREA: Soul (*Kalinowsky*).

Frequent in Japan.

This species has the utricle quite glabrous; the *C. villosa* of Boott, in Perry, *Exped.* ii. (1857) p. 327, with hairy utricles, was perhaps, wholly or partly *C. Duvaliana*, *Franch. et Savat.*

148. *Carex Wallichiana*, Prescott, ex Nees, in *Wight, Contrib.* (1834) p. 129; *Kunth, Enum. Pl.* ii. p. 483; *Boott, Carex*, i. p. 16, t. 18; *Boeck. in Linnæa*, xli. (1877) p. 310 partim; *C. B. Clarke, in Hook. f. Fl. Brit. Ind.* vi. p. 747; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) pp. 87, 99.

*Carex Fedia*, Nees, in *Wight, Contrib.* p. 129; *Kunth, Enum. Pl.* ii. p. 483.

*Carex hirta*, *Boiss. Fl. Orient.* v. p. 431; (var.) *Boeck. in Journ. Linn. Soc., Bot.* xviii. (1881) p. 105; non *Linn.*

YUNNAN: Mosoyu (*Delavay*, 796!), Tapintze (*Delavay*, 1744!), Kiangyu (*Delavay*, 4382!).

Cabul to Tonkin; frequent in North India.

149. *Carex Wilfordii*, *C. B. Clarke*, n. sp.

*Culmi* 1-2-pedales, subrobusti. *Folia* inflorescentiam sæpe superantia,  $\frac{1}{3}$  unc. lata. *Spicæ* 3-4, distantes, usque ad 4 unc. longæ, terminalis mascula. *Utriculi* rostro brevi incluso  $\frac{1}{8}$  unc. longi, fusce virides, obovoidei, 10-12-striati. *Nux* obovoidea, apice obtusius pyramidata, nigrobrunea, erostrata.

*Carex wahuensis*, *Boott, Carex*, iii. p. 111 partim, tt. 352, 353; *Franch. in Nouv. Arch. Mus.* 3me sér., x. (1898) p. 52 partim (var. 3, p. 53) *Palibin, Fl. Koreæ*, iii. p. 26 partim; non *C. A. Meyer*.

*Carex Bongardi*, var.  $\beta$ . robusta, *Franch. et Savat. Enum. Pl. Jap.* ii. pp. 134, 561, saltem pro magnâ parte.

CHEKIANG: Shrub Island (*A. Henry*, 47!); COREA: Port Hamilton (*Wilford*, 703, 707!). *Herb. Kew.*

Japan.

This forms the main part of *C. wahuensis*, Boott (and of Boeck. in Linnæa, xli. (1877) p. 279). The type of *C. wahuensis*, C. A. Meyer (Cyp. Novæ, t. 10), is here, and agrees with t. 10; it is a slender Sandwich Island plant with narrow leaves; the utricle has a linear beak  $\frac{3}{4}$  the length of its body; it is in general aspect as in the utricle very distinct from Wilford's 703, 707.

150. *Carex yunnanensis*, Franch. in *Bull. Soc. Philom. Paris*, 3me sér., vii. (1895) p. 31, et in *Nouv. Arch. Mus.* 3me sér., ix. (1897) pp. 116, 158.

YUNNAN: Tapintze (*Delavay*), Mosoyn (*Delavay*, 4788!). Diagnosed as having a bifid style; but it is always 3-fid.

### 159. GRAMINEÆ.

(By A. B. RENDLE, M.A., D.Sc., F.L.S.)

#### Tribe PANICEÆ.

1. *Paspalum conjugatum*, Berg, in *Act. Helvet.* vii. (1772) p. 129, t. 8; *Steud. Syn. Gram.* p. 21; *Hook. f. Fl. Brit. Ind.* vii. p. 11.

HONGKONG: Mt. Gough (*Ford*, 594!). Mus. Brit.; Herb. Kew.

Widely distributed in the tropics, especially of the New World.

2. *Paspalum distichum*, Linn. *Amæn. Acad.* v. p. 391; *Steud. Syn. Gram.* p. 29; *Hook. f. Fl. Brit. Ind.* vii. p. 12; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 721.

*Paspalum vaginatum*, Sw. *Prodr. Veg. Ind. Occ.* p. 21.

*Paspalum* sp., *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106.

FORMOSA (*Swinhoe*, in herb. *Hance*, 4972 in part!), Takow lagoon growing in a few inches of salt-water (*A. Henry*, 1036!), wet places (*A. Henry*, 1062!), in water (*A. Henry*, 1900!); KWANGTUNG: Pakshá in the extreme south of the province (*Hance*, 4972 in part!); LUCHU ARCHIPELAGO: Okinawa (*Tashiro*, fide *Hackel*). Mus. Brit.; Herb. Kew.

Widely distributed throughout the tropics.

3. *Paspalum scrobiculatum*, *Linn. Mant.* p. 29; *Steud. Syn. Gram.* p. 21; *Benth. Fl. Hongk.* p. 408; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 231; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 721, 723; *Hook. f. Fl. Brit. Ind.* vii. p. 10.

*Paspalum dissectum*, *Linn. Sp. Pl.* ed. 2, p. 81, in part; *Thunb. Fl. Jap.* p. 45.

*Paspalum Thunbergii*, *Kunth, ex Steud. Syn. Gram.* p. 28; *Pilger, in Engl. Bot. Jahrb.* l. c.; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 159; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643.

*Paspalum* sp., *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106.

KIANGSU: Kianang (*Staunton*!); FORMOSA: Takow (*A. Henry, 736, 1164*!); HUPEH (*A. Henry, 3986 A*!); HONGKONG (*Hance, 1111*!), roadside near Pokfoolum and West Point (*Lamont*!); LUCHU ARCHIPELAGO: Okinawa (*Tashiro, fide Hackel*). *Mus. Brit.*; *Herb. Kew.*

Japan.

Widely distributed in the warmer parts of the earth.

1. *Eriochloa polystachya*, *H. B. & K. Nov. Gen. et Sp.* i. p. 95, t. 31.

*Eriochloa annulata*, *Kunth, Rév. Gram.* i. p. 30; *Benth. Fl. Hongk.* p. 409; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267.

*Helopus annulatus*, *Nees, Agrost. Bras.* p. 17, et in *Hook. & Arn. Bot. Beechey's Voy.* p. 232.

*Paspalus annulatus*, *Fluegge, Monogr.* p. 133.

FORMOSA: Takow (*A. Henry, 1024, 1108*!); KWANGTUNG: Macao (*Vachell*!); HONGKONG (*Hance, 5200*!). *Mus. Brit.*; *Herb. Kew.*

Tropics of Old World and Brazil.

2. *Eriochloa villosa*, *Kunth, Rév. Gram.* i. p. 203, t. 13; *Debeaux, Fl. Shanghai*, p. 62, et *Fl. Tchéfou*, p. 155; *Franchet, Pl. David.* i. p. 322; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 49; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 126; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 164; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643, et sér. 2, iii. (1903) p. 501.

*Paspalum villosum*, *Thunb. Fl. Jap.* p. 45, t. 8; *Franchet, in Mém.*

*Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267; *Maxim. Prim. Fl. Amur.* p. 327.

CHIHLI: Jehol (*David*, 2207!), Peking (*Williams*, in herb. *Hance*, 13380!), Taliénwan (*Birnie*!); KIANGSU: Shanghai (*Maingay*, 758!); CHEKIANG: Ningpo Mts. (*Faber*, 285!); KIANGSI: Kiukiang (*Reid*!); HUPEH (*A. Henry*, 677, 4787!), Patung (*Wilson*, 1434!). *Mus. Brit.*; *Herb. Kew.*

Japan.

1. *Isachne albens*, *Trin. Sp. Gram. Ic. t.* 85; *Kunth, Enum. i.* p. 137; *Hook. f. Fl. Brit. Ind.* vii. p. 22.

*Panicum albens*, *Steud. Syn. Gram.* p. 96.

SZECHUEN (*Faber*, 1170!); YUNNAN: Mountain forests, 5000 to 5500 feet, Mengtze (*A. Henry*, 9379 A, 9379 B!). *Mus. Brit.*; *Herb. Kew.*

North India, Malaya.

2. *Isachne altissima*, *Debeaux, Fl. Shanghai*, p. 70.

KIANGSU: Shanghai (*Debeaux*).

I have seen no specimen. Is it a large form of *I. australis*?

3. *Isachne australis*, *R. Br. Prodr.* p. 196; *Kunth, Enum. i.* p. 136; *Benth. Fl. Hongk.* p. 414; *Debeaux, Fl. Shanghai*, p. 70; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 26; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 164; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 50, et in *Bull. Herb. Boiss.* vii. (1899) p. 643, et sér. 2, iii. (1903) p. 501; *Hook. f. Fl. Brit. Ind.* vii. p. 24.

? *Isachne muricata*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 236.

*Panicum patens*, *Osbeck, Resa*, p. 215, non *Linn.* (according to original specimen in *Herb. Mus. Brit.*).

CHINA (*Osbeck*!), East Coast (*Home*!); SHANTUNG: Chefoo (*Forbes*!); KIANGSU: Shanghai (*Maingay*, 565, 567, 695!); FOKIEN: Amoy (*Hance*, 1398 in part!); FORMOSA: Apes' Hill, Takow (*Playfair*, 36! *A. Henry*, 1076, 1920!), Tamsui (*Oldham*, 604!); HUPEH: Ichang (*A. Henry*, 244, 3649, 3520!); SHENSI (?): Mt. Laoysan (*Hugh*!); SZECHUEN: Mt. Uomisan, Tcentosen (*Hugh*!); KWANGTUNG: Canton (*Hance*, 1186, 1398 in part! *Sampson*, 291!); HONGKONG (*Forbes*! *Lamont*! *Seemann*, 658! *Wright*!); COREA (*Wilford*, 925!). *Mus. Brit.*; *Herb. Kew.*

Japan, India, Malaya, Australasia.

The small forms of this species, for example the Takow specimens from Playfair and Henry, approach *I. miliacea*, Roth.

4. *Isachne debilis*, Rendle, n. sp.

*Herba* parva debilis, caule tenui ramosa repente cum nodis radicantibus, ramis sub apice florenti ascendentibus. *Foliorum* vaginæ quam internodia breviores, compressæ, striatæ, pilosæ, præcipue in nodis marginibusque; ligula pilosa; lamina membranacea, lanceolata, acuta, sparse pilosa, venis inconspicuis, 5-9-lineas longa,  $1\frac{1}{2}$ -2 lin. lata. *Panicula* laxè pauciflora ramis filiformibus, spiculis ovalibus,  $\frac{2}{3}$  lin. longis. *Glumæ* steriles virides, membranacæ, ovales, cum marginibus incurvis et apice obtuso, 7-nerves, dorso hirsutulæ, superior quam inferior paullo minor; fertiles quam steriles breviores, dense et breviter pilosæ, suborbiculares, valde concavæ, cum marginibus inflexis paleam in dorso pilosam amplectantibus, nervis inconspicuis, inferior florem ♂, superior paullo minor, florem ♀ fovens.

FORMOSA: Tamsui (*Oldham*, 605!). Mus. Brit.; Herb. Kew.

Nearly allied to *I. miliacea*, Roth, but distinguished by the two very similar hairy fertile glumes.

5. *Isachne firmula*, Büse, in *Miq. Pl. Junghuhn*. p. 379.

*Isachne rigida*, Nees, ex *Steud. Syn. Gram.* p. 95; *Hook. f. Fl. Brit. Ind.* vii. p. 24.

*Isachne Myosotis*, *Benth. Fl. Hongk.* p. 415, non Nees.

*Panicum Rhignon*, *Steud. l. c.*

HONGKONG (*Wilford! Wright!*). Herb. Kew.

India, Malaya, Australia.

I think, judging from the description, that *I. firmula*, Büse, is identical with *I. rigida*, Nees.

Var. *nudiglumis*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 721.

LUCHU ARCHIPELAGO: Mount Yonahadake (*Tashiro*, fide *Hackel*).

Specimen not seen.

1. *Digitaria barbata*, *Willd. Enum. Hort. Berol.* p. 91.

*Panicum barbatum*, *Kunth, Rév. Gram.* i. p. 33; *Benth. Fl. Hongk.* p. 410.

*Panicum heteranthum*, Nees, in *Nov. Act. Nat. Cur.* xix. *Suppl.* 1 (1843) p. 174; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723.

*Paspalum heteranthum*, *Hook. f. Fl. Brit. Ind.* vii. p. 16.

FOKIEN: Amoy (*Hance*, 1379 in part!); FORMOSA: Takow spit (*A. Henry*, 1073!); KWANGTUNG: Pakhoi (*Playfair*, 143!); HONGKONG (*Hance*, 1379 in part!), Happy Valley (*Wright*!); HAINAN (*A. Henry*, 7972!). Mus. Brit.; Herb. Kew.

Malaya.

### 2. *Digitaria Henryi*, Rendle, n. sp.

*Herba*, ut apparet, annua; culmi tenues, e basi decumbente et radicante erecti, subpedales, vix  $\frac{1}{3}$  lin. diametro excedentes, ramosi. *Foliorum* vaginæ glabræ, inferiores laxæ, latæ, suprema cylindrica; ligula scariosa, 1 lin. longa, e vaginæ lateribus continuata; lamina flaccida, anguste lineari-lanceolata, acuta, plana, glabra, marginata, 5-nervis, circa 5 poll. longa, et  $2\frac{1}{2}$  lin. lata. *Spicæ* circa 5, laxiusculæ, ad  $2\frac{1}{2}$  poll. longæ, in axe communi  $1\frac{1}{2}$  poll. longo ordinatæ, suberectæ; rhachis communis tenuis, haud alata; spiculæ lanceolatæ, acutæ, virides, dorso et margine pilosæ,  $1\frac{1}{2}$  lin. longæ. *Gluma* i. deltoidea, minuta, enervis; gl. ii. circa dimidiam gl. iii. æquans, vel paullo superans, obtuse lanceolata, 3-nervis, nervus medianus conspicuus, latera et margines longe pilosæ; gl. iii. 7-nervis, nervi laterales a mediano remoti, inter nervos laterales et marginibus pilosa; gl. fertilis quam iii. paullo brevior, cum margine involuta lanceolata et breviter acuminata, 5-nervis, nervi laterales a mediano remoti, glabra.

FORMOSA: Apes' Hill, Takow (*A. Henry*, 1031!). Mus. Brit.; Herb. Kew.

The character of the spikelet closely recalls that of *Digitaria eriantha*, Steud., but the plant differs in habit, being apparently an annual, with slender delicate culms and flaccid leaves. The arrangement of the spikes is less digitate than in *D. eriantha*.

### 3. *Digitaria formosana*, Rendle, n. sp.

*Herba* minor annua; culmi graciles, cæspitosi, suberecti, plurinodi, 6-14 poll. alti. *Foliorum* vaginæ, ultima excepta, subbreves, omnes glabræ; ligula scariosa, obtusa, 1 lin. longa; lamina lineari-attenuata, basi sparsius et longe pilosa, margine crispulata,  $1\frac{1}{4}$ - $2\frac{1}{2}$  poll. longa. *Spicæ* binæ vel ternæ, digitatæ,

densiusculæ,  $1\frac{1}{4}$ - $2\frac{1}{2}$  poll. longæ, rhachis anguste alata; spiculæ ovali-lanceolatæ, acutæ,  $1\frac{3}{4}$  lin. longæ; cujusve paris alter sessilis, alter pedicellata. *Gluma* i. punctiformis; gl. ii.  $\frac{2}{3}$  longit. gl. iii., trinervis, margine dense pilosa, pili in spicula pedicellata magis patentés; gl. iii. spiculam æquans, 5-nervis, interstitiis sub lente pilosellis, margine dense ciliata, pili in spicula pedicellata patentés velut in *D. sanguinale* var. *ciliare*.

FORMOSA: Apes' Hill, Takow (*A. Henry*, 1941!). Mus. Brit.; Herb. Kew.

A somewhat delicate plant, recalling in habit *Digitaria debilis*, from which it differs in its shorter second glume and the bearded third glume.

4. *Digitaria humifusa*, *Pers. Syn.* i. p. 85.

*Panicum glabrum*, *Gaudin, Agrost. Helvet.* i. p. 22; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643.

*Panicum humifusum*, *Kunth, Rév. Gram.* i. p. 33.

*Paspalum ambiguum*, *DC. Fl. Franç.* iii. p. 16; *Hook. f. Fl. Brit. Ind.* vii. p. 17.

*Digitaria glabra*, *Beauv. Agrost.* p. 160; *Maxim. Prim. Fl. Amur.* p. 328; *Regel, Tent. Fl. Ussur.* p. 171.

SHINGKING (*Ross*, 78, 141!), Kaichow to Kinchow (*Ross*, 190!). Herb. Kew.

Japan, Himalayas, Central and North Asia, Europe.

The last-mentioned specimen approaches *D. longiflora*, *Pers.*

5. *Digitaria longiflora*, *Pers. Syn.* i. p. 85; *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 162.

*Digitaria tenuiflora*, *Beauv. Agrost.* p. 51.

*Panicum parvulum*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, iii. (1835) p. 205; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643.

*Panicum tenuiflorum*, *R. Br. Prodr.* p. 193.

*Paspalum brevifolium*, *Fluegg. Monogr.* p. 150; *Benth. Fl. Hongk.* p. 408; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 159.

*Paspalus chinensis*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 231.

*Paspalum longiflorum*, *Retz. Obs.* iv. p. 15; *Hook. f. Fl. Brit. Ind.* vii. p. 17.

CHINA (*Baird! Fortune*, 128!); SHANTUNG: Chefoo (*Forbes!*); KIANGSU: Shanghai (*Maingay*, 551, 552 in part, 107!); CHE-KIANG (*Staunton!*); FORMOSA: Takow (*A. Henry*, 1939!); HUPEH: Patung (*A. Henry*, 2504 in part!); SHENSI: Mt.



Miaowansan (*Hugh*!); NORTH CENTRAL CHINA: Leunteon, Fenghosan (*Hugh*!); KWANGTUNG: Whampoa (*Robertson*!); HONGKONG (*Hance*, 1070!). Mus. Brit.; Herb. Kew.

Japan, India, and warmer regions of the Old World.

In the plant from Chefoo (*Forbes*) the spikelets are 1 line long.

6. *Digitaria sanguinalis*, Scop. *Fl. Carniol.* ed. 2, i. p. 52; *Maxim. Prim. Fl. Amur.* p. 479.

*Panicum sanguinale*, Linn. *Sp. Pl.* p. 57; *Benth. Fl. Hongk.* p. 410; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Franchet, Pl. David.* ii. p. 142; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 163; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 643, 723.

*Paspalum sanguinale*, Lam. *Illustr.* i. p. 176; *Hook. f. Fl. Brit. Ind.* vii. p. 13; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 126.

CHINA: without locality (*Staunton*!); SHANTUNG (*Maingay*, 70!); KIANGSU: Shanghai (*Maingay*, 552 in part, 553, 670, 694!), Ningpo Mts. (*Faber*, 301!); KIANGSI: Kiukiang (*Reid*!); HUPEH: Patung (*A. Henry*, 2504 in part); KWANGTUNG (*Lamont*!), Canton (*Sampson*, 285!); HONGKONG (*Wilford*, 241! *Lamont*!); LUCHU ARCHIPELAGO (*Beechey*!). Mus. Brit.; Herb. Kew.

Widely distributed in warm countries.

Var. *ciliaris*, Doell, *Rhein. Fl.* p. 126; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 163; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 643, 723.

*Digitaria ciliaris*, Pers. *Syn.* i. p. 85; *Miq. Prol. Fl. Jap.* p. 164.

*Panicum ciliare*, Retz. *Obs.* iv. p. 16; *Debeaux, Fl. Tchéfou*, p. 154; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267.

*Panicum commutatum*, Nees, in *Hook. & Arn. Bot. Beechey's Voy.* p. 232, non in *Linnaea*, vii. (1832) p. 274.

EAST COAST: without locality (*Horne*!); CHIHLI: Peking (*Bretschneider*, 818! *Bushell*! *Williams*, in herb. *Hance*, 1556!); SHINGKING: Kaichow Hills (*Ross*, 309!); CHEKIANG (*Staunton*!); FORMOSA (*Campbell*!), Takow, near sea-beach cliff (*A. Henry*, 727!), South Cape (*Schmüerer*, in coll. *Henry*, 250!); SHENSI: Mt. Kiansan, Huscien-sien, and Kaolinsien (*Hugh*!); KWANGTUNG: Sam Cocks Island near Macao (*Staunton*!), Macao (*Vachell*!). Mus. Brit.; Herb. Kew.

Japan, Europe, and warmer parts of Old World.

Var. *cruciata*, *Hook. f. Fl. Brit. Ind.* vii. p. 14.

HUPEH: Changyang (*A. Henry*, 7508!), Patung (*A. Henry*, 5044!); SZECHUEN (*A. Henry*, 7036!), a young plant with panicle still partly enclosed in the uppermost leaf-sheath, which from the structure of the spikelet may belong to this variety. Mus. Brit.; Herb. Kew.

Himalayas.

Var. *extensa*, *Hook. f. Fl. Brit. Ind.* vii. p. 16; *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 163.

FORMOSA: Apes' Hill, Takow (*A. Henry*, 1128!), Takow (*A. Henry*, 764!); LUCHU ARCHIPELAGO (*Beechey*!). Mus. Brit.; Herb. Kew.

India, Tropical Africa.

Var. *pruriens*, *Hook. f. Fl. Brit. Ind.* vii. p. 15.

KIANGSI: Kiukiang (*Shearer*!); LUCHU ARCHIPELAGO (*Wright*!). Herb. Kew.

India, Malaya.

7. *Digitaria setifolia*, *Stapf, in Dyer, Fl. Cap.* vii. p. 376.

*Panicum fibrosum*, *Hack. in Oest. Bot. Zeitschr.* li. (1901) p. 332.

KWANGTUNG: near Canton (*Hance*, 7425!); HONGKONG: hills on the south side of the island (*Wilford*, 257!). Mus. Brit.; Herb. Kew.

South Africa.

The leaves in the Chinese specimens are broader and less rigid than in the South African, not glaucous and somewhat more sparsely hairy; the hairs, moreover, are finer. There is no difference between the spikelets. It is impossible to separate specifically the plants from the two countries, or to regard them otherwise than as geographical forms. I find MS. notes by General Munro in the Kew Herbarium on the ticket with Wilford's specimen, pointing out its identity with the South African plant.

8. *Digitaria tenuispica*, *Rendle*, n. sp.

*Culmus* repens, radicans, pertenuis, ramosus, rami erecti, florentes ad 9 poll. longi. *Foliorum* vaginæ tenues, glabræ; ligula truncata, vix 1 lin. longa; lamina anguste lineari-lanceolata, flaccida, facie sparse pilosa, margine indurata, ad 2-2½ poll. longa.

*Spicæ* digitatæ, ternæ, pertenuæ, 2-3½ poll. longæ; rhachis anguste viride-alata; spiculæ densiusculæ, 1¼-1½ lin. longæ, anguste lanceolatæ, acutæ, alter cujusque paris pedicellata, pedicellus ¾ lin. longus. *Gluma* i. pæne obsoleta, triangularis; gl. ii. suboblunga, quam gl. iii. circa duplo brevior, 3-nervis, breviter pilosa; gl. iii. spiculæ æquans, anguste lanceolata, acuta, 5-nervis, nervis lateralibus in parte inferiore sparsius pilosis.

KIANGSU: Shanghai (*Maingay*, 554!); KWANGTUNG: Macao (*Buchanan-Hamilton*!). Mus. Brit.; Herb. Kew.

Near the Sumatran *D. pertenuis*, Büse, which, from the description, it apparently resembles in habit, but is distinguished by its lanceolate, not oval, spikelets, shorter second glume, and absence of hairs between the veins on the larger barren glume.

9. **Digitaria ternata**, *Stapf*, in *Dyer*, *Fl. Cap.* vii. p. 376.

*Panicum ternatum*, *Hochst.* in *Flora*, xxiv. *Intelligenzbl.* i. (1841) p. 19; *Steud. Syn. Gram.* p. 40.

*Paspalum ternatum*, *Hook. f. Fl. Brit. Ind.* vii. p. 17.

YUNNAN: Mengtze, cliff at 4700 ft. (*A. Henry*, 11180!). —  
Mus. Brit.; Herb. Kew.

India, Tropical and South Africa.

1. **Panicum acroanthum**, *Steud. Syn. Gram.* p. 87; *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 275; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 162; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 49, et in *Bull. Herb. Boiss.* vii. (1899) p. 644, et sér. 2, iii. (1903) p. 502.

SHANTUNG: Chefoo (*Forbes*, 398!); KIANGSU: Kiangnan (*Staunton*!); SZECHUEN: North Wushan (*A. Henry*, 7051!); LUCHU ARCHIPELAGO (*Döderlein*, fide *Hackel*). Mus. Brit.; Herb. Kew.

Japan, Assam.

2. **Panicum ambiguum**, *Trin. in Mém. Acad. Ptéersb.* sér. 6, iii. p. 243; *Hook. f. Fl. Brit. Ind.* vii. p. 33.

LUCHU ARCHIPELAGO (*Wright*!). Herb. Kew.

India, Ceylon, Malaya, Pacific Islands.

3. **Panicum amplexicaule**, *Rudge, Pl. Gui.* i. p. 21, t. 27.

*Panicum Myurus*, *H. B. & K. Nov. Gen. & Sp.* i. p. 98; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723; *Hook. f. Fl. Brit. Ind.* vii. p. 39; *Benth. Fl. Austral.* vii. p. 480.

Hymenachne Myuros, *Beauv. Agrost.* p. 165, t. 10. fig. 8; *Steud. Syn. Gram.* p. 101.

Hymenachne amplexicaulis, *Nees, Agrost. Brasil.* p. 276; *Steud. Syn. Gram.* p. 102.

FORMOSA: Takow (*A. Henry*, 1950!). Mus. Brit.; Herb. Kew.

India, Malaya, Tropical Australia, South America.

4. **Panicum brevifolium**, *Linn. Sp. Pl.* p. 59, excl. syn.; *Osbeck, Resa*, p. 215.

*Panicum arborescens*, *Linn. Sp. Pl.* p. 59, in part; ? *Osbeck, l. c.* p. 206.

*Panicum ovalifolium*, *Poir. Encycl.* iv. p. 279; *Benth. Fl. Hongk.* p. 413; *Hook. f. Fl. Brit. Ind.* vii. p. 44.

FORMOSA (*Campbell!*); KWANGTUNG: Sichuan, 40 miles west of Canton (*Parry*, in herb. *Hance*, 9391 in part!) (*Sampson*, 418!), Whampoa (*Hance*, 1006 in part!); HONGKONG (*Hance*, 1006 in part, 9391 in part! *Harland*, 684! in herb. *Hance*, *Wilford*, 251!), Little Hongkong Woods (*Lamont!*); HAINAN (*A. Henry*, 8076!). Mus. Brit.; Herb. Kew.

India, Malaya, Tropical Africa.

5. **Panicum colonum**, *Linn. Syst.* ed. 10, p. 870; *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 232; *Benth. Fl. Hongk.* p. 411; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 223; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 160; *Hook. f. Fl. Brit. Ind.* vii. p. 32.

CHEKIANG: Ningpo (*Swinhoe*, in herb. *Hance!*); HUPEH: Ichang (*A. Henry*, 2769, 4286!), Patung (*A. Henry*, 7313!); HONGKONG: Aberdeen (*Lamont!* *Hance*, 1258!); HAINAN (*A. Henry*, 31!). Mus. Brit.; Herb. Kew.

Widely distributed in warm countries.

6. **Panicum Crus-Galli**, *Linn. Sp. Pl.* p. 56; *Osbeck, Resa*, pp. 48, 215; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Benth. Fl. Hongk.* p. 411; *Franchet*, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266, et *Pl. David.* i. p. 322; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Palibin*, in *Act. Hort. Petrop.* xix. (1901) p. 127, et xxi. (1903) p. 10; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106; *Franch. et Savat. in*

*Enum. Pl. Jap.* ii. p. 160; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643, et sér. 2, iii. (1903) p. 501.

*Panicum Crus-Corvi*, *Linn. Syst. Nat.* ed. 10, p. 870; *Lour. Fl. Cochinch.* p. 46.

*Panicum cæsiuum*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 235.

*Panicum hispidulum*, *Hook. & Arn. l. c.* pp. 235, 273.

*Echinochloa Crus-Galli*, *Beauv. Agrost.* p. 53; *Maxim. Primit. Fl. Amur.* p. 479.

*Oplismenus Crus-Galli*, *Dum. Obs. Gram. Fl. Belg.* p. 138; *Kunth, Enum. Pl.* i. p. 143; *Debeaux, Fl. Tchéfou*, p. 154; *Fl. Tientsin*, p. 46; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 274.

CHINA: without locality (*Fortune*, 123!); NORTH CHINA: without locality (*Daniell! Moellendorf*, in herb. *Hance*, 1549!); EAST COAST (*Home!*); CHIHLI: Jehol (*David*, 2114!); KIANGSU: Shanghai (*Maingay*, 569, 570, 668! *Faber!*); CHEKIANG: Ningpo (*Swinhoe*, in herb. *Hance*, 10234 in part!); FORMOSA (*Oldham*, 601, 602, 603!), Takow (*A. Henry*, 1046, 1796, 1960, 2003!), Bankinsing (*A. Henry*, 841!), South Cape (*Schmüerer*, in coll. *A. Henry*, 276!); HUPEH: Ichang (*A. Henry*, 746, 2341, 4336, 7743!); SHENSI: Mt. Kiansan, Husiencien (*Hugh!*), Kaolinsien, Kusan and Hukiascien (*Hugh!*); NORTH CENTRAL CHINA; Leunteon, Fenghosan (*Hugh!*); KWANGTUNG: Canton, hillside near Sunin (*Lamont!*), Whampoa (*Hance*, 10234 in part! 10235!), Macao (*Calléry*, 355! *Vachell*, 43!), Pakhoi (*Playfair!*); HONGKONG: paddy-fields (*Ford*, 598! *Harland*, 698!); COREA: Port Hamilton (*Oldham*, 928!); HOOPER ISLAND (*Oldham*, 1073!). *Mus. Brit.*; *Herb. Kew.*

A weed in the temperate zones of both hemispheres.

Var. *frumentaceum*, *Trimen, Cat. Ceyl. Pl.* p. 104; *Hook. f. Fl. Brit. Ind.* vii. p. 31; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 127.

*Panicum frumentaceum*, *Roxb. Fl. Ind.* p. 102; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266.

*Panicum stagninum*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 133, non *Retz.*

*Oplismenus frumentaceus*, *Kunth, Rév. Gram.* i. p. 45; *Debeaux, Fl. Tchéfou*, p. 154.

CHIHLI: Peking (*Bretschneider*, 810, 811!); SHINGKING (*Ross*, 33½!); SZECHUEN (*Faber*, 1165!); KWEICHAU: Paitzu-Weining (*Bourne!*). *Herb. Kew.*

Cultivated in hotter parts of Asia and Africa.

7. *Panicum decompositum*, *R. Br. Prodr.* p. 191; *Benth. Fl. Austral.* vii. p. 489.

FORMOSA: sea-face of Apes' Hill, Takow, rare, growing more or less in clusters (*A. Henry*, 1080!), South Cape (*Schmürer*, in coll. *Henry*, 255!). Mus. Brit.; Herb. Kew.

Tropical Australia.

8. *Panicum distachyum*, *Linn. Mant.* p. 183; *Kunth, Enum. Pl.* i. p. 91; *Benth. Fl. Hongk.* p. 410; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723; *Hook. f. Fl. Brit. Ind.* vii. p. 37.

FORMOSA: Kelung (*Makino*, fide *Hackel*); HONGKONG: spontaneous (*Hance*, 1983!). Mus. Brit.; Herb. Kew.

India, Malaya, Australia.

9. *Panicum humile*, *Nees, ex Steud. Syn. Gram.* p. 84; *Hance, in Journ. Bot.* xvi. (1878) p. 233; *Hook. f. Fl. Brit. Ind.* vii. p. 48.

KWANGTUNG: Canton (*Sampson*, in herb. *Hance*, 9896 in part!), Whampoa (*Hance*, 9686 in part!), Pakhoi (*Playfair*, 158!). Mus. Brit.; Herb. Kew.

India, Malaya.

10. *Panicum indicum*, *Linn. Mant.* p. 184; *Steud. Syn. Gram.* p. 84; *Benth. Fl. Hongk.* p. 413; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 644, et sér. 2, iii. (1903) p. 502.

CHINA: without locality (*Staunton!* *Fortune!*); FORMOSA: Bankinsing Mts. (*A. Henry*, 1547!); HUPEH: Ichang (*A. Henry*, 2750, 4312!), Nanto (*A. Henry*, 4423!); HONGKONG (*Hance*, 1072! *Wilford*, 240! *Wright!*). Mus. Brit.; Herb. Kew.

Japan, and tropics of Old World.

Var. *angustum*, *Hook. f. Fl. Brit. Ind.* vii. p. 42.

*Panicum angustum*, *Trin. Ic. Gram.* t. 334.

KWANGTUNG (*Tate!*), Pakhoi (*Playfair*, 141!). Herb. Kew.  
India.

Var. *contractum*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 275; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 49; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 163.

LUCHU ARCHIPELAGO: Amami Osima (*Döderlein*, fide *Hackel*).  
Japan.

I have seen no specimen.

11. *Panicum khasianum*, *Munro, ex Hook. f. Fl. Brit. Ind.* vii. p. 54.

YUNNAN: Mengtze, 5500 ft., in a wood (*A. Henry, 9379!*).  
Herb. Kew.

Himalayas and Khasia Hills.

12. *Panicum maximum*, *Jacq. Ic. Pl. Rar.* i. t. 13; *Steud. Syn. Gram.* p. 72; *Benth. in Hook. Niger Fl.* p. 560; *Hook. f. Fl. Brit. Ind.* vii. p. 49.

HONGKONG: cultivated (*Hance, 716!*). Mus. Brit.

An African grass which is cultivated in many parts of India.

13. *Panicum miliaceum*, *Linn. Sp. Pl.* p. 58; *Steud. Syn. Gram.* p. 77; *Lour. Fl. Cochinch.* p. 59; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 233; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Maxim. Prim. Fl. Amur.* p. 479; *Debeaux, Fl. Shanghai*, p. 69, *Fl. Tchéfou*, p. 155, *et Fl. Tientsin*, p. 48; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266; *Palibin, in Act. Hort. Petrop.* xxi. (1903) p. 10; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 275; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 163; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 644; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106; *Hook. f. Fl. Brit. Ind.* vii. p. 45.

NORTH CHINA (*Daniell!*); CHILHI: Peking, cult. (*Bretschneider, 804!*); SHANTUNG: Chefoo (*Forbes, 1069! Maingay, 80!*), Tsien Mountains (*Faber, 1531!*); HUPEH (*A. Henry, 6611!*); WEST CHINA: Chiensi (*Wilson, 1667!*); COREA (*Oldham, 926!*).  
Mus. Brit.; Herb. Kew.

Widely cultivated in warm countries.

14. *Panicum montanum*, *Roxb. Fl. Ind.* i. p. 313; *Kunth, Enum. Pl.* i. p. 126; *Benth. Fl. Hongk.* p. 412; *Hook. f. Fl. Brit. Ind.* vii. p. 53.

KWANGTUNG: Canton (*Hance, 9905!*); HONGKONG (*Wright!*).  
Mus. Brit.; Herb. Kew.

Hotter hill-districts of India southwards to Ceylon; Malaya.

15. *Panicum nodosum*, *Kunth, Enum. Pl.* i. p. 97; *Benth. Fl. Hongk.* p. 412; *Hook. f. Fl. Brit. Ind.* vii. p. 43.

*Panicum multinode*, *Presl, Rel. Haenk.* i. p. 303; *Steud. Syn. Gram.* p. 59.

HONGKONG (*Wilford, 254!*). Herb. Kew.

India to Malacca and Ceylon; Malaya.

16. *Panicum patens*, *Linn. Sp. Pl.* p. 58; *Kunth, Enum. Pl.* i. p. 126; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 721, 723; *Hook. f. Fl. Brit. Ind.* vii. p. 57.

*Panicum accrescens*, *Trin. Ic. Gram.* t. 88.

*Panicum radicans*, *Retz. Obs.* iv. p. 18.

FOKIEN: Amoy (*Hance*, 1400!); FORMOSA: Bankinsing Mountains (*A. Henry*, 1545!); HONGKONG: Race-course, Happy Valley (*Lamont*!); LUCHU ARCHIPELAGO (*Tashiro*, fide *Hackel*). *Mus. Brit.*; *Herb. Kew.*

India, Malaya, Pacific Islands.

17. *Panicum proliferum*, *Lam. Encycl.* iv. p. 747; *Steud. Syn. Gram.* p. 71; *Hook. f. Fl. Brit. Ind.* vii. p. 50.

*Panicum natans*, *Koen. ex Trin. Gram. Pan.* p. 237.

*Panicum paludosum*, *Roxb. in Hort. Bengal.* p. 8 (nomen), et *Fl. Ind.* i. p. 307; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723.

FORMOSA: Takow, in ditch (*A. Henry*, 1948!). *Mus. Brit.*; *Herb. Kew.*

India, Tropical Africa, Atlantic States of North America, and Central America.

18. *Panicum prostratum*, *Lam. Illustr.* i. p. 171; *Steud. Syn. Gram.* p. 62; *Hook. f. Fl. Brit. Ind.* vii. p. 33; *Benth. Fl. Austral.* vii. p. 476.

FORMOSA: Takow (*A. Henry*, 1020, 1020 A, 1919!), Apes' Hill (*Playfair*, 22!); KWANGTUNG: Whampoa (*Hance*, 5694!). *Mus. Brit.*; *Herb. Kew.*

India, Malaya, Tropical Australia; West Indies.

19. *Panicum psilopodium*, *Trin. Gram. Panic.* p. 217; *Steud. Syn. Gram.* p. 83; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 233; *Hook. f. Fl. Brit. Ind.* vii. p. 46.

CHINA: without locality (*Beechey*!); KWANGTUNG: around Macao (*Millett*, fide *Nees*). *Herb. Kew.*

India to Burma, Malacca, and Ceylon.

20. *Panicum repens*, *Linn. Sp. Pl.* ed. 2, p. 87; *Kunth, Enum. Pl.* i. p. 103; *Benth. Fl. Hongk.* p. 412; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 644; *Hook. f. Fl. Erit. Ind.* vii. p. 49.



*Panicum ischæmoides*, *Retz. Obs.* iv. p. 17; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 233; *Hook. & Arn. l. c.* p. 273.

FORMOSA (*Campbell! Playfair, 172, 195!*), Apes' Hill, sea-face in wet place (*A. Henry, 724!*), Takow, moist place (*A. Henry, 1171!*); HONGKONG (*Hance, 1186! Wright!*); LUCHU ARCHIPELAGO (*Beechey!*). Mus. Brit.; Herb. Kew.

Japan, India, Malaya, and Mediterranean region.

21. *Panicum sarmentosum*, *Roxb. Hort. Bengal.* p. 8 (nomen), *et Fl. Ind.* i. p. 308; *Benth. Fl. Hongk.* p. 412; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 133; *Hook. f. Fl. Brit. Ind.* vii. p. 54.

*Panicum concinnum*, *Nees, in Hook. Kew Journ.* ii. (1850) p. 97; *Steud. Syn. Gram.* p. 78.

*Panicum incomtum*, *Trin. Gram. Pan.* p. 200, *et Sp. Gram. Ic.* t. 232.

*Panicum micrognostum*, *Steud. Syn. Gram.* p. 78.

*Panicum vacillans*, *Steud. l. c.* p. 75.

HONGKONG (*Harland! Wilford, 242! Wright!*); CHINA: without locality (*Seemann, 2397!*). Mus. Brit.; Herb. Kew.

India, Malaya to Philippine Islands, and Borneo.

22. *Panicum trypheron*, *Schult. Mant.* ii. p. 244; *Hook. f. Fl. Brit. Ind.* vii. p. 47; *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 181.

CHIHLI: Salt plains, Tientsin (*Fortune!*); KWANGTUNG: Whampoa and Macao (*Hance, 1944 in part!*); HONGKONG (*Hance, 1944 in part!*). Mus. Brit.; Herb. Kew.

India, Malaya, Tropical Africa.

23. *Panicum villosum*, *Lam. Illustr.* i. p. 173; *Kunth, Enum. Pl.* i. p. 98; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 644; *Hook. f. Fl. Brit. Ind.* vii. p. 34.

*Panicum cocospermum*, *Steud. Syn. Gram.* p. 62.

*Panicum nilagiricum*, *Steud. l. c.*

FOKIEN: near Amoy (*Sampson, in herb. Hance, 8223 in part!*); FORMOSA (*Playfair!*); KWANGTUNG (*Ford, 210!*), Canton and Macao (*Hance, 8223 in part!*), Whampoa (*Wright!*), Wong Mau Islands, Chinese Sea (*Sampson, in herb. Hance, 8223 in part!*). Mus. Brit.; Herb. Kew.

India (hill-regions), Burma, Tonkin.

1. *Axonopus cimicinus*, *Beauv. Agrost.* p. 12; *Hook. f. Fl. Brit. Ind.* vii. p. 64.

*Panicum cimicinum*, *Retz. Obs.* iii. p. 9.

HONGKONG: Happy Valley woods (*Lamont*!). Mus. Brit. India, Malaya.

2. *Axonopus semialatus*, *Hook. f. Fl. Brit. Ind.* vii. p. 64; *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 184.

*Panicum semialatum*, *R. Br. Prodr.* p. 192; *Benth. Fl. Austral.* vii. p. 472.

KWANGTUNG: Canton (*Sampson*, 280!), Pakwan Hills near Canton (*Hance*, 9900!), Swatow (*Perry*!); KWANGTUNG (*A. Henry*, 39!). Mus. Brit.; Herb. Kew.

India, Malaya, Australia, Tropical Africa.

1. *Ichnanthus pallens*, *Munro, in Benth. Fl. Hongk.* p. 414; *Hook. f. Fl. Brit. Ind.* vii. p. 60; *Doell, in Martius, Fl. Bras.* ii. pt. 2, p. 290.

*Panicum pallens*, *Sw. Prodr.* p. 23, et *Fl. Ind. Occ.* i. p. 164; *Kunth, Enum.* i. p. 89.

FORMOSA: Tamsui (*Oldham*, 646!); KWANGTUNG: Canton (*Sampson & Hance*, in herb. *Hance*, 2676!); HONGKONG: Little Hongkong woods (*Lamont*! *Wright*!). Mus. Brit.; Herb. Kew.

Tropics of both worlds.

1. *Setaria Forbesiana*, *Hook. f. Fl. Brit. Ind.* vii. p. 81; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223.

*Setaria macrostachya*, var. *Schultzii*, *Benth. Fl. Austral.* vii. p. 493.

*Panicum Forbesianum*, *Nees, ex Steud. Syn. Gram.* p. 98.

*Chamæraphis setosa*, var. *Schultzii*, *Kuntze, Rev. Gen.* ii. p. 769.

KIANGSU: Shanghai (*Faber*, 281!); HUPEH: Ichang (*A. Henry*, 4246!), Nanto and mountains to north (*A. Henry*, 4642!); CHINA: no locality (*Macartney*!). Mus. Brit.; Herb. Kew.

Northern India.

I have followed Sir Joseph Hooker in regarding this as distinct from the American and Australian *S. macrostachya*, but the two species are very closely allied.

2. *Setaria glauca*, Beauv. *Agrost.* p. 51; *Kunth, Enum. Pl.* i. p. 149; *Maxim. Primit. Fl. Amur.* pp. 330, 479; *Debeaux, Fl. Shanghai*, p. 68, *Fl. Tchéfou*, p. 155, et *Fl. Tientsin*, p. 47; *Franchet, Pl. David.* i. p. 323, ii. p. 142; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 128; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 645; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106.

*Panicum glaucum*, *Linn. Syst. Nat.* ed. 10, p. 870; *Benth. Fl. Hongk.* p. 411; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 234; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 161.

*Panicum penicillatum*, *Nees, Agrost. Bras.* p. 242, et in *Hook. & Arn. Bot. Beechey's Voy.* p. 234; *Hook. & Arn. l. c.* p. 273.

KIANGSU: Shanghai (*Maingay*, 562!); CHUSAN ISLAND: Tin-Hae (*Leclancher*!); FORMOSA (*Campbell! Oldham*, 608!), Takow, in ditch (*A. Henry*, 1109!), Apes' Hill (*A. Henry*, 1904!); HUPEH: Ichang (*A. Henry*, 18, 2263!); SHENSI: Kusan and Kaolinsien (*Hugh*!); SOUTH CHINA: without locality (*Seemann*, 2395!); YUNNAN: Mengtze, 4800 ft. (*A. Henry*, 9613!), and grass mountains 5500 ft. (*A. Henry*, 10994!); KWANGTUNG (*Sampson*, 289!), Lienchau (*Lamont*!), Macao (*Abel*!), Pakhoi (*Playfair*, 149!), Whampoa (*Wright*!); HONGKONG (*Lamont! Hance*, 847! *Wilford*, 262! *Wright*!); HAINAN (*A. Henry*, 8687!). *Mus. Brit.*; *Herb. Kew.*

Japan, and widely distributed in temperate and tropical regions.

3. *Setaria italica*, Beauv. *Agrost.* p. 51; *Kunth, Enum.* i. p. 153; *Debeaux, Fl. Shanghai*, p. 69, *Fl. Tchéfou*, p. 155, et *Fl. Tientsin*, p. 47; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 645.

*Panicum italicum*, *Linn. Sp. Pl.* p. 56; *Lour. Fl. Cochinch.* p. 46; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 234; *Hook. & Arn. l. c.* p. 273; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 162; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70 (var. *germanicum*, *Willd.*).

NORTH CHINA: without locality (monstrous specimen) (*Daniell*!); CHIHLI: Peking (*Bretschneider*, 806-7!); SHING-

KING : Moukden (*Ross!*) ; SHANTUNG (*Maingay, 69!*), Chefoo (*Forbes!*) ; FORMOSA : Bankinsing, cultivated in the mountains (*A. Henry, 571!*) ; EAST COAST (*Home!*) ; HUPEH : Patung (*A. Henry, 885, 2565 A, 5054!*), mountains near Ichang (*A. Henry, 4240!*) ; SZECHUEN (*Faber, 1189!*), Kweichau (*Faber, 1190!*) ; HONGKONG : cultivated (*Hance, 848!*) ; HAINAN (*A. Henry, 8286!*) ; KOREA (*Oldham!*). Mus. Brit. ; Herb. Kew.

Cultivated from Southern Europe to Japan and elsewhere in temperate and tropical countries.

4. *Setaria mauritiana*, *Spreng. Syst. i. p. 305* ; *Rendle, in Cat. Afr. Pl. Welw. ii. p. 187.*

*Panicum palmaefolium*, *Koen. in Naturforsch. xxiii. (1788) p. 208.*

*Panicum plicatum*, *Lam. Illustr. p. 171 (1791)* ; *Kunth, Enum. Pl. i. p. 94* ; *Benth. Fl. Hongk. p. 411* ; *Pilger, in Engl. Bot. Jahrb. xxix. (1900) p. 223* ; *A. Henry, in Trans. Asiat. Soc. Jap. xxiv. Suppl. (1896) p. 106* ; *Hook. f. Fl. Brit. Ind. vii. p. 55.*

*Panicum excurrans*, *Trin. Ic. Gram. t. 89* ; *Kunth, Enum. Pl. i. p. 94* ; *Benth. Fl. Hongk. p. 412* ; *Hack. in Bull. Herb. Boiss. vii. (1899) p. 644.*

*Panicum nervosum*, *Roxb. Fl. Ind. i. p. 314.*

SHINGKING : Chienshan (*Ross, 443!*) ; FOKIEN : Amcy (*Hance, 1397!*) ; FORMOSA (*Oldham, 600!*), Bankinsing Mountains (*A. Henry, 1570!*), South Cape (*A. Henry, 1238!*), Siko Hill (*A. Henry, 1400!*) ; HUPEH : Ichang (*A. Henry, 4264, 4265, 4316, 4320!*), Nanto (*A. Henry, 4450, 4630!*) ; SZECHUEN : Yangtze-kiang, Wushan gorge (*Faber, 1164!*) ; YUNNAN : Mengtze, 4600 — to 5500 ft. (*A. Henry, 9496, 9496 A, 10013, 10013 A, 10972, 10999!*) ; KWANGTUNG : Canton (*Sampson, 632! Hance, 2133 in part!*) ; HONGKONG (*Lamont! Hance 2133 in part! Faurie, 15863!*) ; KWANGSI : Lungchow (*Morse, 368!*). Mus. Brit. ; Herb. Kew.

Japan, India, Malaya, Tropical Africa.

5. *Setaria viridis*, *Beauv. Agrost. p. 51* ; *Kunth, Enum. Pl. i. p. 151* ; *Maxim. Primit. Fl. Amur. pp. 330 & 479* ; *Debeaux, Fl. Tientsin, p. 47* ; *Franchet, Pl. David. ii. p. 323* ; *Pilger, in Engl. Bot. Jahrb. xxix. (1900) p. 223* ; *A. Henry, in Trans. Asiat. Soc. Jap. xxiv. Suppl. (1896) p. 106* ; *Palibin, in Act. Hort. Petrop. xix. (1901) p. 128* ; *Hack. in Bull. Herb. Boiss. vii. (1899) p. 645.*

*Panicum viride*, *Linn. Syst. Nat.* ed. 10, p. 870; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 235; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267.

CHIHLI: Peking (*Bushell!*), Tientsin (*Fortune!*), mountains west of Peking (*Bretschneider*, 815!); SHINGKING: Taliénwan (*Birnie!*); SHANTUNG: Chefoo (*Forbes!*); KIANGSU (*Maingay*, 561, 761!); CHEKIANG: Ningpo (*Swinhoe*, in herb. *Hance!*); FORMOSA: Tamsui (*Oldham*, 607!), Takow (*A. Henry*, 1104!), Apes' Hill (*Playfair*, 37!), South Cape (*Schmüerer*, in coll. *A. Henry*, 969!), Pescadores (*Swinhoe!*); HUPEH: Ichang (*A. Henry*, 191, 2565 A!), Patung (*A. Henry*, 5054 A!); NORTH CENTRAL CHINA: Leunteon (*Hugh!*); SHENSI: Kusan, Kaolin-sien, Paoki, Mt. Kiansan and Mt. Lao-y-san (*Hugh!*); SZECHUEN (*Faber*, 1182!), Lofaushan (*Faber*, 18!); KWANGTUNG: Canton (*Sampson*, 230!); LUCHU ARCHIPELAGO (*Wright!*). *Mus. Brit.*; *Herb. Kew.*

Widely distributed in temperate and subtropical regions of the Old World.

1. *Oplismenus compositus*, *Beauv. Agrost.* p. 54; *Kunth, Enum. Pl.* i. p. 141; *Franchet, Pl. David.* i. p. 323; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 223; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 106; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 721, 723; *Hook. f. Fl. Brit. Ind.* vii. p. 66.

*Oplismenus loliaceus*, *Beauv. l. c.*; *Hack. l. c.* p. 721.

*Panicum compositum*, *Linn. Sp. Pl.* p. 57; *Steud. Syn. Gram.* p. 44; *Benth. Fl. Hongk.* p. 411.

*Orthopogon compositus*, *R. Br. Prodr.* p. 194.

FORMOSA: Bankinsing (*A. Henry*, 1557!), South Cape (*Schmüerer*, in coll. *A. Henry*, 232!); HUPEH (*A. Henry*, 4349!); SHENSI: Paokisan, Mt. Miaowansan (*Hugh!*); KWANGTUNG: Canton (*Lamont! Hance*, 1010 in part!); HONGKONG (*Harland*, 714, in herb. *Hance*, 1010 in part! *Forbes*, 486!); YUNNAN: Mengtze woods, 5500 feet (*A. Henry*, 10971!); LUCHU ARCHIPELAGO (*Tashiro*, fide *Hackel*). *Mus. Brit.*; *Herb. Kew.*

Japan.

Widely distributed in the warmer parts of both Old and New Worlds.

2. *Oplismenus undulatifolius*, Beauv. *Agrost.* p. 54; Kunth, *Enum. Pl.* i. p. 139; Pilger, in *Engl. Bot. Jahrb.* xxix. (1900) p. 223; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 654; Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 645, et sér. 2, iii. (1903) p. 502; Hook. f. *Fl. Brit. Ind.* vii. p. 66.

*Panicum undulatifolium*, Arduino, *Animad. Bot. Spec. alter.* p. 14, t. 4; Maxim. *Prim. Fl. Amur.* p. 479.

CHIHLI: Peking, imperial garden (Williams, in herb. Hance, 13487!), in the mountains (David!); HUPEH: Patung (A. Henry, 4782, 7277!), Fang (A. Henry, 666!). Mus. Brit.; Herb. Kew.

Japan, Southern Europe to India, Tropical Africa.

1. *Pennisetum compressum*, R. Br. *Prodr.* p. 195; Steud. *Syn. Gram.* p. 102.

*Pennisetum chinense*, Nees, ex Steud. *Syn. Gram.* p. 108.

*Pennisetum japonicum*, Trin. in Spreng. *Neue Entdeck.* ii. p. 76, et *Sp. Gram. Ic.* t. 19; Maxim. *Primit. Fl. Amur.* p. 479; Franchet, *Pl. David.* ii. p. 324; Palibin, in *Act. Hort. Petrop.* xix. (1901) p. 128; Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 645, et sér. 2, iii. (1903) p. 502.

*Gymnothrix japonica*, Kunth, *Enum. Pl.* i. p. 158; Benth. *Fl. Hongk.* p. 413; Debeaux, *Fl. Tchéfou*, p. 156; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 164.

? *Alopecurus hordeiformis*, Osbeck, *Resu*, p. 232.

CHIHLI: Peking (Bushell!), mountains west of Peking (Bretschneider, 827!), Tachiaosze (Moellendorf, in herb. Hance, 1270 in part!); SHINGKING: Kaichow to Kinchow (Ross, 292!); SHANSI and SHENSI (Shockley!); KIANGSU: Kianang (Staunton!); CHEKIANG: Chusan Island (no collector), Ningpo (Faber!); EAST COAST (Home!); FOKIEN: Amoy and French Island (Hance, 1270 in part!); HUPEH: Patung (A. Henry, 2540!), Ichang (A. Henry, 97, 2364!); SHENSI: Miaowansan, Paoki, and Kusan (Hugh!); YUNNAN: Mengtze, 4500 feet (A. Henry, 10170!); KWANGTUNG: Canton (Macartney! Sampson, 290!), Lippas Island, Danes' Island, and Macao (Vachell!), Whampoa (Robertson! Lamont!), Pakhoi (Playfair, 154!); HONGKONG (Hance, 1270 in part! Seeman, 233!), Aberdeen (Lamont!); HAINAN (A. Henry, 7950!); COREA (Perry!);

CHINA: without locality (*Bladh! Fortune, 12!*). Mus. Brit; Herb. Kew.

Japan, Burma, Tonkin; Australia.

Var. *viridescens*, *Rendle*.

*Gymnothrix japonica*, var. *viridescens*, *Miq. in Ann. Mus. Bot. Lugd.-Bat. ii. p. 276.*

*Pennisetum japonicum*, var. *viridescens*, *Palibin, in Act. Hort. Petrop. xix. (1901) p. 128.*

COREA: Kyongkwi, Inchon and Soul (*Gottsche, fide Palibin*).  
Japan.

2. *Pennisetum flaccidum*, *Griseb. in Nachr. Ges. Wiss. Götting. ii. (1868) p. 86; Franchet, Pl. David. p. 323; Hook. f. Fl. Brit. Ind. vii. p. 84.*

*Gymnothrix hordeiformis*, *Hance, in Ann. Sci. Nat. sér. 5, v. (1866) p. 250, non Nees.*

CHIHLI: Peking, walls of the town (*Williams, in herb. Hance, 3386 in part!*), Jehol (*David, 2050!*); YUNNAN: Mengtze (*A. Henry, 9619, 9619 A!*); SZECHUEN: Tongolo (*Soulié, 695!*). —  
Mus. Brit.; Herb. Kew.

Himalayas, Central Asia.

3. *Pennisetum typhoideum*, *Rich. in Pers. Syn. i. p. 72; Hook. f. Fl. Brit. Ind. vii. p. 82; Rendle, in Cat. Afr. Pl. Welw. ii. p. 191; Stapf, in Fl. Cap. vii. p. 432.*

*Pennisetum spicatum*, *Roem. & Schult. Syst. ii. p. 499.*

*Holcus spicatus*, *Linn. Syst. ed. 10, p. 1305.*

*Panicum spicatum*, *Roxb. Fl. Ind. i. (1832) p. 283.*

*Penicillaria spicata*, *Willd. Enum. Hort. Berol. p. 1037; Kunth, Enum. i. p. 165.*

CHIHLI: Peking, Imperial garden (*Bretschneider, in herb. Hance, 15910!*). Mus. Brit.

Cultivated in India and Tropical Africa.

1. *Chamæraphis spinescens*, *Poir. Encycl. ii. p. 189; var. Brunoniana, Hook. f. Fl. Brit. Ind. vii. p. 62.*

*Panicum asperum*, *Koen. in Naturf. xxiii. (1788) p. 209.*

*Panicum spinescens*, *R. Br. Prodr. p. 193; Kunth, Enum. Pl. i. p. 131.*

KWANGTUNG: in rather deep ponds around Canton (*Sampson, 367!*). Mus. Brit.; Herb. Kew.

India, Tonkin, Malaya, Australia.

1. *Thysanolaena Agrostis*, Nees, in *Edinb. New Phil. Journ.* xviii. (1835) p. 180; *Hook. f. Fl. Brit. Ind.* vii. p. 61.

*Thysanolaena acarifera*, Arn. et Nees, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 181; *Benth. Fl. Hongk.* p. 417; *Hack. in Bull. Herb. Boiss.* vii. (1889) p. 723.

FORMOSA: Bankinsing (*A. Henry*, 407!); YUNNAN: Red River Valley (*A. Henry*, 10950!); HONGKONG (*Hance*, 1187! *Faurie*, 15862!). Mus. Brit.; Herb. Kew.

Northern India, Malaya, Mauritius.

1. *Thuarea sarmentosa*, Pers. *Syn.* i. p. 110; *Munro*, in *Seemann, Bot. Voy. 'Herald,'* p. 423; *Benth. Fl. Hongk.* p. 415; *Hook. f. Fl. Brit. Ind.* vii. p. 91.

*Ornithocephalochloa arenicola*, Kurz, in *Journ. Bot.* xiii. (1875) p. 332, t. 171.

HONGKONG (*Hance*, 10230!); SOUTH CHINA SEA: Pulo Redang (*Jensen!*), Pratas Island (*Wilford*, 451!). Mus. Brit.; Herb. Kew.

Ceylon, Cochin China, Malaya, North Australia, Pacific Islands.

1. *Spinifex squarrosus*, Linn. *Mant.* p. 300; *Steud. Syn. Gram.* p. 113; *Benth. Fl. Hongk.* p. 415; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 50; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Hook. f. Fl. Brit. Ind.* vii. p. 63.

FORMOSA (*Oldham*, 606!), Tamsui sand-hills (*Hancock*, 18!); KWANGTUNG: Macao (*Abel!* *Vachell!* *Tate!* *Hance*, 865 in part!); HONGKONG: sandy sea-shore (*Hance*, 865 in part! *Forbes*, 610!). Mus. Brit.; Herb. Kew.

India, Ceylon, Malaya.

1. *Phænospërma globosa*, *Munro*, ex *Benth. in Journ. Linn. Soc., Bot.* xix. (1881) p. 59; *Franchet, Pl. David.* i. p. 326; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222.

CHINA: without locality (*Faber!*); KIANGSI: Kiukiang (*Shearer!*); HUPEH: Ichang (*A. Henry*, 626, 3966, 3968!), Western Hupeh (*Wilson*, 530!), Nanto and mountains to north (*A. Henry*, 1943, 2073!), Patung (*A. Henry*, 4743!); SZECHUEN: near Tachienlu (*Pratt*, 426!) (*A. Henry*, 8911!); YUNNAN: Mengtze, wooded cliffs, 6000 ft. (*A. Henry*, 11299!). Mus. Brit.; Herb. Kew.



1. *Arundinella anomala*, *Steud. Syn. Gram.* p. 116; *Debeaux, Fl. Tchéfou*, p. 156; *Franchet, in Mém. Soc. Sci. Nat. Cherb.* xxiv. (1884) p. 267, et *Pl. David.* i. p. 325; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 127; *Pilger, in Engl. Jahrb.* xxix. (1900) p. 222; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 276; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 165; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 643, et sér. 2, iii. (1903) p. 501.

*Panicum mandshuricum*, *Maxim. Prim. Fl. Amur.* p. 328; *Hance, in Journ. Bot.* vii. (1869) p. 41; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 338.

*Panicum Williamsii*, *Hance, in Ann. Sci. Nat.* sér. 5, v. (1866) p. 250.

*Chalynochlamis anomala*, *Franchet*, apud *Debeaux, Fl. Tientsin*, p. 71.

CHIHLI: Peking (*Bushell! Williams, in herb. Hance, 11458 in part! 12864! Bretschneider, 828-829!*), Jehol (*David, 1901!*); SHINGKING (*Ross, 81, 469!*), Kaichow to Kinchow (*Ross, 255, 275, 290!*), Kaichow Hills (*Ross, 320!*), Changpeishan (*James!*); SHANTUNG (*Maingay, 74!*), Chefoo (*Forbes!*); KIANGSU: Kianang (*Staunton!*); CHEKIANG: Ningpo Mts. to Chefoo Mts. (*Faber, 297!*), Ningpo (*Swinhoe, in herb. Hance, 11458 in part!*); KIANGSI: Kiukiang (*Shearer!*); HUPEH: Patung (*A. Henry, 4845!*), Ichang (*A. Henry, 431, 2265, 2272, 4284, 4290!*); SHENSI: Kusan (*Hugh!*), Kisanshien (*Hugh!*); ? HUNAN: Sungatechi (*Maack!*); KWANGTUNG: Canton (*Sampson, in herb. Hance, 12928!*); COREA: Soul (*Sontag!*). *Mus. Brit.; Herb. Kew.*

Japan, North Asia.

Var. *depauperata*, *Rendle*.

*Herba* quam in specie minor, culmis gracilibus, 16 poll. vel brevioribus. *Folia* brevia, angusta, convoluta, demum setacea, 1½-2½ poll. longa, ad ½ lin. diametro. *Panicula* angusta, 3-3½ poll. longa, 3-6 lin. lata; ramis paucis, brevibus, appressis, infimis ad 1 poll. longis.

HUPEH: Ichang (*A. Henry, 1059, 4290!*). *Herb. Kew.*

2. *Arundinella nepalensis*, *Trin. Gram. Pan.* p. 62, et *Sp. Gram. Ic.* t. 268; *Benth. Fl. Hongk.* p. 416.

*Arundinella miliacea*, *Nees, in Hook. Kew Journ.* ii. (1850) p. 102; *Munro, in Seemann, Bot. Voy. 'Herald,'* p. 423.

? *Arundinella glabra*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 237 (plant not seen).

HUPEH: Ichang (*A. Henry*, 2264!); YUNNAN: Mengtze, 4600 ft. (*A. Henry*, 9139!); KWANGTUNG (*Ford*, 214!), Canton (*Sampson*, 1049!); HONGKONG (*Hance*, 8111 in part! *Faurie*, 15852!); LAPPAS ISLAND (no collector). Mus. Brit.; Herb. Kew.

India, Malaya, Australia.

Sir Joseph Hooker (*Fl. Brit. Ind.* vii. p. 73) unites this species with *A. brasiliensis*, Raddi. The spikelets in Raddi's type at the British Museum are, however, conspicuously smaller than anything I have seen in the Asiatic plants, and I prefer to regard the latter as forming a distinct species.

3. *Arundinella setosa*, *Trin. Gram. Pan.* p. 63; *Benth. Fl. Hongk.* p. 416; *Hook. f. Fl. Brit. Ind.* vii. p. 70; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723.

*Arundinella stricta*, *Nees, in Hook. Kew Journ.* ii. (1850) p. 102.

*Danthonia neuroelytrum*, *Steud. Syn. Gram.* p. 245.

*Milium cimicinoïdes*, *Roxb. ex Hook. f. Fl. Brit. Ind.* vii. p. 70.

CHINA: without locality (*Fortune*, 5!); FOKIEN: Amoy (*Hance*, 1282 in part!); FORMOSA: Takow (*A. Henry*, 771!), Apes' Hill (*A. Henry*, 1033, 1107!); YUNNAN: Mengtze (*A. Henry*, 10518, 10518 A!), Szemao Mts., 5000 ft. (*A. Henry*, 13193!); KWANGTUNG (*Ford*, 211!), Canton (*Sampson*, 1167!); HONGKONG (*Hance*, 1282 in part! *Lamont*! *Wright*!). Mus. Brit.; Herb. Kew.

India, Ceylon, Philippines.

4. *Arundinella sinensis*, *Rendle*, n. sp.

*Herba* perennis, rhizomate duro, 1½–2 lin. crasso, culmis simplicibus, 2-pedalibus, nodis glabris, internodiis 6, inferioribus 1–1¼ lin. crassis. *Foliorum* vaginæ cylindricæ, internodis haud æquantes, margine et ore pilosæ, aliter plus minus scabride-pilosæ vel superiores glabrescentes; ligula e pilis 1 lin. longis constituta; lamina linearis, basi paullo angustata et superne in apice acuto gradatim attenuata, rigidula, cum faciebus striatis et margine scabrida, 6–14 poll. longa, 2–4½ lin. lata. *Panicula*, basi nondum evoluta, pedalis, rhacheos scabride-angulati internodiis inferioribus circa 1 poll. longis, ramis ad nodum quemque 1–3, fastigiatis, ad 8 poll. longis, complanatis, scabridis, laxè spiculatis; spiculæ 3 lin. longæ, anguste lanceolatæ, acuminatæ, glabræ, alter pedicellum scabridulum apice sparse barbatur dimidia,

alter eo plus triplo excedens. *Gluma* i. lanceolata acuminata, cum nervis tribus validis, quorum mediano scabridulo, et tribus incompletis instructa, 3 lin. longa; gl. ii. primo æquans et ei similis, 5-nervis; gl. iii. ovata, 3-nervis cum nervulis 4 debilis superadditis,  $1\frac{2}{5}$  lin. longa, florem ♂ celans; gl. iv. oblonga, 1 lin. longa, apice in setam debilem imperfectam 3 lin. longam cum setulis lateralibus duabus prolongata.

KWANGTUNG (*Ford*, 201, 212!), Lofaushan (*Ford*, 125!).  
Herb. Kew.

Approaches *A. setosa*, from which it is distinguished by its larger panicle and subequal multinerved barren glumes.

#### Tribe ZOYSIÆ.

1. *Tragus racemosus*, *Scop. Introd.* p. 73; *Franchet, Pl. David.* i. p. 327; *Hook. f. Fl. Brit. Ind.* vii. p. 97.

*Tragus tcheliensis*, *Debeaux, Fl. Tientsin*, p. 48.

*Lappago racemosa*, *Honck. Syn.* i. p. 440; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70.

*Cenchrus racemosus*, *Linn. Sp. Pl.* p. 1049; *Staunton, Embass. Chin.* ii. p. 346.

CHIHLI: Pechili (*Staunton!*), Jehol (*David*, 2069!), Peking (*Williams*, in herb. *Hance*, 10522! *Bushell!*); SHENSI: Kaolin-sien (*Hugh!*). Mus. Brit.; Herb. Kew.

Warmer parts of both hemispheres.

1. *Perotis latifolia*, *Ait. Hort. Kew.* i. p. 85; *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 202; *Hook. f. Fl. Brit. Ind.* vii. p. 98; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107.

*Perotis longiflora*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 247; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 133.

*Perotis patula*, *Nees, ex Steud. Nom.* ed. 2, ii. p. 306.

*Saccharum spicatum*, *Linn. Sp. Pl.* p. 54.

*Alopecurus hordeiformis*, *Lour. Fl. Cochinch.* p. 60.

FOKIEN: Amoy (*Hance*, 1384 in part!); FORMOSA: Takow spit (*A. Henry*, 1043, 1110!); KWANGTUNG: Macao (*Hance*, 1384 in part! *Vachell*, 38!), Pakhoi (*Playfair*, 151!); HONG-KONG (*Hance*, 1384 in part! *Wright!*); CHINA: without locality (*Fortune*, 89! *Meyen!*). Mus. Brit.; Herb. Kew.

Tropical Asia and Africa.

1. *Zoysia pungens*, Willd. in *Ges. Naturf. Fr. N. Schrift.* iii. (1801) p. 441; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 271; A. Henry, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; Palibin, in *Act. Hort. Petrop.* xix. (1901) p. 128; Hook. f. *Fl. Brit. Ind.* vii. p. 99.

SHINGKING: Taliénwan (*Birnie!*); SHANTUNG: Chefoo (*Perry!*); FORMOSA: Takow (*A. Henry*, 1798, 1798 A, 1906!); COREA: East (*Perry!*), Port Chusan (*Wilford*, 942!); COREAN ARCHIPELAGO (*Oldham*, 951, 1015!); LUCHU ARCHIPELAGO (*Wright!*). Mus. Brit.; Herb. Kew.

Tropical Asia, Australia, Mascarene Islands.

2. *Zoysia sinica*, Hance, in *Journ. Bot.* vii. (1869) p. 168, et in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 134.

*Zoysia pungens*, Munro, in *Seemann, Bot. Voy. 'Herald,'* p. 424, non Willd.; Benth. *Fl. Hongk.* p. 418.

*Zoysia macrostachya*, Franch. et Savat. *Enum. Pl. Jap.* ii. p. 608; Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 642.

KIANGSI (*Faber!*); FOKIEN: Amoy (*Sampson*, in herb. Hance, 10155 in part! *De Grijs*, *ibid.*!); KWANGTUNG: Pakhoi (*Playfair*, 142!); HONGKONG (*Hance*, 10155 in part!). Mus. Brit.; Herb. Kew.

Japan.

#### Tribe ORYZEÆ.

1. *Oryza sativa*, Linn. *Sp. Pl.* p. 333; Steud. *Syn. Gram.* p. 3; Osbeck, *Resa*, p. 218; Thunb. *Fl. Japon.* p. 147; Lour. *Fl. Cochinch.* p. 214; Franch. et Savat. *Enum. Pl. Japon.* ii. p. 155; Debeaux, *Fl. Shanghai*, p. 67; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 266, et *Pl. David.* i. p. 324; Hack. *Bull. Herb. Boiss.* vii. (1899) p. 645; Palibin, in *Act. Hort. Petrop.* xix. (1901) p. 128; A. Henry, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107.

*Oryza communissima*, *O. præcox*, *O. montana*, *O. glutinosa*, Lour. *Fl. Cochinch.* p. 215.

NORTH CHINA: without locality, cult. (*Daniell!*); SHANTUNG (*Maingay*, 68!); FOKIEN: Amoy (*Hance!*); FORMOSA (*A. Henry*); HONGKONG: cult. (*Hance*, 706! *Urquhart*, 243!). Mus. Brit.; Herb. Kew.

Tropical Asia.

1. *Zizania aquatica*, *Linn. Sp. Pl.* p. 991 (excluding reference to *Sloane, Jam. & Hist.*); *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 502.

*Hydropyrum latifolium*, *Griseb. in Ledeb. Fl. Ross.* iv. p. 466; *Turcz. in Bull. Soc. Nat. Mosc.* xxix. (1856) pt. 1, p. 2; *Maxim. Prim. Fl. Amur.* p. 479; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 156; *Hack. Bull. Herb. Boiss.* vii. (1899) p. 646.

CHIHLI: Peking (*Swinhoe, in herb. Hance, 6496 in part!*); KWANGTUNG: Canton, cult. (*Hance, 6496 in part!*). Mus. Brit.

Japan, Tonkin; Eastern North America.

1. *Leersia hexandra*, *Sw. Prodr.* p. 21; *Hook. f. Fl. Brit. Ind.* vii. p. 94; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 646, 723; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 135.

KIANGSU: Kianang (*Staunton!*); FOKIEN: Amoy (*Hance, 1383!*); FORMOSA: Takow (*A. Henry, 1170, 2034!*); HUPEH: Ichang (*A. Henry, 4249, 2783!*); KWANGTUNG: Canton, edges of rice-fields (*Sampson, 312!*). Mus. Brit.; Herb. Kew.

Widely distributed in the tropics.

#### Tribe MAYDEÆ.

1. *Coix Lachryma-Jobi*, *Linn. Sp. Pl.* p. 972; *Hook. f. Fl. Brit. Ind.* vii. p. 100.

*Coix agrestis*, *Lour. Fl. Cochinch.* p. 551; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 273; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 638.

*Coix chinensis*, *Tod. Ind. Sem. Hort. Bot. Panorm.* 1861, p. 5, et *Fl. Sic. Exsicc.* no. 918.

*Coix Lachryma*, *Linn. Syst.* ed. 10, p. 1261; *Franchet, in Mém. Soc. Sci. Nat. Cherb.* xxiv. (1884) p. 266; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Lour. Fl. Cochinch.* p. 551; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 156; *Thunb. Fl. Jap.* p. 37; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 638.

CHIHLI: Peking (*Bretschneider, 798!*); KIANGSI: Kiukiang (*Reid, 45!*); FORMOSA: Bankinsing (*A. Henry, 51!*); HUPEH: Nanto and mountains to north (*A. Henry, 2635 A!*); YUNNAN (*A. Henry, 10137!*); KWANGTUNG (*Ford, 195!*), mainland

opposite Hongkong (*Harland*, in herb. *Hance*, 402!), Macao (*Calléry*, 496!). Mus. Brit.; Herb. Kew.

Warmer parts of the world.

1. *Zea Mays*, *Linn. Sp. Pl.* p. 971; *Debeaux*, *Fl. Tientsin*, p. 46; *Hook. & Arn. Bot. Beechey's Voy.* p. 347; *Palibin*, in *Act. Hort. Petrop.* xix. (1901) p. 129; *Lour. Fl. Cochinch.* p. 550; *Thunb. Fl. Jap.* p. 37; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 156; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Hook. f. Fl. Brit. Ind.* vii. p. 102.

CHEKIANG: Ningpo Mts. (*Faber*, 299!); SZECHUEN (*Faber*, 1174!); HONGKONG: cultivated (*Hance*, 1333!). Mus. Brit.; Herb. Kew.

Widely cultivated.

#### Tribe ANDROPOGONEÆ.

1. *Imperata arundinacea*, *Cyrill.*, var. *Koenigii*, *Hack. Mon. Androp.* p. 94, et in *Engl. Bot. Jahrb.* vi. (1884) p. 50, et in *Bull. Herb. Boiss.* vii. (1899) p. 639, et sér. 2, iii. (1903) p. 501.

*Imperata arundinacea*, *Cyrill. Pl. Rar.* ii. p. 27, t. 11, in part; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 72; *Maxim. Prim. Fl. Amur.* p. 479; *Franchet, Pl. David.* i. p. 327; *A. Henry*, in *Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 107; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 221; *Palibin*, in *Act. Hort. Petrop.* xix. (1901) p. 129; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 188; *Hook. f. Fl. Brit. Ind.* vii. p. 106.

*Imperata arundinacea*, var. *pedicellata*, *Debeaux, Fl. Tchéfou*, p. 165; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272.

*Imperata Koenigii*, *Beauv. Agrost.* p. 165; *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* pp. 240, 273, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 183.

*Imperata pedicellata*, *Steud. in Flora*, xxix. (1846) p. 22.

*Saccharum spicatum*, *Thunb. Fl. Jap.* p. 42; *Lour. Fl. Cochinch.* p. 53.

CHIHLI: Peking (*Bretschneider*, 824!); EAST COAST (*Home*!); SHANTUNG: Chefoo (*Carmichael*, in herb. *Forbes*!); KIANGSI: Shanghai (*Maingay*, 374, 564! *Forbes*!); FORMOSA (*Oldham*, 619!), Takow (*Playfair*, 100!), South Cape (*Schmürer*, in coll. *A. Henry*, 955!); HUPEH: Ichang (*A. Henry*, 1136!); SHENSI: Kusan (*Hugh*!), Kisansien (*Hugh*!); SZECHUEN: Chungching (*Bourne*, 116!); YUNNAN: Mengtze (*A. Henry*, 7506!); KWANG-TUNG: Ookaisa (*Tate*!); HONGKONG (*Hance*, 577! *Lamont*!);

*Wilford*, 244!); COREA: Chemulpo (*Carles*, 43! *Perry*!); LUCHU ARCHIPELAGO (*Döderlein*, fide *Hackel*). *Mus. Brit.*; *Herb. Kew*.

The species is generally distributed throughout the warmer parts of the earth.

1. *Miscanthus japonicus*, *Anderss. in Öfvers. Vetensk.-Akad. Förhandl.* xii. (1855) p. 166; *Hack. Mon. Androp.* p. 107, et in *Bull. Herb. Boiss.* vii. (1899) p. 639; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 221.

*Saccharum polydactylon*, var. *α*, *Thunb. Fl. Jap.* p. 43.

*Saccharum japonicum*, *Thunb. in Trans. Linn. Soc.* ii. (1794) p. 328, in part.

*Eulalia densa*, *Munro, in Seem. Bot. Voy. 'Herald,'* p. 424.

KIANGSU: Kianang (*Staunton*!), Shanghai (*Maingay*, 575!); Kiangsi: Kiukiang (*Forbes*, 74!); FORMOSA: Takow (*A. Henry*, 1823, 1823 A, 1690, 1692, 1693!), Apes' Hill (*A. Henry*, 1689, 1691!), Kelung (*Ford*, 68!), Tamsui (*Walters*, 45!), South Cape (*A. Henry*, 1332!); KWANGTUNG: Whampoa (*Hance*, 485 in part!); HONGKONG (*Hance*, 485 in part! 557! *Champion*! *Ford*, 596!). *Mus. Brit.*; *Herb. Kew*.

Japan, Malaya, and Pacific Islands.

It seems impossible to distinguish between extreme forms of this species and of *M. sinensis*, *Anderss.*

2. *Miscanthus nepalensis*, *Hack. Mon. Androp.* p. 104; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 221; *Hook. f. Fl. Brit. Ind.* vii. p. 107.

SZECHUEN: Mt. Omei, 4200 feet (*Faber*, 1152!), Moupin (*David*!); LAPPAS ISLAND (herb. *Munro*!). *Herb. Kew*.

Mountains of Northern India.

3. *Miscanthus sacchariflorus*, *Benth. & Hook. f. Gen. Pl.* iii. 2, p. 1125; *Hack. Mon. Androp.* p. 102, et in *Bull. Herb. Boiss.* vii. (1899) p. 639; *Franchet, Pl. David.* i. p. 327; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 243.

*Imperata sacchariflora*, *Maxim. Prim. Fl. Amur.* p. 331.

? *Imperata eulalioides*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 289; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 188.

CHIHLI: Peking (*Williams*, in herb. *Hance*, 11451! *Bretschneider*, 823!), Jehol (*David*!); KIANGSU: Shanghai (*Maingay*, 577!), Kianang (*Staunton*!); HUPEH: Patung (*A. Henry*, 5123!), Changyang (*A. Henry*, 7507!); SHENSI: Kaolinsien (*Hugh*!), Mt. Kiansan, Husiencien (*Hugh*!), Mt. Laoyan, Tciuzsien (*Hugh*!). Mus. Brit.; Herb. Kew.

North Asia, Japan.

4. *Miscanthus sinensis*, *Anderss.* in *Öfvers. Vetensk.-Akad. Förhandl.* xii. (1855) p. 166; *Hack. Mon. Androp.* p. 105, et in *Bull. Herb. Boiss.* vii. (1889) p. 639, et sér. 2, iii. (1903) p. 501.

*Saccharum polydactylon*, var.  $\beta$ , *Thunb. Fl. Jap.* p. 43.

*Saccharum japonicum*, *Thunb.* in *Trans. Linn. Soc.* ii. (1794) p. 328, in part.

*Erianthus japonicus*, *Beauv. Agrost.* p. 14, in part; *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 242, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 184.

*Eulalia densa*, *Baker & Moore*, in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 388, non *Munro*.

*Eulalia japonica*, *Trin.* in *Mém. Ac. Sci. Pétersb.* sér. 6, ii. p. 333, in part; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 189; *Franchet*, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272.

*Ripidium japonicum*, *Trin. Fund. Agrost.* p. 169, in part.

SHINGKING (*Ross*, 44!); FORMOSA: Takow (*A. Henry*, 1173, 1173 A, 1175, 1185!); HUPEH: Patung (*A. Henry*, 5065!), Ichang (*A. Henry*, 93!); SHENSI: Kusan (*Hugh*!), Mt. Kiansan, Husiencien (*Hugh*!); SZECHUEN: Tachou (*Rockhill*!); YUNNAN: Mengtze (*A. Henry*, 10014!); KWANGTUNG: Macao, Canton (*Sampson*, 973!); HONGKONG (*Lamont*! *Reeves*! *Urquhart*, 251! *Hind*!); COREA (*Perry*!); LUCHU ARCHIPELAGO (*Wright*!); CHINA: without locality (*Macartney*!). Mus. Brit.; Herb. Kew.

Japan, Tonkin, Celebes, Borneo, Philippines.

Var. *purpurascens*, *Rendle*.

*Miscanthus purpurascens*, *Anderss. l. c.* p. 167; *Hack. Mon. Androp.* p. 106; *Palibin*, in *Act. Hort. Petrop.* xix. (1901) p. 129; *Komarow*, *ibid.* xx. (1901) p. 244.

SHANTUNG (*Maingay*, 77!), Chefoo (*Forbes*!). Herb. Kew.

Japan.

I regard this merely as a variety of *M. sinensis*. Hackel himself says (*l. c.* p. 107): "cui fortasse melius subspeciei numero subjiciendus."



1. *Saccharum arundinaceum*, *Retz. Obs.* iv. p. 14; *Hack. Monogr. Androp.* p. 117.

*Saccharum procerum*, *Roxb. Fl. Ind.* i. p. 243; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 134.

*Saccharum Sara*, *Roxb. l. c.* p. 244; *Hook. & Arn. in Bot. Beechey's Voy.* p. 240.

CHEKIANG (*Staunton!*); FOKIEN: Amoy (*Hance, 1406 in part!*); HUPEH (*A. Henry, 4227!*); KWANGTUNG (*Staunton!*), Whampoa (*Hance, 1406 in part!*), Pakhoi (*Playfair, 160!*); CHINA: without locality (*Fortune, 9!*). Mus. Brit.; Herb. Kew.  
India, Ceylon, China.

2. *Saccharum Narenga*, *Ham. in Wall. List*, no. 8856; *Hack. Mon. Androp.* p. 119; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222; *Hook. f. Fl. Brit. Ind.* vii. p. 120.

*Saccharum porphyrocomum*, *Hack. Mon. Androp.* p. 120.

*Eriochrysis Narenga*, *Nees, ex Steud. Syn. Gram.* p. 411.

*Eriochrysis porphyrocoma*, *Hance, in Journ. Bot.* xiv. (1876) p. 294.

FORMOSA: Takow Plain and Lake Hill (*A. Henry, 2036!*); HUPEH: Ichang (*A. Henry, 154!*); KWANGTUNG: Canton (*Staunton! Lamont! Nevin, in herb. Hance, 19285 in part!*), by River Lienchau (*Galbraith, in herb. Hance, 19285 in part!*). Mus. Brit.; Herb. Kew.

India, Burma.

3. *Saccharum officinarum*, *Linn. Sp. Pl.* p. 54; *Hack. Monogr. Androp.* p. 111; *Benth. Fl. Hongk.* p. 419; *Lour. Fl. Cochinch.* p. 52; *Thunb. Fl. Japon.* p. 42; *Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107.

*Saccharum sinense*, *Roxb. Fl. Ind.* i. p. 244; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 241.

FORMOSA: Bankinsing Plain, "the variety which is generally cultivated in South Formosa" (*A. Henry, 1688!*); KWANGTUNG: Canton (*Macartney!*), Whampoa (*Hance, 6556!*); HONGKONG (*Harland, 683!*). Mus. Brit.; Herb. Kew.

Widely cultivated in the tropics.

4. *Saccharum spontaneum*, *Linn. Mant.* ii. p. 183; *Lour. Fl. Cochinch.* p. 52; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222.

Subsp.  $\alpha$ . *indicum*, *Hack. Mon. Androp.* p. 114.

FORMOSA: Takow (*A. Henry, 1077, 1077 A!*); HUPEH: Ichang (*A. Henry, 34, 2266, 4222!*); KWANGTUNG: Canton (*Galbraith, in*

herb. *Hance*, 19304! *Lamont*!); CHINA: without locality (*Fortune*, 6!). Mus. Brit.; Herb. Kew.

India, Central Asia, Malaya.

1. *Erianthus formosanus*, *Stapf*, in *Kew Bull.* 1898, p. 228.

*Erianthus fastigiatus*, *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 107.

FORMOSA: Apes' Hill (*Playfair*, 314!). Herb. Kew.

2. *Erianthus fulvus*, *Nees*, ex *Steud. Syn. Gram.* p. 409; *Hack. Mon. Androp.* p. 147; *Hook. f. Fl. Brit. Ind.* vii. p. 123.

*Saccharum rufipilum*, *Steud. l. c.*

HUPEH: Patung (*A. Henry*, 5115!); SHENSI: Paokisan, near Mt. Miaowansan (*Hugh*!); YUNNAN: Mengtze, grass mountains, 4600–6000 feet (*A. Henry*, 9818, 9818 A, 10325!). Mus. Brit.; Herb. Kew.

Hills of Northern India.

3. *Erianthus Hookeri*, *Hack. Mon. Androp.* p. 142; *Hook. f. Fl. Brit. Ind.* vii. p. 125.

YUNNAN: without locality (*Anderson*!), Mengtze, 4600 feet (*A. Henry*, 9660!). Herb. Kew.

Himalayas.

4. *Erianthus pollinioides*, *Rendle*, n. sp.

*Culmi* erecti, arundinacei,  $4\frac{1}{2}$  ped. alti, basibus 2 lin. crassi, teretes, glabri, simplices, foliosi, basibus cum vaginis fibrosis marcidis vestiti; internodia 7. *Foliorum* vaginæ glabræ, stramineæ, suprema excepta internodia excedentes; ligula brevis, membranacea, vix semilinea longa, pilosa; lamina elongata, anguste linearis, ad apicem acutum attenuata, rigidula, usque ad 3 ped. longa, medio  $2\frac{1}{2}$  lin. lata, cum nervo mediano valido albido lineata. *Panicula* eximie purpurascens, subcorymbosa, 8 poll. longa, superne 5 poll. lata; rhachis communis 4 poll. longa; racemi ad 30, 5 poll. longi vel breviores, sessiles, erecto-patuli, graciles, dense et longe purpureo-pilosi; articuli pedicellis æquantes sed magis robusti,  $\frac{1}{10}$  poll. longi, velut pedicellis cum pilis purpureis ad 3 lin. longis dense vestiti. *Spicula sessilis* uniflora, lanceolata, læte brunnea,  $1\frac{1}{2}$  lin. longa; gluma i. cum callo brevi ( $\frac{1}{8}$  lin.), lanceolata, apice anguste truncata, cum nervis utrinque marginalibus binis, dorso pilis purpureis plus duplo longioribus

vestita; gl. ii. primæ subæquans sed paullo angustior et superne magis membranacea, cum carina et nervis lateralibus duobus; gl. iii. paullo brevior, anguste lanceolata, hyalina, supra medium ciliata cum marginibus involutis, vacua; gl. iv. minor,  $\frac{4}{5}$  lin. longa, lineari-oblonga, nervo excepto hyalina, apice integro, cum arista simplici, debili,  $3\frac{1}{2}$  lin. longa, instructa. *Spicula pedicellata sessili similis, vix  $1\frac{1}{2}$  lin. longa.*

FORMOSA: Takow (*A. Henry*, 1143!). Mus. Brit.; Herb. Kew.

A well-marked species near the Indian *E. fastigiatus*, but distinguished by its larger, gracefully spreading, purple-haired racemes, and the hairs on the spikelet more than twice as long as the glume.

1. *Spodiopogon cotulifer*, *Hack. Mon. Androp.* p. 187, et in *Bull. Herb. Boiss.* vii. (1899) p. 641, et sér. 2, iii. (1903) p. 501; *Hook. f. Fl. Brit. Ind.* vii. p. 108; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 129.

*Andropogon cotuliferum*, *Thunb. Fl. Jap.* p. 41.

*Eccoilopus andropogonoides*, *Steud. Syn. Gram.* p. 124.

*Eulalia cotulifera*, *Munro, ex Miq. Ann. Mus. Bot. Lugd.-Bat.* ii. p. 289; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 189.

CHEKIANG: Ningpo Mountains (*Faber*, 3, 1734!); FORMOSA: Takow (*A. Henry*, 1045!), Apes' Hill (*Playfair*, 240, 309! *A. Henry*, 1022!), Bankinsing Mountains (*A. Henry*, 1558!); LUCHU ARCHIPELAGO (*Wright*!). Mus. Brit.; Herb. Kew.

Japan; Mountains of North India.

2. *Spodiopogon formosanus*, *Rendle*, n. sp.

*Planta* perennis; culmus erectus, subvalidus, subcompressus, in parte inferiore unilateraliter sulcatus, basi  $1\frac{2}{3}$  lin. crassus. *Folii* vagina glabra, membranacea, ad internodium applicita; ligula brevis, pilosa; lamina anguste lanceolata, acuta, plana, glabra, 8 poll. longa,  $\frac{1}{2}$  poll. lata. *Panicula* e vagina folii supremi breviter (c. 1 poll.) exserta, subovata, 7 poll. longa, ad 2 poll. lata; rhachis flexuosa; rami ad nodos fasciculati, flexuose ascendentopatuli, ad nodum infimum usque ad  $3\frac{1}{2}$  poll. longi, in parte inferiore, interdum usque ad medium, nudi, superne breviter ramulosi et densius spiculiferi; spiculæ sæpissime pedicellatæ, pedicelli glabri, inæquales, quam spiculis sæpius breviores; spiculæ bilineales vel paullo longiores, lanceolatæ, sparsius pilosæ.

*Gluma* i. ovata, apice breviter mucronata, margine ciliata, 9-nervis, ad  $2\frac{1}{3}$  lin. longa; gl. ii. 7-nervis, primæ subæquilonga sed latior, marginibus subhyalinis involutis, apice mucronata; gl. iii. hyalina, enervis, late elliptica, superne rotundata et ciliolata,  $1\frac{1}{2}$  lin. longa, cum palea specie vacua subæquilonga sed angustiore; gl. iv. oblonga, hyalina, vix 2 lin. longa, apice bifida, arista breve imperfecta, ad  $2\frac{1}{2}$  lin. longa sed sæpe vix e spicula exserta; palea quam gluma paullo brevior florem ♂ celans; antheræ rubro-brunneæ,  $1\frac{1}{3}$  lin. longæ.

FORMOSA: planted by savages in mountains near Bankinsing, the grain good to eat (*A. Henry*, 76!). Herb. Kew.

Described from a single specimen which includes the base of the stem and the upper portion bearing a single leaf and the panicle. The plant is evidently closely allied to *S. sibiricus*, in some respects suggesting a cultivated form of that species. It differs in its more open panicle, the almost complete absence of sessile spikelets, the less hairy character of the spikelets, the almost complete disappearance of the awn on the fertile glume, and the shape of the glume of the empty (or perhaps male) floret.

### 3. *Spodiopogon sagittifolia*, Rendle, n. sp.

*Planta* perennis; culmus basi innovationibus obsessus, erectus, teres, glaberrimus, ad  $3\frac{1}{2}$ -pedalis altus, basi vix 1 lin. diametro excedens, 3-4-nodus. *Foliorum* vaginae glabræ, ore pilosæ, infima excepta internodiis breviores; ligula ad 3 lin. longa, scariosa, obtusa, vel apice fimbriata, dorso pilosa; lamina petiolata; petiolus tenuis, in facie superiore complanatus, in foliis inferioribus ad 5 poll. longus, in superioribus deminuens, in supremis pæne obsoletus, pilosulus vel glabrescens; lamina linearis, superne ad apicem acutum attenuata, basi angustata et sagittata, plana, in facie superiore glabra, inferiore pilosa ad 10 poll. longa et  $4\frac{1}{2}$  lin. lata. *Panicula* ovoidea, ad  $4\frac{1}{2}$  poll. longa,  $2\frac{1}{2}$  poll. lata, rhacheos nodi pilosuli, rami filiformes verticillati (interdum cum paucis interpositis), patuli,  $1\frac{1}{2}$  poll. longi vel breviores, apice clavati et spiculas binas gerentes; spiculæ 3 lin. longæ, conformes, altera sessilis, altera pedicellata; pedicellus filiformis, 2 lin. longus, marginibus purpureo-ciliatus. *Glumæ* i. et ii. æquales, dorso pilosæ, virides, membranaceæ, prima 10-nervis, anguste ovata, obtusa,  $2\frac{3}{4}$  lin. longa; secunda primæ similis sed cum basi latiore, et 8-nervis, apice emarginata,

margine ciliolata; gl. iii. oblonga, apice biloba, 1-nervis, hyalina, margine et in parte superiore dorsali ciliolata, vix  $2\frac{1}{2}$  lin. longa, florem ♂ fovens; palea glumæ æqualis; gl. iv. fertilis, oblonga, basi angustata, valde bipartita, vix  $2\frac{1}{2}$  lin. longa, segmentis tenuibus, linearibus, acutis, hyalinis, arista e sinu orta, perfecta, 10 lin. longa; palea ovata,  $2\frac{1}{4}$  lin. longa; antheræ ut in flore masculo lineares sed paullo longiores ( $2\frac{1}{4}$  lin.), lodiculæ oblique cuneatæ.

YUNNAN: Mengtze, grass mountains, 5500 ft. (*A. Henry*, 10997!), wood, 4600 ft. (*A. Henry*, 11256!). Mus. Brit.; *Herb. Kew.*

A very striking species characterized by the reduction of the raceme to one pair of spikelets, and the petiolate sagittate leaves. In the reduction of the spikelets it resembles the Japanese *S. depauperatus*, Hack. (in *Bull. Herb. Boiss.* vii. p. 641), while in *S. albidus* the lamina has a well-marked petiole, and shows a tendency to become petiolate in *S. cotulifer*.

4. *Spodiopogon sibiricus*, *Trin. Fund. Agrost.* p. 192, et in *Bunge, Enum. Pl. Chin. Bor.* p. 73; *Hack. Mon. Androp.* p. 183, et in *Bull. Herb. Boiss.* vii. (1899) p. 641, et sér. 2, iii. (1903) p. 501; *Maxim. Prim. Fl. Amur.* p. 331; *Franchet, Pl. David.* i. p. 327, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 101; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222; *Komarow, in Act. Hort. Petrop.* xx. (1903) p. 245.

*Andropogon sibiricus*, *Steud. Syn. Gram.* p. 398.

CHIHLI: Peking (*Bushell! Williams & Bretschneider*, in herb. *Hance*, 11429!), Puhuashan (*Bretschneider*, in herb. *Forbes!*), mountains west of Peking (*Bretschneider!*), Jehol (*David*, 2056!); SHINGKING (*Ross*, 28!); SHANTUNG: Chefoo (*Forbes! Maingay*, 81!); HUPEH (*A. Henry*, 6717, 6520!); SHENSI: Mt. Kiansan, Huscenscien (*Hugh!*), Kusan, Kisanshien (*Hugh!*), Mt. Thuikiotsuen (*Hugh!*); KWANGTUNG (*Ford*, 196!), Taiwan (*Nevin*, in herb. *Hance*, 20156!). Mus. Brit.; *Herb. Kew.*

North Asia, Japan.

1. *Pollinia articulata*, *Trin. in Mém. Ac. Sci. Pétersb.* sér. 6, iv. (1836) p. 90; *Hack. Mon. Androp.* p. 153; *Hance, in Journ. Bot.* xviii. (1880) p. 302; *Hook. f. Fl. Brit. Ind.* vii. p. 109.

FOKIEN : Amoy (*Hance*, 1378 in part !); KWANGTUNG : Whampoa (*Hance*, 1378 in part !), Canton (*Sampson*, 478 !). Mus. Brit. ; Herb. Kew.

India, Malaya, Australia.

2. *Pollinia cantonensis*, *Rendle*, n. sp.

*Culmi* sejuncti, foliosi, ad 14 poll. longi, solum exstant, internodiis, ad 7, gracilibus, ad semilineam latis, levibus, nodis glabris, plus minus geniculatis. *Foliorum* vaginæ tenues, glabræ, ore sparse pilosulæ, quam internodia paullo breviores ; ligula membranacea, breviter ciliata, vix linea longa ; lamina anguste lanceolata, acuta, basi valde attenuata, 2-3 poll. longa,  $2\frac{1}{2}$ -4 lin. lata, leviter pilosa, cum nervo mediano albo-lineato et utrinque nervis 3-4 percursa. *Inflorescentia* vix evoluta, racemi ad 4, rhacheos internodia circa  $\frac{1}{4}$  poll. longa ; articuli racemorum haud sulcati, quam spicula lanceolata  $\frac{1}{4}$  breviores, in utraque margine pilis quam articuli  $\frac{2}{3}$  brevioribus instructi. *Spicula sessilis* biflora,  $2\frac{1}{2}$  lin. longa ; gluma i. membranacea, vix acuta, in carinis et nervis 2 dorsalibus pilosa, dorso leviter sulcata,  $2\frac{1}{4}$  lin. longa ; gl. ii. primæ æquans, mutica ; flos inferior masculus inter gl. primam paleamque ( $1\frac{3}{4}$  lin. longam) gl. iii. deficientis inclusus, antheris tribus  $1\frac{1}{3}$  lin. longis ; gl. iv.  $\frac{1}{2}$  lin. longa, hyalina, oblonga, apice integra, cum arista debili geniculata 2 lin. longa, palea ovata vix  $\frac{1}{2}$  lineam excedens, antheræ tres, 1 lineam paullo excedentes. *Spicula pedicellata* sessili similis, pedicellus quam rhachis angustior et paullo brevior, similiter pilosus.

KWANGTUNG : Canton (*Sampson*, in herb. *Hance*, 14923 in part !). Mus. Brit.

A distinct species of the subgenus *Leptatherum*, perhaps nearest to *P. vagans*, Nees, from which it is distinguished by the membranous hairy outermost glume, and imperfectly shortly awned fertile glume of the spikelet.

Described from a few flowering culms mixed with *Pollinia imberbis* from Canton (*Sampson*, in herb. *Hance*).

3. *Pollinia ciliata*, *Trin.* in *Mém. Ac. Sci. Pétersb.* sér. 6, ii. (1832) p. 306 ; *Hack. Mon. Androp.* p. 176 (subsp. c. *Wallichiana*), et in *Bull. Herb. Boiss.* vii. (1899) p. 723 ; *Hook. f. Fl. Brit. Ind.* vii. p. 116 ; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 186.

*Pollinia tenuis*, *Benth.* in *Fl. Hongk.* p. 421, non *Trin.*

*Pollinia Wallichiana*, *Nees, ex Steud. Syn. Gram.* p. 410.

FORMOSA: Takow (*A. Henry*, 1150!); KWANGTUNG: Whampoa (*Hance*, 5231 in part!); HONGKONG (*Hance*, 5231 in part! *Wilford*, 61!); LUCHU ARCHIPELAGO (*Wright*!). Mus. Brit.; Herb. Kew.

India, Malaya.

Var. *breviaristata*, *Rendle*, n. var. *Spicæ* numerosæ densæ; rhacheos internodia utrinque ciliata, pili  $\frac{1}{3}$  internodii longitudine haud æquantes. *Glumæ* i. et ii. subobtusæ; gl. iv. parva, integra; arista 3 lin. longa, perfecta, columna ejus subulæ æquans.

CHINA: without locality (*Mulgrave*!); HUPEH: Ichang (*A. Henry*, 3133!); YUNNAN: Mengtze, 5000 ft. (*A. Henry*, 11412!). Mus. Brit.; Herb. Kew.

4. *Pollinia Cumingii*, *Nees, in Hook. Kew Journ.* ii. (1850) p. 98; *Hack. Mon. Androp.* p. 167 (excl. var. *fulva*); *Hook. f. Fl. Brit. Ind.* vii. p. 114.

CHINA: without locality (*Fortune*, 4!); FOKIEN: Amoy (*Hance*, 1390!). Mus. Brit.; Herb. Kew.

North India, Malaya.

[? *Pollinia glabrata*, *Trin.*, fide *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

FORMOSA: Takow (*Playfair*, fide *A. Henry*).

*P. glabrata* is a plant from the Society Islands; there is no Formosan specimen either at the British Museum or Kew.]

5. *Pollinia imberbis*, *Nees*, var. *Willdenowiana*, *Hack. Mon. Androp.* p. 178; *Hook. f. Fl. Brit. Ind.* vii. p. 117.

*Pollinia imberbis*, *Nees, ex Steud. Syn. Gram.* p. 410 ampl., in part; *Hance, in Journ. Bot.* xvi. (1878) p. 233; *Komarow, in Act. Hort. Petrop.* xx. (1903) p. 249.

*Pollinia japonica*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 190, non *Miquel*.

KWANGTUNG: Canton (*Sampson*, in herb. *Hance*, 14923 in part!). Mus. Brit.; Herb. Kew.

Japan, North India, Java.

6. *Pollinia monantha*, *Nees, ex Steud. Syn. Gram.* p. 410; *Hack. Mon. Androp.* p. 174; *Hook. f. Fl. Brit. Ind.* vii. p. 116.

YUNNAN: Mengtze, 4500 ft. (*A. Henry*, 9210!). Herb. Kew.  
North India, Burmah.

The Yunnan specimen shows a remarkable variation as regards glume iii. of the spikelet. It may be absent or minute (smaller than the fertile gl. iv.) or form an important 3-nerved structure reaching 1 line in length with a weak geniculate awn  $1\frac{2}{3}$  line long.

7. *Pollinia nuda*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, ii. (1832) p. 307; *Hack. Mon. Androp.* p. 178, et in *Bull. Herb. Boiss.* vii. (1899) p. 641; *Hook. f. Fl. Brit. Ind.* vii. p. 117.

*Pollinia japonica*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 290.

*Leptatherum japonicum*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 190.

CHINA: without locality (*Seemann!*); LUCHU ARCHIPELAGO (*Wright!*). Mus. Brit.; Herb. Kew.

Japan, North India; South Africa.

8. *Pollinia pallens*, *Hack. Mon. Androp.* p. 156; *Hook. f. Fl. Brit. Ind.* vii. p. 110.

YUNNAN: near Tapintze (*Delavay*, 1793, in herb. Paris. *fide* Hackel).

North India.

I have seen no specimen.

9. *Pollinia phæothrix*, *Hack. Mon. Androp.* p. 168; *Hook. f. Fl. Brit. Ind.* vii. p. 112.

HUPEH: Ichang (*A. Henry*, 289!). Herb. Kew.

South India.

10. *Pollinia quadrinervis*, *Hack. Mon. Androp.* p. 158, et in *Bull. Herb. Boiss.* vii. (1899) pp. 640, 721; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 130; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222; *Hook. f. Fl. Brit. Ind.* vii. p. 110.

*Pollinia villosa*, *Munro, in Benth. Fl. Hongk.* p. 420, non *Spreng.*

*Pollinia villosa*, *Munro*, var. *chifuensis*, *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 271. (Probably belongs here. I have not seen the plant, and the diagnosis is inadequate.)

*Erianthus tristachyus*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 241, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 183.

FOKIEN: Amoy (*Hance*, 1381!); YUNNAN: Mengtze, grass mountains, 7000 ft. (*A. Henry*, 10052!); KWANGTUNG: Macao (*Vachell*, 351!); HONGKONG: Mount Victoria (*Lamont!*);



LUCHU ARCHIPELAGO (*Tashiro*, fide *Hackel*). Mus. Brit.; Herb. Kew.

Japan, North India.

Var. *latifolia*, *Rendle*, n. var. *Planta* quam in specie robustior, culmis quadripedalibus, ad  $2\frac{1}{2}$  lineas crassis prope basin. *Folia* lineari-lanceolata, rigidula, glabra, subtus pruinosa, in culmo 8-10 poll. longa, 6-7 lin. lata.

YUNNAN: Mengtze, grass mts., 6000 ft. (*A. Henry*, 10993!). — Mus. Brit.; Herb. Kew.

11. *Pollinia speciosa*, *Hack. Mon. Androp.* p. 159; *Hook. f. Fl. Brit. Ind.* vii. p. 113.

*Erianthus speciosus*, *Debeaux, Fl. Tchéfou*, p. 166; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272.

HUPEH: Ichang (*A. Henry*, 2796, 4374!); KWANGTUNG: Canton (*Sampson*, 297!). Herb. Kew.

North India.

12. *Pollinia velutina*, *Hack. Mon. Androp.* p. 169; *Hook. f. Fl. Brit. Ind.* vii. p. 113.

HUPEH (*A. Henry*, 4374!), Ichang (*A. Henry*, 2796!). Mus. Brit.; Herb. Kew.

North India.

1. *Pogonatherum saccharoideum*, *Beauv. Agrost.* p. 176, t. xi. fig. 7; *Hook. f. Fl. Brit. Ind.* vii. p. 141; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222; *Hack. Mon. Androp.* p. 193 (var. *α. genuinum*).

*Pogonatherum polystachum*, *Rœm. et Sch. Syst. Veg.* ii. p. 497.

*Saccharum paniceum*, *Lam. Encycl. Méth.* i. p. 595, t. 40. fig. 3.

HUPEH: Ichang (*A. Henry*, 1142!), Patung (*A. Henry*, 574, 3487!); YUNNAN (*A. Henry*, 11695!), Mengtze, grass hills, 5000 feet (*A. Henry*, 10360!); KWANGSI: Lungchow, common on slopes of hills (*Morse*, 496!). Mus. Brit.; Herb. Kew.

India, Malaya.

Var. *β. monandrum*, *Hack. Mon. Androp.* p. 193, et in *Bull. Herb. Boiss.* vii. (1899) p. 641.

*Pogonatherum crinitum*, *Kunth, Enum.* i. p. 478; *Miq. in Ann. Mus.*

*Bot. Lugd.-Bat.* ii. p. 289; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 189; *Hook. f. Fl. Brit. Ind.* vii. p. 141.

*Andropogon crinitus*, *Thunb. Fl. Jap.* p. 40, t. 7.

*Homoplitis crinita*, *Trin. Fund. Agrost.* p. 166.

*Pogonopsis tenera*, *Presl, Reliq. Hænk.* i. p. 333, t. 46.

*Pogonatherum polystachyum*, *Nees, in Hook. et Arn. Bot. Beechey's Voy.* p. 238, non *Rœm. et Sch.*

*Pogonatherum refractum*, *Nees, in Hook. et Arn. l. c.* p. 239, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 182; *Hook. et Arn. l. c.* p. 273.

*Pogonatherum saccharoideum*, *Beauv.; Benth. Fl. Hongk.* p. 421; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

CHINA: without locality (*Meyen!*); FORMOSA (*Oldham, 621!* *Campbell! Playfair, 250!*), Bankinsing (*A. Henry, 382!*), Takow, on rock in streamlet (*A. Henry, 762!*); SZECHUEN (*Faber, 1180!*), Min River (*Faber, 1155!*); KWANGTUNG: Canton (*Lamont!*), Macao (*Buchanan!*); HONGKONG (*Hance, 1033! Wright! Wilford, 62! Harland!*), Aberdeen and Mt. Victoria (*Lamont!*); LUCHU ARCHIPELAGO (*Wright! Beechey!*). *Mus. Brit.; Herb. Kew.*

Japan, India, Malaya.

I believe that there is only one species of this genus, of which *P. rufobarbatum*, *Griff.*, distinguished as a second species by *Hackel*, forms part. It is often difficult to say to which of *Hackel's* varieties a given specimen belongs, and both seem to be distributed over the whole area of the species. *Bentham* (*Journ. Linn. Soc., Bot.* xix. p. 67) says the genus "is a single tropical and subtropical Asiatic species."

1. *Apocopsis Wightii*, *Nees, ex Steud. Syn. Gram.* p. 377; *Hook. f. Fl. Brit. Ind.* vii. p. 142.

*Apocopsis Wightii*, subsp. *a. genuinus*, var. *γ. Wightii*, *Hack. Mon. Androp.* p. 259.

*Apocopsis Wightii*, *Munro, in Proc. Amer. Acad.* iv. (1860) p. 363, et in *Benth. Fl. Hongk.* p. 421.

*Andropogon courtallumensis*, *Steud. Syn. Gram.* p. 377.

KWANGTUNG: Whampoa (*Hance, 7432* in part!), Canton (*Sampson, 296!*), Lofausban (*Ford, 134* in part!); HONGKONG (*Hance, 7432* in part!), Cumsingmoon (*Wright! Ford, 134* in part!). *Mus. Brit.; Herb. Kew.*

India, Ceylon.

1. *Dimeria falcata*, *Hack. Mon. Androp.* p. 85.

*Dimeria fuscescens*, *Munro, in Benth. Fl. Hongk.* p. 419, non *Trin.*

FOKIEN: Amoy (*Hance, 1385 in part!*); KWANGTUNG: Canton (*Sampson, 1051!*); HONGKONG (*Hance, 1385 in part! Lamont!*).  
Mus. Brit.; Herb. Kew.

Tonkin.

2. *Dimeria sinensis*, *Rendle, n. sp.*

*Herba* annua; culmi erecti, filiformes, simplices (vel interdum cum ramo racemifero altero e nodo secundo orto), 7-nodi, 6–11 poll. alti. *Foliorum* vaginæ, ut laminæ præsertim in marginibus, tuberculato-ciliatæ, ad nodos barbatae; ligula perbrevis, truncata; laminæ planæ, lineares, acutæ, in parte culmi inferiore ad  $1\frac{1}{2}$  poll. longæ et  $\frac{3}{4}$  lin. latæ, superne breviores, summa rudimentaria. *Racemus* solitarius terminalis, ultra folium ultimum plus minus elatus, densus,  $1-1\frac{1}{4}$  poll. longus, circa 1 lin. latus, rhachi subflexuosa, dorso in parte inferiore sulco tenui percursa, marginibus anguste alata et breviter ciliata; spiculæ conniventes,  $1\frac{1}{2}$  lin. longæ; pedicelli brevissimi dorso breviter barbati; callus brevissimus cum pilis ipsi subæquantibus instructus. *Gluma* i. linearis, obtusa, basi subangustata, in margine, sub apice excepta, et carina dense pilosa, vix  $1\frac{1}{2}$  lin. longa; gl. ii. quam prima duplo latior, rhomboideo-oblonga, obtusa, basi angustata, margine membranacea, carina lata, pilosa,  $1\frac{1}{2}$  lin. longa,  $\frac{1}{2}$  lin. lata; gl. iii. angusta, hyalina, enervis, apice obtuso pilosa, margine sparse pilosa, vix  $\frac{1}{2}$  lin. longa; gl. iv. oblonga, basi angustata, hyalina, uninervis, apice obtusa,  $1\frac{1}{6}$  lin. longa; arista perfecta,  $6\frac{1}{2}$  lin. longa, columna vix 3 lin. longa.

KWANGTUNG: Canton (*Hance, 1385 in part!*). Mus. Brit.

Has the habit of the Ceylon species *Dimeria pusilla*, Thwaites, but is at once distinguished by its remarkably blunt spikelets.

1. *Arthraxon breviaristatus*, *Hack. Mon. Androp.* p. 350; *Hook. f. Fl. Brit. Ind.* vii. p. 144.

YUNNAN: Kimise (*Delavay, 1811 in herb. Paris. fide Hackel*).  
Khasia Hills.

I have seen no specimen.

2. *Arthraxon ciliaris*, *Beauv. Agrost.* p. 111, t. xi. fig. 6; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 134; *Franchet,*

*Pl. David.* i. p. 328, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 187; *Hook. f. Fl. Brit. Ind.* vii. p. 145.

Subsp. *a.* **Langsdorffii**, var. *α.* genuinus, *Hack. Mon. Androp.* p. 355, et in *Bull. Herb. Boiss.* vii. (1899) p. 642, et sér. 2, iii. (1903) p. 501; *Komarow, in Act. Hort. Petrop.* xx. (1903) p. 248.

*Arthraxon japonicus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 288.

*Arthraxon Langsdorffianus*, *Hochst. in Flora*, xxxix. (1856) p. 188.

*Andropogon lasiocoles*, *Steud. Syn. Gram.* p. 383.

*Andropogon amplexifolius*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, ii. p. 274.

*Deyeuxia japonica*, *Spreng. Syst.* i. p. 254.

*Lasiolytrum hispidum*, *Steud. in Flora*, xxxi. (1846) p. 18.

*Lasiolytrum hirtum*, *Steud. Syn. Gram.* p. 12.

*Lasiolytrum japonicum*, *Steud. ex Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 288.

*Pleuroplitis Langsdorffii*, *Trin. Fund. Agrost.* p. 175, t. xvi.

*Pleuroplitis Langsdorffii*, varr. *japonica*, *typica*, et *breviseta*, *Regel, in Bull. Acad. Sci. Pétersb.* x. (1866) pp. 373, 374.

*Luçæa Langsdorffiana*, *Steud. Syn. Gram.* p. 413.

CHINA: East Coast (*Home!*); SHINGKING (*Ross, 64!*), Kai-chow to Kinchow (*Ross, 182!*); SHANGTUNG: Chefoo (*Forbes, 405! Maingay, 65!*); KIANGSU: Shanghai (*Maingay, 659! Faber, 304!*); KIANGSI: Kiukiang (*Shearer!*); FOKIEN: Amoy (*Hance, 1396 in part!*); SZECHUEN: Mt. Thaepeisan (*Hugh!*); YUNNAN: Mengtze, 4700 feet (*A. Henry, 9137 A!*); KWANG-TUNG: Whampoa (*Hance, 1396 in part!*). *Mus. Brit.*; *Herb. Kew.*

Japan.

Subsp. *b.* **submuticus**, *Hack. Mon. Androp.* p. 356.

KIANGSU: Shanghai (*Seemann!*); HUPEH: Patung (*A. Henry, 4806!*); LUCHU ARCHIPELAGO (*Wright!*). *Herb. Kew.*

North India.

A species with numerous varieties which is widely spread through the warmer parts of the Old World.

3. **Arthraxon cuspidatus**, *Hochst. in Flora*, xxxix. (1856) p. 188; var. **micans**, *Hack. Mon. Androp.* p. 353; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222.

*Arthraxon micans*, *Hochst. l. c.*; *Franchet, Pl. David.* ii. p. 147.

*Arthraxon ciliaris*, *Beaur.*, group *c*, *Hook. f. Fl. Brit. Ind.* vii. p. 146.

SZECHUEN: Moupin (*David, fide Franchet*).

India, Mauritius.

I have seen no specimen.

4. *Arthraxon lanceolatus*, *Hochst. in Flora*, xxxix. (1856) p. 188; *Hack. Mon. Androp.* p. 347; *Hook. f. Fl. Brit. Ind.* vii. p. 143.

CHIHLI: Mountains west of Peking (*Bretschneider, in herb. Forbes!*); KIANGSU: Kiangnan (*Staunton!*); HUPEH (*A. Henry, 2282!*); YUNNAN: Mengtze, 4700 feet (*A. Henry, 9137!*).  
Mus. Brit.; Herb. Kew.

India; Abyssinia; Nubia.

5. *Arthraxon nodosus*, *Komarow, in Act. Hort. Petrop.* xviii. (1901) p. 448.

COREA: Czaschin, Kenge district (*Komarow*).

I have seen no specimen.

1. *Rottboellia compressa*, *Linn. f. Suppl.* p. 114; *Hack. Mon. Androp.* p. 286, et in *Bull. Herb. Boiss.* vii. (1899) p. 723; *Hook. f. Fl. Brit. Ind.* vii. p. 153; *Komarow, in Act. Hort. Petrop.* xx. (1903) p. 247.

*Hemarthria compressa*, *R. Br. Prodr.* p. 207; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 287; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 186; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272.

Var. *genuina*, *Hack. Mon. Androp.* p. 286; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108, sub *Hemarthria fasciculata*, *Kunth*.

FORMOSA: Takow (*A. Henry, 1896!*), Apes' Hill (*Playfair, 123!*); YUNNAN: Mengtze, 4500 feet (*A. Henry, 10169!*);  
KWANGTUNG: Canton (*Sampson et Hance, in herb. Hance, 1862 in part!*). Mus. Brit.; Herb. Kew.

India.

Var. *fasciculata*, *Hack. Mon. Androp.* p. 286.

*Hemarthria fasciculata*, *Kunth, Rév. Gram.* i. p. 153; *Benth. Fl. Hongk.* p. 418.

HONGKONG (*Hance, 1862 in part!*). Mus. Brit.

Mediterranean region to North India, South Africa, Madagascar, Tropical and Subtropical America.

Var. *japonica*, *Hack. l. c.* p. 288, et in *Bull. Herb. Boiss. sér. 2, iii.* (1903) p. 501.

CHIHLI (*Staunton!*); SHINGKING (*Ross, 55!*); SHANTUNG:  
LINN. JOURN.—BOTANY, VOL. XXXVI. 2 B

Chefoo (*Faber*, 290! *Forbes*, 399!). Mus. Brit.; Herb. Kew.

Japan.

Scarcely distinguishable from var. *fasciculata*.

2. *Rottboellia exaltata*, *Linn. f. Suppl.* p. 114; *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 201; *Hack. Mon. Androp.* p. 293; *Hook. f. Fl. Brit. Ind.* vii. p. 156; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

*Stegosia cochinchinensis*, *Lour. Fl. Cochinch.* p. 51.

FOKIEN: Amoy (*Hance*, 1395 in part!); FORMOSA: Takow (*A. Henry*, 1127 A!), Apes' Hill (*A. Henry*, 1127! *Playfair*, 194!); KWANGTUNG: Whampoa (*Hance*, 1395 in part!), Macao (*Bladh*! *Ford*, 203!). Mus. Brit.; Herb. Kew.

Widely distributed in the tropics.

3. *Rottboellia latifolia*, *Steud. in Flora*, xxxi. (1846) p. 21, et *Syn. Gram.* p. 361; *Hack. Mon. Androp.* p. 281, et in *Bull. Herb. Boiss.* vii. (1899) p. 641, et sér. 2, iii. (1903) p. 501; *Debeaux, Fl. Shanghai*, p. 71.

*Rottboellia foliata*, *Steud. Syn. Gram.* p. 361.

*Ischæmum latifolium*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 291; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 193.

CHINA: without locality (*Fortune*, A 29!); East Coast (*Home*!). Mus. Brit.; Herb. Kew.

Japan.

4. *Rottboellia mollicoma*, *Hance, in Journ. Bot.* ix. (1871) p. 134; *Hack. Mon. Androp.* p. 297.

KWANGTUNG: Whampoa (*Hance*, 7558!), Lofaushan (*Ford*, 130!). Mus. Brit.; Herb. Kew.

Tonkin.

A very distinct species with the habit of *R. glandulosa*, *Trin.*, but at once distinguished by the sessile spikelet at each node, the back of which is conspicuously tessellated as in *R. nigrescens*, *Thwaites*; it differs from the latter in the perfectly free stalk of the rudimentary pedicelled spikelet.

1. *Ophiurus monostachyus*, *J. S. Presl, in C. B. Presl, Rel. Hænk.* i. p. 330; *Hack. Mon. Androp.* p. 318; *Benth. Fl. Hongk.* p. 417.

*Ophiurus undatus*, Nees, in *Hook. Kew Journ.* ii. (1850) p. 100.

FORMOSA: Takow (*A. Henry*, 1201!); HONGKONG (*Hance*, 1248! *Harland*!). Mus. Brit.; Herb. Kew.

Philippine Islands; Tonkin.

1. *Manisuris granularis*, *Linn. f. Nov. Gram. Gen.* p. 37, figs. 4-7; *Hack. Mon. Androp.* p. 314; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

FORMOSA: Apes' Hill, Takow (*A. Henry*, 1865!), Tamsui (*A. Henry*, 1478!); KWANGTUNG: Whampoa (*Hance*, 5022! *Ford*, 205!); HAINAN (*A. Henry*, 8120!). Mus. Brit.; Herb. Kew.

Tropics of both hemispheres.

1. *Eremochloa leersioides*, *Hack. Mon. Androp.* p. 264; *Hook. f. Fl. Brit. Ind.* vii. p. 140.

*Eremochloa falcata*, *Hack. l. c.* p. 263.

*Ischæmum leersioides*, *Munro*, in *Proc. Amer. Acad.* iv. (1864-65) p. 363; *Benth. Fl. Hongk.* p. 425; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 135; *Pilger*, in *Engl. Jahrb.* xxix. (1900) p. 222.

*Ischæmum pectinatum*, *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 201 (quoad plantam chinensem).

*Nardus ciliaris*, *Linn. Sp. Pl.* p. 53; *Osbeck, Resa*, p. 220.

KWANGTUNG (*Tate! Ford*, 198!), Sam Cocks Island near Macao (*Staunton!*), Canton (*Sampson*, 303!), Whampoa (*Wright!*), Pakhoi (*Playfair*, 147!); HONGKONG (*Hance*, 1251! *Lamont! Forbes*, 497! *Wilford*, 249! *Seemann*, 223! *Faurie*, 15849!). Mus. Brit.; Herb. Kew.

Pegu, Tonkin.

This species is very near *Eremochloa zeylanica*, of which it is probably only a less robust variety, with shorter, more falcate spikes and narrower, less obtuse leaves.

2. *Eremochloa ophiuroides*, *Hack. Mon. Androp.* p. 261.

*Ischæmum ophiuroides*, *Munro*, in *Proc. Amer. Acad.* iv. (1864-65) p. 363; *Benth. Fl. Hongk.* p. 425.

KIANGSU: Kiangnan (*Staunton!*); CHEKIANG: Ningpo Mts. (*Faber*, 1532!); HUPEH (*A. Henry*, 4148!); KWANGTUNG: Whampoa (*Wright!*), Macao (*Tate!*), Pakhoi (*Playfair*, 142 in part!); HONGKONG (*Hance*, 1249!). Mus. Brit.; Herb. Kew.

3. *Eremochloa zeylanica*, *Hack. Mon. Androp.* p. 263; *Hook. f. Fl. Brit. Ind.* vii. p. 140.

HUPEH (*A. Henry*, 3502, 569!). Mus. Brit.; Herb. Kew.  
Ceylon.

1. *Ischæmum angustifolium*, *Hack. ex Oliver, in Hook. Icon. Pl.* (1888) t. 1773, et in *Mon. Androp.* p. 241, et in *Bull. Herb. Boiss.* vii. (1899) p. 723; *Hook. f. Fl. Brit. Ind.* vii. p. 129; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222.

*Pollinia eriopoda*, *Hance, in Journ. Bot.* iv. (1866) p. 173.

FORMOSA: Apes' Hill (*Swinhoe, in herb. Hance*, 12585! *A. Henry*, 1056!); HUPEH (*A. Henry*, 568!); YUNNAN: Mengtze, 5000 ft. (*A. Henry*, 10390!). Mus. Brit.; Herb. Kew.

India, North to Central; Philippines.

2. *Ischæmum antephoroides*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* iii. p. 193; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 193; *Hack. Mon. Androp.* p. 216, et in *Bull. Herb. Boiss.* vii. (1899) p. 641.

*Ischæmum barbatum*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 291, non *Retz.*

SHANTUNG: Chefoo (*Carmichael, in herb. Forbes*, 1074! *Main-gay*, 66!). Mus. Brit.; Herb. Kew.

Japan.

3. *Ischæmum aristatum*, *Linn. Sp. Pl.* p. 1049; *Hook. f. Fl. Brit. Ind.* vii. p. 126.

Subsp. *imberbe*, *Hack. Mon. Androp.* p. 203.

FORMOSA: Takow (*A. Henry*, 1807, 2046!); HUPEH: Ichang (*A. Henry*, 2299, 4242!); KWANGTUNG: Canton (*Sampson*, 304!). Mus. Brit.; Herb. Kew.

India, Malaya.

Subsp. *barbatum*, *Hack. l. c.* p. 204; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 222.

*Ischæmum barbatum*, *Retz. Obs.* vi. p. 35; *Benth. Fl. Hongk.* p. 425; *Franchet, in Mém. Soc. Sc. Nat. Cherbourg*, xxiv. (1884) p. 273.

*Meoschium lodiculare*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 246, et in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 195.

*Meoschium Meyenianum*, *Nees, in Nov. Act. l. c.* p. 197.

CHINA: without locality (*Staunton!*); KWANGTUNG: Macao



(*Tate*!), Pakhoi (*Playfair*!); HONGKONG (*Hance*, 1257! *Harland*! *Hind*! *Seemann*, 232! *Wilford*! *Wright*! *Champion*, 545!), Mt. Victoria (*Lamont*!). Mus. Brit.; Herb. Kew.

India, Malaya.

4. *Ischæmum aureum*, *Hack. Mon. Androp.* p. 224.

*Spodiopogon aureus*, *Hook. & Arn. Bot. Beechey's Voy.* p. 273.

LUCHU ARCHIPELAGO (*Beechey*, 672!). Herb. Kew.

5. *Ischæmum ciliare*, *Retz. Obs.* vi. p. 36; *Hook. f. Fl. Brit. Ind.* vii. p. 134, *et in Trim. Fl. Ceyl.* v. p. 216; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273; *Debeaux, Fl. Tchéfou*, p. 163; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 723.

*Ischæmum ciliare*, var. *genuinum*, *Hack. Mon. Androp.* p. 226.

*Spodiopogon obliquivalvis*, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 185; *Benth. Fl. Hongk.* p. 426, including var. *villosus*, *Benth.*

*Spodiopogon villosus*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 242.

CHINA: without locality (*Staunton*! *Reeves*!); KIANGSI (*Staunton*!); FOKIEN: Amoy (*Hance*, 1257!); FORMOSA: Tamsui (*Oldham*, 623!); KWANGTUNG: Whampoa (*Robertson*! *Hance*, 1526 in part!), hills near Canton (*Sampson*, 371!), Pakhoi (*Playfair*, 155!), Okkaiisa, sandy banks near sea (*Tate*!), Cape Syngmun (*Wright*!); HONGKONG: Mt. Victoria and Govt. Gardens (*Lamont*! *Forbes*, 496! *Hance*, 1526 in part! *Tate*!). Mus. Brit.; Herb. Kew.

India, Ceylon, Malaya.

I cannot arrange the Chinese specimens under the subvarieties of the var. *genuinum* (which corresponds with the species) proposed by Hackel. I find villous forms with scrobiculate first glumes, &c.

6. *Ischæmum eriostachyum*, *Hack. Mon. Androp.* p. 218.

*Ischæmum barbatum*, *Debeaux, Fl. Tchéfou*, p. 163, non *Retz.*, fide *Hackel, l. c.*

SHANTUNG: Chefoo (*Debeaux, Wawra* fide *Hackel*).

Japan.

I have seen no specimen.

7. *Ischæmum laxum*, *R. Br. Prodr.* p. 205; *Hack. Mon. Androp.* p. 243; *Hook. f. Fl. Brit. Ind.* vii. p. 136.

— YUNNAN: Mengtze, 4600 ft. (*A. Henry*, 9618, 9618 A!).  
Herb. Kew.

Cape Verde Is., Tropical Africa, Arabia, Tropical Asia to  
Australia.

8. *Ischæmum rugosum*, *Salisb. Ic. Stirp. Rar.* i. t. 1; *Hack. Mon. Androp.* p. 206, et in *Bull. Mus. Herb. Boiss.* vii. (1899) p. 723; *Hook. f. Fl. Brit. Ind.* vii. p. 127.

*Ischæmum segetum*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, ii. p. 294; *Hance, in Journ. Bot.* xvi. (1878) p. 233.

*Meoschium rugosum*, *Nees, in Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 200.

FORMOSA: Takow (*A. Henry*, 2001!); KWANGTUNG: Whampoa (*Hance*, 9665!), by river Lienchau (*Nevin*, in herb. *Hance*, 20150!), Pakhoi (*Playfair*, 153!). Mus. Brit.; Herb. Kew.

India, Malaya.

9. *Ischæmum Sieboldi*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 291; *Franch. et Savat. Enum. Pl. Japon.* ii. p. 193; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273; *Hack. Mon. Androp.* p. 217, et in *Bull. Herb. Boiss.* vii. (1899) p. 641, et sér. 2, iii. (1903) p. 501; *Debeaux, Fl. Tchéfou*, p. 163.

CHINA: East Coast (*Home*!); SHANTUNG: Chefoo (*Forbes*, 384!); KIANGSU: Shanghai (*Fabre-Tonnerre*, in herb. *Hance*, 10156! *Maingay*, 557, 667!). Mus. Brit.; Herb. Kew.

Japan.

1. *Heteropogon contortus*, *Beauv. ex Roem. et Schult. Syst.* ii. p. 836; *Nees, in Hook. & Arn. in Bot. Beechey's Voy.* p. 241.

*Andropogon contortus*, *Linn. Sp. Pl.* p. 1045; *Hack. Mon. Androp.* p. 585, et in *Bull. Herb. Boiss.* vii. pp. 721, 723; *Hook. f. Fl. Brit. Ind.* vii. p. 199.

*Heteropogon hirtus*, *Pers. Syn.* ii. p. 533; *Benth. Fl. Hongk.* p. 424; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

FORMOSA: Apes' Hill, Takow (*A. Henry*! *Playfair*, 289!); HUPEH: Ichang (*A. Henry*, 442, 4143!); KWANGTUNG: Canton (*Sampson*, 302!); HONGKONG (*Hance*, 555! *Wright*!), Mount Victoria (*Lamont*!), Aberdeen and Little Hongkong (*Wilford*, 258!). Mus. Brit.; Herb. Kew.

Generally distributed in the warmer parts of the earth.

1. *Sorghum fulvum*, Beauv. *Agrost.* p. 164.

*Andropogon serratus*, Thunb. *Fl. Jap.* p. 41; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 192.

*Andropogon serratus*, var. *genuinus*, Hack. *Mon. Androp.* p. 521, et in *Bull. Herb. Boiss.* vii. (1899) p. 642, et sér. 2, iii. (1903) p. 501; Hook. f. *Fl. Brit. Ind.* vii. p. 185.

*Andropogon tropicus*, Spreng. *Syst. Veg.* i. p. 287; Benth. *Fl. Hongk.* p. 423; Miq. in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 290; Debeaux, *Fl. Tientsin*, p. 74; Franchet, *Pl. David.* i. p. 328; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 192.

*Holcus fulvus*, R. Br. *Prodr.* p. 199.

FOKIEN: Amoy (*Hance*, 1394 in part!); KIANGSU: Chinkiang (*Carles*, 501!); KIANGSI: Kiukiang (*Shearer*!); HUPEH: Ichang (*A. Henry*, 28, 4234!); KWANGTUNG: Canton (*Hance*, 1394 in part! *Sampson*, 476!). *Mus. Brit.*; *Herb. Kew.*

India, Malaya, Japan, Australia.

Var. *nitidum*, Hack. *Mon. Androp.* p. 521; Hook. f. *Fl. Brit. Ind.* vii. p. 185.

*Andropogon nitidus*, Kunth, *Rév. Gram.* i. p. 166.

*Sorghum nitidum*, Pers. *Syn.* p. 101.

FORMOSA: Tamsui (*Oldham*, 624!), Takow plain (*A. Henry*, 1895!). *Mus. Brit.*; *Herb. Kew.*

India, Malaya.

2. *Sorghum halepense*, Pers. *Syn.* i. p. 101.

*Andropogon halepensis*, Brot. *Fl. Lusit.* i. p. 89; Hook. f. *Fl. Brit. Ind.* vii. p. 182; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273.

*Andropogon Sorghum*, subsp. *halepensis*, Hack. *Mon. Androp.* p. 501.

KWANGTUNG: Canton, by river Lienchau (*Nevin*, in herb. *Hance*, 4879 in part!), banks of North River, Canton (*Sampson*, in herb. *Hance*, 4879 in part!); HAINAN: Nartaisee (herb. *Ford*, 484! *A. Henry*, 8295!). *Mus. Brit.*; *Herb. Kew.*

Most warm countries.

3. *Sorghum vulgare*, Pers. *Syn.* i. p. 101; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

*Andropogon Sorghum*, Brot. *Fl. Lusit.* i. p. 88; Komarow, in *Act. Hort. Petrop.* xx. (1901) p. 250.

*Andropogon Sorghum*, subsp. *sativus*, var. *vulgaris*, Hack. *Mon. Androp.* p. 515; Stapf, in Hook. f. *Fl. Brit. Ind.* vii. p. 184.

FORMOSA: Bankinsing, cultivated (*A. Henry*, 807!); South Cape (*Schmüerer*, in herb. *A. Henry*, 257!). Mus. Brit.; Herb. Kew.

Widely cultivated in warm countries.

Var. *bicolor*, *Hack. l. c.* p. 519.

CHIHLI: Peking (*Bretschneider*, 799–800!). Herb. Kew.

Var. *nervosum*, *Hack. l. c.* p. 513; *Stapf*, in *Hook. f. Fl. Brit. Ind.* vii. p. 184.

CHIHLI: Peking, cultivated (*Bretschneider*, in herb. *Hance*, 1349 in part!); FOKIEN: Amoy, cultivated (*Hance*, 1349 in part!); CHINA: East Coast (*Home*!); HAINAN (*A. Henry*, 8757!). Mus. Brit.; Herb. Kew.

India.

Var. *obovatum*, *Hack. l. c.* p. 514; *Stapf*, *l. c.*

KIANGSU: Shanghai (*Faber*, 277 in part!); CHEKIANG: Ningpo (*Faber*, 277 in part!); KWANGTUNG: waste ground near Canton (*Hance*, 5544!). Mus. Brit.; Herb. Kew.

Madeira, Mediterranean Region, India, Japan.

1. *Chrysopogon aciculatus*, *Trin. Fund. Agrost.* p. 188; *Benth. Fl. Hongk.* p. 424; *Hance*, in *Ann. Sci. Nat. sér. 5*, v. (1866) p. 253; *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 238; *Hook. & Arn. l. c.* p. 273; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

*Andropogon aciculatus*, *Retz. Obs.* v. p. 22; *Hack. Mon. Androp.* p. 562, et in *Bull. Herb. Boiss.* vii. (1899) pp. 721, 723.

*Centrophorum chinense*, *Trin. Fund. Agrost.* p. 106, t. 5.

*Rhaphis trivialis*, *Lour. Fl. Cochinch.* p. 553.

FORMOSA (*Campbell*!), Tamsui (*Oldham*, 622!), Takow (*A. Henry*, 1034!), Apes' Hill (*Playfair*, 33!); KWANGTUNG: Macao (*Vachell*, 49!), Pakhoi (*Playfair*, 152!), Lofaushan (*Ford*!); HONGKONG (*Hance*, 383! *Tate*! *Wright*!), Little Hongkong (*Wilford*, 326!), Happy Valley and West Point (*Lamont*!); LUCHU ARCHIPELAGO (*Lay & Collie*!). Mus. Brit.; Herb. Kew.

Mauritius, Tropical Asia, Australia, Polynesia.

2. *Chrysopogon sinensis*, *Rendle*, n. sp.

*Planta* perennis; culmi ascendentes, demum erecti, ramosi, 12–15 poll. alti, vix semilineam crassit. excedentes, plurinodi.

*Foliorum* vaginæ glabræ, in parte culmi inferiore breviores, ad  $2\frac{1}{4}$  poll. long., internodiis sæpe æquantes vel excedentes; ligula in seriem ciliorum brevissimorum soluta; laminæ sæpius conduplicatæ, lineares, obtusiusculæ, ad  $2\frac{3}{4}$  poll. longæ, vix 1 lin. crassit. superantes, rigidiusculæ, basi in margine sparse barbatae, ceterum glabræ. *Panicula* breviter exserta, demum subpyramidalis, ad 6 poll. longa; rami inæquales, septeni vel pauciores, demum patentes, filiformes, sub apice sensim incrassati, cum ramulo terminale spiculam singulam fertilem gerente, oblique articulati, disjuncti cicatricem ovalem ferrugineo-ciliatam exhibentes; ramulus 2 lin. longus, ferrugineo-pilosus. *Spicula sessilis*  $2\frac{1}{2}$  lin. longa, glabra; gluma i. elliptico-oblonga, binervis, apice breviter bifida et hirsutula; gl. ii. primam subæquans, oblonga, compressa, 3-nervis, dorso rotundata et coriacea, marginibus hyalina, cum arista imperfecta dorsali 6 lin. longa; gl. iii. anguste linearis hyalina, margine ciliolata, quam prima et secunda paullo brevior; gl. fertilis anguste linearis, 3-nervis, hyalina, 2 lin. longa, arista perfecta  $1\frac{1}{4}$  poll. longa, columna brunnea semipollicaris; antheræ lineares  $1\frac{1}{2}$  lin. longæ. *Spicula pedicellata* 3 lin. longa, pedicellus ferrugineo-pilosus, vix 2 lin. long.; gluma i. lanceolata, apice obsolete aristulata, 6-7-nervis; gl. ii. lanceolata, acuta, 3-nervis, primam æquans; gl. iii. & iv. hyalinæ, margine ciliatæ, gl. iii. vix 3 lin. longa, binervis, gl. iv. anguste elliptica, 3-nervis, quam tertia paullo brevior, antheræ vix 2 lin. longæ.

KWANGTUNG: Pakhoi (*Playfair*, 144!); HAINAN: roadside leading from Hoihau to Kingchau (*Sampson*, in herb. *Hance*, 3453!). Mus. Brit.; Herb. Kew.

Near *Andropogon Trinii*, Steud., but distinguished by the longer partial inflorescence, the much elongated oblique base by which the latter is attached to the branch, the longer stalk of the partial inflorescence, &c.

1. *Andropogon annulatus*, *Forsk. Fl. Æg.-Arab.* p. 173; *Hack. Mon. Androp.* p. 570; *Hook. f. Fl. Brit. Ind.* vii. p. 196.

Var. *genuinus*, *Hack. l. c.* p. 572.

HUPEH: Ichang (*A. Henry*, 200, 1090, 3528!); HONGKONG (*Ford*, 102!). Mus. Brit.; Herb. Kew.

Arabia to India, Africa, Australia.

Var. *Bladhii*, *Hack. Mon. Androp.* p. 572; *Hook. f. Fl. Brit. Ind.* vii. p. 197.

*Andropogon Bladhii*, *Retz. Obs.* ii. p. 27; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 243.

KWANGTUNG: Canton, very abundant (*Sampson*, 1057!). *Herb. Kew.*

India.

2. *Andropogon apricus*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, iv. (1836) p. 83; var. *indicus*, *Hack. Mon. Androp.* p. 457.

*Andropogon apricus*, *Hook. f. Fl. Brit. Ind.* vii. p. 169.

YUNNAN: Mengtze, barren clay hills, 4600 ft. (*A. Henry*, 10995!). *Herb. Kew.*

The species occurs in India, West Tropical Africa, and Tropical America.

3. *Andropogon assimilis*, *Steud. Syn. Gram.* p. 397; *Hook. f. Fl. Brit. Ind.* vii. p. 179.

*Andropogon glaucopsis* et *A. subrepens*, *Steud. l. c.*

*Andropogon montanus*, *Hack. Mon. Androp.* p. 490, non *Roxb.*

*Chrysopogon pictus*, *Hance, in Ann. Sci. Nat.* sér. 5, v. (1866) p. 252.

CHIHLI: Peking (*Williams*, in herb. *Hance*, 10154 in part!); CHEKIANG: Changking (*Faber*!); FOKIEN (*De Grijs*, in herb. *Hance*, 10154 in part!); HUPEH: Ichang (*A. Henry*, 4220, 4177!); SZECHUEN: Suifu, Yangtzekiang (*Faber*, 1169!); YUNNAN: Mengtze, 4600–4800 ft. (*A. Henry*, 9138, 9615, 9617!), Szemao, 5500 ft. (*A. Henry*, 13414!); KWANGTUNG: Canton (*Lamont*!), White Cloud Hills (*Sampson*, 1902 in part!); LUCHU ARCHIPELAGO (*Wright*!). *Mus. Brit.*; *Herb. Kew.*

India, Java, Timor.

Scarcely distinct from *A. micranthus*, *Kunth.*

4. *Andropogon brevifolius*, *Swartz, Prodr. Veg. Ind. Occ.* p. 26; *Hack. Mon. Androp.* p. 363, et in *Bull. Herb. Boiss.* vii. (1899) p. 642, et sér. 2, iii. (1903) p. 501; *Benth. Fl. Hongk.* p. 423; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 135; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 191; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 291; *Hook. f. Fl. Brit. Ind.* vii. p. 165; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273.

SHANTUNG: Chefoo (*Forbes*, 395!), Chefoo Mountains (*Faber*,

288 !); KWANGTUNG: Whampoa (*Hance*, 6401 !), Canton (*Sampson*, 1409 !). Mus. Brit.; Herb. Kew.

Japan; tropics of both worlds.

5. *Andropogon caricosus*, *Linn. Sp. Pl.* ed. 2, p. 1480; *Hack. Mon. Androp.* p. 567; *Hook. f. Fl. Brit. Ind.* vii. p. 196.

YUNNAN: near Tapintze (*Delavay*, fide *Hackel*).

India, Timor, Mauritius.

I have seen no specimen.

6. *Andropogon Delavayi*, *Hack. Mon. Androp.* p. 404.

YUNNAN: Kimise (*Delavay*, 1800), meadows at Kipinkay near Tali (*Delavay*, 2242).

I have seen no specimen.

7. *Andropogon fastigiatus*, *Sw. Prodr. Veg. Ind. Occ.* p. 26; *Hack. Mon. Androp.* p. 393; *Hook. f. Fl. Brit. Ind.* vii. p. 167.

*Andropogon apricus*, *Trin.*, var. *chinensis*, *Hack. Mon. Androp.* p. 457.

*Homœatherum chinense*, *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 239.

KWANGTUNG: Macao (*Vachell*, fide *Nees*).

Widely spread in the tropics of both hemispheres.

I have not seen the specimen.

8. *Andropogon formosanus*, *Rendle*, n. sp.

*Herba* late ramosa; rami multinodi repentis ad nodos geniculatos radicanter, demum adscendentes et floriferi, radicibus valde elongatis, internodiis teretibus, ad  $\frac{2}{3}$  lin. crass., glabris. *Foliorum* vaginæ cylindricæ ad internodia breviora arcte applicitæ, glabræ (sub ore quidem sparsius pilosulæ); ligula membranacea, ovata, obtusa, brunneo-tincta, pilosula, vix 1 lin. longa; lamina linearis, apice acuminato-setacea, basi valde angustata, plana, glabra, costa media præcipue in facie inferiore albida et conspicua,  $2\frac{1}{2}$ –4 poll. longa,  $2\frac{1}{2}$ –3 lin. lata. *Panicula* terminalis, ramis 3–6, fastigiatis, sæpe subdigitatis, tenuibus, 1–1 $\frac{1}{4}$  poll. longis, axem communem duplo vel pluries excellentibus; racemi fragiles, articuli pedicellique tenues, anguste clavati, haud sulcati, articuli circa  $1\frac{2}{3}$  lin. longi, glabri, vel

marginē exteriore, in parte racemi superiore, piloso, pedicelli  $\frac{1}{2}$  ad  $\frac{3}{4}$  articularum longitudini æquantes, utroque margine cum pilis patentibus instructi. *Spicula sessilis* lineari-lanceolata acuta, 2–2 $\frac{1}{2}$  lin. longa; gluma i. dorso subtiliter tuberculata, et sulcata, apice breviter acuminata et biaristulata, marginibus implicatis superne ciliolatis, callo brevissimo (vix  $\frac{1}{8}$  lin. longo) barbato; gl. ii. cymbiformis, primam subæquans, apice leviter aristata, arista vix 2 lin. longa; gl. iii. minuta ( $\frac{1}{3}$  lin. longa), hyalina, late elliptica, obtusa; gl. iv. minuta (vix  $\frac{1}{4}$  lin. longa), hyalina, breviter oblonga, apice in aristam debilem, imperfectam, purpureo-tinctam, 6–7 lin. longam, producta; palea absens; caryopsis anguste ovalis, pallide brunnea, glabra, 1 lin. longa. *Spicula pedicellata* quam sessilis paullo minor, ceterum similis, mox tabescens.

FORMOSA: on cliff, Takow (*A. Henry*, 1167!). Herb. Kew.

Near *A. vagans*, Rendle, resembling in habit the more delicate form of that species. It is distinguished by the narrower racemes and spikelets, the awned second glume, and the larger but weak and imperfect awn of the fertile glume.

Var. *minor*, Rendle, n. var. Rhacheos articuli pedicelli et spiculæ quam in specie breviores; *spicula sessilis* cum callo brevissimo vix 1 $\frac{1}{2}$  lin. longa; gluma i. obtusa.

HONGKONG: Climbing in hedges near Aberdeen (*Hance*, 10746 in part!). Mus. Brit.; Herb. Kew.

9. *Andropogon fragilis*, R. Br. *Prodr.* p. 202.

Var. *sinensis*, Rendle, n. var. *Planta* annua quam typus minus robusta, cum spathis florentibus paullo angustioribus. *Spicula sessilis* gluma i. dorso ad medium pilosa, non velut in specie glabra.

KWANGTUNG: French Island, Whampoa (*Hance*, 1388(?)!). Herb. Kew.

The type occurs in Queensland and North Australia.

The species is included by Hackel as a variety ( $\delta$ ) of *Andropogon brevifolius*, from which, however, it is, I think, quite distinct. Brown founded his species on plants collected during Captain Cook's first voyage by Banks and Solander at Endeavour River. Professor Hackel has probably not seen the original specimens at the British Museum; they differ from *A. brevi-*



*folius* in their more robust habit, subacute foliage-leaves, the broad spathe-like sheath enveloping the inflorescence (which suggested to the discoverers a more appropriate species-name than that adopted by Brown), and the stouter hairy racemes. Hackel describes the rhachis-joints as glabrous, whereas they bear a row of white hairs on the outer edge, the hairs being almost as long as the joint.

10. *Andropogon hirtiflorus*, *Kunth, Rév. Gram.* ii. p. 569, t. 198; *Hack. Mon. Androp.* p. 371; *Hook. f. Fl. Brit. Ind.* vii. p. 167.

*Andropogon Pseudograya*, *Steud. Syn. Gram.* p. 365; *Hack. Mon. Androp.* p. 370.

*Andropogon zeylanicus*, *Arn. ex Benth. Fl. Hongk.* p. 423.

FOKIEN: Amoy (*Hance*, 1388 in part!); KWANGTUNG: Whampoa (*Hance*, 1388 in part!), Cape Syngmun (*Wright*!); HONGKONG (*Lamont*!). *Mus. Brit.*; *Herb. Kew.*

India, Ceylon, Tonkin; Madagascar, Tropical and South Africa; America.

11. *Andropogon intermedius*, *R. Br. Prodr.* p. 202; *Hack. Mon. Androp.* p. 485; *Hook. f. Fl. Brit. Ind.* vii. p. 175; *Pilger, in Engl. Jahrb.* xxix. (1900) p. 222.

*Andropogon punctatus*, *Roxb. Fl. Ind.* i. p. 264; *Trin. Sp. Gram. Ic.* t. 328; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 135; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 242.

*Andropogon Vachellii*, *Nees, l. c.* p. 243; *Benth. Fl. Hongk.* p. 423; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

SHANTUNG (*Maingay*, 76!); FOKIEN: Amoy (*Seemann*, 547! *Hance*, 1078!); FORMOSA: Takow (*A. Henry*, 1039, 1094, 1152!), Apes' Hill (*Playfair*, 102!); HUPEH (*A. Henry*, 2373!), Ichang (*A. Henry*, 4096, 2766, 2374!); YUNNAN: Mengtze, 4800 ft. (*A. Henry*, 9614!); KWANGTUNG (*Staunton*! *Sampson*, 299!), Macao (*Hance*, 1407 in part! *Vachell*!); HONGKONG (*Hance*, 1407 in part! 10229! *Harland*, 801 in herb. *Hance*, 1078!), roadsides at Aberdeen (*Wilford*, 248!), ravine, Mt. Victoria (*Lamont*!). *Mus. Brit.*; *Herb. Kew.*

Caucasus, Tropical Asia to Australia, Pacific Islands, Tropical and South Africa.

The specimens vary considerably in size of panicle, length of panicle-branches, amount of secondary branching, size of

spikelets, and presence or absence of a pit on the outer glume of the sessile spikelet; also in the presence or absence of hairs at the mouth of the sheath. I prefer to consider them all under the species rather than to attempt to assign them to the varieties proposed by Hackel; in such case it would be necessary to make other varieties.

12. *Andropogon Ischæmum*, *Linn. Sp. Pl.* p. 1047; *Hack. Mon. Androp.* p. 474, et in *Bull. Herb. Boiss*, sér. 2, iii. (1903) p. 501; *Trin. in Bunge, Enum. Pl.* p. 73; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273; *Hook. f. Fl. Brit. Ind.* vii. p. 171.

CHIHLI: Peking (*Bushell! Bretschneider*, 822!); SHINGKING: Bay of Taliénwan (*Swinhoe, in herb. Hance*, 3452!); SHANTUNG: Chefoo (*Forbes*, 404!); KIANGSI: Kiukiang (*Reid*, 9!); FOKIEN: Amoy (*Hance*, 1389, 3452 in part!); FORMOSA: Apes' Hill (*Playfair*, 272!), Takow (*A. Henry*, 761!); SHENSI: Kusan (*Hugh!*); KWANGTUNG: Whampoa (*Hance*, 3452 in part!). *Mus. Brit.*; *Herb. Kew.*

South and Central Europe, Central to North Asia, Northwest India, and in many parts of the tropics and subtropics of both hemispheres.

13. *Andropogon micranthus*, *Kunth, Rév. Gram.* i. p. 165, et *Enum.* i. p. 504; *Hack. Mon. Androp.* p. 488, et in *Bull. Herb. Boiss.* vii. (1899) p. 642, et sér. 2, iii. (1903) p. 501; *Hook. f. Fl. Brit. Ind.* vii. p. 178.

*Andropogon capilliflorus*, *Steud. Syn. Gram.* p. 397 (fide *Hackel*); *Franch. et Savat. Enum. Pl. Jap.* ii. p. 192; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 50.

*Andropogon montanus*, *Benth. Fl. Hongk.* p. 423, non *Roxb.*

CHINA: without locality (*Fortune*, 13!); SHINGKING: Kaichow to Kinchau (*Ross*, 278!); SHANTUNG: Chefoo (*Forbes! Main-gay*, 83!); CHEKIANG (*Staunton!*), Ningpo (*Faber*, 283!); HUPEH: Ichang (*A. Henry*, 26, 2373, 4285!), Changlo (*A. Henry*, 7815!); SZECHUEN: Omei, 3000 ft. (*Faber*, 1177!); YUNNAN: Mengtze, grass mountains, 5500 ft. (*A. Henry*, 9615 A!); KWANGTUNG: Canton (*Lamont*), Whampoa (*Hance*, 1078 in part! *Sampson*, 289!); HONGKONG (*Hance*, 1078 in part!). *Mus. Brit.*; *Herb. Kew.*

Japan; Old World tropics.

variety of *A. annulatus* and as occurring also in India. The Whampoa specimen is a more delicate plant, with slenderer stem and smaller narrower leaves than the one from Canton.

15. *Andropogon yunnanensis*, *Hack. Mon. Androp.* p. 440.

YUNNAN: Mosoyn (*Delavay*, 1782, fide *Hackel*).

I have seen no specimen.

1. *Cymbopogon Nardus*, *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 155.

*Andropogon Nardus*, *Linn. Sp. Pl.* p. 1046; *Hack. Mon. Androp.* p. 601; *Hook. f. Fl. Brit. Ind.* vii. p. 205; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

Forma culta, *Hook. f. Fl. Brit. Ind.* vii. p. 206; subsp. *a. genuinus*, *Hack. l. c.* p. 602.

HONGKONG: cultivated (*Hance*, 255!). Mus. Brit.

Cultivated in Tropical Asia.

Subsp. *hamatulus*, *Hack. Mon. Androp.* p. 606.

*Andropogon hamatulus*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 244.

FORMOSA: Takow (*A. Henry*, 1009, 1921!), Apes' Hill, Takow (*A. Henry*, 1021!); KWANGTUNG: Danes Island, below Canton (*Vachell*, 416! *Herb. Cantab.*); HONGKONG (*Champion*, 541!); CHINA: without locality (*Millett*!). Herb. Kew.

Philippines.

Subsp. *marginatus*, var. *Goeringii*, *Hack. Mon. Androp.* p. 607, *et in Bull. Herb. Boiss.* vii. p. 642, et sér. 2, iii. (1903) p. 501; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 30.

*Andropogon Goeringii*, *Steud. in Flora*, xxxi. (1846) p. 22.

*Andropogon Schoenanthus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 290; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 191.

CHINA: without locality (*Fortune*, 99!); SHANGTUNG (*Main-gay*, 79!), Chefoo (*Forbes*!); KIANGSU: Shanghai (*Faber*, 296!), Fengwangshan (*Forbes*, 494!); CHEKIANG: Chusan and Ningpo (*Home*!), Ningpo Mts. (*Faber*, 291!); FORMOSA (*Swinhoe*!), South Cape (*A. Henry*, 1309!), Apes' Hill, Takow (*Playfair*, 35, 193, 237!); KWANGTUNG (*Ford*, 204!), Lienchau, Canton (*Lamont*!); HONGKONG: Mount Victoria (*Lamont*! *Forbes*, 493! *Seemann*, 754!); LUCHU ARCHIPELAGO (*Wright*!). Mus. Brit.; Herb. Kew.

Japan.

Var. *villosulus*, *Hack. Mon. Androp.* p. 490; *Hook. f. Fl. Brit. Ind.* vii. p. 179.

*Andropogon villosulus*, *Steud. Syn. Gram.* p. 397.

YUNNAN (*A. Henry*, 11256!); HONGKONG (*Sampson*, 1902 in part!). *Herb. Kew.*

North India.

14. *Andropogon vagans*, *Rendle*, n. sp.

*Planta* divaricata, culmi demum erecti, teretes, in speciminibus incompleti, ad 20 poll. longi et 1 lin. crassi, internodia superiora vaginas superantia, striata, et plus minus pilosula. *Foliorum* vaginæ ad internodia arcte applicitæ, glabræ, superne autem pilosulæ, nodis puberulis; ligula truncata ad  $\frac{3}{4}$  lin. longa; laminæ anguste lineari-lanceolatæ, basi valde angustatæ, superne acuminatæ pæne ad aristulam attenuatæ, 3-6 poll. longæ, 3-8 lin. latæ, planæ, herbacæ, dorso puberulæ, et nervo mediano albo-lineatæ. *Racemi* 4-10, dense fastigiati, 2-4 $\frac{3}{4}$  poll. longi, axem communem rarius  $\frac{1}{2}$  poll. excedens; spiculæ densæ, rhacheos articuli et pedicelli in marginibus pilosi, haud canaliculati, hi paullo breviores, 1 lin. longi. *Spicula sessilis* cum callo brevissimo barbato 1 $\frac{1}{2}$  lin. longa, calli pili haud medium glumæ attingentes; gluma i. ovali-oblonga, obtusa, marginibus implicatis, carinulis superne ciliatis, dorso binervis, glabra et per longitudinem sulcata; gl. ii. cymbiformis, primam subæquans, mutica, carina in parte superiore scabridula; gl. iii. ovata, hyalina, enervis, longitudine vix plus quam  $\frac{1}{2}$  lin.; gl. iv. brevissima, hyalina, breviter bilobulata, arista e sinu perfecta, 3 lin. longa. *Spicula pedicillata* ♀ sessili similis, sed minor (1 $\frac{1}{4}$  lin. longa) lanceolata et cum arista brevior (vix 2 lineas long. excedens).

A distinct species in the alliance of *A. Ischænum*, but distinguished by its habit, puberulous internodes, denser less hairy inflorescence, bisexual stalked flower, absence of a channel on the pedicel, &c.

KWANGTUNG: Canton, ravine of the White Cloud Hill (*Sampson*, 300!), Whampoa (*Hance*, 10746 in part!). *Mus. Brit.*; *Herb. Kew.*

Both the plants cited were distributed as *Andropogon Bladhii*, Retz., of which I have not seen the type. It is, however, described as having unawned and neuter stalked spikelets and long-awned sessile spikelets, and is generally regarded as a

# NOTICE.

---

*The Society's Apartments will be closed  
for cleaning during the month of  
AUGUST.*

# LINNEAN SOCIETY OF LONDON.

---

In future all Publications for the LINNEAN SOCIETY should be addressed

**THE LIBRARIAN,**

**Linnean Society,**

**Burlington House,**

**Piccadilly,**

**LONDON, W.**

**It is desirable that all parcels should be sent direct by post, as far as possible, in order to avoid delay in transmission.**

# INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

## BOTANY

has been undertaken by the Council of the Linnean Society, and they appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

The volume for 1901, in two parts, has been published; the volume for 1902 is now being prepared for press.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.

---

VOL. XXXV. is still in course of issue, the Parts already published being Nos. 242-247. Part 248 will complete the volume.

VOL. XXXVI., of which Parts 249-253 are already published, is reserved for the completion of the Chinese Flora.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."



JULY 25.

Price 6s.

# THE JOURNAL

OF

# THE LINNEAN SOCIETY.

VOL. XXXVI.

BOTANY.

No. 254.

## CONTENTS.

Page

An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.—

PART XIX. . . . . 377

---

See Notice on last page of Wrapper.

---

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE, PICCADILLY, W.;

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1904.

12

# LINNEAN SOCIETY OF LONDON.

## LIST OF THE OFFICERS AND COUNCIL.

Elected 24th May, 1904.

### PRESIDENT.

Prof. W. A. Herdman, D.Sc., F.R.S.

### VICE-PRESIDENTS.

Frank Crisp, LL.B., B.A.	Albert C. Seward, F.R.S.
Dr. A. Günther, F.R.S.	Prof. S. H. Vines, F.R.S.

### TREASURER.

Frank Crisp, LL.B., B.A.

### SECRETARIES.

D. H. Scott, M.A., Ph.D., F.R.S.	Rev. T. R. R. Stebbing, M.A., F.R.S.
----------------------------------	--------------------------------------

### COUNCIL.

V. H. Blackman, M.A.	Dr. W. G. Ridewood, F.Z.S.
Charles Baron Clarke, M.A., F.R.S.	G. S. Saunders, F.E.S.
Frank Crisp, LL.B., B.A.	Dukinfield H. Scott, M.A., Ph.D., F.R.S.
Prof. J. B. Farmer, M.A., D.Sc., F.R.S.	A. C. Seward, M.A., F.R.S.
Dr. A. Günther, F.R.S.	Dr. Otto Stapf.
Prof. W. A. Herdman, F.R.S.	Rev. T. R. R. Stebbing, M.A., F.R.S.
B. Daydon Jackson, Esq.	Prof. Sydney H. Vines, D.Sc., F.R.S.
F. G. Parsons, F.Z.S.	

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*; the former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meets at 4 P.M., as required during the Session. The Members for 1904-1905, in addition to the officers, are:—

Herbert Druce, F.Z.S.	A. B. Rendle, M.A., D.Sc.
Antony Gepp, M.A.	A. G. Tansley, M.A.
W. Botting Hemsley, F.R.S.	Prof. H. Marshall Ward, F.R.S.
Dr. G. Henderson.	Dr. A. Smith Woodward, F.R.S.
P. Chalmers Mitchell, M.A., F.Z.S.	

2. **Cymbopogon Schœnanthus**, *Spreng. Pl. Min. Cogn. Pug.* ii. p. 14.

*Andropogon Schœnanthus*, *Linn. Sp. Pl.* p. 1046; *Lour. Fl. Cochinch.* p. 793; *Debeaux, Fl. Tchef.* p. 162; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 273.

Var. **cæsius**, *Hack. Mon. Androp.* p. 610; *Hook. f. Fl. Brit. Ind.* vii. p. 205.

*Andropogon cæsius*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 244.

*Andropogon Martini*, *Benth. Fl. Hongk.* p. 424, non *Roab.*

KWANGTUNG: Macao (*Abel, Vachell!*), Whampoa (*Robertson, Wright! Millett!*), Lofaushan (*Ford, 127!*); HONGKONG: wild (*Hance, 254!*), Cumsingmoon (*Wright!*); HAINAN (*A. Henry, 8287!*). *Mus. Brit.*; *Herb. Kew.*

Arabia, Afghanistan, India.

1. **Themeda gigantea**, *Hack. Mon. Androp.* p. 670.

*Anthistiria gigantea*, *Cav. Icon.* v. p. 35; *Hook. f. Fl. Brit. Ind.* vii. p. 216.

Subsp. **caudata**, *Hack. Mon. Androp.* p. 676; *Hook. f. Fl. Brit. Ind.* vii. p. 217.

*Androscepia gigantea*, var. *armata*, *Anderss. in Nov. Act. Soc. Sci. Upsal.* ser. 3, ii. (1856) p. 248.

*Anthistiria caudata*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 245.

*Anthistiria ciliata*, *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108.

CHINA: without locality (*Staunton!*); FORMOSA: Bankinsing Mountains (*A. Henry, 1518!*); YUNNAN: Mengtze Mountains, 5500 ft. (*A. Henry, 9459!*); KWANGTUNG: near Canton (*Hance, 6489!*), Canton, common on the hills, sometimes 8 ft. high (*Sampson, 276!*). *Mus. Brit.*; *Herb. Kew.*

Mountains of North India.

Subsp. **villosa**, *Hack. Mon. Androp.* p. 675; *Hook. f. Fl. Brit. Ind.* vii. p. 217.

*Anthistiria villosa*, *Lam. Illustr.* t. 841, fig. 3; *Poir. Encycl. Suppl.* i. p. 396.

HUPEH: So Tunghu (*A. Henry, 7696!*), Ichang (*A. Henry, 980!*); HAINAN: Nartaisee (herb. *Ford, 483!*). *Herb. Kew.*

India, Malaya.

2. **Themeda triandra**, *Forsk. Fl. Æg.-Arab.* p. 178.

*Themeda Forskalii*, *Hack. Mon. Androp.* p. 659.

*Anthistiria ciliata*, *Retz. Obs.* iii. p. 11 (non *Linn. f.*), *Baker & Moore,*

in *Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Franchet, Pl. David.* i. p. 328.

*Anthistiria Forskalii*, *Kunth, Rév. Gram.* p. 162.

*Anthistiria imberbis*, *Retz. Obs.* iii. p. 11; *Hook. f. Fl. Brit. Ind.* vii. p. 211.

Var. **major**, subvar. **japonica**, *Hack. Mon. Androp.* p. 662, et in *Bull. Herb. Boiss.* vii. (1899) p. 642, et sér. 2, iii. (1903) p. 501; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 131; *Komarow, ibid.* xx. (1901) p. 251.

*Andropogon ciliatum*, *Thunb. Fl. Jap.* p. 40.

*Anthistiria arguens*, *Franchet, Pl. David.* i. p. 328 (non *Willd.*), et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 272.

*Anthistiria arguens*, *Willd.*, var. *japonica*, *Anderss. in Nov. Act. Soc. Sci. Upsal.* ser. 3, ii. (1856) p. 236; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 191.

*Anthistiria japonica*, *Willd. Sp. Pl.* iv. p. 901; *Trin. in Bunge, Enum. Plant. Chin. Bor.* p. 73.

CHINA: without locality (*Seemann! Bradley!*); CHIHLI: mountains west of Peking (*Bretschneider, 838!*), Peking (*Williams, in herb. Hance, 12758! David, 593! Bushell!*); SHINGKING: Chienshan (*Ross, 470!*), Kaichow to Kinchow (*Ross, 277!*); SHANTUNG: Chefoo (*Forbes! Hance, in herb. Forbes, 1444 in part! Maingay, 71!*); CHEKIANG: Chusan and Ningpo (*Home!*); KIANGSI: Kiukiang (*Reid, 7!*); KIANGSU: Chinkiang (*Carles, 502, 503!*), Shanghai (*Seemann!*); HUPEH: Ichang (*A. Henry, 959, 4091!*); SHENSI: Leunteon, Fenghosan (*Hugh!*), Kusan, Kisanshien (*Hugh!*), Mt. Laoysan (*Hugh!*); YUNNAN: Mengtze, grass mountains, 5500 ft. (*A. Henry, 10996!*); COREA: Soul (*Sontag!*). Mus. Brit.; Herb. Kew.

The species is widely distributed in the warmer parts of the Old World.

1. ***Pseudanthistiria heteroclita***, *Hook. f. Fl. Brit. Ind.* vii. p. 219.

*Andropogon heteroclitus*, *Nees, Fl. Afr. Austr.* p. 115; *Hack. Mon. Androp.* p. 400.

*Anthistiria heteroclita*, *Roxb. Fl. Ind.* i. p. 249; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 135.

HONGKONG (*Hance, 9667!*). Mus. Brit.; Herb. Kew.  
India.

1. *Germainia capitata*, *Bal. & Poitr. in Bull. Soc. Hist. Nat. Toulouse*, vii. (1873) p. 344; *Hack. Mon. Androp.* p. 686.

*Anthistiria caudata*, *Hance, in Journ. Bot.* xvi. (1878) p. 234, non *Nees*.

KWANGTUNG: Pakwan Mountains above Canton (*Hance*, 14392!), White Cloud Hills, Canton (*Sampson*, 1162!). Mus. Brit.; Herb. Kew.

Cochin-China.

1. *Apluda mutica*, *Linn. Sp. Pl.* p. 82; *Benth. Fl. Hongk.* p. 422; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 721, 723.

*Apluda varia*, subsp. *mutica*, *Hack. Mon. Androp.* p. 198; *Hook. f. Fl. Brit. Ind.* vii. p. 150.

CHINA: without locality (*Bradley!* *Macartney!* *Willett!*); FOKIEN: Amoy (*Hance*, 1003!); FORMOSA: Tamsui (*Oldham*, 625!), Takow (*A. Henry*, 743, 1114!); KWANGTUNG: Lienchau, Canton (*Lamont!*), Whampoa (*Wright!*), Canton (*Sampson*, 292!), Macao and adjacent islands (*Vachell*, 42!), hills, Pybus Island (*Wright!*); HONGKONG (*Hance*, ex herb. *Harland*, 588, 1003! *Ford*, 597!), Causeway Bay (*Ford*, 600!); HAINAN (*A. Henry*, 52!); LUCHU ARCHIPELAGO: Okinawa (*Tashiro*, fide *Hack. in Bull. Herb. Boiss.* vii. p. 721). Mus. Brit.; Herb. Kew.

India, Malaya, Australia, Pacific Islands.

Var. *aristata*, *Hack. Mon. Androp.* p. 199; *Hook. f. Fl. Brit. Ind.* vii. p. 151.

*Apluda aristata*, *Linn. Amœn. Acad.* iv. p. 303.

YUNNAN: Mengtze Ravine, 4700 ft. (*A. Henry*, 9616!). — Herb. Kew.

Socotra, India, Malaya.

#### Tribe PHALARIDÆ.

1. *Phalaris arundinacea*, *Linn. Sp. Pl.* p. 55; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 69; *Franch. Pl. David.* i. p. 328; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 157; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 646, et sér. 2, iii. (1903) p. 502; *Hook. f. Fl. Brit. Ind.* vii. p. 221; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 263.

CHIHLI: Jehol (*David*, in herb. *Hance*, 3574!); CHEKIANG:

Ningpo, Shanghai (*Faber*!); SHINGKING: between Tungcheshien and Maoershan, Yaloo River (*James*!). Mus. Brit.; Herb. Kew.  
Japan; North temperate regions.

1. *Anthoxanthum Hookeri*, *Rendle*.

*Ataxia Hookeri*, *Griseb. in Goett. Nachr.* 1868, p. 77.

*Hierochloe Hookeri*, *Maxim. in Bull. Acad. Pétersb.* xii. (1888) p. 930;  
*Hook. f. Fl. Brit. Ind.* vii. p. 223.

SZECHUEN: West Szechuen and Tibetan frontier (*Pratt*, 889!);  
YUNNAN: Mengtze, grass mountains, at 5000 feet (*A. Henry*,  
10517!). Mus. Brit.

North India (Himalayas).

I agree with Dr. Stapf in regarding Robert Brown's genus *Ataxia* as more closely allied to *Anthoxanthum* than to *Hierochloe*.

1. *Hierochloe borealis*, *Roem. et Schult. Syst.* ii. p. 513; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 131; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 157; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 646, et sér. 2, iii. (1903) p. 502; *Regel, Tent. Fl. Ussur.* p. 169; *Maxim. Prim. Fl. Amur.* p. 322.

*Hierochloe Bungeana*, *Trin. in Mém. Acad. Pétersb.* sér. 5, vi. II. (1839) p. 82.

CHEKIANG: Tientai Mountains (*Faber*, 73!). Herb. Kew.  
Japan, North Asia, and North temperate regions.

2. *Hierochloe glabra*, *Trin. in Spreng. Neue Entdeck. Pflanz.* ii. (1821) p. 66.

*Hierochloe dahurica*, *Trin. in Mém. Acad. Pétersb.* sér. 5, v. II. (1839) p. 80; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 91; *Maxim. Prim. Fl. Amur.* p. 322; *Turcz. Fl. Baic.-Dahur.* ii. p. 296; *Franchet, Pl. David.* i. p. 329, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267.

*Hierochloe borealis*, *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388, non *Roem. et Schult.*

CHIHLI: Mt. Puhuashan, west of Peking (*Bretschneider*, in herb. *Forbes*, 484! and in herb. *Hance*!), Jehol (*David*, in herb. *Hance*!); SHINGKING: hills near Takushan (*Ross*, 56!), Ccrean Gate (*Ross*, 422!), Peiling (*Webster*, 190!). Mus. Brit.; Herb. Kew.

North Asia.

## Tribe AGROSTIDÆ.

1. *Aristida Adscensionis*, *Linn. Sp. Pl.* p. 82; *Steud. Syn. Gram.* p. 139; *Hook. f. Fl. Brit. Ind.* vii. p. 224.

*Aristida cœrulescens*, *Desf. Fl. Atlant.* i. p. 109, t. 21. fig. 2; *Franchet, Pl. David.* i. p. 329.

*Aristida vulpioides*, *Hance, in Ann. Sci. Nat. sér. 5, v.* (1866) p. 251.

CHIHLI: Peking (*Williams, in herb. Hance, 11465!*), between Peking and Jehol (*Staunton!*); NORTH CHINA (*Archer!*). *Mus. Brit.*; *Herb. Kew.*

India, and widely distributed in the warmer parts of the earth.

2. *Aristida chinensis*, *Munro, in Proc. Amer. Acad.* iv. (1860) p. 363; *Benth. Fl. Hongk.* p. 427.

KWANGTUNG: Lienchau, Canton (*Lamont!*), Whampoa (*Wright! Hance, 1241 in part!*); HONGKONG (*Hance, 1241 in part! Harland!*), Cumsingmoon (*Wright!*). *Mus. Brit.*; *Herb. Kew.*

3. *Aristida Cumingiana*, *Trin. et Rupr. in Mém. Acad. Pétersb. sér. 6, vii.* (1843) p. 141; *Steud. Syn. Gram.* p. 140; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 135; *Hook. f. Fl. Brit. Ind.* vii. p. 224.

*Aristida capillacea*, *Cav. Icon.* v. p. 43, t. 468. fig. 1, non *Lam.*

*Chætaria trichodes*, *Nees, in Hook. Kew Journ.* ii. (1850) p. 101.

KWANGTUNG: dry grassy slopes near Canton (*Sampson, 1194!*); HONGKONG: steep rocks near sea between Aberdeen and Hongkong (*Sampson, in herb. Hance, 15057! Lamont!*). *Mus. Brit.*; *Herb. Kew.*

North India, Philippines.

1. *Stipa Bungeana*, *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70, et in *Mém. Acad. Pétersb. sér. 6, vii.* (1843) p. 61; *Steud. Syn. Gram.* p. 129; *Franchet, Pl. David.* i. p. 330.

CHIHLI: Kalgan, by the Great Wall (*Williams, in herb. Hance, 13831 in part!*), Peking (*Bretschneider, 832!*). *Herb. Kew.*

2. *Stipa capillata*, *Linn. Sp. Pl.* ed. 2, p. 116; *Steud. Syn. Gram.* p. 130; *Franchet, Pl. David.* i. p. 330; *Turcz. Fl. Baic Dahur.* ii. p. 303; *Hook. f. Fl. Brit. Ind.* vii. p. 230.

CHIHLI: near Peking (*David, 501!*). *Herb. Kew.*

Himalaya and temperate Asia; Europe.

3. *Stipa consanguinea*, *Trin. et Rupr. e descript. in Mém. Acad. Pétersb. sér. 6, vii. (1843) p. 78; Steud. Syn. Gram. p. 130.*

SZECHUEN: Tongolo (*Soulié, 341!*). Herb. Kew.  
Central Asia.

4. *Stipa* (§ *Lasiagrostis*) *Henryi*, *Rendle, n. sp.*

*Herba* perennis, rhizomate valido 4 lin. crasso, glabra; culmi et axes foliosi conferti numerosi, illi ad 2 ped. alti, basi vaginis scarioso-membranaceis sæpe tribus circumdati; internodia elongata gracilia tria, basi  $\frac{1}{2}$  ad  $\frac{4}{5}$  lin. crass. *Foliorum* vaginæ internodia sæpe vix æquantes, striatæ, sæpe purpureo-tinctæ; ligula obsoleta; lamina linearis, basi angustata et apice longe attenuata, plana vel interdum siccitate convoluta, plus quam pedalis longa et 2 lin. lata, sæpius autem angustior, cum nervo mediano in facie superiore prominente. *Panicula* subelongata, ad 11 poll. longa, rami capillares ascendentes, ad 5 poll. longi, plures in nodis rhacheos distantibus lateraliter fasciati, ramuli pauci-spiculiferi; pedicellus spiculam æquans vel superans; spiculæ  $1\frac{2}{3}$ – $1\frac{3}{4}$  lin. longæ, obtuse lanceolatæ. *Gluma* i. membranacea translucens, cum nervis tribus interdum duobus viridibus ad apicem haud porrectis, lineata, elliptico-ovata obtusa, spiculam æquans; gl. ii. primæ similis at paullo brevior; gl. florens subcoriacea, anguste ovalis (quum convoluta), aperta late oblonga, trinervis, apice breviter bilobulata, dorso minute puberula, cum callo breve (vix  $\frac{1}{8}$  lin.) et breviter piloso,  $1\frac{1}{2}$  lin. longæ; arista debilis, continua, 5 lin. longa, puberula; palea quam gluma florens paullo brevior; lodiculæ 2, oblongæ, basi angustatæ,  $\frac{1}{4}$  ad  $\frac{3}{5}$  lin. longæ; antheræ 3, lineares, haud apice barbatae, 1 lin. longæ.

HUPEH: Ichang (*A. Henry, 3444!*); SZECHUEN: South Wushan (*A. Henry, 5530!*), Yangtzekiang, Kweichau (*Faber, 1160!*). Herb. Kew.

Near *S. splendens*, Trin., but differing in its less robust habit, obsolete ligule, subequal barren glumes, and glabrescent fertile glume. There are occasionally traces of bearding on the anther-tips.

5. *Stipa mongholica*, *Turcz. ex Trin. in Bull. Sci. Acad. Pétersb. i. (1836) p. 67; Steud. Syn. Gram. p. 132; Hook. f. Fl. Brit. Ind. vii. p. 229.*



*Lasiagrostis mongholica*, *Trin. et Rupr. in Mém. Acad. Pétersb. sér. 6, vii. (1843) p. 87.*

*Ptilagrostis mongolica*, *Griseb. in Ledeb. Fl. Ross. iv. p. 447; Turcz. Fl. Baic.-Dahur. ii. i. p. 302.*

SHENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

Himalaya, North and Central Asia; Western North America.

6. *Stipa sibirica*, *Lam. Illustr. i. p. 158; Steud. Syn. Gram. p. 129; Hance, in Journ. Linn. Soc., Bot. xiii. (1873) p. 91; Franchet, Pl. David. i. p. 330, et in Mém. Soc. Sci. Nat. Cherbourg, xxiv. (1884) p. 268; Hack. in Bull. Herb. Boiss. sér. 2, iii. (1903) p. 502; Maxim. Prim. Fl. Amur. p. 325; Turcz. Fl. Baic.-Dahur. p. 302; Hook. f. Fl. Brit. Ind. vii. p. 231.*

*Stipa pekinensis*, *Hance, in Journ. Bot. xv. (1877) p. 268; Franchet, Pl. David. i. p. 331.*

CHIHLI: Kupehkan (*Hance, 13831 in part!*), Peking (*Bretschneider, 834! Williams, in herb. Hance, 13831 in part!*); SHANTUNG: Chefoo (*Forbes! Faber, 1529!*), (*Maingay, 75!*); CHEKIANG: Kaichow to Kinchow (*Ross, 276!*). Mus. Brit.; Herb. Kew.

Himalaya, North Asia; Western North America.

[*Stipa Aliciæ*, *Kanitz, in Math. Naturw. Ber. Ungarn, iii. (1884) p. 11 (nomen).*]

I have seen no specimen.

1. *Oryzopsis obtusa*, *Stapf, in Hook. Icon. t. 2393; Filger, in Engl. Bot. Jahrb. xxix. (1900) p. 224.*

HUPEH: near Ichang (*A. Henry, 3507!*), Nanto and mountains to northward (*A. Henry, 3896!*). Mus. Brit.; Herb. Kew.

1. *Milium effusum*, *Linn. Sp. Pl. p. 61; Steud. Syn. Gram. p. 34; Maxim. Prim. Fl. Amur. p. 325; Turcz. Fl. Baic.-Dahur. ii. p. 300; Miq. in Ann. Mus. Bot. Lugd.-Bat. ii. p. 274; Franch. et Savat. Enum. Pl. Jap. ii. p. 160; Hack. in Bull. Herb. Boiss. vii. (1899) p. 647; Komarow, in Act. Hort. Petrop. xx. (1901) p. 368; Loureiro, Fl. Cochinch. p. 49; Hook. f. Fl. Brit. Ind. vii. p. 235.*

SHINGKING: Chienshan (*Ross, 575!*); HUPEH: Patung dis-

trict (*A. Henry*, 4032, 5413 !); SZECHUEN (*A. Henry*, 5661 !), Mt. Omei (*A. Henry*, 1161 !). Mus. Brit.; Herb. Kew.

Japan, Cochin China; Himalaya; Europe; Eastern North America.

1. *Crypsis aculeata*, *Ait. Hort. Kew.* i. p. 48; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 91; *Franchet, Pl. David.* i. p. 331, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 267; *Debeaux, Fl. Tchef.* p. 157; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 270.

CHIHLI: Peking (*David*, in herb. *Hance*!). Mus. Brit.  
Europe; temperate Asia; North Africa.

1. *Phleum alpinum*, *Linn. Sp. Pl.* p. 59; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 647; *Hook. f. Fl. Brit. Ind.* vii. p. 236; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 271.

SHENSI: Mt. Miaowansan (*Hugh*!). Mus. Brit.

North Asia, Japan, Himalayas; Arctic and Alpine regions.

2. *Phleum asperum*, *Jacq. Collect.* i. p. 110; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 647; *Hook. f. Fl. Brit. Ind.* vii. p. 237.

SHENSI: Paoki, Chifengsan (*Hugh*!), plain of Hukiascien (*Hugh*!), Laoshien and Kasan (*Hugh*!), Mt. Laoysan, Teuizscien (*Hugh*!), Mt. Kanysan (*Hugh*!); KIANGSU: Shanghai (*Maingay*, 566, 716, 718 !). Mus. Brit.; Herb. Kew.

Japan, Himalaya, Western Asia, Europe.

1. *Alopecurus æqualis*, *Sobol. Fl. Petropol.* (1799) p. 16.

*Alopecurus aristulatus*, *Michx. Fl. Bor.-Am.* i. p. 43 (1803); *Hook. f. Fl. Brit. Ind.* vii. p. 238.

*Alopecurus fulvus*, *Smith, Engl. Bot.* xxi. t. 1467 (1805); *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648, et sér. 2, iii. (1903) p. 502; *Franchet, Pl. David.* i. p. 329; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 131; *Komarow, ibid.* xx. (1901) p. 271.

*Alopecurus geniculatus*, *Linn. Sp. Pl.* p. 60 in part; *Benth. Fl. Hongk.* p. 408; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 69; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Thunb. Fl. Jap.* p. 49; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 277; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 158.

SHINGKING: West of Sarhoo (*Ross*, 361!), near Moukden (*James*!); KIANGSU: Chinkiang (*Maries*!), Shanghai (*Faber*!); CHEKIANG: Taihoo Lake (*Forbes*, 483!); FORMOSA (*Oldham*, 599!); HUPEH: Ichang (*A. Henry*, 3475!), Patung (*A. Henry*, 4733!); HONGKONG (*Lamont*! *Wright*! *Hance*, 1087! *Wilford*, 259!); CHINA: without locality (*Staunton*!). Mus. Brit.; Herb. Kew.

Japan; North temperate regions.

2. *Alopecurus agrestis*, *Linn. Sp. Pl.* ed. 2, p. 89; *Steud. Syn. Gram.* p. 149; *Benth. Fl. Hongk.* p. 407; *Hook. f. Fl. Brit. Ind.* vii. p. 239; *Ledeb. Fl. Ross.* iv. p. 465.

KIANGSU: Shanghai (*Maingay*, 719!). Herb. Kew.

Europe; Western and Northern Asia; India.

3. *Alopecurus japonicus*, *Steud. Syn. Gram.* p. 149; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 277; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 158; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648.

*Alopecurus malacostachyus*, *A. Gray, in Perry's Exped. Japan*, ii. p. 328.

*Alopecurus agrestis*, *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224, non *Linn.*

HONGKONG (*Hance*, 1153!); HUPEH (*A. Henry*, 7869!), Patung (*Wilson*, 1661!). Mus. Brit.; Herb. Kew.

Japan.

1. *Muehlenbergia Hugelii*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, vi. (1841) p. 293; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 647, et sér. 2, iii. (1903) p. 502.

*Muehlenbergia viridissima*, *Nees, ex Steud. Syn. Gram.* p. 178; *Hook. f. Fl. Brit. Ind.* vii. p. 259; *Franchet, Pl. David.* ii. p. 143; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224.

KIANGSI: Kiukiang (*Shearer*!); HUPEH: Chiensbih (*A. Henry*, 7410!), Fang (*A. Henry*, 6720!), Ichang (*A. Henry*, 4289!), Patung (*A. Henry*, 2513!). Mus. Brit.; Herb. Kew.

Japan, North Asia, Himalaya, Tibet, Java.

2. *Muehlenbergia japonica*, *Steud. Syn. Gram.* p. 422; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 647, sér. 2, iii. (1903) p. 502; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 165.

COREA: Roadsides near Fusan (*Faurie*, 805, fide *Hackel*).

Japan.

I have seen no specimen.

1. *Polypogon littoralis*, *Smith*, *Compend. Fl. Brit.* p. 13; *Steud. Syn. Gram.* p. 184; *Benth. Fl. Hongk.* p. 427; *Debeaux, Fl. Shanghai*, p. 70; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 131; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 167; *Hook. f. Fl. Brit. Ind.* p. 246.

7 — KIANGSU: Shanghai (*Faber*, 306! *Maingay*, 720!); FOKIEN (*Grijs*, in herb. *Hance*, 6681 in part!); SHENSI: Kusan (*Hugh*!); HUPEH: Ichang (*A. Henry*, 190!); YUNNAN: Mengtze, beside streamlet at 4500 feet (*A. Henry*, 10556!), ditto in dry rice-field at 4500 feet (*A. Henry*, 10246!), ditto on rocks in streamlet at 4600 feet (*A. Henry*, 10246 A!); KWANGTUNG: Canton (*Parry*, in herb. *Hance*, 6681 in part!); HONGKONG (*Harland*, 711!); COREA: Port Hamilton (*Wilford*, 702!); CHINA: East Coast (*Home*!). Mus. Brit.; Herb. Kew.

Japan, Himalaya; temperate and subtropical regions.

2. *Polypogon monspeliensis*, *Desf. Fl. Atl.* i. p. 67; *Steud. Syn. Gram.* p. 184; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Franchet, in Pl. David.* i. p. 332; *Kanitz, in Math. Naturw. Ber. Ungarn*, iii. (1884) p. 8; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 167; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648; *Hook. f. Fl. Brit. Ind.* vii. p. 245.

KIANGSU: Fengwangshan (*Forbes*, 488!), Woosing River, Shanghai (*Perry*!); FOKIEN (*Grijs*, in herb. *Hance*, 3449!); FORMOSA: Tamsui (*Oldham*, 610!); CHINA: no locality (*Oldham*, 936!). Mus. Brit.; Herb. Kew.

Japan, India, and widely spread in tropical and temperate regions.

1. *Garnotia patula*, *Munro, ex Benth. Fl. Hongk.* p. 416.

*Garnotia drymeia*, *Hance, in Ann. Sci. Nat. sér. 4*, xviii. (1862) p. 233.

*Berghausia patula*, *Munro, in Proc. Amer. Acad.* iv. (1860) p. 362.

*Miquelia barbulata*, *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 178.

HONGKONG (*Hance*, 1009 in part! *Harland*, 692, in herb. *Hance*, 1009 in part! *Wright*!); KWANGTUNG: Canton (*Sampson*, 1783! and in herb. *Hance*, 8135!). Mus. Brit.; Herb. Kew.

Var. *mutica*, *Rendle*.

*Garnotia stricta*, *Benth.* *Fl. Hongk.* p. 416, non *Brongn.*

*Berghausia mutica*, *Munro*, in *Proc. Amer. Acad.* iv. (1860) p. 362.

HONGKONG (*Wright*!). Herb. Kew.

1. *Sphærocaryum elegans*, *Nees*, ex *Steud.* *Nom.* ed. 2, ii. p. 620; *Hook. f.* *Fl. Brit. Ind.* vii. p. 246.

*Graya elegans*, *Nees*, ex *Steud.* *Syn. Gram.* p. 119.

*Isachne pulchella*, *Roth*, *Nov. Pl. Sp.* p. 58; *Benth.* *Fl. Hongk.* p. 414.

*Panicum pulchellum*, *Spreng.* *Syst.* i. p. 322.

SZECHUEN: Mt. Uomisan, Tcentosen (*Hugh*!); KWANGTUNG: swamps near Canton (*Simpson*, 1203!); HONGKONG (*Harland*, 718, in herb. *Hance*, 1005!), waste places, Government Gardens (*Lamont*!). Mus. Brit.; Herb. Kew.

Hills of Northern India, Ceylon, Mergui, Malacca, Singapore.

1. *Sporobolus diandrus*, *Beauv.* *Agrost.* p. 26; *Hook. f.* *Fl. Brit. Ind.* vii. p. 247.

*Agrostis diandra*, *Retz.* *Obs.* v. p. 19.

*Vilfa diandra*, *Steud.* *Syn. Gram.* p. 155.

FORMOSA (*Campbell*!), Apes' Hill, summit 1100 feet (*A. Henry*, 1106!), Takow (*Playfair*! *A. Henry*, 1044!), South Cape (*Schmü-  
rer*, in coll. *A. Henry*, 625!); KWANGTUNG: Macao (*Abel*!), Pakhoi (*Playfair*!). Mus. Brit.; Herb. Kew.

India, Malaya.

2. *Sporobolus Hancei*, *Rendle*, n. sp.

*Herba* perennis cæspitosa, glabra, 8-13 poll. alta; culmus erectus vel basi geniculato-ascendens, 3-nodus, basi ad  $\frac{2}{5}$  lin. crassus, a foliorum vaginis pæne ad paniculæ basin arcte velatus. *Folia* basalia linearia, superne attenuata, plus minus convoluta, ad 5 poll. longa, vix rigidula; caulina ad  $1\frac{2}{3}$  poll. longa, e basi ad apicem acutum attenuata, marginibus involutis, rigidula;

ligula obsoleta. *Panicula*  $2\frac{1}{4}$  ad  $3\frac{3}{4}$  poll. longa, ramis brevibus (ad  $\frac{1}{2}$  poll. longis), tenuibus, flexuosis, e basi spiculiferis, interdum suberectis paniculam densam ad  $\frac{1}{4}$  poll. crassam formantibus, interdum subpatentibus; rachis et ramuli leves; spiculæ lanceolatae, 1 lin. longæ, sub medio purpureo-tinctæ, breviter pedicellatæ. *Gluma* i. lanceolata, acuta, enervis, quam secunda et tertia paullo brevior ( $\frac{7}{8}$  lin. long.); gl. ii. ovata, subacuta, uninervis, 1 lin. long.; gl. iii. ovata, subobtusata, uninervis, secundam æquans; palea paullo brevior; antheræ 3, atro-purpureæ,  $\frac{1}{2}$  lin. long. *Semen* ellipticum, leve, rubrum,  $\frac{1}{2}$  lin. longum.

FOKIEN: Hills, Amoy (*Sampson*, in herb. *Hance*, 13055 in part!); KWANGTUNG: Canton (*Hance*, 13055 in part!). Mus. Brit.; Herb. Kew.

Near *Sporobolus orientalis*, Kunth, but differing in habit, especially in the absence of stolons, in the sometimes effuse panicle, and the relatively longer acute lowermost barren glume.

3. *Sporobolus indicus*, *R. Br. Prodr.* p. 170; *Benth. Fl. Hongk.* p. 426; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Hack. in Bull. Herb. Boiss. sér. 2, iii.* (1903) p. 502; *Hook. f. Fl. Brit. Ind.* vii. p. 247.

*Sporobolus elongatus*, *Franchet, Pl. David.* i. p. 333, non *R. Br.*; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 166.

*Agrostis indica*, *Linn. Sp. Pl.* p. 63; *Loureiro, Fl. Cochinch.* p. 49; *Osbeck, Resa*, p. 215.

*Vilfa elongata*, *Beauv. Agrost.* p. 181; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 248; *Hook. & Arn. l. c.* p. 274.

*Vilfa indica*, *Steud. Syn. Gram.* p. 162.

KIANGSU: above Changchau, Yangtzekiang (*Faber*, 1183!); FORMOSA: Apes' Hill, Takow (*A. Henry*, 1030!); HUPEH (*A. Henry*, 4287!), Ichang (*A. Henry*, 889!), Patung (*A. Henry*, 2510!); KWANGTUNG: Whampoa (*Robertson*!), Canton (*Sampson*, 306!); HONGKONG: Mt. Victoria (*Lamont*! *Hance*, 1114! *Wright*!); LUCHU ARCHIPELAGO (*Wright*!); CHINA: without locality (*Seemann*! *Reeves*!). Mus. Brit.; Herb. Kew.

Japan, India, and tropics of both hemispheres.

4. *Sporobolus japonicus*, *Maxim. MS.*

*Sporobolus ciliatus*, *Presl*, var. *japonicus*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648.

*Agrostis japonica*, *Steud. Syn. Gram.* p. 171; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 277; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 167.

KIANGSI: Kiukiang, Nankangfu (*Reid!*). Herb. Kew.

Japan.

5. *Sporobolus virginicus*, *Kunth, Rév. Gram.* i. p. 67; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 108; *Hook. f. Fl. Brit. Ind.* vii. p. 249.

*Agrostis virginica*, *Linn. Sp. Pl.* p. 63.

FOKIEN: Amoy, on the sea-shore (*Sampson, in herb. Hance, 12295!*); FORMOSA: Takow (*A. Henry, 1055, 1067, 1844!*), Apes' Hill (*Playfair, 150!*). Mus. Brit.; Herb. Kew.

Ceylon, Java, Australia, Tropical and South Africa, and America.

1. *Agrostis alba*, *Linn. Sp. Pl.* p. 63; *Steud. Syn. Gram.* p. 167; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 649; *Hook. f. Fl. Brit. Ind.* vii. p. 254; *Franchet, Pl. David.* ii. p. 142.

SZECHUEN: Moupin (*David, fide Franchet*).

Japan, Himalaya, Central Asia, Europe; introduced in America.

I have seen no specimen.

2. *Agrostis canina*, *Linn. Sp. Pl.* p. 62; *Steud. Syn. Gram.* p. 165; *Maxim. Primit. Fl. Amur.* p. 325; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 276; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 649, et sér. 2, iii. (1903) p. 502; *Hook. f. Fl. Brit. Ind.* vii. p. 255.

FORMOSA: Tamsui (*Oldham, 609!*); HUPEH: Ichang (*A. Henry, 245, 3464!*). Herb. Kew.

Japan, Himalaya, and North temperate regions generally.

In Dr. Henry's specimen no. 245 the pale is one-third the length of the fertile glume, and the callus bears very short lateral tufts of stiffish hairs, scarcely  $\frac{1}{10}$  of a line long.

3. *Agrostis Hugoniana*, *Rendle, n. sp.*

*Herba* parva, cæspitosa, perennis, rhizomate tenui fragili, radicibus dense fibrosis; culmi 3-4½ poll. longi, subcompressi, basi ad 1½ lin. crassi, 2-nodi, internodium inferius breve, basi vaginis membranaceis marcidis obtectum, superius elongatum, vaginam

laxam glabram sæpius excedente. *Foliorum* ligula scariosa, truncata vel obtusa,  $\frac{3}{4}$  ad 1 lin. longa; lamina culmi vix 2 poll. longa, 1 vel  $1\frac{1}{2}$  lin. lata, linearis superne attenuata, glabra, innovationum longior, ad  $4\frac{1}{2}$  poll. *Panicula* densa, lineari-oblongata vel lineari-oblonga,  $1\frac{1}{2}$ - $2\frac{1}{2}$  in. longa ad  $\frac{1}{2}$  in. lata; rami erecti, terni (ad nodos superiores bini et solitarii), laterales e basi ramulosi, ut ramuli pedicellique leves; pedicellus quam spicula brevior; spiculæ  $1\frac{3}{4}$  lin. longæ, lanceolatæ, pallide virides sæpe purpureo-tinctæ. *Glumæ* steriles subæquales, ovato-lanceolatæ, acutæ, 1-nerves, carina exterioris supra basin hirsutula; gl. fertilis paullo brevior, perlucida ( $1\frac{1}{2}$  lin. longa), cum venis 5 inconspicuis; palea brevis, circiter  $\frac{1}{4}$  lin. longa; anthera oblonga,  $\frac{1}{2}$  lin. longa; processus rhachillæ absens; callus cum pilis brevissimis (vix  $\frac{1}{4}$  lin.) instructus.

SHENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

This species recalls the Northern Indian *Agrostis inæquiglumis*, Griseb., but is of more robust habit, with much larger spikelets, broader leaves, &c.

4. *Agrostis micrantha*, *Steud. Syn. Gram.* p. 170; *Hook. f. Fl. Brit. Ind.* vii. p. 256.

*Agrostis alba*, *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388, non *Linn.*

SHINGKING: Chienshan (*Ross*, 428, 581!); HUPEH: Patung (*A. Henry*, 4722, 4723!); YUNNAN: Mengtze forest, at 5500 ft. (*A. Henry*, 11257!). Mus. Brit.; Herb. Kew.

Himalaya.

5. *Agrostis myriantha*, *Hook. f. Fl. Brit. Ind.* vii. p. 257.

HUPEH (*A. Henry*, 4036, 4698!). Mus. Brit.; Herb. Kew.

Himalaya.

6. *Agrostis perennans*, *Tuck. in Amer. Journ. Sci.* ser. 2, xlv. (1843) p. 44; *A. Gray, Man. Bot.* ed. 5, p. 611; *Steud. Syn. Gram.* p. 165; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 132; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 277; *Franch. et Savat. Enum. Pl. Jap.* p. 166; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648.

HUPEH: Patung (*A. Henry*, 4034, 4697!); COREA: Soul (*Sontag!*). Mus. Brit.; Herb. Kew.

Japan; Eastern United States.



7. *Agrostis scabra*, Willd. *Sp. Pl.* i. p. 370; *Steud. Syn. Gram.* p. 164; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 648, et sér. 2, iii. (1903) p. 502.

*Agrostis laxiflora*, R. Br. in *Richards, App. Frankl. Voy.* p. 3; *Franchet, Pl. David.* i. p. 332; *Regel, Tent. Fl. Ussur.* p. 170; *Turez. Fl. Baic.-Dahur.* ii. p. 305; *Maxim. Prim. Fl. Amur.* p. 325.

CHIHLI: environs of Peking (*David, fide Franchet*); COREA: meadows near Kanouento (*Faurie, fide Hackel*).

Japan, North Asia; North America.

I have seen no specimen.

8. *Agrostis vulgaris*, *With.*, var. *ligularis*, *Franchet, Pl. David.* ii. p. 142.

SZECHUEN: Moupin (*David, fide Franchet*).

I have seen no specimen.

1. *Calamagrostis epigejos*, *Roth, Tent. Fl. Germ.* i. p. 34; *Steud. Syn. Gram.* p. 187; *Franchet, Pl. David.* i. p. 331, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 268; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 224; *Regel, Tent. Fl. Ussur.* p. 170; *Maxim. Primit. Fl. Amur.* p. 324; *Turez. Fl. Baic.-Dahur.* ii. p. 310.

Var. *densiflora*, *Ledeb. Fl. Altaic.* i. p. 87; *Franchet, Pl. David.* i. p. 332; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 649, et sér. 2, iii. (1903) p. 502; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 132; *Komarow, ibid.* xx. (1901) p. 276.

*Arundo epigejos*, *Linn. Sp. Pl.* p. 81; *Loureir. Fl. Cochinch.* p. 45.

CHIHLI: Jehol (*David, 1953!*), mountains near Tachiaosze (*Moellendorff!*), Peking (*Bretschneider, 847!*); SHINGKING: Moukden to Yaloo River (*Webster!*); HUPEH (*A. Henry, 7371!*). Patung (*Wilson, 1656! A. Henry, 4722!*), Ichang (*A. Henry, 3967!*), Changyang (*A. Henry, 6213!*); SHENSI: Kusan, Kisan-shien (*Hugh!*); KWANGTUNG: Lofaushan, 3100 feet (*Ford, 17!*); COREA (*Veitch!*); COREAN ARCHIPELAGO (*Oldham, 943!*). Mus. Brit.; Herb. Kew.

Japan and temperate Asia; Europe; South Africa.

2. *Calamagrostis Langsdorffii*, *Trin. De Gram. Unifl. &c.* p. 225, tab. iv. fig. 10; *Franchet, Pl. David.* i. p. 332.

*Calamagrostis villosa*, *Mutel*, var. *Langsdorffii*, *Hack. in Sommer, Fl. Ob. infer.* p. 107, et *Bull. Herb. Boiss.* vii. (1899) p. 650.

CHIHLI: environs of Pohnashan (*David, fide Franchet*).

Japan.

I have seen no specimen.

3. *Calamagrostis Onoei*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 198; *Franchet, Pl. David.* i. p. 332; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 650,

CHIHLI: borders of Lanho, Jehol (*David, fide Franchet*).

Japan.

I have seen no specimen.

4. *Calamagrostis villosa*, *Mutel, Fl. Fr.* p. 41; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 650, et sér. 2, iii. (1903) p. 502; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 277.

COREA: damp grassy places below Kanouento Mountains (*Faurie, 850, fide Hackel*).

Japan.

I have seen no specimen.

1. *Deyeuxia effusiflora*, *Rendle, n. sp.*

*Folia* . . . . . *Panicula* diffusa, ad 14 poll. longa, rhachis scabridula, nodi inferiores distantes, rami 4 ad 5 poll. longi, tenues, flexuosi, subpatentes vel superiores ascendentes, ad nodum quemque sæpe quini lateraliter fasciculati, interdum e basi divisi, sæpius ad  $\frac{1}{4}$ – $\frac{1}{2}$  longitudinis nudi, ut ramuli longi capillares et pedicelli scabriduli; pedicelli quam spiculæ breviores; spiculæ purpureo-tinctæ, lanceolatae,  $1\frac{3}{4}$  lin. long. *Glumæ* steriles rigidulæ, scaberulæ; gl. i. lanceolata, obtusa, uninervis, nervo in  $\frac{2}{3}$  superiore hirsutulo; gl. ii. paullo brevior, elliptica, obtusa, 3-nervis; gl. fertilis quam steriles paullo brevior, ovata, inconspicue 4-nervis, apice perlucida 4-denticulata; arista basalis, geniculata,  $2\frac{1}{2}$  lin. longa; pili in callo laterales vix  $\frac{1}{2}$  lin. longi; palea quam gluma paullulo brevior; processus rhachillæ barbatus cum pilis vix 1 lin. longus.

SZECHUEN (*Rockhill*!). *Herb. Kew.*

Approaches *D. hupehensis*, mihi, in the characters of the spikelets, which, however, are less rigid and not scaberulous in that species, but differs strikingly in the large, laxly-branching, spreading panicle.

2. *Deyeuxia elatior*, *Hook. f. Fl. Brit. Ind.* vii. p. 266.

*Calamagrostis scabrescens*, var. *elatior*, *Griseb. in Goett. Nachr.* 1868, p. 79.

CHIHLI: between Peking and Jehol (*Staunton*!). Mus. Brit. Khasia Hills.

Differs from the Indian specimens in the much shorter hairs on the callus.

3. *Deyeuxia Henryi*, *Rendle*, n. sp.

*Culmus* elatus 5-pedalis, basi vix 2 lin. crass., 3-nodus. *Foliorum* vaginæ elongatæ, 9-15 poll. longæ (vagina suprema inferiores excedens), quam internodia breviores, sublaxæ, marginibus et in striis longitudinalibus breviter pilosæ; ligula ovata, obtusa, 4-4½ lin. longa; lamina linearis, basi in nervo mediano valido plicata, superne longe attenuata, bipedalis, 3 lin. lata, utrinque scabridula. *Panicula* vix pedalis, oblanceolata, densiuscula at superne laxior, culmus sub panicula, rhachis, et rami scabriduli; rami sæpius quini, ad nodos inferiores ascendentes, supra paniculæ medium subpatentes, infra medium ramulosi, ramuli et pedicelli breves scabriduli; spiculæ 2 lin. longæ, purpureo-tinctæ. *Gluma* i. secundam paullulo excedens, ovato-lanceolata, acuta, uninervis, in nervo scabridula; gl. ii. late ovata obtusiuscula, 3-nervis, glabra; gl. fertilis steriles subæquans, late ovata, 4-nervis, apice truncata denticulata, arista basalis, geniculata, 3 lin. longa, pili laterales in callo minus quam 1 lin. longi; palea quam gluma paullo brevior; antheræ purpureæ, 1 lin. longæ; processus rhachillæ barbatus, cum pilis 1¼ lin. longus.

HUPEH: Fang (*A. Henry*, 6724!). Mus. Brit.; Herb. Kew.

This species has the general habit of *Deyeuxia sylvatica*, but is distinguished by its blunt spikelets, subequal glumes, and form of panicle.

4. *Deyeuxia Hugoniana*, *Rendle*, n. sp.

*Herba* perennis, cæspitosa; culmi aggregati, ad 10 pollicares longi, cum internodiis basalibus brevibus tenuibus cylindricis, ad nodum supremum autem geniculati, tum erecti internodio elongato ad 8 poll. longo compresso, viride, glabro sed sub panicula scabridulo. *Folia* erecta, vaginæ glabræ, internodia inferiora tegentes, suprema circa 2 poll. longa, ad internodii basin arcte applicita; ligula brevis, obtusa, rotundata, glabra, ½ lin. longa; lamina linearis, superne attenuata, acutiuscula, plana vel con-

duplicata, glabra, folii supremi  $3\frac{1}{2}$  ad 4 poll. longa, ad 1 lin. lata. *Panicula* laxa, sublanceolata vel ovata, 3 ad 4 poll. longa,  $\frac{1}{2}$  ad  $1\frac{1}{4}$  poll. lata; rami ad quemque nodum bini, demum sæpius patentés, flexuosi, in parte  $\frac{1}{2}$  ad  $\frac{1}{3}$  inferiore nudi, ramus longior ad  $1\frac{1}{2}$  poll. longus, sparse ramulosus, rami, velut rhachi, ramulis et pedicellis scabridi; spiculæ vix 2 lin. longæ, purpureo-tinctæ. *Glumæ steriles* lanceolatæ, acutæ; gl. i.  $1\frac{1}{4}$  ad  $1\frac{1}{2}$  lin. longa, infra medium 3-nervis, carina supra medium scabridula; gl. ii. quam prima paullo longior, 3-nervis, nervi laterales medium superantes, carina in parte  $\frac{2}{3}$  superiore scabridula; gl. fertilis ovata (quam aperta) obtusiuscula, gl. secundam subæquans, imperfecte 5-nervis, carina in parte  $\frac{2}{3}$  superiore scabridula in parte inferiore pilosa; callus breviter pilosus, palea quam gluma paullo brevior; antheræ  $\frac{2}{3}$  lin. longæ; rhachillæ processus glaber vel glaberulus, vix 1 lin. longus, interdum glumæ rudimentum minutum gerens.

SHENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

A very distinct species characterized by the absence of an awn on the fertile glume, and the glabrous continuation of the rhachilla.

5. *Deyeuxia hupehensis*, *Rendle*, n. sp.

*Herba* cæspitosa; culmi aggregati, tenues, glabri, 2-nodi, geniculati, pedales, basi  $\frac{1}{2}$ - $\frac{2}{3}$  lin. crassi. *Folia* rigidula, suberecta; vaginæ striatæ, haud scabridæ, margine superiore et ore breviter pilosæ, internodia haud æquantes; ligula brevis,  $\frac{1}{2}$ -1 lin. longa; lamina linearis, basi attenuata, superne tenuiter acuminata, induplicata, in facie superiore pilosula, dorso scabridula, ad 10 poll. longa, et 2 lin. lata. *Panicula* vix semipedalis, lanceolata, vix densa, 1 poll. lata; rhachis tenuis, levis; rami numerosi, fasciculati, in  $\frac{1}{3}$  inferiore ramulosi, erecto-patentes, ramuli capillares, ut pedicelli minute scabriduli; pedicellus spiculam subæquans vel paullo brevior; spiculæ pallide virides, purpureo-tinctæ, vix 2 lin. longæ. *Glumæ* membranaceæ; gl. i. lanceolata, acuta, 1-nervis, margine ciliolata, nervo scabridula; gl. ii. paullo brevior, elliptica, obtusa vel breviter et abrupte subacuta, nervis tribus parallelis viridibus lineata; gl. fertilis secundam subæquans, 4-nervis, apice denticulata, arista basalis, geniculata, vix 3 lin. longa, pili laterales in callo  $\frac{1}{4}$  lin. longi, palea quam gluma paullo vel vix brevior; antheræ purpuræ, 1 lin. longæ; rhachillæ processus barbatus, cum pilis  $\frac{3}{4}$  lin. longus.

HUPEH (*A. Henry*, 6726!). Mus. Brit.; Herb. Kew.

This species seems nearest to *Calamagrostis hakonensis*,

Franchet & Savat., but differs in habit, having much fewer stem-nodes, also in the longer ligule, the long-exserted awn of the fertile glume, and the very short hairs on its callus.

6. *Deyeuxia moupinensis*, *Pilger, in Engl. Jahrb.* xxix. (1900) p. 225.

*Calamagrostis moupinensis*, *Franchet, Pl. David.* ii. p. 144.

SZECHUEN: Moupin (*David, fide Franchet*).

I have seen no specimen.

7. *Deyeuxia pulchella*, *Hook. f. Fl. Brit. Ind.* vii. p. 268.

*Calamagrostis pulchella*, *Griseb. in Goett. Nachr.* 1868, p. 78.

SZECHUEN: Tachienlu (*Pratt, 553!*). Mus. Brit.; Herb. Kew.

Himalaya, Tibet.

8. *Deyeuxia sachalinensis*, *Rendle*.

*Calamagrostis sachalinensis*, *Fr. Schmidt, Reisen in Amurl.* p. 203; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 650, et sér. 2, iii. (1903) p. 502.

COREA: near Naipiang (*Faurie, 874, fide Hackel*).

North Asia.

I have not seen the specimen.

9. *Deyeuxia scabrescens*, *Munro, ex Duthie, Grasses N.W. India*, p. 31; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Hook. f. Fl. Brit. Ind.* vii. p. 267.

*Calamagrostis scabrescens*, *Griseb. in Goett. Nachr.* 1868, p. 79; *Franchet, Pl. David.* ii. p. 143.

SZECHUEN: Tongolo (*Soulié, 263!*), summit of Mt. Omei (*Faber, 1168!*). Herb. Kew.

Himalayas.

10. *Deyeuxia sylvatica*, *Kunth, Rév. Gram.* i. p. 77, et *Enum.* i. p. 243; *Hook. f. Fl. Brit. Ind.* vii. p. 266.

*Calamagrostis arundinacea*, *Roth, Tent. Fl. Germ.* i. p. 33; *Hance, in Journ. Bot.* xvi. (1878) p. 234; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 652, et sér. 2, iii. (1903) p. 502; *Regel, Tent. Fl. Ussur.* p. 169; *Maxim. Fl. As. Or. Fragm.* p. 70.

*Calamagrostis sylvatica*, *DC. Fl. Franç.* v. p. 253; *Ledeb. Fl. Ross.* iv. p. 426; *Turcz. Fl. Baic.-Dah.* ii. p. 314; *Maxim. Prim. Fl. Amur.* p. 323.

CHIHLI: Mt. Miaofengshan (*Moellendorff, in herb. Hance, 17528 in part!*); HUPEH: Fang (*A. Henry, 6727!*), Patung (*A. Henry, 2564, 5052, 7411!*); SHENSI: Mt. Laoyuan (*Hugh!*), Mt. Miaowansan (*Hugh!*), Mt. Kiansan, Husiencen (*Hugh!*);

SZECHUEN: North Wushan, 4000-7000 feet (*A. Henry*, 7043!); KWANGTUNG: near River Lienchan (*Nevin*, in herb. *Hance*, 17528 in part!). Mus. Brit.; Herb. Kew.

Himalayas, Japan, Temperate Asia, Europe.

Var. *borealis*, *Rendle*, var. nov. Culmi ad 2-pedales, basi 1 lin. crass., 3-nodi. Foliorum vaginæ glabræ, striatæ, vix scabridulæ, sublaxæ; ligula ovata, 2-5 lin. longa; lamina ut in var. *ligulata* sed minus scabridula. Panicula densiuscula, ramis brevibus ascendentibus, lanceolata, 8-9 poll. longa, ad 7 lin. lata; spiculæ 3 lin. longæ. Gluma i. secundam paullo ( $\frac{1}{4}$ - $\frac{1}{2}$  lin.) excedens; gl. fertilis c.  $2\frac{1}{2}$  lin. longa, arista basalis c. 4 lin. longa, pili laterales in callo  $\frac{1}{2}$  lin. longi; palea 2 lin. longa; rhachilla cum pilis c.  $1\frac{1}{2}$  lin. longa.

SHANTUNG: Chefoo (*Forbes*, 397! *Maingay*, 63!). Herb. Kew.

Near var. *ligulata*, but distinguished by its less strict and laxer panicle and larger spikelets.

Var. *brachytricha*, *Hack.* in *Bull. Herb. Boiss.* vii. (1899) p. 652.

*Calamagrostis brachytricha*, *Steud. Syn. Gram.* p. 189; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 91; *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 169.

CHIHLI: Peking (*Williams*, in herb. *Hance*, 12793 in part!); SHINGKING (*Ross*, 130!), Newchang (*Shearer*!); HUPEH, 7000-9500 ft. (*A. Henry*, 6792!), Ichang (*A. Henry*, 2745!), Tsien Mountains (*Faber*, 1528!). Mus. Brit.; Herb. Kew.

Japan.

I find it difficult to distinguish this variety from the type-species.

Var. *collina*, *Rendle*.

*Deyeuxia collina*, *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 225.

*Calamagrostis collina*, *Franchet, Pl. David.* ii. p. 143.

SZECHUEN: Moupin, on sunny hills (*David*, fide *Franchet*).

I have not seen a specimen of this plant, but do not find in the description anything by which to separate it specifically from *Deyeuxia sylvatica*. The author says that it has the habit of *Calamagrostis sciuroides*, from which it differs in having the culms smooth below the panicle, the tuft of hairs on the callus shorter, equalling scarcely a third of the length of the flower,

and the sheaths glabrous at the top. The ligule is short, the fertile glume and palea are equal in length and hardly shorter than the barren glumes; the spikelets are barely 2 lines long.

Var. *hirsuta*, Hack. in *Bull. Herb. Boiss*, sér. 2, iii. (1903) p. 502.

COREA: near Fusan (*Faurie*, 851, fide *Hackel*).

I have not seen the specimen.

Var. *latifolia*, Rendle, var. nov. *Planta* robusta; culmus 3-pedalis, basi vix 2 lin. crass., 3-nodus. *Foliorum* vaginæ pilosæ; ligula brevis obtusa, ad  $1\frac{1}{2}$  lin. longa; lamina elongata, lineari-lanceolata, 2-pedalis et ultra,  $\frac{1}{2}$  poll. lata, in parte inferiore plicata, angustata, superne in apicem aristiformem attenuata; in facie superiore scabridula, cum nervo mediano præcipue in parte inferiore eminente. *Panicula* pedalis, stricta, densa, ad  $\frac{3}{4}$  poll. lata; spiculæ  $2\frac{1}{2}$ – $2\frac{3}{4}$  lin. longæ. *Glumæ* steriles æquales, gl. i. uninervis, ii. trinervis, venis conspicuis, hirsutulis; gl. fertilis  $2\frac{1}{4}$ – $2\frac{1}{2}$  lin. longa, cum arista basali  $3\frac{1}{2}$  lin. longa, palea paullo brevior, pili laterales in callo  $\frac{1}{2}$  lin. longi, rhachillæ continuatio cum pilis insidentibus  $1\frac{1}{4}$  lin. longa.

CHIHLI: Peking (*Bretschneider*, 851–2! *Moellendorff*, in herb. *Hance*, 12793 in part!). Mus. Brit.; Herb. Kew.

Characterized by its robust habit, very long broad leaves with hairy sheaths and short ligules, and long slender dense panicles. Evidently very near *Calamagrostis robusta*, Franchet et Savat., which from the description resembles it in habit, size of spikelets, &c., but is described as having a ligule nearly 1 cm. long, and a very minute oblong glabrous (?) rudiment of a second flower.

Var. *laxiflora*, Rendle, var. nov. *Folia* innovationum e vaginis marcidis rufescentibus dense involutis orientia, anguste linearia, superne longe attenuata, plana, sesquipedalia, 1 lin. lata, rigidula, marginibus scabridulis; folia culmi linearia, scabridula; ligula brevis, 1 lin. long. *Panicula* 7–9 poll. long., pyramidata, rami ad nodi inferiora quini patentes, in parte inferiore nudi, superne laxiflori; spiculæ  $2\frac{1}{4}$  lin. long. *Gluma* i. secundam paullo excedens; gl. fertilis 2 lin. longa, paleam paullo ( $\frac{1}{4}$  lin.) excedens, arista basalis 2 lin. longa, rhachilla cum pilis  $\frac{3}{4}$  lin. longa; pili laterales in callo  $\frac{1}{2}$  lin. longi.

HUPEH: Ichang (*A. Henry*, 978!). Herb. Kew.

Characterized by the long narrow linear leaves of the sterile shoots, which are surrounded at the base by a dense covering of reddish sheath-remains, and the habit of the inflorescence with its pyramidal outline and laxly-spreading lower branches.

Var. *ligulata*, Rendle, var. nov. Culmus  $2\frac{1}{2}$ -pedalis, parte inferiore  $1\frac{1}{2}$  lin. crass., 3-nodus. Foliorum vaginae glabræ, scabridulo-striatæ; ligula ovata, ad  $3\frac{1}{2}$  lin. longa; lamina linearis, sub medio angustata et plicata, superne longe attenuata, ad sesquipedalis longa,  $2\frac{1}{2}$  lin. lata, utrinque scabrida. Panicula stricta, densa, 9 poll. longa; spiculæ 2- $2\frac{1}{2}$  lin. longæ. Gluma i. secundam paullo excedens; gl. fertilis sterilibus c.  $\frac{1}{2}$  lin. brevior (sæpe 2 lin. longa); arista basalis  $2\frac{1}{2}$  lin. longa; pili laterales in callo c. 1 lin. longi; palea glumam subæquans; rhachilla cum pilis paleam subæquans vel paullo brevior.

HUPEH (*A. Henry*, 5021!). Mus. Brit.; Herb. Kew.

Characterized by its long ligule, its dense strict panicle, and by having the rhachilla-continuation with its hair-tuft about equal to the pale, and the lateral tufts of hairs on the callus barely half the length of the glume.

Var. *sciuroides*, Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 652.

*Calamagrostis sciuroides*, Franch. et Savat. *Enum. Pl. Jap.* ii. p. 600.

CHIHLI: hills in neighbourhood of Jehol (*David*, 2182!), dry hills and edges of fields in Peking district (*Williams*, in herb. *Hance*, 12793 in part!). Mus. Brit.; Herb. Kew.

Japan.

Distinguished from the species by its longer, more rigid and scabrid spikelets, conspicuous nerves on the upper barren glume, and longer hairs on callus of fertile glume.

#### Tribe AVENÆÆ.

1. *Eriachne pallescens*, R. Br. *Prodr.* p. 184; *Steud. Syn. Gram.* p. 237; *Benth. Fl. Austr.* vii. p. 630.

*Eriachne Hookeri*, Munro (nomen), in *Journ. Linn. Soc., Bot.* vi. (1862) p. 42; *Hance*, in *Ann. Sci. Nat.* sér. 5, v. (1866) p. 251.

*Eriachne chinensis*, *Hance*, in *Ann. Sci. Nat.* sér. 4, xv. (1861) p. 228, and in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 136.

*Aira seminibus hirsutis*, aristis terminalibus, flore longioribus, *Osbeck, Resa*, p. 220.



KWANGTUNG: French Island (*Osbeck*, in herb. *Linn.*!), hills north of Canton (*Hance*!), hills; Danes Island, Whampoa (*Hance*, 7424 in part! *Ford*, 207!); HONGKONG: tops of mountains (*Hance*, 7424 in part!). Mus. Brit.; Herb. Kew.

India, Malaya, Australia.

1. *Cœlachne pulchella*, *R. Br. Prodr.* p. 187; var. *simpliciuscula*, *Hook. f. Fl. Brit. Ind.* vii. p. 271.

*Cœlachne pulchella*, *Benth. Fl. Hongk.* p. 430.

*Panicum simpliciusculum*, *Wight & Arn. ex Steud. Syn. Gram.* p. 96.

SZETCHUEN: Tcentosen, Mt. Uomisan (*Hugh*!); KWANGTUNG: swampy ground near Canton (*Sampson*, 1144!), pools at the foot of Mt. Pakwan, outside Cantou (*Sampson*, in herb. *Hance*, 8142!), rice-field Kowlung, mainland opposite Hongkong (*Lamont*!); HONGKONG: ditch, Little Hongkong (*Lamont*! *Wilford*, 253! *Fuber*, 279!). Mus. Brit.; Herb. Kew.

India, Ceylon, Malaya, Madagascar.

1. *Deschampsia cæspitosa*, *Beauv. Agrost.* p. 91, t. xviii. fig. 3; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 702; *Maxim. Prim. Fl. Amur.* p. 323; *Turcz. Fl. Baic.-Dahur.* ii. p. 318; *Hook. f. Fl. Brit. Ind.* vii. p. 273.

*Aira cæspitosa*, *Linn. Sp. Pl.* p. 64; *Steud. Syn. Gram.* p. 219; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 283.

SZETCHUEN: Tongolo (*Soulié*, 703, 732!). Herb. Kew.

Japan, Himalayas, and temperate and cold regions of Northern and Southern hemispheres; alpine in Tropics.

1. *Trisetum flavescens*, *Beauv. Agrost.* p. 88; *Steud. Syn. Gram.* p. 226; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 702; *Turcz. Fl. Baic.-Dahur.* ii. p. 319; *Maxim. Prim. Fl. Amur.* p. 323; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 283.

*Avena flavescens*, *Linn. Sp. Pl.* p. 80; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Hook. f. Fl. Brit. Ind.* vii. p. 279.

SHENSI: Miaowansan (*Hugh*!); HUPEH: Patung (*A. Henry*, 4696, 4027, 2558!); COREAN ARCHIPELAGO: Tsusima Islands, Strait of Corea (*Wilford*, 804!), Port Chusan (*Wilford*, 938!). Mus. Brit.; Herb. Kew.

Japan; Europe, Temperate Asia, North Africa.

Var. *macranthum*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703, et sér. 2, iii. (1903) p. 503.

COREA: Soul (*Faurie*, 807), hills near Chinampo (*Faurie*, 883), Mt. Kanouento (*Faurie*, 896), all *fide* Hackel.

Japan.

I have seen no specimen.

## 2. *Trisetum Henryi*, *Rendle*, n. sp.

*Planta* robusta, elata, culmus 4-5-pedalis, basi plus quam 2 lin. crassus, 8-nodus, internodia glabra polita. *Foliorum* vaginæ, superioribus exceptis, internodia æquantes vel superantes, striatæ, plus minus scabridæ, marginibus et ore, interdum quoque superficie, piliferæ; ligula brevissima breviter pilosa; lamina anguste lineari-lanceolata, apice attenuata ad 1 ped. longa, et  $\frac{1}{2}$  poll. lata, scabridula, cum nervo mediano dorso prominulo in facie superiore interdum sparse pilifera. *Panicula* ovalis, densa, 8 poll. longa, ad  $3\frac{1}{2}$  poll. diametro; rami ad nodum quemque terni vel plurimi, multi-ramulosi, ad paniculæ basin ascendentes, superne curvulopatentes, rami ramulique super basin nudam spiculiferi velut rhachide leves; spiculæ pedicellos sæpius duplo excedentes, bifloræ, iis *T. flavescens* similes sed argenteo-fulvæ, vix 3 lin. longa. *Gluma* i. lanceolata, acuta, 1-nervis,  $1\frac{1}{2}$  lin. longa; gl. ii. ovali-lanceolata, acuta, 3-nervis,  $2\frac{1}{2}$  lin. longa, carina ut in gl. i. supra medium minute scabridula; gl. iii. (plicata) anguste lanceolata, apice acuminata bifida, 3-nervis, dorso scabridula,  $2\frac{1}{2}$  lin. longa, arista tenuis, geniculata, circa tertiam partem ab apice inserta, 3 lin. long.; palea 2 lin. long., carinæ minute scabridulæ; antheræ  $1\frac{1}{2}$  lin. long.; rhachilla inter flores et super florem alterum, breviter pilosa.

HUPEH: Fang (*A. Henry*, 6643, 6706!). Mus. Brit.; Herb. Kew.

Closely allied to *Trisetum flavescens*, but a taller, more robust plant, with denser panicle and silvery-brown coloured spikelets. Except in the somewhat less acute lower barren glume, I find no difference in the details of the spikelet from those of *T. flavescens*.

## 3. *Trisetum subspicatum*, *Beauv. Agrost.* p. 88; *Steud. Syn. Gram.* p. 225; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703.

*Avena subspicata*, *Clairv. Man. Herb.* p. 17; *Hook. f. Fl. Brit. Ind.* vii. p. 278.

SHENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

Japan, Himalayas, and generally distributed in alpine and frigid regions.

1. *Avena Delavayi*, *Hack. in Oest. bot. Zeitschr.* lii. (1902) p. 189.

YUNNAN: "Mountain pastures of Heechanmen (Lankong) at 10,000 feet" (*Delavay*, fide *Hackel*).

I have seen no specimen.

2. *Avena fatua*, *Linn. Sp. Pl.* p. 80; *Steud. Syn. Gram.* p. 230; *Benth. Fl. Hongk.* p. 430; *Hook. & Arn. Bot. Beechey's Voy.* p. 274; *Thunb. Fl. Jap.* p. 54; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Franch. & Sav. Enum. Pl. Jap.* ii. p. 173; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703; *Hook. f. Fl. Brit. Ind.* vii. p. 275.

KIANGSU: Shanghai (*Faber*, 276!); FOKIEN: Foochow, spontaneous (*Gregory*, in herb. *Hance*, 1266 in part!); SZECHUEN: Tongolo (*Soulié*, 249!); HONGKONG (*Hance*, 1266 in part!). Mus. Brit.; Herb. Kew.

Europe, Temperate Asia, North Africa.

Var. *glabrata*, *Stapf, MS. in herb. Kew*; glumæ fertiles glabræ; carum callis breviter et dense hirsutis; rhachis hirsuta.

HUPEH: Ichang, wild at side of wheat-fields &c. (*A. Henry*, 3474!); COREA: Chemulpo (*Carles*, 164!). Mus. Brit.; Herb. Kew.

No. 3474 of Henry is referred by Pilger (in *Engl. Bot. Jahrb.* xxix. p. 225) to *Avena strigosa*, Schreb.

3. *Avena nuda*, *Linn. Amœn. Acad.* iii. p. 401; *Steud. Syn. Gram.* p. 230; *Host, Gram. Austr.* iii. t. 43.

*Avena nuda*, *Linn.*, var. *chinensis*, *Cat. Gor.*, ex *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 71, probably belongs here.

HUPEH (*A. Henry*, 6098!), cultivated oats, Yenmeh of the Chinese, Ichang (*A. Henry*, 3573!). Mus. Brit.; Herb. Kew.

Cultivated.

4. *Avena pratensis*, *Linn. Sp. Pl.* p. 80; *Steud. Syn. Gram.* p. 231; *Franchet, Pl. David.* i. p. 333; *Maxim. Prim. Fl. Amur.* p. 323; *Hook. f. Fl. Brit. Ind.* vii. p. 276.

CHIHLI: Environs of Puhuashan (*David, fide Franchet*).  
Himalayas and Temperate Asia, Europe.  
I have not seen the specimen.

5. *Avena sativa*, *Linn. Sp. Pl.* p. 79; *Steud. Syn. Gram.* p. 230; *Franchet, Pl. David.* i. p. 333; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 287; *Thunb. Fl. Jap.* p. 54; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Franch. & Savat. Enum. Pl. Jap.* ii. p. 173; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703; *Regel, Tent. Fl. Ussur.* p. 169; *Maxim. Prim. Fl. Amur.* p. 323; *Hook. f. Fl. Brit. Ind.* vii. p. 275.

COREAN ARCHIPELAGO (*Oldham, 941!*). Herb. Kew.  
Cultivated.

#### Tribe CHLORIDÆ.

1. *Microchloa setacea*, *R. Br. Prodr.* p. 208; *Steud. Syn. Gram.* p. 202; *Benth. Fl. Hongk.* p. 428; *Hook. f. Fl. Brit. Ind.* vii. p. 283.

*Nardus indica*, *Linn. f. Suppl.* p. 105; *Lour. Fl. Cochinch.* p. 44.

*Rottbœllia setacea*, *Roxb. Corom. Pl.* ii. p. 17, t. 132.

—YUNNAN: rocky mountains 6500 ft., Mengtze (*A. Henry, 10998!*); KWANGTUNG: turfy places at the edges of bare rocks, Fuungkwongshan near Whampoa (*Hance, 13057 in part!*), forms turf on damp lichen-covered rocks, Kulangsu Island, opposite Amoy (*Sampson, in herb. Hance, 13057 in part!*). Mus. Brit.; Herb. Kew.

India; tropics of Old and New Worlds.

1. *Cynodon Dactylon*, *Pers. Syn.* i. p. 85; *Benth. Fl. Hongk.* p. 428; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Franchet, Pl. David.* i. p. 334, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 269; *Debeaux, Fl. Tchefou*, p. 159; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 109; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 172; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703; *Hook. f. Fl. Brit. Ind.* vii. p. 288.

KIANGSU: Shanghai (*Maingay, 556!*); FORMOSA (*Campbell!*), Tamsui (*Oldham, 613!*), Takow (*A. Henry, 1038!*), Apes' Hill (*Playfair, 32!*); HUPEH (*A. Henry, 478!*), Ichang (*A. Henry, 1365!*); SHENSI: Kaolinsien (*Hugh!*), Kisansien (*Hugh!*);

KWANGTUNG: Macao (*Abel!*); HONGKONG: spontaneous (*Hance*, 1069!), Aberdeen (*Lamont!*). Mus. Brit.; Herb. Kew.

India, Japan, and widely distributed in the warmer parts of the earth.

1. *Enteropogon gracilior*, *Rendle*, n. sp.

*Herba* perennis, culmi tenues e basi sæpe repente ascendentes demum erecti, ramosi, floriferi ad sesquipedales, plurinodes, internodia  $\frac{1}{3}$  ad  $\frac{1}{2}$  lin. lata, glabra, inferiora a foliorum vaginis sæpius obtecta, superiora sæpius vaginas paullo excedentia, supremum valde exsertum. *Foliorum* vaginæ cylindricæ, internodiis arcte circumvolutæ, glabræ, ore albido-pilosæ, robustiores dorso carinulatæ; ligula brevissima, pilis mollibus albidis 1 lineis longis instructa; lamina glauca, rigidula, anguste linearis, superne attenuata, plana vel sæpe plus minus convoluta, 2-8 poll. longa, ad  $1\frac{1}{3}$  lin. lata, glabra, dorso scabridula. *Spica* linearis, densa, ad 5 poll. longa, 1 lin. lata; spiculæ sessiles, in seriebus duabus ordinatæ,  $2\frac{1}{4}$  lin. longæ, unifloræ cum flosculo abortivo. *Glumæ* steriles hyalinæ, lanceolatæ, uninerves (nervi virides), persistentes, inferior obtusa,  $\frac{2}{3}$  lin. longa, superior acuta pæne aristulata, vix 2 lin. longa, nervo valido; gl. fertilis, 2 lin. longa, cum marginibus ad nervos laterales inflexis elliptica, 3-nervis, nervi validi, dorso in parte superiore scabriduli, medianus ex apice bilobulato in aristam debilem gluma duplo longiorem prolongatus; callus breviter pilosus; palea quam gluma paullo brevior, carinulis superne scabridulis; flosculus abortivus,  $1\frac{1}{3}$  lin. longus, cum aristula debile,  $\frac{1}{2}$  lin. longa, glumam fertilem paullo superante.

FORMOSA: Takow (*A. Henry*, 1119!), Apes' Hill (*A. Henry*, 1940!). Mus. Brit.; Herb. Kew.

A well-marked species, resembling *Enteropogon melicoides* in habit but less robust, and differing in the slenderer shorter spike, and smaller, not clustered spikelets.

1. *Chloris barbata*, *Sw.* *Fl. Ind. Occ.* i. p. 200; *Steud. Syn. Gram.* p. 204; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 250; *Benth. Fl. Hongk.* p. 429; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 109; *Hook. f. Fl. Brit. Ind.* vii. p. 292.

FORMOSA: Takow, on cliff and near sea-level (*A. Henry*, 726,

1023!), Apes' Hill (*Playfair*, 53!), Pescadores (*Swinhoe*!); KWANGTUNG: British Kowloon (*Ford*, 599!), Macao (*Seemann*!); HONGKONG (*Hance*, 1943!), sea-beach at Saywan (*Wilford*, 246!). Mus. Brit.; Herb. Kew.

India and tropics generally.

2. *Chloris incompleta*, *Roth*, *Nov. Sp.* p. 60; *Steud. Syn. Gram.* p. 207; *Hook. f. Fl. Brit. Ind.* vii. p. 290.

*Chloris digitata*, *Steud. l. c.*

*Gymnopogon digitatus*, *Nees*, *ex Steud. Nom.* ed. 2, i. p. 713; *A. Henry*, in *Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 109.

FORMOSA: Takow (*Campbell*! *A. Henry*, 1151!), Apes' Hill (*Playfair*, 308!). Mus. Brit.; Herb. Kew.

India, Malaya.

3. *Chloris virgata*, *Sw. Fl. Ind. Occ.* i. p. 203; *Steud. Syn. Gram.* p. 205; *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 503; *Hook. f. Fl. Brit. Ind.* vii. p. 291.

*Chloris caudata*, *Trin. in Bunge, Enum. Pl. Chin.* p. 70; *Debeaux, Fl. Tchefou*, p. 158; *Franchet, Pl. David.* i. p. 334, *et in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 268; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 288; *Palibin, ibid.* xxi. (1903) p. 209.

CHIHLI: Tientsin (*Fortune*!), mountains west of Peking (*Bretschneider*!), near Peking (*Williams*, in herb. *Hance*, 11462 in part!) (*Bushell*! *Staunton*! *Pierson*, in herb. *Hance*, 11462 in part!); SHANTUNG: Chefoo (*Forbes*, 400! *Maingay*, 82!); KIANGSU: Hai (*Faber*, 1534!); SHENSI: Kaolinsien (*Hugh*!); NORTH CENTRAL CHINA: Leunteon, Fenghosan (*Hugh*!); HONGKONG (*herb. Munro*, 752!). Mus. Brit.; Herb. Kew.

India and warmer parts of both hemispheres.

1. *Tripogon chinensis*, *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 503.

*Catapodium unilaterale*, *Hance, in Ann. Sci. Nat.* sér. 5, v. (1866) p. 251, non *Griseb.*

*Nardurus filiformis*, var. *chinensis*, *Franchet, Pl. David.* i. p. 339.

CHIHLI (*Staunton*!), Peking (*Bretschneider*, 840! *Bushell*! *Williams*, in herb. *Hance*, 11444!), Sanyu, on the hills (*David*, 2065!); FORMOSA: Apes' Hill, Takow (*A. Henry*, 1064!). Mus. Brit.; Herb. Kew.

The plant cited by Franchet in his 'Catalogue des Plantes de

Tché-fou' (in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. p. 271) as *Festuca filiformis*, Nees, ex Steud. *Syn. Gram.* p. 302, probably belongs here.

Var. *coreensis*, *Hack. l. c.*

COREA: dry hills near Chinampo (*Faurie*, 803, fide *Hackel*).

I have not seen authoritative specimens of the forms which Hackel includes under this species, but judging from his description I think the plants cited above must be placed here. They show a wider variation in size than indicated by Hackel. In Staunton's specimen the culms reach 2 feet in length, while the Formosa plants are not more than 6 inches high.

1. *Beckmannia erucæformis*, *Host, Gram. Austr.* iii. p. 5, t. 6; *Franchet, Pl. David.* i. p. 322; *Maxim. Prim. Fl. Amur.* p. 479; *Debeaux, Fl. Shanghai*, p. 68; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 70; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 134; *Komarow, ibid.* xx. (1901) p. 289.

CHIHLI: Peking (*David*, in herb. *Hance*, 4952! *Bretschneider*, 814!), Jehol (*David*, 1893!); SHINGKING (*Ross*, 44!), Moukden (*Webster*, 107!), between Moukden and Tungcheshien, and between Tungcheshien and Mäorshan, Yalu River (*James*!); KIANGSU: Fengwangshan (*Forbes*, 485!), Shanghai (*Maingay*, 725!), Chapu (*Oldham*, 18!); SZECHUEN: Tongolo (*Soulié*, 248!). *Mus. Brit.*; *Herb. Kew.*

Japan, North Asia, Asia Minor, Eastern Europe, North America.

1. *Eleusine coracana*, *Gaertn. Fruct.* i. p. 8, t. i. fig. 11; *Steud. Syn. Gram.* p. 211; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 249; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 172; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703; *Hook. f. Fl. Brit. Ind.* vii. p. 294.

*Cynosurus coracanus*, *Linn. Syst. Nat.* ed. 11, p. 875; *Thunb. Fl. Jap.* p. 51.

FORMOSA (*Oldham*, 615!); YUNNAN: South Mengtze, mountain forests at 5000 ft., side of streamlet (*A. Henry*, 10962!), Szemao, 4500 ft. (*A. Henry*, 12431!); KWANGTUNG: Lienchau, Canton (*Lamont*!); HAINAN (*A. Henry*, 8284!). *Mus. Brit.*; *Herb. Kew.*

Cultivated in the tropics of the Old World, probably originating from *Eleusine indica*.

2. *Eleusine indica*, *Gaertn. Fruct.* i. p. 8; *Steud. Syn. Gram.* p. 211; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 249; *Benth. Fl. Hongk.* p. 429; *Franchet, Pl. David.* i. p. 334, et in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 268; *Debeaux, Fl. Tchefou*, p. 158; *Hook. f. Fl. Brit. Ind.* vii. p. 293; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 171; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703, et sér. 2, iii. (1903) p. 503.

*Cynosurus indicus*, *Linn. Sp. Pl.* p. 72; *Lour. Fl. Cochinch.* p. 59; *Thunb. Fl. Jap.* p. 52.

KIANGSU: Hai city (*Faber*, 1539!), Shanghai (*Maingay*, 549!); FORMOSA (*Campbell*!), Takow (*A. Henry*, 1124!), Apes' Hill (*A. Henry*, 1013!); NORTH CENTRAL CHINA: Leunteon, Fenghosan (*Hugh*!); SHENSI: in and near water in the plain of Hukiascien (*Hugh*!); SZECHUEN: Omei plain (*Faber*, 1166!); KWANGTUNG: Macao (*Abel*!), Lienchau, Canton (*Lamont*!), Shamien, Canton (*Sampson*, 309!); HONGKONG: wet place, Happy Valley (*Lamont*!); CHINA: without locality (*Seemann*!). *Mus. Brit.*; *Herb. Kew.*

Warmer parts of the Old World.

1. *Dactyloctenium ægyptiacum*, *Willd. Enum. Hort. Berol.* p. 1029; *Steud. Syn. Gram.* p. 212; *Benth. Fl. Hongk.* p. 429.

*Dactyloctenium mucronatum*, *Willd. l. c.*; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 250.

*Eleusine ægyptia*, *Desf. Fl. Atlant.* i. p. 85; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv., *Suppl.* (1896) p. 109 (*ægyptiaca*); *Hook. f. Fl. Brit. Ind.* vii. p. 295 (*ægyptiaca*).

*Cynosurus ægyptius*, *Linn. Sp. Pl.* p. 72; *Lour. Fl. Cochinch.* p. 59; *Osbeck, Resa*, p. 232.

FORMOSA: Takow (*A. Henry*, 755, 1087!), Apes' Hill (*Playfair*, 110!), South Cape (*Schmürer, in coll. A. Henry*, 258!), without locality (*Campbell*!); KWANGTUNG: Whampoa (*Robertson*!); HONGKONG: Robinson road, Victoria (*Lamont*! *Wright*!). *Mus. Brit.*; *Herb. Kew.*

India, and warmer parts of the Old World.



1. *Leptochloa chinensis*, *Nees, in Syll. Pl. Nov. Ratisbon.* i. (1824) p. 5; *Steud. Syn. Gram.* p. 209; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 249; *Benth. Fl. Hongk.* p. 430; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 704; *Hook. f. Fl. Brit. Ind.* vii. p. 299.

*Leptochloa tenerrima*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279, non *Kunth*; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 171.

*Poa chinensis*, *Linn. Sp. Pl.* p. 69; *Osbeck, Resa*, p. 206; *Lour. Fl. Cochinch.* p. 54.

SHANTUNG (*Staunton*!); KIANGSU: Kianang (*Staunton*!); CHEKIANG: Ningpo Mts. (*Faber*, 282!); KIANGSI: Kiukiang (*Shearer*!); FOKIEN: Amoy (*Hance*, 1292 in part!); FORMOSA: Tamsui (*Oldham*, 614!); KWANGTUNG: Lienchau, Canton (*Lamont*!); HONGKONG (*Hance*, 1292 in part!). Mus. Brit.; Herb. Kew.

Japan, and tropics of Asia, Africa, and Australia.

2. *Leptochloa filiformis*, *Roem. & Schult. Syst.* ii. p. 580; *Steud. Syn. Gram.* p. 209; *Hook. f. Fl. Brit. Ind.* vii. p. 298.

Var. *humilior*, *Presl, Rel. Hænk.* i. p. 288; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 249.

*Leptochloa tenerrima*, *Roem. & Schult. l. c.* p. 581, probably belongs here.

KIANGSU: Shanghai (*Maingay*, 563!); SZECHUEN: Yangtze-kiang, Min River (*Faber*, 1156!); KWANGTUNG: Macao (*Abel*!); HONGKONG: Government new gardens (*Lamont*!) (*Hance*, 1292 in part!). Mus. Brit.; Herb. Kew.

Tropics of Asia, Africa, and America.

#### Tribe FESTUCEÆ.

1. *Enneapogon brachystachyus*, *Stapf, in Dyer, Fl. Capens.* vii. p. 654.

*Pappophorum brachystachyum*, *Jaub. & Spach, Illustr. Pl. Or.* iv. t. 324; *Steud. Syn. Gram.* p. 201; *Hook. f. Fl. Brit. Ind.* vii. p. 302.

*Pappophorum boreale*, *Griseb. in Ledeb. Fl. Ross.* iv. p. 404; *Turcz. Fl. Baic.-Dahur.* ii. p. 317; *Steud. Syn. Gram. l. c.*; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 92.

CHIHLI: on the road near Kalgan (*Williams, in herb. Hance*, 14583!). Mus. Brit.; Herb. Kew.

Tropical and North Africa to North-west India and Central Asia. A variety in South Africa.

1. *Arundo Donax*, *Linn. Sp. Pl.* p. 81; *Lour. Fl. Cochinch.* p. 45; *Steud. Syn. Gram.* p. 193; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 705; *Hook. f. Fl. Brit. Ind.* vii. p. 302.

*Arundo bengalensis*, *Retz. Obs.* v. p. 20; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv., *Suppl.* (1896) p. 100.

*Arundo bifaria*, *Retz. Obs.* iv. p. 21.

*Amphidonax bifaria*, *Nees, ex Steud. Syn. Gram.* p. 197; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 171.

YUNNAN: 8-10 ft. high, Mengtze at 4600 ft. (*A. Henry, 9660 A!*); KWANGTUNG: by North River, Canton (*Nevin and Graves, in herb. Hance, 19303 in part!*), Amoy (*Hance, 1405!*), Whampoa (*Hance, 5501!*). Mus. Brit.; Herb. Kew.

Mountains of India; Europe, Northern Africa, Temperate Asia.

Var. *coleotricha*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 724.

FORMOSA: Tamsui (*Makino, fide Hackel*).

I have seen no specimen.

2. *Arundo formosana*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 724.

FORMOSA: Takow, Apes' Hill (*A. Henry, 1153, 1153 A!* *Playfair, 299!*), Bankinsing mountains (*A. Henry, 1153 B!*), Tamsui, sandhills (*Hancock, 8!*). Mus. Brit.; Herb. Kew.

Var. *gracilis*, *Hack. l. c.*

FORMOSA: Tamsui (*A. Henry, 1462!*). Herb. Kew.

I have not seen Professor Hackel's specimens and have referred the above to his species and variety on account of a general agreement with the descriptions. There is, however, one serious difference. Hackel describes the sterile glumes in his species as one-nerved, a remarkable character, whereas in the specimens which I have seen the sterile glumes are three-nerved as in other species of the genus. The spikelets are moreover two- to three-flowered, and show a wider range in size than is indicated in Hackel's description, which was based on a single specimen. The spikelets in the species are 2-4-flowered, and  $2\frac{3}{4}$ -4 lines long, the lower sterile glume ranges from  $2\frac{1}{2}$ - $2\frac{3}{4}$  lines, the upper from  $2\frac{3}{4}$ -3 lines, and the lowest fertile glume from 2- $2\frac{1}{2}$  lines long, with an awn  $1\frac{1}{4}$ - $1\frac{1}{2}$  lines long. The spikelets

in the variety are 2-flowered, scarcely  $2\frac{1}{2}$  lines long, the two sterile glumes a little less and a little more than  $1\frac{1}{2}$  lines respectively, the lower fertile glume  $1\frac{3}{4}$  lines long with an awn 1 line long.

1. *Neyraudia madagascariensis*, *Hook. f. Fl. Brit. Ind.* vii. p. 305; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225.

*Arundo madagascariensis*, *Kunth, Rév. Gram.* i. p. 273, t. 48; *Benth. Fl. Hongk.* p. 427.

*Arundo Henslowiana*, *Hook. & Arn. Bot. Beechey's Voy.* p. 248.

*Arundo Reynaudiana*, *Kunth, l. c.* p. 275, t. 49.

KIANGSI: Kiukiang (*Shearer!*); FOKIEN: Amoy (*Hance, 1407!*); HUPEH: Ichang gorge, River Yangtze (*Maries!*), Ichang (*A. Henry, 94, 4317!*); YUNNAN: Mengtze, 4600 ft. (*A. Henry, 10063!*), 7 ft. high, Szemao, 4000 ft. (*A. Henry, 13544!*); KWANGTUNG (*Staunton!*), Lienchau, Canton (*Lamont!*), Whampoa (*Hance, 1490 in part!*), face of city walls, Canton (*Sampson, 308!*), Macao (*Gaudichaud!*); HONGKONG: 3-4 ft. high, Robinson Road (*Hughes! Hance, 1490 in part! Champion! Wilford, 250!*); CHINA: South (*Seemann, 2396!*), without locality (*Seemann! Macartney!*). *Mus. Brit.*; *Herb. Kew.*

India, Malaya; Tropical Africa and Madagascar.

1. *Phragmites communis*, *Trin. Fund. Agrost.* p. 134; *Steud. Syn. Gram.* p. 195; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 268, et *Pl. David.* ii. p. 334; *Debeaux, Fl. Tchefou*, p. 157, et *Fl. Tientsien*, p. 49; *Ledeb. Fl. Alt.* i. p. 88; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 289; *Turczan. Fl. Baic.-Dahur.* ii. p. 315; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 170; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 704, et sér. 2, iii. (1903) p. 504; *Hook. f. Fl. Brit. Ind.* vii. p. 303.

*Phragmites longivalvis*, *Steud. Syn. Gram.* p. 196; *Debeaux, Fl. Tientsien*, p. 49.

*Arundo Phragmites*, *Linn. Sp. Pl.* i. p. 81; *Maxim. Prim. Fl. Amur.* p. 321; *Regel, Tent. Fl. Ussur.* p. 168; *Thunb. Fl. Jap.* p. 55.

CHIHLI: Peking (*Williams, in herb. Hance, 3472 in part!*); SHINGKING: Newchang (*Webster, 313!*), between Tungcheshien and Mäorshan, Yalu River (*James!*); KIANGSI (*Staunton!*); FOKIEN: near Amoy (*Hance, 3472 in part!*); FORMOSA: Tamsui

(*Oldham*, 611, 612 !), side of creek Takow (*A. Henry*, 1149 !); HUPEH: Ichang (*A. Henry*, 79 !); SHENSI: Kaolinsien (*Hugh* !); KWANGTUNG: near Canton (*Macartney* !), banks of the Canton River (*Sampson*, 1050 !). Mus. Brit.; Herb. Kew.

Widely distributed, especially in temperate regions.

2. *Phragmites Karka*, *Trin. ex Steud. Nom.* ii. p. 324; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Hack. in Bull. Herb. Boiss.* vii. (1899) pp. 704, 723; *Hook. f. Fl. Brit. Ind.* vii. p. 304.

*Phragmites Roxburghii*, *Steud. l. c.*; *Benth. Fl. Hongk.* p. 428; *Debeaux, Fl. Shanghai*, p. 70; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 278; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 170.

*Arundo Roxburghii*, *Kunth, Rév. Gram.* i. p. 79; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv., *Suppl.* (1896) p. 109.

SHANTUNG (*Maingay*, 67 !); KIANGSU: Shanghai (*Maingay*, 578 !); FORMOSA: Takow (*A. Henry*, 1205 !), Takow spit (*A. Henry*, 1208 !); YUNNAN: Mengtze, moist places, 4500 ft. (*A. Henry*, 9209 !), ravine 4600 ft. (*A. Henry*, 9209 A !); KWANGTUNG: growing plentifully and gregariously and reaching a height of 15 feet on the banks of the Canton River (*Hance*, 6492 !). Mus. Brit.; Herb. Kew.

Widely distributed in the warmer parts of the world.

1. *Kœleria cristata*, *Pers. Syn.* i. p. 97; *Steud. Syn. Gram.* p. 292; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 72; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 270, et *Pl. David.* i. p. 335; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 132; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 283; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 179; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 703, et in sér. 2, iii. (1903) p. 505; *Hook. f. Fl. Brit. Ind.* vii. p. 308; *Maxim. Prim. Fl. Amur.* p. 322; *Turcz. Fl. Baic.-Dahur.* ii. p. 338.

SHINGKING: Taliénwan Bay (*Swinhoe, in herb. Hance*, 3611 in part !); CHIHLI: Peking (*Bretschneider*, 848 !); SHANTUNG: Chefoo (*King, in herb. Hance*, 3611 in part! *Perry* !); KIANGSU: Taihu Lake (*Forbes*, 498 !); HUPEH: Ichang (*A. Henry*, 3505 !); SHENSI: Kusan (*Hugh* !), Mt. Laoysan, Teuizscien (*Hugh* !); COREA: East (*Perry* !). Mus. Brit.; Herb. Kew.

Widely distributed in temperate regions.

1. *Diplachne fusca*, Beauv. *Agrost.* p. 163; *Hook. f. Fl. Brit. Ind.* vii. p. 329.

*Uralepis fusca*, Steud. *Syn. Gram.* p. 247.

SHANTUNG (*Staunton!*); FOKIEN: Amoy (*Hance*, 1399!); FORMOSA: Takow, rice-field (*A. Henry*, 1959!), Takow spit (*A. Henry*, 1822, 1822 A, 1846!). Mus. Brit.; Herb. Kew.

Warmer regions of Old World.

2. *Diplachne serotina*, Link, *Hort. Berol.* i. p. 155; var. *chinensis*, Maxim. *Fl. As. Or. Fragm.* p. 70; *Franchet, Pl. David.* i. p. 334; Komarow, in *Act. Hort. Petrop.* xx. (1901) p. 291; Palibin, *ibid.* xxi. (1903) p. 209.

*Diplachne sinensis*, Hance, in *Journ. Bot.* viii. (1870) p. 76; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 270.

*Molinia serotina*, Mert. & Koch, *Deutschl. Fl.* i. p. 585.

CHIHLI: hills to the west of Peking (*Williams*, in herb. *Hance*, 12572!), Peking (*Bretschneider*, 853, 854!); SHING-KING: Kaichow to Kinchow (*Ross*, 263!); SHANTUNG: Chefoo (*Forbes!*). Mus. Brit.; Herb. Kew.

Var. *aristata*, Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 704, et sér. 2, iii. (1903) p. 504.

SHENSI: Kaolinsien (*Hugh!*), Mt. Laoyan (*Hugh!*); COREA: Hills near Pyengyang (*Faurie*, 878, fide *Hackel*). Mus. Brit.

Japan.

3. *Diplachne squarrosa*, Richter, *Plant. Europ.* p. 72.

*Molinia squarrosa*, Trin. in *Ledeb. Ic. Pl. Ross.* t. 227; *Ledeb. Fl. Ross.* iv. p. 396.

Var. *longe-aristata*, Rendle, n. var. Culmi tenues, dense cæspitosi, ad 1½-pedales, in parte superiore valde flexuosi. Spiculæ 4¼ lin. longæ; rhachilla glabra, sub floris autem utriusque articulationem breviter pilosa. Glumæ steriles lanceolatae; gl. i. 1⅛ lin. longa; gl. ii. 2 lin. longa, acuminata; gluma fertilis inferior, 2¾ lin. longa, cum arista terminali 3¼ lin. longa, lanceolata, in carinæ parte superiore et arista scabridula, 3-nervis, inter nervum lateralem et marginem utrinque sparse pilosula; palea 2½ lin. longa, carinæ conspicuæ aristuliferæ (aristula 1½ lin. longa); gluma floris superioris 2⅛ lin. longa, breviter aristata.

SHENSI: Kaolinsien (*Hugh!*), Mt. Kiansan, Husiensien (*Hugh!*). Mus. Brit.

Distinguished from the species by its longer awned, less hairy fertile glume.

The species occurs in South-east Russia, and Central and Northern Asia.

1. *Molinia Fauriei*, *Hack. in Bull. Herb. Boiss. sér. 2, iii.* (1903) p. 504.

COREA: by streams near Naipiang (*Faurie*, 804, *fide Hackel*).  
I have seen no specimen.

2. *Molinia japonica*, *Hack. in Bull. Herb. Boiss. vii.* (1899) p. 704, et *sér. 2, iii.* (1903) p. 504.

COREA: hills near Ouensan (*Faurie*, 810, *fide Hackel*).

Japan.

I have seen no specimen.

1. *Eragrostis amabilis*, *Wright & Arn. ex Nees, in Hook & Arn. Bot. Beechey's Voy.* p. 251; *Steud. Syn. Gram.* p. 265; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 317.

*Eragrostis unioloides*, *Nees, ex Steud. Syn. Gram.* p. 264; *Benth. Fl. Hongk.* p. 432; *Huck. in Bull. Herb. Boiss. vii.* (1899) p. 724.

*Poa unioloides*, *Retz. Obs. v.* p. 19.

KWANGTUNG: Whampoa (*Robertson!*), Pakhoi (*Playfair*, 132!); Lofaushan (*Ford*, 128!); HONGKONG: Race-course, Happy Valley (*Lamont! Forbes! Harland*, 695, in herb. *Hance*, 1007!); CHINA: without locality (*Baird!*). Mus. Brit.; Herb. Kew.

Tropical Asia and Tropical Africa.

2. *Eragrostis elegantula*, *Steud. Syn. Gram.* p. 266; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 318.

FORMOSA: path in rice-field, east plain, Takow (*A. Henry*, 1820!); KWANGTUNG: Lienchau, Canton (*Lamont!*), near top of White Cloud Hills, Canton (*Sampson*, 1055!), Pakhoi (*Playfair*, 159!); HONGKONG: Lam Tong Island (*Wright!*) (*Wilford*, 247!); HAINAN: without locality (*A. Henry*, 8072!); KWANGSI: Lungchow (*Morse*, 367!). Mus. Brit.; Herb. Kew.

India.

3. *Eragrostis elongata*, *Jacq. Eclog. Gram.* p. 3, t. 3; *Steud. Syn. Gram.* p. 264; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 319.

*Eragrostis atrovirens*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 725 (? of Trinius).

*Eragrostis bahiensis*, *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 136 (non *Schrad.*).

*Eragrostis Brownei*, *Nees, ex Steud. Nom.* ed. 2, p. 562, et in *Hook. & Arn. Bot. Beechey's Voy.* p. 253; *Benth. Fl. Hongk.* p. 432; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 281.

*Eragrostis orientalis*, *Nees, in Nov. Act. Nat. Cur.* xix., *Suppl.* i. (1843) p. 205, non *Trin.*; *Benth. Fl. Hongk.* p. 432; *A. Henry, in Trans. Asiatic Soc. Jap.* xxiv., *Suppl.* (1896) p. 109.

*Eragrostis Willdenoviana*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 252.

*Eragrostis zeylanica*, *Nees, in Nov. Act. Nat. Cur.*, tom. cit. p. 204; *Benth. Fl. Hongk.* p. 433.

CHEKIANG: without locality (*Staunton!*); FORMOSA: Takow (*A. Henry, 1019!*), Apes' Hill (*A. Henry, 1872! Playfair, 106!*); KWANGTUNG: abundant near Canton (*Sampson, 1796!*), Whampoa (*Wright! Hance, 11063 in part!*); HONGKONG (*Hance, 1004 in part, 11063 in part! Harland, 694, in herb. Hance, 1004 in part! Seemann, 749!*), dry sunny places (*Hance, 11064!*); Victoria Race-course, Happy Valley, and roadside, West Point (*Lamont!*), Little Hongkong and Aberdeen (*Wilford, 261!*); LUCHU ARCHIPELAGO: hill-sides (*Wright!*). *Mus. Brit.*; *Herb. Kew.*

Tropical Asia; Australia.

4. *Eragrostis ferruginea*, *Beauv. Agrost.* p. 71; *Steud. Syn. Gram.* p. 267; *Debeaux, Fl. Tchefou*, p. 161; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Franchet, Pl. David.* i. p. 335, et ii. p. 146; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 294; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 177; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 705, et sér. 2, iii. (1903) p. 505; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 324.

*Eragrostis orientalis*, *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 71.

*Poa ferruginea*, *Thunb. Fl. Jap.* p. 50; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 270.

*Poa orientalis*, *Franchet, l. c.*

SHINGKING: Kaichow to Kinchow (*Ross, 296!*); KIANGSU:

Shanghai (*Maingay*, 548! *Seemann*! *Tonnerre*, in herb. *Hance*, 8133!); EAST COAST (*Home*!); KIANGSI: Nauk'angfu, Kiukiang (*Reid*!), Kiukiang (*Shearer*!); HUPEH: Ichang (*A. Henry*, 2743, 4288!); SHENSI: Mt. Thaepeisan (*Hugh*!), Mt. Miao-wansan (*Hugh*!), Kisanshien (*Hugh*!); SZECHUEN: Mt. Omei (*Faber*, 1179!); YUNNAN: Mengtze, grass mountains to north, — at 5000 ft. (*A. Henry*, 10602!); CHINA: without locality (*Macartney*!). Mus. Brit.; Herb. Kew.

Japan, Himalayas.

5. *Eragrostis geniculata*, *Nees*, in *Nov. Act. Nat. Cur.* xix. *Suppl.* i. (1843) p. 203; *Benth. Fl. Hongk.* p. 433.

*Eragrostis cylindrica*, *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 251.

*Eragrostis Nevinii*, *Hance*, in *Journ. Bot.* xviii. (1880) p. 302.

FOKIEN: Amoy (*Hance*, 1380 in part!); KWANGTUNG: Canton (*Sampson*, 316, 317! and in herb. *Hance*, 20602 in part!; *Hance*, 1380 in part!), by North River at Tsingiin (*Nevin*, in herb. *Hance*, 20602 in part!), Pakhoi (*Playfair*, 145!); HONGKONG: Mt. Victoria (*Lamont*! *Wright*!); HAINAN (*A. Henry*, 7970!). Mus. Brit.; Herb. Kew.

*Hance's Eragrostis Nevinii* (herb. 20602) is a robust form with broader leaves, somewhat laxer panicle, and the keels of the pale more shortly haired than in the type. I do not think it is specifically distinct.

6. *Eragrostis harpachnoides*, *Hack.* in *Oest. Bot. Zeitschr.* lii. (1902) p. 306.

YUNNAN: roadsides at Peelongtau near Tapintze (*Delavay*, fide *Hackel*).

I have not seen a specimen.

7. *Eragrostis Hugoniana*, *Rendle*, n. sp.

*Herba* glauca, rhizomate brevi perennis; culmi ramosi, cæspitiosi, ascendentes, nodis inferioribus geniculati, 4-5-nodi, teretes, ad 11 poll. alti et  $\frac{2}{3}$  lin. crassi. *Foliorum* vaginæ internodiis sæpe breviores, laxæ, marginibus sparse pilosæ, ore barbatae; ligula ad lineam breviter et dense pilosam reducta; lamina rigidula, lineari-acuminata, plana, margine glandulosa et sæpe involuta, facies superior in venis scabridula, ad 5 poll. longa, et vix



2 lin. lata. *Panicula* ellipsoidea, aut ob ramos demum patentez ovoidea, ad 6 poll. longa et  $5\frac{1}{2}$  poll. diametro; rhachis, rami, ramuli et pedicelli leves, pedicelli ad medium originemque, rami ramulique ad eorum origines glanduliferi; rami solitarii, inferiores  $2\frac{1}{2}$  ad 3 poll. longi, mox supra basin ramulosi, ramuli demum divaricati; spiculæ compactæ, olivaceæ, lineari-lanceolatæ, 10-18 flores,  $2\frac{1}{2}$  ad 6 lin. longæ quam pedicellæ longiores. *Glumæ* steriles inæquales, inferior lanceolata, subacuta, quam gl. superior ovata subacuta,  $\frac{3}{4}$  lin. longa, paullo brevior; fertiles ovatæ, obtusiusculæ, 1 lin. longæ, nervis lateralibus 2 prominulis, e basi versus apicem rhachillæ deciduæ; paleæ glumis paullo breviores, persistentes; antheræ  $\frac{1}{10}$  lin. longæ. *Fructus* oblongus,  $\frac{1}{2}$  lin. longus.

SHENSHI: Kaolinsien (*Hugh!*). Mus. Brit.

Approaches the Southern Indian and Ceylon species, *Eragrostis Willdenoviana*, Nees, which it recalls in general habit, but has a more compact spikelet, with less blunt flowering-glumes and less unequal barren glumes.

8. *Eragrostis interrupta*, Beauv., var. *tenuissima*, Stapf, in *Hook. f. Fl. Brit. Ind.* vii. p. 316.

*Eragrostis interrupta*, Beauv.; Pilger, in *Engl. Bot. Jahrb.* xxix. (1900) p. 225.

*Eragrostis japonica*, Trin. in *Mém. Acad. Pétersb.* sér. 6, i. (1830) p. 450; Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 705.

*Eragrostis tenella*, Benth. *Fl. Hongk.* p. 431; Debeaux, *Fl. Tchefou*, p. 161; Franchet, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 270, et *Pl. David.* i. p. 335; Miq. in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; Franch. et Savat. *Enum. Pl. Jap.* ii. p. 177.

*Eragrostis tenuissima*, Schrad. ex Nees, *Fl. Afr. Austr.* p. 410.

*Poa japonica*, Thunb. *Fl. Jap.* p. 51.

KIANGSU: Shanghai (*Maingay*, 559!), Chiukiang, Nank'angfu (*Reid!*); CHEKIANG: Ningpo Mountains (*Faber!*); FOKIEN: in rice-fields, Amoy (*Hance*, 1291, 3376 in part!); EAST COAST (*Home!*); FORMOSA (*Oldham*, 616!), Takow plain, in field (*A. Henry*, 2018!), Tamsui (*A. Henry*, 1450!); HUPEH: Ichang (*A. Henry*, 137, 2220!), Changyang (*A. Henry*, 7738!); SOUTH CHINA (*Petersen*, in herb. *Hance*, 3376 in part!); KWANGTUNG: Lienchau, Canton (*Lamont!*), North River, Canton (*Ford*, 7!), Lantao (*Ford*, 14!), Whampoa (*Robertson!* *Hance!*). Mus. Brit.; Herb. Kew.

Japan; Tropical Asia.

7 9. *Eragrostis major*, *Host, Gram. Austr.* iv. p. 14, t. 24; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 133; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 706, et sér. 2, iii. (1903) p. 505; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 320.

*Eragrostis megastachya*, *Link, Hort. Berol.* i. p. 187; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 251; *Debeaux, Fl. Tchefou*, p. 160; *Franchet, Pl. David.* i. p. 335; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 177; *Ledeb. Fl. Ross.* iv. p. 382; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 294.

*Eragrostis poæoides*, *Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, i. (1830) p. 404.

*Poa megastachya*, *Koel. Descr. Gram.* p. 181; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 269.

CHINA: without locality (*Fortune*, 3!); CHIHLI: Peking (*Bretschneider*, 859! *Bushell*!); SHINGKING: Newchang (*Webster*, 311!) (*Ross*, 44!); SHANTUNG: Chefoo (*Stuhlmann* in herb. *Hance*, 3514 in part!); KIANGSU: Shanghai (*Maingay*, 547, 741!); KIANGSI: Kiukiang (*Shearer*!); FOKIEN: without locality (*De Grijs*, in herb. *Hance*, 3514 in part!); HUPEH: Ichang (*A. Henry*, 172!); CHINA: East Coast (*Home*!); FORMOSA (*Campbell*!), Apes' Hill, Takow (*A. Henry*, 1207!); NORTH CENTRAL CHINA: Leunteon, Fenghosan (*Hugh*!); SHENSI: Paoki, Miaowansan (*Hugh*!), Kusan, Kisansien (*Hugh*!); KWANGTUNG (*Ford*, 216!). Mus. Brit.; Herb. Kew.

Japan, warmer parts of Asia; Mediterranean region, Tropical and South Africa, warmer parts of America.

10. *Eragrostis Makinoi*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 725.

FORMOSA: Shinchiku (*Makino*, fide *Hackel*).

I have seen no specimen.

11. *Eragrostis minor*, *Host, Gram. Austr.* iv. p. 15; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 724, et sér. 2, iii. (1903) p. 505; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 321.

*Eragrostis poæoides*, *Beauv. Agrost.* p. 162; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 293; *Ledeb. Fl. Ross.* iv. p. 381; *Turcz. Fl. Baic.-Dahur.* ii. p. 329.

CHIHLI: near Peking (*Bushell*!); NORTH CENTRAL CHINA: Leunteon, Fenghosan (*Hugh*!); SZECHUEN: Yangtzekiang (*Faber*, 1131!); KWANGTUNG: Lienchau (*Lamont*!). Mus. Brit.; Herb. Kew.

South Europe, North Africa to Abyssinia, Temperate Asia, and Northern India.

12. *Eragrostis pilosa*, Beauv. *Agrost.* p. 71; *Steud. Syn. Gram.* p. 263; *Benth. Fl. Hongk.* p. 432; *Debeaux, Fl. Tchefou*, p. 159; *Franchet, Pl. David.* i. p. 335 (et var. *imberbis*); *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 176; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 706, et sér. 2, iii. (1903) p. 505; *Ledeb. Fl. Ross.* iv. p. 382; *Turcz. Fl. Baic.-Dahur.* ii. p. 329; *Maxim. Prim. Fl. Amur.* p. 320; *Regel, Tent. Fl. Ussur.* p. 168; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 292; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 323.

*Eragrostis verticillata*, Beauv. *l. c.*; *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 253.

*Poa pilosa*, *Linn. Sp. Pl.* p. 68; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 269.

SHINGKING: Newchang plain (*Faber*, 1530!), Kaichow to Kinchow (*Ross*, 291!); SHANTUNG (*Maingay*, 72, 216!); KIANGSU: Shanghai (*Maingay*, 698!); HUPEH (*A. Henry*, 5001!); SHENSI: Paoki, Mt. Miaowansan (*Hugh*!), Mt. Thaepeisan (*Hugh*!); SZECHUEN: Mt. Omei (*Faber*, 1172!); KWANGTUNG: Whampoa (*Hance*, 3573! *Wright*!), Canton (*Sampson*, 315!). *Mus. Brit.*; *Herb. Kew.*

Japan; Southern Europe, and widely distributed in the warmer parts of the earth.

13. *Eragrostis pilosissima*, *Link, Hort. Berol.* i. p. 189; *Steud. Syn. Gram.* p. 280; *Benth. Fl. Hongk.* p. 432.

*Eragrostis Millettii*, *Nees, in Hook. & Arn. Bot. Beechey's Voy.* p. 252.

KWANGTUNG: Pakwan Mountains, near Canton (*Sampson*, in herb. *Hance*, 1311!), Lofaushan (*Ford*, 133!), British Kowlung (*Ford*, 593!), Whampoa (*Wright*!), Macao (*Vachell*!); HONGKONG (*Hance*, 1311! *Harland*, 1007! *Ford*, in herb. *Forbes*, 489!). *Mus. Brit.*; *Herb. Kew.*

Japan.

14. *Eragrostis stenophylla*, *Hochst. ex Miq. Anal. Bot. Ind.* ii. p. 27; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 318.

HONGKONG: hill-sides in small tufts and along mountain rivulets in crevices of rocks (*Wright*!). *Herb. Kew.*

Tropical Asia and Africa.

15. *Eragrostis tenella*, *Roem. et Schult. Syst.* ii. p. 576; *Steud. Syn. Gram.* p. 315.

*Eragrostis tenella*, var. *plumosa*, *Stapf*, in *Hook. f. Fl. Brit. Ind.* vii. p. 315.

*Eragrostis plumosa*, *Link, Hort. Berol.* i. p. 192; *Steud. l. c.*; *Nees*, in *Hook. & Arn. Bot. Beechey's Voy.* p. 254; *Benth. Fl. Hongk.* p. 431; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 724.

*Poa plumosa*, *Retz. Obs.* iv. p. 20.

FORMOSA (*Campbell!*); KWANGTUNG: Canton (*Sampson*, 313!), Pakhoi (*Playfair*, 148!); HONGKONG: on walls (*Hance*, 1291!) (*Harland!*); CHINA; without locality (*Macartney! Millett!*). *Mus. Brit.*; *Herb. Kew.*

Tropical Asia, Africa.

1. *Diarrhena japonica*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 603; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 707, et sér. 2, iii. (1903) p. 505.

*Oncea japonica*, *Franch. et Savat. loc. cit.* p. 178.

COREA: woods near Ouensan (*Faurie*, 835, ex *Hackel*).

Japan.

1. *Melica Gmelini*, *Turcz. ex Trin. in Mém. Acad. Sci. Pétersb.* sér. 6, i. (1831) p. 368; *Steud. Syn. Gram.* p. 289; *Franchet, Pl. David.* i. p. 336; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 296; *Maxim. Prim. Fl. Amur.* p. 322; *Regel, Tent. Fl. Ussur.* p. 169; *Ledeb. Fl. Ross.* iv. p. 398.

CHILI: near top of Mt. Puhuashan, at 6000–7000 ft. (*Moellendorff*, in herb. *Hance*, 3605!). *Mus. Brit.*

North Asia.

2. *Melica nutans*, *Linn. Sp. Pl.* p. 66; *Steud. Syn. Gram.* p. 291; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 178; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 706; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 297; *Maxim. Prim. Fl. Amur.* p. 322; *Regel, Tent. Fl. Ussur.* p. 169; *Turcz. Fl. Baic.-Dahur.* ii. p. 336; *Hook. f. Fl. Brit. Ind.* vii. p. 330.

SHINGKING: between Moukden and Tungehshien (*James!*); CHEKIANG OF SHANTUNG (*Faber*, 289 or 291!). *Herb. Kew.*

Japan, North and Central Asia, Himalayas, Europe.

*Faber's* specimen (the locality of which is uncertain owing to

a mixing of two labels) resembles specimens collected in Japan by the 'Challenger' Expedition in having whitish-green spikelets.

3. *Melica Onœi*, *Franch. et Savat. Enum. Pl. Jap.* ii. p. 603; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 707, et sér. 2, iii. (1903) p. 505.

CHEKIANG: Ningpo Mountains (*Faber*, 12, 1733!); HUPEH: Fang (*A. Henry*, 6641, 6716!); SZECHUEN: North Wushan, 4000-7000 feet (*A. Henry*, 7034!); COREA: hills near Ouensan (*Faurie*, 809, ex *Hackel*). Mus. Brit.; Herb. Kew.

Japan.

I have seen no authenticated specimen of this species; the plants referred to it agree fairly with the author's description, though the spikelets in specimens collected by Henry and Faber are slightly larger.

4. *Melica Radula*, *Franchet, Pl. David.* i. p. 336.

SHENSI: Kusan (*Hugh*!), Mt. Laoysan, Teuizscien (*Hugh*!). Mus. Brit.

I have seen no authenticated specimen of this species; Father Hugh's plants agree fairly well with Franchet's description.

5. *Melica scabrosa*, *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 72; *Steud. Syn. Gram.* p. 291; *Franchet, Pl. David.* i. p. 336, et in *Mém. Soc. Sci. Nat. Cherb.* xxiv. (1884) p. 270; *Kanitz, in Math. Naturw. Ber. Ungarn*, iii. (1885) p. 10; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Komarow, in Act. Hort. Petrop.* xx. (1901) p. 297; *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 505.

CHIHLI: mountain west of Peking (*Bretschneider*, in herb. *Forbes*, 1984!); SZECHUEN: Wushan gorge, Yangtzekiang (*Faber*, 1178!). Mus. Brit.; Herb. Kew.

The plant from Nanchuan referred by Pilger (in *Engl. Bot. Jahrb.* xxix. p. 225) to *Melica scabra*, Linn., belongs here.

1. *Centotheca lappacea*, *Desv. in Nouv. Bull. Soc. Philom.* ii. (1810) p. 189; *Steud. Syn. Gram.* p. 116; *Benth. in Hook. Niger Fl.* p. 56; *Hance, in Journ. Linn. Soc., Bot.* xiii. (1873) p. 136; *Rendle, in Cat. Afr. Pl. Welw.* ii. p. 228.

*Holcus latifolius*, *Osbeck, Resa*, p. 247; *Linn. Syst.* ed. 10, p. 1305.

FORMOSA: Bankinsing (*A. Henry*, 99!); KWANGTUNG: White

Cloud hills, Canton (*Sampson*, 1556 !), mainland opposite Hongkong (*Harland*, in herb. *Hance*, 2131 in part !); HONGKONG (*Hance*, 2131 in part !), Happy Valley woods (*Lamont* !), ravine leading up to 900 yards range (*Urquhart*, 250 !). Mus. Brit.; Herb. Kew.

Tropical Asia to Polynesia and Tropical Australia; Tropical Africa.

Var. *inermis*, *Rendle*, n. var. *Folia* speciei ad 7 poll. longa et  $1\frac{1}{2}$  poll. lata. *Panicula* triangulari-ovata, densa,  $3\frac{3}{4}$  poll. longa et lata, rami horizontaliter patentés, secundi, spiculis nutantibus; spiculæ  $\frac{1}{4}$  poll. longæ, 3-4-floræ. *Glumæ* florentes rarissime setam solitariam sub apice gerentes.

FORMOSA: Bankinsing (*A. Henry*, 417 !). Herb. Kew.

At first sight this has the appearance of a distinct species, owing to the habit of the inflorescence with crowded secund branches and nodding spikelets. But, except for the practically complete absence of the characteristic recurved bristles of the flowering glumes, the spikelets do not differ from those of the species—in which, moreover, there is considerable variation in the number of bristles on the glumes.

1. *Lophatherum gracile*, *Brongn.* in *Duperrey, Voyage, Bot.* p. 50, t. 8; *Steud. Syn. Gram.* p. 300; *Benth. Fl. Hongk.* p. 433; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 707; *Franchet, Pl. David.* i. p. 337; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Hook. f. Fl. Brit. Ind.* vii. p. 331.

*Lophatherum elatum*, *Zoll. et Mor. in Mor. Syst. Verz.* p. 102; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 179; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 50.

*Lophatherum japonicum*, *Steud. Syn. Gram.* p. 300.

*Lophatherum Lehmanni*, *Nees, ex Steud. l. c.*; *A. Henry, in Trans. Asiat. Soc. Jap.* xxiv. *Suppl.* (1896) p. 109.

CHEKIANG: Ningpo Mountains (*Faber*, 284 !); FOKIEN: Amoy (*Hance*, 1013 in part !); FORMOSA (*Oldham*, 618 !), Bankinsing Mountains (*A. Henry*, 1647 !); HUPEH: West (*Wilson*, 533 !), Nanto and mountains to the northward (*A. Henry*, 4635 !); SZECHUEN: Mt. Omei (*Faber*, 1184 !); KWANGTUNG: wood at White Cloud Monastery, Canton (*Sampson*, 1166 !); HONGKONG: Happy Valley woods (*Lamont* ! *Ford*, 595 ! *Hance*, 1013 in part ! *Wright* !); LUCHU ARCHIPELAGO (*Wright* !). Mus. Brit.; Herb. Kew.

Japan, Hill country of India and Ceylon, Malaya.

2. *Lophatherum sinense*, Rendle, n. sp.

*Herba*; culmus (e specimine unico quod ex internodiis culmi superioribus tribus constat) teres, 1 lin. crassus, internodium supremum, ad paniculæ basin, 14 poll. longum. *Foliorum* vaginæ internodia haud æquantes, striatæ, sparsius scabridulo-pilosæ; ligula obsoleta; lamina vix petiolata, lanceolata, acuminata, flaccida, glabra, tesselata, ad 6 poll. longa,  $\frac{3}{4}$  poll. lata. *Panicula* 8 poll. longa, rami distantes 5, rigide patento-ascendentes, infimus 2 poll. longus; spiculæ late lanceolatæ,  $3\frac{1}{2}$  lin. longæ. *Glumæ* steriles dorso rotundatæ, obtusæ, 5-nerves, glabræ, inferior late oblonga, 2 lin. longa, superior ovata,  $2\frac{1}{4}$  lin. longa; fertilis latissime ovata, mucronata, 7-nervis, glabra, cum mucrone ( $\frac{1}{2}$  lin.) 3 lin. longa,  $2\frac{2}{3}$  lin. lata; palea  $2\frac{1}{2}$  lin. longa. *Caryopsis* ovoidea,  $1\frac{1}{2}$  lin. longa,  $\frac{2}{3}$  lin. lata.

KIANGSI: Kiukiang (*Shearer*!). Herb. Kew.

Distinguished from *Lophatherum gracile* by the broader spikelet, very broad fertile glume, and broad ovoid grain.

1. *Æluropus littoralis*, *Parl. Fl. Ital.* i. p. 461; *Ledeb. Fl. Ross.* iv. p. 369; var. *sinensis*, *Debeaux, Fl. Tientsin*, p. 50.

CHIHLI: sands at Peiho, Tientsin (*Debeaux*).

The species occurs in the Mediterranean region and North Asia.

1. *Dactylis glomerata*, *Linn. Sp. Pl.* p. 71; *Steud. Syn. Gram.* p. 297; *Franchet, Pl. David.* i. p. 146; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 225; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 179; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 708; *Hook. f. Fl. Brit. Ind.* vii. p. 335.

HUPEH: Hsingshan (*A. Henry*, 6530 A!), Patung (*A. Henry*, 4673!); SZECHUEN: Tachienlu (*Pratt*, 121!); YUNNAN: Mengtze, grass mountains to south, 5000 feet (*A. Henry*, 10740!). Mus. Brit.; Herb. Kew.

Europe, North Africa, West and North Asia, and Himalayas.

1. *Briza elegans*, *Osbeck, Resa*, p. 246.

KWANGTUNG: near Canton (*Osbeck*).

I have not seen this specimen. It is described as an exceedingly fine grass, which grew near the highest plantations, having

oblong spikelets and keeled glumes. It may perhaps be a synonym of *Eragrostis major*, Host.

2. *Briza minor*, *Linn. Sp. Pl.* p. 70; *Steud. Syn. Gram.* p. 282; *Franchet, Pl. David.* i. p. 337; *Hack. in Engl. Bot. Jahrb.* vi. (1884) p. 50; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 178; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 708.

CHIHLI: environs of Jehol (*David, ex Franchet*).

Europe, North Africa, Western Asia; widely distributed as an alien.

1. *Poa acroleuca*, *Steud. Syn. Gram.* p. 256; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Franchet, Pl. David.* ii. p. 145; *Palibin, in Act. Hort. Petrop.* xix. (1901) p. 133; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 175; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 710, et sér. 2, iii. (1903) p. 506.

CHINA, without locality (*Faber!*); SHINGKING: Peiling (*Webster, 99!*); KIANGSU: Chinkiang (*Maries!*); HUPEH (*A. Henry, 5792!*); SHENSI: Paoki, Chifengsan (*Hugh!*), Hansunfu, Mt. Leansan (*Hugh!*), Mt. Laoysan, Teuizscien (*Hugh*); SZECHUEN: South Wushan (*A. Henry, 5685!*); COREA: Port Hamilton (*Wilford, 714!*); LUCHU ARCHIPELAGO (*Wright!*). *Mus. Brit.*; *Herb. Kew.*

Japan.

2. *Poa annua*, *Linn. Sp. Pl.* p. 68; *Steud. Syn. Gram.* p. 250; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 279; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 174; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 708; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 345; *Turcz. Fl. Baic.-Dahur.* ii. p. 327.

KIANGSU: Shanghai (*Maingay, 580!*); KIANGSI: Kiukiang (*Faber, 1150!*); FORMOSA (*Oldham, 617!*); HUPEH: Patung (*Wilson, 1646!* *A. Henry, 3746, 5425!*), Ichang (*A. Henry, 824!*). *Mus. Brit.*; *Herb. Kew.*

Japan, mountains of India, Temperate Asia, Europe. Widely colonised in other countries.

Var. *nepalensis*, *Griseb. Goett. Nachricht.* 1868, p. 75; *Stapf,*



in *Hook. f. Fl. Brit. Ind.* vii. p. 346; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226.

*Poa nepalensis*, *Wall. ex Duthie, Grasses N.W. India*, p. 40 (nomen); *Franchet, Pl. David.* ii. p. 145.

SZECHUEN: Moupin (*David, ex Franchet*).

Western Himalayas.

3. *Poa attenuata*, *Trin. in Bunge, Verz. Suppl. Fl. Altaic.* p. 9; *Steud. Syn. Gram.* p. 254; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 340; *Turez. Fl. Baic.-Dahur.* ii. p. 324.

*Poa nemoralis*, var., *Hance, in Journ. Bot.* xviii. (1880) p. 303.

CHIHILI: near the top of Mt. Puhuashan, 6000–7000 feet (*Moellendorff, in herb. Hance, 20314!*). Mus. Brit.; Herb. Kew.

Himalayas, Central and North Asia.

A small form of the species.

Var. *vivipara*, *Rendle, n. var.* *Herba* humilis; culmi (cum panicula) 5–7-pollicares. *Folia* plana vel plicata, lineari-acuminata, ad  $2\frac{3}{4}$  poll. longa; ligula lata, obtusa vel abrupte acuta,  $1\frac{1}{4}$  lin. longa; lamina suprema vaginam excedens. *Panicula*  $1\frac{3}{4}$ – $2\frac{1}{4}$  poll. longa, lineari-oblongata, densa; rami ad quemque nodum gemini, scabridi, suberecti, sub apice pauci-spiculiferi; spiculæ 2 lin. longæ, ovato-lanceolatæ, viviparæ, violaceæ. *Glumæ* steriles cum carina sub apice solum scabridulæ; callus glumæ fertilis breviter et sparse pilosus, carina et nervi marginales ad basin tantum pilosa; antheræ  $\frac{1}{8}$  poll. longæ.

SIENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

#### 4. *Poa Faberi*, *Rendle, n. sp.*

*Herba* perennis, glabra, rhizomate duro; culmi sesquipedales, conferti, graciles, basi breviter decumbentes et cum vaginis scariosis brunneis circumdati. *Folia* caulina tria, nodum supremum sub medio culmi positum; vaginæ striatæ, subscabridulæ; ligula acuta, ad 2 lin. longa; lamina plana, linearis, basi angustata, superne longe attenuata, scabridula, nervis primariis tribus; folii supremi vagina  $4\frac{1}{3}$  poll. longa, lamina  $3\frac{1}{2}$  poll. longa,  $\frac{3}{4}$  lin. lata; folii secundi vagina 3 poll. longa, lamina 4 poll. longa. *Panicula* lanceolata, densa,  $4\frac{1}{2}$  poll. longa, axis et rami tenues, scabriduli; rami flexuosi in parte inferiore nudi, superne sparse et breviter ramulosi, dense et stricte spiculiferi, ad nodum infe-

riorem 5, 1-2 poll. longi; spiculæ oblanceolatae et obovatae, 4-floræ, rigidulæ,  $2\frac{1}{2}$  lin. longæ, pedicellos scabridulos excedentes. *Glumæ* steriles acutæ, lanceolatae, 3-nerves, in carina nervisque lateralibus scabridæ; superior latior et longior (vix  $1\frac{1}{2}$  lin. longa) quam inferior; florentes ovatae obtusæ, apice breviter membranaceæ, 5-nerves, carina scabrida et velut nervis marginalibus ad medium pilosa, pedicellus contexto-lanatus; gl. infima  $1\frac{1}{2}$  lin. longa; palea  $1\frac{1}{4}$  lin. longa; carinis ciliolatis; antheræ  $\frac{1}{2}$  lin. longæ.

KIANGSU: Shanghai (*Faber*, 293!); SHENSI: Mt. Laoysan, Tcuizscien (*Hugh*!), Kaolinsien (*Hugh*!), Hansunfu, Mt. Leansan (*Hugh*!). Mus. Brit.; Herb. Kew.

A distinct species of the *Poa nemoralis* group, differentiated by its long acute ligule, dense narrow panicle with quinate branches which are free from spikelets below, and minute anthers.

Var. *ligulata*, *Rendle*, n. var. *Planta* gracilis, bipedalis; ligula elongata, acuta, ad 5 lin. longa. *Panicula* gracilis; spiculæ bifloræ.

SZETCHUEN: Fou, Yangtzekiang (*Faber*, 1159!). Herb. Kew.

5. *Poa gracillima*, *Rendle*, n. sp.

*Herba* humilis, ut apparet cæspitosa, glabra, 4 poll. alta; culmi tenues, compressi. *Folia* caulina 3, supremum basi paniculæ amplectans; vagina suprema quam inferiores longior (10 lin. longa) laminæ æquans, lamina folii secundi longior ( $1\frac{1}{4}$  poll. longa); vaginæ strictæ, compressæ, striatæ; ligula vix  $\frac{1}{3}$  lin. longa, obtusa; lamina linearis, superne longe attenuata, plana, flaccida,  $\frac{3}{4}$  lin. lata. *Panicula*  $1\frac{3}{4}$  poll. longa, rhachis capillaris, flexuosa, levis, nodis 5, rami distantes, ascendentes, solitarii, tenuissimi, scabriduli, parte inferiore (ad dimidiam) nudi, superne 5-1-spiculiferi; rami duo infimi semipollicares, interdum cum ramulo breve 2-spiculifero. *Spiculæ* ellipticæ, 2-3-floræ,  $1\frac{1}{4}$  lin. longæ; pedicello scabridulo spiculam æquante. *Glumæ* subrigidulæ, steriles, lanceolatae, acutæ, cum carina scabridula, inferior uninervis,  $\frac{1}{2}$  lin. longa, superior latior, trinervis,  $\frac{7}{8}$  lin. longa; fertiles omnino glabræ, plicatæ, oblongo-lanceolatae, obtusæ, in parte  $\frac{1}{5}$  superiore membranaceæ, albidæ, 5-nerves, in carina scabridulæ, inferior 1 lin. longa, palea paullo brevior, carinis in parte superiore scabridulis; antheræ vix  $\frac{1}{4}$  lin. longæ.

SZECHUEN: Mt. Omei (*Faber*, 1185!). Herb. Kew.

A very small delicate specimen, suggesting somewhat a dwarf form of *Poa annua*, but differing in the habit of the panicle and the complete absence of hair on the fertile glumes, which are also less membranous than in *P. annua*.

6. *Poa Mariesii*, *Rendle*, n. sp.

*Planta* subrobusta; culmi flexuosi, erecti, sub medio  $1\frac{1}{2}$  lin. crassi, internodiis levibus striatis. *Folia* culmi 4, supremum basin paniculæ vaginans; vaginæ laxæ, striatæ, glabræ, suprema quam inferiores paullo longior (circa 5 poll. longa) quam ejus lamina paullo brevior; ligula brevis, truncata, 1 lin. longa; lamina linearis, superne longe attenuata, plana, flaccida, multinervis, glabra, 3 lin. lata,  $5-6\frac{2}{3}$  poll. longa; suprema quam secunda paullo brevior. *Panicula* subpyramidalis, flexuosa,  $8\frac{1}{2}$  poll. longa,  $2-2\frac{1}{2}$  poll. lata, ramis geminis, axis in parte superiore solum subscabridulus ceterum levis, rami in parte superiore subscabriduli, ramuli scabriduli; rami ad nodum infimum 3 poll. et  $2\frac{1}{8}$  poll. longi, superne gradatim deminuentes, in parte inferiore ( $\frac{1}{3}-\frac{1}{2}$ ) nudi, superne (cum ramulis) subdense spiculiferi; spiculæ ellipticæ, 5-6-floræ, 3 lin. longæ. *Glumæ* steriles lanceolatæ, acutæ, apice membranaceæ et albidæ, inferior  $1\frac{1}{2}$  lin. longa, uninervis, superior 3-nervis,  $1\frac{5}{8}$  lin. longa, quam inferior latior et minus acuta; fertiles firmæ, duplatae lanceolatæ, obtusæ vel subacutæ, apice et margine membranaceæ et albidæ, 5-nerves, carina (ad  $\frac{3}{4}-\frac{4}{5}$  longitudinis) ut nervus marginalis (ad  $\frac{1}{2}-\frac{3}{4}$  longit.) dense et regulariter pilosa, pilis  $\frac{1}{4}$  lin. longa; infima  $2\frac{1}{3}$  lin. longa; carinæ paleæ ciliolatæ; antheræ  $\frac{1}{2}$  lin. longæ; pedicellus breviter contortuplicato-lanatus.

KIANGSU: Chiukiang and River Yangtze (*Maries!*). Herb. Kew.

A well-marked species of the *Poa pratensis* group, characterized by its robust habit, large broad leaves, the upper blade being nearly as long as that of the leaf at the second node, the large panicle, the 4-5-flowered spikelets, the very hairy keels and marginal nerves of the fertile glumes, and the small anthers.

7. *Poa nemoralis*, *Linn. Sp. Pl.* p. 69; *Steud. Syn. Gram.* p. 251; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 174; *Hack. in Bull. Herb. Boiss.*

vii. (1899) p. 710; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 341; *Turcz. Fl. Baic.-Dahur.* ii. p. 327.

SHENSI: Mt. Miaowansan (*Hugh!*). Mus. Brit.

Var. *ligulata*, *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 341.

HUPEH (*A. Henry*, 6807!). Mus. Brit.; Herb. Kew.

Himalayas.

The specimen, which is in fruit, is not in a very satisfactory condition for determination.

Var. *mongolica*, *Rendle*, n. var. *Paniculæ* rami tantum in parte superiore ramulosi et spiculiferi. *Glumæ* fertiles in carina ad  $\frac{2}{3}$  longit. longius pilosæ, in nervo marginale ad  $\frac{2}{5}$  longit.

SHINGKING: between Moukden and Tungcheshien (*James!*). Herb. Kew.

The geminate branches of the panicle, which are bare and unbranched for half or more than half their length, give a characteristic appearance to the inflorescence. The keel and marginal nerves of the fertile glumes bear more and longer hairs than usual in the species. There is a scanty tuft of wool on the short pedicel.

8. *Poa pratensis*, *Linn. Sp. Pl.* p. 67; *Steud. Syn. Gram.* p. 251; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Franchet, Pl. David.* i. p. 337; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 174; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 708, et sér. 2, iii. (1903) p. 506; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 339; *Turcz. Fl. Baic.-Dahur.* ii. p. 328.

CHIHLI: Jehol (*David*, 1807!); SHINGKING: Chienshan (*Ross*, 586!), Jaoling (*Ross*, 383!), between Moukden and Tungcheshien (*James!*); SHENSI: Kusan (*Hugh!*); HUPEH: Patung (*A. Henry*, 4026!); COREA: East Corea (*Perry!*). Mus. Brit.; Herb. Kew.

Japan, Himalayas, and North temperate regions.

Var. *angustifolia*, *Gaud. Agrost. Helvet.* i. p. 214; *Kunth, Enum.* i. p. 353; *Franchet, Pl. David.* i. p. 337; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 340; *Turcz. Fl. Baic.-Dahur.* ii. p. 328.

SHENSI: Paoki, Chifengsan (*Hugh!*). Mus. Brit.

North temperate regions.

9. *Poa prolixior*, Rendle, n. sp.

*Herba* perennis?, glabra; culmi tenuissimi, 27 poll. alti, vix  $\frac{1}{3}$  lin. crassi, basi ascendentes et geniculati, 4-nodi; nodum supremum  $\frac{1}{4}$  culmi longit. a basi locatum. *Foliorum* vaginæ ut culmi scabridule striatæ, strictæ, quam internodia breviores, inferæ albido-membranaceæ quam suprema breviores; ligula lata, superne fimbriata, ad  $1\frac{1}{2}$  lin. longa; laminæ angusto-lineares ad apicem acutum attenuatæ, suprema quam inferiores longior,  $3\frac{1}{2}$ – $4\frac{1}{3}$  poll. longa,  $\frac{1}{2}$  lin. lata, ejus vaginæ vix æquans vel eam paullo excedens. *Panicula* stricta, flexuosa, interdum secunda, ad 3 poll. longa; rami gemini erecti vel suberecti, in parte inferiore (circa  $\frac{1}{3}$ ) nudi, scabriduli ut in ramulis brevibus pedicellisque, in parte superiore sublaxiter spiculiferi; spiculæ lanceolatæ,  $1\frac{1}{2}$  lin. longæ, pedicellos excedentes, 2-floræ. *Glumæ* firmæ, steriles lanceolatæ, acutæ, trinerves, cum carina scabridula, et marginibus anguste membranaceis,  $1\frac{1}{4}$  lin. longæ; fertiles duplicatæ oblongæ vel oblongo-lanceolatæ, sub apice (ad  $\frac{1}{5}$  longit.) membranaceæ, 5-nerves, nervo marginale ad basin sparse piloso, carina ad medium pilosa, callus sparsius contortuplicato-lanatus; inferior sterilibus subæquans vel paullo longior; palea in carinis supra medium minute ciliolata; antheræ  $\frac{1}{2}$  lin. longæ.

HUPH: Patung, in grass, 6000 ft. (*A. Henry*, 6049!); SZECHUEN: Mt. Omei (*Faber*, 1147!). Herb. Kew.

A distinct species of the *Poa nemoralis* group, characterized by its very slender elongated stem, developed ligule, strict inflorescence, and small few-flowered spikelets.

10. *Poa spondylodes*, *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 71; *Steud. Syn. Gram.* p. 256; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Franchet, Pl. David.* i. p. 338; *Palibin, in Act. Horti Petrop.* xix. (1901) p. 133; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 280; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 175.

*Poa linearis*, *Trin. l. c.* and its var. *subdiantha*, *Franchet, in Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 269, may belong here.

*Poa strictula*, *Steud. l. c.* p. 426.

*Poa palustris*, *Linn.*, var. *strictula*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 710, et sér. 2, iii. (1903) p. 506.

CHIHLI: Jehol (*David*, in herb. *Hance*, 15612 in part!), Peking (*Bushell*!), mountains near Peking (*Bretschneider*, in herb. *Hance*, 15612 in part!); SHINGKING: Sarhoo (*Ross*, 337!), East of Fooling (*Ross*, 359!), Chienshan (*Ross*, 526!), Liao River, west of Liaoyang (*Ross*, 11!); KIANGSU: Shanghai (*Perry*! *Forbes*, 1073!); HUPEH: Patung (*Wilson*, 1640!); SHENSI: Mt. Laoysan, Tcuizscien (*Hugh*!), Paoki, Chifengsan (*Hugh*!); COREA: East (*Perry*!), Soul (*Sontag*!); COREAN ARCHIPELAGO (*Oldham*, 946!). Mus. Brit.; Herb. Kew.

Japan.

11. *Poa viridula*, *Palibin*, in *Act. Horti Petrop.* xix. (1901) p. 134.

COREA: Soul, Mt. Yisan (*Sontag*!). Mus. Brit.

1. *Glyceria aquatica*, *Wahlenb. Fl. Goth.* p. 18; *Smith, Engl. Fl.* p. 116; *Franchet, Pl. David.* i. p. 338; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 711, et sér. 2, iii. (1903) p. 506; *Ledeb. Fl. Ross.* iv. p. 392; *Turez. Fl. Baic.-Dahur.* ii. p. 334; *Regel, Tent. Fl. Ussur.* p. 168.

CHIHLI: Jehol, banks of streams (*David*, in herb. *Hance*, 3617!); SHINGKING: between Tungcheshien and Maorshan, Yalu River (*James*!); CHEKIANG: Ningpo Mountains (*Faber*, 1735!). Mus. Brit.; Herb. Kew.

Japan, North and Central Asia; Europe.

2. *Glyceria distans*, *Wahlenb. Fl. Upsal.* p. 36; *Hook. f. Fl. Brit. Ind.* vii. p. 347.

*Atropis distans*, *Griseb. in Ledeb. Fl. Ross.* iv. p. 388; *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 506; *Kantz, in Math. u. Naturw. Ber. aus Ungarn*, iii. (1884) p. 9.

CHIHLI: Peking (*David*! *Bretschneider*, 887!); COREA: wet places near Chinampo (*Faurie*, 865, 873, fide *Hackel*). Herb. Kew.

Europe, North and Central Asia (Western Tibet).

The plant from Peking collected by David is probably the one referred by Franchet (*Pl. David.* i. p. 338) to *Glyceria convoluta*, Fries, which in the 'Flora of British India' is regarded as a variety (var. *convoluta*) of *G. distans*.

3. *Glyceria procumbens*, Dumort. *Obs. Gram. Fl. Belg.* p. 145.

*Festuca procumbens*, Kunth, *Rév. Gram.* p. 129.

*Poa procumbens*, Curt. *Fl. Lond.* fasc. vi. p. 11.

KIANGSU: Shanghai (*Maingay*, 579!). Herb. Kew.

Europe; probably an introduction in China.

1. *Festuca gigantea*, Vill. *Hist. Pl. Dauph.* ii. p. 110; Kunth, *Enum. Pl.* i. p. 407; Hack. in *Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 506; Komarow, in *Act. Horti Petrop.* xx. (1901) p. 311; Ledeb. *Fl. Ross.* iv. p. 354; Maxim. *Prim. Fl. Amur.* p. 318; Stapf, in *Hook. f. Fl. Brit. Ind.* vii. p. 353.

SHINGKING: Chienshan (*Ross*, 444!). Herb. Kew.

Japan, North Asia, Temperate Himalaya; Europe.

A depauperate form.

2. *Festuca modesta*, Steud. *Syn. Gram.* p. 316; Stapf, in *Hook. f. Fl. Brit. Ind.* vii. p. 354.

HUPEH: South Patung (*A. Henry*, 5345!), Patung (*Wilson*, 1643!). Mus. Brit.; Herb. Kew.

Himalayas.

Wilson's specimen has a longer ligule than usual in the species, reaching sometimes a quarter of an inch or more in length; but I find considerable variation in this character both in the Indian and Chinese specimens.

3. *Festuca ovina*, Linn. *Sp. Pl.* p. 73; var. *vulgaris*, Koch, *Syn.* ii. p. 812; Hack. *Mon. Festuc. Europ.* p. 86, et in *Bull. Herb. Boiss.* vii. (1899) p. 713, et sér. 2, iii. (1903) p. 506; Palibin, in *Act. Horti Petrop.* xix. (1901) p. 35.

SHENSI: Mt. Miaowansan (*Hugh*!); HUPEH: Hsingshan, 10,000 ft. (*A. Henry*, 6966!); COREA: dry places on Mt. Namsan near Soul (*Faurie*, 879, fide *Hackel*). Mus. Brit.; Herb. Kew.

Europe, North Asia.

4. *Festuca parvigluma*, Steud. *Syn. Gram.* p. 305; Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 713, et sér. 2, iii. (1903) p. 506; Palibin, in *Act. Horti Petrop.* xix. (1901) p. 35.

CHEKIANG: Ningpo Mountains (*Faber*, 300!); HUPEH: Chien-shih (*A. Henry*, 5792!); COREA: Port Hamilton (*Wilford*, 701!). Herb. Kew.

Japan.

1. *Bromus asper*, *Murr. Prodr. Stirp. Gotting.* p. 42; *Steud. Syn. Gram.* p. 320; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 358; *Ledeb. Fl. Ross.* iv. p. 356.

KIANGSU: Shanghai (*Maingay*, 708 !); CHEKIANG: Ningpo Mountains (*Faber*, 1736 !); HUPEH: Ichang (*A. Henry*, 3560 !), Hsingshan, 8000–9000 ft. (*A. Henry*, 6994 !); HONGKONG: Putoy Isle, sand of sea-shore (*Faber*, 18 !). *Mus. Brit.*; *Herb. Kew.*

Europe, North Africa, West and North Asia, and Hill Country of India.

2. *Bromus japonicus*, *Thunb. Fl. Jap.* p. 51; *Steud. Syn. Gram.* p. 326; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 283; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 181; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 714, et sér. 2, iii. (1903) p. 506; *Palibin, in Act. Horti Petrop.* xix. (1901) p. 135; *Hance, in Journ. Bot.* xvi. (1878) p. 234.

*Bromus patulus*, *Mert. & Koch, in Röhl. Deutschl. Fl.* i. p. 685; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 361.

SHANTUNG: Chefoo (*Swinhoe! Perry!*); KIANGSI: Kiukiang (*Forbes*, 18 !), Shanghai (*Maingay*, 352 ! *Faber*, 289 !), Woosing River, Shanghai (*Perry!*), Nanking and Wan City (*Faber*, 1158 !), Chapu (*Oldham*, 19 !); HUPEH: Ichang (*A. Henry*, 4033, 3523 !), Patung (*A. Henry*, 4732 !), West Patung (*Wilson*, 1649 !); SHENSI: Paoki, Chifengsan (*Hugh!*), Kusan (*Hugh!*), Kaolinsien (*Hugh!*), Mt. Laoysan, Teuizscien (*Hugh!*), Hukiascien (*Hugh!*); SZECHUEN (*Parker, in herb. Hance*, 17893 in part !); COREA: Port Chusan (*Wilford*, 929 !), East Corea (*Perry!*). *Mus. Brit.*; *Herb. Kew.*

Japan, Temperate Asia; Europe.

3. *Bromus pauciflorus*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 713, et sér. 2, iii. (1903) p. 506.

*Festuca pauciflora*, *Thunb. Fl. Jap.* p. 52.

*Festuca remotiflora*, *Steud. Syn. Gram.* p. 315; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226.

*Schedonorus remotiflorus*, *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 283; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 181; *Franchet, Pl. David.* i. p. 339, ii. p. 146.

SZECHUEN: Moupin (*David, fide Franchet*).

Japan.



4. *Bromus squarrosus*, *Linn. Sp. Pl.* p. 76; *Steud. Syn. Gram.* p. 325; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Ledeb. Fl. Ross.* iv. p. 363.

SZECHUEN: Nanchuan (*Bock, fide Pilger*).

West and North Asia; Europe.

5. *Bromus tectorum*, *Linn. Sp. Pl.* p. 77; *Steud. Syn. Gram.* p. 318; *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 359.

SZECHUEN: Tachienlu (*Pratt, 198!*). *Mus. Brit.*; *Herb. Kew.*

Europe, North Africa, Western and Northern Asia, Himalayas.

1. *Brachypodium silvaticum*, *Beauv. Agrost.* p. 155; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 286; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 185; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 714; *Baker & Moore, in Journ. Linn. Soc., Bot.* xvii. (1879) p. 388; *Hook. f. Fl. Brit. Ind.* vii. p. 363; *Ledeb. Fl. Ross.* iv. p. 346.

SHINGKING: Chienshan (*Ross, 580!*); HUPEH: Fang, 7000 ft. (*A. Henry, 6745!*), Patung (*A. Henry, 4721!*); SZECHUEN: Tongolo (*Soulié, 246!*); YUNNAN: Mengtze, woods at 4500 ft. (*A. Henry, 11255!*). *Mus. Brit.*; *Herb. Kew.*

Japan, North Asia, mountains of India; Europe.

#### Tribe HORDEÆ.

1. *Agropyrum caninum*, *Beauv. Agrost.* p. 146; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 714.

*Triticum caninum*, *Schreb. Spicil. Fl. Lips.* p. 51; *Steud. Syn. Gram.* p. 344; *Trin. in Bunge, Enum. Pl. Chin. Bor.* p. 72; *Maxim. Prim. Fl. Amur.* p. 478.

CHINA: East Coast (*Home!*); CHIHLI: mountains west of Peking (*Bretschneider, 856!*), Peking (*Bushell!*); KIANGSU: Shanghai (*Perry!*); HUPEH: Patung (*A. Henry, 4726!*), Ichang (*A. Henry, 197, 798!*); SZECHUEN (*A. Henry, 5873!*); COREA: East (*Perry!*); COREAN ARCHIPELAGO: Tsusima Islands (*Wilford, 779!*). *Mus. Brit.*; *Herb. Kew.*

Japan, Europe, North America.

2. *Agropyrum ciliare*, *Franchet, Pl. David. i. p. 341*; *Palibin, in Act. Horti Petrop. xix. (1901) p. 135*; *Komarow, ibid. xx. (1901) p. 315*.

*Agropyrum semicostatum*, *Nees, var. ciliare, Hack. in Bull. Herb. Boiss. sér. 2, iii. (1903) p. 506*.

*Triticum ciliare*, *Trin. in Bunge, Enum. Pl. Chin. Bor. p. 72*; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg, xxiv. (1884) p. 271*; *Maxim. Prim. Fl. Amur. p. 478*.

*Brachypodium chinense*, *S. Moore, in Journ. Bot. xiii. (1875) p. 230*.

CHIHLI: mountains west of Peking (*Bretschneider, 835!*), hills about Peking (*Williams, in herb. Hance, 13838 in part!*), dry hills, Jehol (*David, in herb. Hance, 13838 in part!*), Mt. Pohuashan (*Moellendorff, in herb. Hance, 13838 in part!*); SHANTUNG: Chefoo (*Perry!*); KIANGSI: Kiukiang (*Shearer!*); HUPEH: Ichang (*A. Henry, 3486, 201!*); SHENSI: Kusan (*Hugh!*); SHINGKING: Moukden to Laioyang (*Ross, 341!*), Chienshan (*Ross, 598!*); COREA: Soul (*Sontag!*), East Corea (*Perry!*). Mus. Brit.; Herb. Kew.

North Asia.

3. *Agropyrum repens*, *Beauv. Agrost. p. 146*; *Hook. f. Fl. Brit. Ind. vii. p. 370*.

*Agropyrum pseudo-agropyrum*, *Franchet, Pl. David. i. p. 340*; *Hack. in Bull. Herb. Boiss. sér. 2, iii. (1903) p. 507*; *Komarow, in Act. Horti Petrop. xx. (1901) p. 315*.

*Triticum pseudo-agropyrum*, *Griseb. in Ledeb. Fl. Ross. iv. p. 343*; *Maxim. Prim. Fl. Amur. p. 318*; *Franchet, in Mém. Soc. Sci. Nat. Cherbourg, xxiv. (1884) p. 271*.

*Triticum chinense*, *Trin. in Bunge, Enum. Pl. Chin. Bor. p. 72*; *Maxim. l. c. p. 478*.

*Triticum repens*, *Linn. Sp. Pl. p. 86*.

*Elymus pseudo-agropyrum*, *Trin. in Bull. Soc. Nat. Mosc. 1838, p. 105 (nomen), et in Turcz. Fl. Baic.-Dahur. ii. p. 350*.

CHIHLI: Jehol, in dry places (*David, in herb. Hance, 15614 in part!*), Peking (*Bretschneider, in herb. Hance, 15614 in part!*); KIANGSU: Shanghai (*Forbes!*). Mus. Brit.; Herb. Kew.

North Asia, Himalayas; widely distributed in temperate regions.

4. *Agropyrum semicostatum*, *Nees, ex Steud. Syn. Gram. p. 346*; *var. transiens, Hack. in Bull. Herb. Boiss. sér. 2, iii. (1903) p. 507*.

COREA: sand of the river Pyengyang (*Faurie*, 841, fide *Hackel*).

5. *Agropyrum strigosum*, *Franchet, Pl. David.* i. p. 340.

*Triticum strigosum*, *Less. in Linnæa*, ix. (1834) p. 170; *Maxim. Prim. Fl. Amur.* p. 478.

CHIHLI: Jehol (*David*, 1878, fide *Franchet*).

1. *Lepturus repens*, *R. Br. Prodr.* p. 207; *Steud. Syn. Gram.* p. 357; *Hook. f. Fl. Brit. Ind.* vii. p. 365.

FORMOSA: Takow (*A. Henry*, 756, 1174, 1202!). Mus. Brit.; Herb. Kew.

Ceylon, Malaya, Pacific Islands, Australia, Mascarene Is., South Africa.

1. *Elymus arenarius*, *Linn. Sp. Pl.* p. 83; var. *coreensis*, *Hack. in Bull. Herb. Boiss. sér. 2*, iii. (1903) p. 507.

COREA: Ouensan (*Faurie*, 843, fide *Hackel*).

The species occurs in Europe including Arctic, North Asia and Japan.

2. *Elymus dahuricus*, *Trin. in Bull. Soc. Nat. Mosc.* 1838, p. 105 (nomen), et *Fl. Altaic.* ii. p. 348; *Steud. Syn. Gram.* p. 348; *Franchet, Pl. David.* i. p. 342, et ii. p. 146; *Komarow, in Act. Horti Petrop.* xx. (1901) p. 318; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226; *Ledeb. Fl. Ross.* iv. p. 331; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 715.

CHIHLI: neighbourhood of Peking (*David*, 562! *Bretschneider*, 846!); HUPEH: Patung (*A. Henry*, 4709!); SZECHUEN: Moupin (*David*, fide *Franchet*). Herb. Kew.

Himalayas, Central Asia, North Asia.

Var. *cylindricus*, *Franchet, Pl. David.* i. p. 342.

CHIHLI: neighbourhood of Peking (*David*, fide *Franchet*).

3. *Elymus dasystachys*, *Trin. in Ledeb. Fl. Altaic.* i. p. 120; *Steud. Syn. Gram.* p. 350; *Franchet, Pl. David.* i. p. 341; *Komarow, in Act. Horti Petrop.* xx. (1901) p. 320; *Hook. f. Fl. Brit. Ind.* vii. p. 374; *Ledeb. Fl. Ross.* i. p. 333.

SHENSI: in the mountains (*David*, in herb. *Hance*, 7960!). Mus. Brit.

Central and North Asia, Himalayas.

4. *Elymus excelsus*, Turcz. in *Bull. Soc. Nat. Mosc.* p. 105 (nomen), et *Fl. Altaic.* ii. p. 349; *Steud. Syn. Gram.* p. 348; *Hack. in Bull. Herb. Boiss.* sér. 2, iii. (1903) p. 507; Komarow, in *Act. Horti Petrop.* xx. (1901) p. 319; *Ledeb. Fl. Ross.* iv. p. 331.

COREA: Ouensan (*Faurie*, 844, fide *Hackel*).

North Asia.

5. *Elymus sibiricus*, *Linn. Sp. Pl.* p. 83; *Steud. Syn. Gram.* p. 348; *Franchet, Pl. David.* i. p. 341; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 715; Komarow, in *Act. Horti Petrop.* xx. (1901) p. 317; *Ledeb. Fl. Altaic.* i. p. 123, et *Fl. Ross.* iv. p. 330; *Regel, Tent. Fl. Ussur.* p. 167; *Maxim. Prim. Fl. Amur.* p. 317; *Hook. f. Fl. Brit. Ind.* vii. p. 373.

SHINGKING: Laoyehling and other hills near Moukden (*James!*);

SHENSI: Mt. Miaowansan (*Hugh!*). *Mus. Brit.*; *Herb. Kew.*

Himalayas, Central and North Asia; America.

1. *Asprella Duthiei*, *Stapf, in Hook. f. Fl. Brit. Ind.* vii. p. 375, et in *Hook. Icon. Pl.* t. 2525; *Pilger, in Engl. Bot. Jahrb.* xxix. (1900) p. 226.

HUPÈH: Chienshih (*A. Henry*, 5918!). *Herb. Kew.*

Western Himalaya.

#### Tribe BAMBUSEÆ.

1. *Arundinaria densiflora*, *Rendle*, n. sp.

*Frutex*; internodia glabra. *Foliorum* vaginæ, in ramis floriferis, glabræ, apice breviter marginatæ et ore fimbriatæ; ligula brevis, truncata; lamina lineari-lanceolata, acuta,  $2\frac{1}{2}$ -4 poll. longa, ad (vix)  $\frac{3}{4}$  poll. lata, in petiolum brevem attenuata, margine subaspera, venæ secundariæ utrinque 5, interspatiis minute et regulariter tessellatis. *Racemi* parvi, densi, pauci-(circa 5-) spiculiferi, in ramulis florentibus approximati, axis brevis cum bracteis sursum majoribus imbricatis indutus; spiculæ sæpe 5-6-floræ, purpureo-tinctæ, subsessiles. *Glumæ* imbricatæ, rhachillæ articuli crassi clavati puberuli, gl. steriles, solitariae vel binæ et inæquales, fertilibus similes sed breviores; gl. fertiles late ovatæ, cuspidatæ, dorso vix carinatæ, cum venulis secundariis utrinque 4 prominulis, dorso hirsutulæ, inferior  $4\frac{1}{2}$  lin.

longa; palea subæquilonga vel paullo longior, dorso et in carinis binis hirsutula, apice bilobulata; lodiculæ 3, obovatæ, superne rotundatæ et ciliatæ,  $1\frac{1}{2}$  lin. longæ; antheræ 3,  $2\frac{1}{2}$  lin. longæ; ovarium glabrum, stylus brevis, stigmata 3 plumosa.

KIANGSU: Kiangsu hills (*Faber*, 19!); CHEKIANG: hills near Huchan, Taihoo Lake (*Carles*, 227!). Mus. Brit.; Herb. Kew.

A distinct species characterized by the short crowded groups of spikelets, closely arranged on the slender flowering shoots which are more or less densely associated and overtopped by leafy shoots.

## 2. *Arundinaria Faberi*, *Rendle*, n. sp.

*Frutex* bipedalis; internodia in parte superiore florifera glabra,  $\frac{2}{3}$  ad  $\frac{1}{3}$  poll. longa, ramuli floriferi fasciculati; vaginæ aphyllæ internodia excedentes, glabræ, margine et apice ut in foliaceis. *Foliorum* vaginæ strictæ, sese involventes, glabræ, margine ciliolatæ, apice breviter marginatæ et cum auriculis fimbriatis instructæ; ligula brevis, truncata; lamina anguste lanceolata,  $\pm$  acuminata, ad 3 poll. longa,  $\frac{1}{3}$  poll. lata, in petiolum brevem attenuata, glabra, margine aspera, subter asperula, venæ secundariæ utrinque 3, interspatiis minute et regulariter tessellatis. *Racemi* terminales, lineares, semipedales, rhachis gracilis ultra foliæ ultimæ vaginam longe (circa 4 poll.) exserta, glabra; spiculæ paucæ, multifloræ (sæpe 6–10) ad  $2\frac{1}{2}$  poll. longæ, atro-purpureæ, pedicellis tenuibus flexuosis. *Glumæ* steriles valde inæquales, inferior lanceolata, acuta,  $1\frac{3}{4}$  lin. longa, uninervis, et dorso carinata, superior, ovata, trinervis, carinata, abrupte acuta,  $3\frac{1}{2}$  lin. longa; rhachillæ articuli clavati, margine et apice ciliolati; glumæ florentes dissitæ, ovatæ, acuminatæ, dorso carinatæ cum venulis secundariis utrinque 3 prominentibus, dorso hirsutulæ, inferior  $\frac{1}{2}$  poll. longa; palea brevior, bidentula, in carinis et dorso sub apice hirsutula; lodiculæ ovatæ, obtusæ, venulosæ; stigmata 3 plumosa.

SZETCHUEN: above Chingking, Yangtzekiang (*Faber*, 1119!). Herb. Kew.

Approaches the Himalayan species *Arundinaria racemosa*, Munro, but differs in the longer-exserted panicles and the large upper barren glume.

3. *Arundinaria flexuosa*, Hance, in *Journ. Bot.* xiv. (1876) p. 339.

KWANGTUNG: Danes Island, Whampoa (Nevin, in herb. Hance, 19336!). Mus. Brit.

4. *Arundinaria Hindsii*, Munro, in *Trans. Linn. Soc.* xxvi. (1868) p. 31.

HONGKONG (*Hinds!*). Herb. Kew.

Known only from leaf-specimens.

5. *Arundinaria linearis*, Hack. in *Bull. Herb. Boiss.* vii. (1899) p. 721.

LUCHU ARCHIPELAGO (*Tashiro, fide Hackel*).

6. *Arundinaria nitida*, Mitford, in *Gard. Chron.* ser. III. xviii. (1895) p. 186, t. 33, et *Bamboo Garden*, p. 73 (nomen); Stapf, in *Kew Bull.* 1896, p. 20; Pilger, in *Engl. Bot. Jahrb.* xxix. (1900) p. 226.

*Arundinaria khasiana*, Bean, in *Gard. Chron.* ser. III. xv. (1894) p. 301, non *Munro*.

HUPEH: Fang district, on cliffs, 6000–9500 ft. (*A. Henry*, 6832!), mountains, Chienshin (*Wilson*, 954!), West Hupeh (*Wilson*, 1040 A!); SZECHUEN: North (*Potanin!*), mountains, South Wushan (*Wilson*, 1040!). Mus. Brit.; Herb. Kew.

7. *Arundinaria sinica*, Hance, in *Ann. Sci. Nat.* sér. 4, xviii. (1862) p. 235, et *Journ. Linn. Soc., Bot.* xiii. (1873) p. 137.

*Arundinaria longiramea*, Munro, in *Trans. Linn. Soc.* xxvi. (1868) p. 19.

*Arundinaria Wightii*, Benth. *Fl. Hongk.* p. 434.

HONGKONG (*Ford*, in herb. *Forbes!* Hance, 1945! *Wright!*), Mount Victoria and Happy Valley Woods (*Lamont!*). Mus. Brit.; Herb. Kew.

8. *Arundinaria sparsiflora*, Rendle, n. sp.

*Frutex* sexpedalis; culmi teretes, leves, in specimine  $1\frac{1}{2}$  lin. crassi; rami fasciculati, in ramulis pluribus brevibus foliiferis et floriferis soluti; foliiferi juveniles (qui solum suppetunt)  $1\frac{1}{2}$  poll. longi, a vaginis pupurascentibus velati. *Foliorum* vaginæ striatæ cum venulis transversalibus sparsis, ore cum fimbriis paucis, mox deciduis instructæ; ligula brevis, truncata; lamina lanceo-

lata, acuminata, brevissime petiolata, ad  $1\frac{1}{2}$  poll. longa,  $2\frac{1}{2}$  lin. lata, glabra, venæ secundariæ 4 a venulis transversalibus creberrimis conjunctæ, margine scabridula. *Racemi* laterales breves,  $1-1\frac{1}{3}$  poll. longi, a bracteis sursum majoribus e basi vaginati; bracteæ 3 inferiores, 2-9 lin. longæ, membranaceæ (infima carinata), 2 sequentes foliaceæ cum lamina  $\frac{1}{2}-1$  poll. longa, ceteræ (ad 5) sursum breviores,  $2\frac{1}{2}-6$  lin. longæ, membranaceæ et sine lamina, quæque spiculam fovens (spiculæ sæpe 3-floræ, glumæ steriles 2, vel in spiculis superioribus 1, fertilibus similes sed breviores), rhachis glabra, cum glumis fertilibus pluribus (sæpe 6 quarum suprema  $\pm$  abortiva est) terminata. *Glumæ* fertiles adultæ late lanceolatae, breviter aristatae, dorso hirsutulæ, cum venulis secundariis 5 prominulis,  $5\frac{1}{2}-6$  lin. longæ, paleæ c.  $4\frac{1}{2}$  lin. longæ, binerves, in carinis et dorso hirsutulæ, apice bidentulæ; lodiculæ 3, falcate ovatae, superne ciliolatae, vix 1 lin. longæ; antheræ 3, 3 lin. longæ; ovarium glabrum, stylus brevis, stigmata tria plumosa.

ИУРЕН: Hsingshan, throughout conifer woods on mountains, 8000 to 9500 ft. (*A. Henry*, 6938!). Mus. Brit.; Herb. Kew.

A distinct species belonging to the section *Thamnocalamus*, and suggesting the Japanese *Arundinaria vaginata*, Hack. (in Bull. Herb. Boiss. vii. p. 717), which I have not seen, but differing in the details of arrangement of the inflorescence and other points.

#### 9. *Arundinaria Wilsoni*, Rendle, n. sp.

*Frutex*; internodia teretia, glabra, ad  $1\frac{2}{3}$  lin. crassa; vaginæ culmi internodia (in parte saltem superiore) excedentes, chartaceæ, glabræ, ramis floriferis arcte circumvolutæ; ligula brevis, truncata. *Foliorum* vaginæ strictæ, arcte convolutæ, glabræ; lamina oblongo-lanceolata, acuminata, basi in petiolum brevem attenuata, ad 5 poll. longa, 11 lin. lata, superne glabra, margine aspera, subter pilosula, venæ secundariæ utrinque 4-6. *Rami floriferi* ad nodos culmi ultimos fastigiati, rarius solitarii; paniculæ parvæ, effusæ, ad  $2\frac{3}{4}$  poll. longæ, et 3 poll. latæ, pedicelli velut rhachi glabri, graciles, flexuosi, sæpius patentés, interdum nutantes; spiculæ purpurascens, sæpius 5-7-floræ, ad 1 poll. longæ; glumæ steriles distantes, inæquales, ovatae, obtusæ vel subacutæ, carinatae, in marginibus præcipue superioribus puberulæ, inferior 5-nervis,  $1\frac{1}{2}$  lin. longa, superior

7-nervis,  $2\frac{2}{3}$  lin. longa; florentes dissitæ (rhachillæ articuli pilosuli), late ovatæ, obtusæ vel breviter acutæ basi rhachillam amplectantes, super basin carinatae, 7-nerves, pilosulæ præcipue ad marginem, gl. infima 4 lin. longa, palea brevior ( $3\frac{1}{2}$  lin. longa) obtusa, in carinis et ad latera puberula; lodiculæ ovatæ, sub apice fimbriatæ; ovarium glabrum; stylus brevis in stigmata tria plumosa solutus. *Caryopsis* oblonga, brunnea, glabra,  $1\frac{2}{3}$  lin. longa, cum basi styli persistente coronata.

HUPEH: Fang, forms scrub on top of highest mountains, 7700 to 9600 ft. (*Wilson*, 1887!). Herb. Kew.

Approaches *Arundinaria elegans*, Kurz, from further India, but differs in the effuse panicle, the markedly unequal barren glumes, and the blunter flowering glumes.

1. *Fargesia spathacea*, *Franchet*, in *Bull. Soc. Linn. Paris*, 1893, p. 1067; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 226.

SZECHUEN: mountains in Tchenkeoutin district (*Farges*, fide *Franchet*).

1. *Phyllostachys bambusoides*, *Sieb. & Zucc.* in *Abh. Akad. Wiss. Münch., Math.-phys. Class.* iii. (1843) p. 746, t. 5. fig. 3; *Munro*, in *Trans. Linn. Soc.* xxvi. (1870) p. 36; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 227; *Miq.* in *Ann. Mus. Bot. Lugd.-Bat.* ii. p. 284; *Franch. et Savat. Enum. Pl. Jap.* ii. pp. 182, 605; *Hack.* in *Bull. Herb. Boiss.* vii. (1899) p. 718; *Gamble*, *Bamb. Brit. Ind.* p. 27, et in *Hook. f. Fl. Brit. Ind.* vii. p. 386.

CHINA: without locality (*De Montigny*!); FOKIEN: Amoy (*De Grijs*, in herb. *Hance*, 2073!); HUPEH: Nanto (*A. Henry*, 3867, 4405, 4420, 7900! *Wilson*, 230!). Mus. Brit.; Herb. Kew.

Japan, Assam.

2. *Phyllostachys congesta*, *Rendle*, n. sp.

*Suffrutex* vel *frutex*; internodia semiteretia, glabra. *Foliorum* vaginæ (in ramulis floriferis) glabræ, striatulæ, ad sesquipollicares, ore fimbriis ad 3 lin. longis instructæ; ligula brevis, truncata, ciliolata; lamina lanceolata, acuta, ad  $3\frac{1}{4}$  poll. longa et  $\frac{2}{3}$  poll. lata, basi rotundata, breviter petiolata, margine scabridula, venæ secundariæ utrinque 4-6, subter vix prominulæ, interspatiis



minute tessellatis. *Paniculæ* multispiculiferæ, congestæ, capituliformes, circa  $\frac{3}{4}$  poll. longæ, ramulos foliiferos terminantes et a vaginis spathiformibus foliorum ultimorum (qui transitum ad bracteas monstrant) primo obsessæ; bracteæ primariæ subfoliaceæ, superiores breviores, anguste lanceolatae, acutæ; spiculæ 3-5-floræ (flos suprema imperfecta), rhachilla glabra. *Gluma* sterilis bracteis superioribus subconformis, angusta, e basi ad apicem elongatum subpungentem attenuata, glumam fertilem infimam subæquans, superne carinulata et sparsius hirsutula; glumæ fertiles oblongæ apice longius pungentes, dorso rotundato hirsutæ, multinerves, infima sæpius semipollicaris; palea paullo brevior (e. g. 5 lin. longa), cum carinis prominulis, dorso superne hirsuta, apice bidentula; lodiculæ inæquales, 2 majores obovatae, apice ciliatæ, vix  $1\frac{1}{2}$  lin. longæ, tertia conformis sed duplo minor; stamina 3; ovarium pedicellatum, stylus diu persistens superne in stigmata 3 plumosa divisus.

HUPEH: Patung (*A. Henry*, 4828! "used for paper-making"), Changyang (*A. Henry*, 6238!), Ichang (*A. Henry*, 1698!), West Hupeh (*Wilson*, 230 A!). Herb. Kew.

The species is characterized by the short dense capitate panicles terminating the leafy shoots. No. 1698 is described by Dr. Henry as a "very small bamboo in ditch"; No. 6238 as 10 ft. high.

### 3. *Phyllostachys Faberi*, Rendle, n. sp.

*Frutex*; internodia glabra. *Foliorum* vaginæ glabræ, striatulae, ore longe fimbriatæ, fimbriæ ad 5 lin. longæ; ligula truncata,  $1\frac{1}{2}$  lin. longa; lamina lineari-lanceolata, acuta vel acuminata, in ramulis sterilibus ad 6 poll. longa et 8 lin. lata, in petiolum bilineum (vel brevior) attenuata, margine altero scabridula, altero levis, venæ secundariæ utrinque 4-6, interspatiis minute tessellatis. *Rami florentes* elongati, aphylli, internodia unilateraliter complanata, levia; ramuli primarii solitarii subpedales, ad originem et ad nodos quosque distantes ramulos secundarios solitarios ad 4 poll. longos gerentes, internodia ramulique sursum deminuti; ramuli ultimi tenues spiculiferi; bracteæ magnæ, membranaceæ, spathiformes, ad  $1\frac{1}{3}$  poll. longæ; spiculæ trifloræ (flos supremus autem imperfectus) distantes. *Gluma* sterilis vaginæformis, sæpe cum ligula obsoleta terminata, 1 poll. vel longior; fertilis inferior gluma sterili ab internodio bilineo clavato puberulo sejuncta, lineari-lanceolata,

acuta, multinervis, dorso rotundata et scabridula, pollicaris vel paullo longior; palea quam gluma paullo vel vix brevior, angusta, multinervis, dorso in parte superiore præcipue in carinis geminis approximatis scabridula, apice bidentula; lodiculæ lineari-lanceolatæ, acutæ,  $2\frac{2}{3}$  lin. longæ; ovarium pedicellatum, napiforme; stylus longus diu persistens.

CHEKIANG: Ningpo Mountains (*Faber*!). Herb. Kew.

Resembles *Phyllostachys Stauntonii*, Munro, in the prominent ligule and long fimbriæ of the leaf-sheath, but differs in the much coarser-looking inflorescence.

4. *Phyllostachys flexuosa*, A. & C. Rivière, in *Bull. Soc. Acclimat.* sér. 3, v. (1878) p. 758.

Stated by the authors to have been received by the Paris Société d'Acclimatation from the cold parts of China. It is apparently near *Phyllostachys Nevinii*, Hance.

I have seen only leaf-specimens from plants grown in Kew Gardens.

5. *Phyllostachys Henryi*, Rendle, n. sp.

*Frutex* 20-pedalis alta; rami quadrangulares, glabri. *Foliorum* vaginæ in ramulis floriferis glabræ, leves, stramineæ,  $\frac{3}{4}$  poll. longæ, ore fimbriatæ (fimbriæ  $1\frac{1}{2}$  lin. longæ); ligula brevis, truncata; lamina anguste lanceolata, acuta,  $1\frac{1}{2}$ – $2\frac{1}{2}$  poll. longa,  $\frac{1}{4}$  ad vix  $\frac{1}{2}$  poll. lata, in petiolum brevem attenuata, margine scabridula, venæ secundariæ utrinque 4, interspatiis minute tessellatis. *Paniculæ* densæ, pollicares, ramulos ultimos terminantes et a foliis supremis geminis approximatis velut cum spatha vaginata; rhachis pilosula, brevis, ramulosa, bracteæ inferiores subfoliaceæ cum lamina breve acuta instructæ, superiores membranaceæ, lanceolatæ, acutæ; spiculæ circa 5, sessiles, bifloræ, flos superior imperfecta. *Gluma* sterilis inter bracteam præcedentem et glumam fertilem forma intermedia, lanceolata, acuta, multinervis, superne sparsius hirsutula; fertilis 7 lin. longa, lanceolata, apice pungens, dorso rotundato hirsutula, cum venulis parallelis utrinque 6 haud prominulis; palea paullo brevior (6 lin. longa), multinervis, carinis approximatis, dorso minute hirsutula, apice bidentula; lodiculæ 3, subæquales, lanceolatæ, acutæ,  $1\frac{3}{4}$  lin. longæ; ovarium pedicellatum; stylus in stigmata 3 plumosa superne divisus.

HUPEH: Nanto (*A. Henry*, 6338!). Mus. Brit.; Herb. Kew.

Near *Phyllostachys Stauntoni*, Munro, from which it differs in the shorter ligule and shorter fimbriæ at the mouth of the sheath, and in the solitary dense panicles situated at the end of short leafy branches.

6. *Phyllostachys heteroclada*, *Oliver*, in *Hook. Icon. Pl.* t. 2288; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 227.

KIANGSI: Kiukiang (*Shearer*!), native name Man-tsze; HUPEH: West (*Wilson*, 332, 332 A!), Nanto and mountains to northward (*A. Henry*, 1932, 3828!); SZECHUEN (*A. Henry*, 8833!), West Szechuen and Tibetan frontier chiefly near Tachienlu, 9000–13,500 ft. (*Pratt*, 384!). Herb. Kew.

7. *Phyllostachys montana*, *Rendle*, n. sp.

*Frutex*. *Folia* papyracea; vaginæ subcompressæ, glabræ, densiter striatæ, apice pulvinatæ; ligula  $\frac{1}{2}$  lin. longa, truncata, longe fimbriata; lamina lineari-oblonga ad apicem acuminatum attenuata, 8 poll. longa, vix 1 poll. lata, margine utrinque scabridula, subter sparse pilosula, venæ secundariæ utrinque 6–7, venulæ inconspicuæ infrequenter et irregulariter transverse junctæ; petiolus brevissimus. *Inflorescentia* aphylla, ramosa, effusa; paniculæ in ramulis ultimis brevibus densius aggregatæ, iis *P. Veitchianæ* similes sed interdum minus congestæ; bracteæ quoque et glumæ velut in *P. Veitchiana*.

SZECHUEN: Mount Omei, 3000 ft. (*Faber*, 1116!). Herb. Kew.

Closely resembles *Phyllostachys Veitchiana*, mihi, in the character of the panicles, but the inflorescence is more diffuse. It also differs in the texture of the leaves which are thinner and papery, the long-fimbriated ligule, and the very short petiole. I am unable to describe the florets, as they have been eaten by a grub.

8. *Phyllostachys nana*, *Rendle*, n. sp.

*Suffrutex* pedalis; rhizoma 2–3 lin. crassum; culmus basi subteres ad  $\frac{1}{4}$  poll. crassus, nodis basalibus subapproximatis ramosus; rami diffuse ramulosi; internodia unilateraliter complanata, brunneo-maculata. *Foliorum* vaginæ pollicares superne ampliatae.

et sparse puberulæ, margine ciliolatæ, ore longe fimbriatæ (fimbriæ  $\frac{1}{4}$  poll. longæ); ligula brevis, truncata, ciliolata; lamina late linearis vel lineari-lanceolata, ad apicem breviter acuminatum attenuata, ad 5 poll. longa et 7 lin. lata basi in petiolum brevem (ad 3 lin. longum) subito attenuata, subter pilosula, margine utroque scabridula, venæ secundariae utrinque 5, interspatiis minute tessellatis. *Paniculæ* in ramulis primo foliiferis laterales vel ramulos breves terminantes, ad sesquipollicares, axis basi cum bracteis ovatis, glabris, subcoriaceis, brevibus sed sursum majoribus, vaginatus; bracteæ superiores ramulos subtendentes, membranaceæ, longiores (ad  $\frac{3}{4}$  poll. longæ), multinerves, hirsutulæ, cum lamina rudimentaria instructæ; ramuli breves congesti a spiculis 2-3-floris terminati. *Gluma* sterilis bracteæ supremæ similis sed brevior (5 lin. longa), apice subpungens; glumæ fertiles oblongæ, superne angustatæ, apice pungentes, dorso rotundatæ, hirsutulæ, imperfecte 9-nerves, infima  $\frac{2}{3}$  poll. longa, suprema florem imperfectum fovens; rhachilla puberula; palea quam gluma paullo brevior, bidentula, dorso hirsutula, plurinervis, carinis 2 vix prominulis; lodiculæ 3 lanceolatæ vel ovatæ, obtusæ, superne ciliolatæ,  $1\frac{3}{4}$  lin. longæ; stamina, ovarium et stylus generis.

HUPEH: Ichang (*A. Henry*, 3278!). Herb. Kew.

The panicle and spikelets resemble those of *Phyllostachys nigra*, Munro, but the habit is very different.

9. *Phyllostachys Nevinii*, *Hance*, in *Journ. Bot.* xiv. (1876) p. 295.

KWANGTUNG: Danes Island, Whampoa (*Nevin*, in herb. *Hance*, 19307!). Mus. Brit.; Herb. Kew.

Var. *hupehensis*, *Rendle*, n. var. A typo recedit fimbriis ad vaginarum ora longioribus (sæpe  $2\frac{1}{2}$  lin. longis) et spiculis densius hirsutulis.

HUPEH: Ichang (*A. Henry*, 3318!). Mus. Brit.; Herb. Kew.

10. *Phyllostachys nidularia*, *Munro*, in *Gard. Chron.* (1876) ii. pp. 773, 774.

*Phyllostachys nigra*, *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 227, non *Munro*.

CHEKIANG: Taiboo Lake (*Forbes*!); KIANGSI: Kiukiang

(*Shearer! Maries!*); HUPEH: Ichang (*A. Henry*, 450, 1322, 7868, 7871!), Nanto (*Wilson*, 10! *A. Henry*, 3823!), Changyang (*A. Henry*, 3671, 7809!); SZECHUEN: Chungking, Yangtzekiang (*Faber*, 1117!); KWANGTUNG: Macao (*Calléry*, 767!). Mus. Brit.; Herb. Kew.

Japan.

11. *Phyllostachys nigra*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 38; *A. & C. Rivière*, in *Bull. Soc. Acclimat. sér. 3*, v. (1878) p. 709.

Var. *Henonis*, *Stapf*, in *Herb. Kew.*

*Phyllostachys Henonis*, *Mitford*, *Bamboo Garden*, p. 149; *Stapf*, in *Hook. Icon. Pl.* t. 2614; *Pilger*, in *Engl. Bot. Jahrb.* xxix. (1900) p. 227.

SZECHUEN: Nanchuan, planted? (*Bock von Rosthorn*, 1207, fide *Pilger*).

Japan.

12. *Phyllostachys quadrangularis*, *Rendle*.

*Bambusa quadrangularis*, *Fenzi*, in *Bull. Soc. Tosc. Ort.* v. (1880) p. 401.

KIANGSU: Shanghai (*Macgowan!*); FOKIEN: Wuiking (*Bourne!*). Herb. Kew.

Japan.

The two plants cited are leaf-specimens which are thus referred in Herb. Kew.

13. *Phyllostachys Stauntoni*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 37; *Franchet*, in *Mém. Soc. Sci. Nat. Cherbourg*, xxiv. (1884) p. 270.

CHIHLI: Peking, cultivated (*Skatschkow!*); KANGSU: Kiang (*Staunton!*). Mus. Brit.; Herb. Kew.

14. *Phyllostachys Veitchiana*, *Rendle*, n. sp.

*Frutex* ramis semiteretibus. *Folia* in ramulis floriferis paniculas superantibus posita; vaginæ sursum ampliatae, pollicares, glabræ, ore fimbriatæ, fimbriæ  $\frac{1}{4}$  poll. longæ; ligula brevissima, truncata; lamina lineari-oblonga ad oblongo-lanceolata, superne in apicem acuminatum attenuata, ad  $3\frac{1}{2}$  poll. longa et vix  $\frac{2}{3}$  lin. lata, glabra, basi in petiolum vix  $\frac{1}{4}$  poll. longum subito attenuata, margine unico tantum scabridula, venæ secundariæ utrinque 5,

interspatiis subtiliter tessellatis. *Rami* florentes ad nodos distantes gemini vel terni fasciculati, tenues ( $\frac{1}{2}$  lin. crassit. vix excedentes) super basin iterum iterumque ramulosi, ramuli ascendo-patentes, cum internodiis brevibus, paniculas sessiles laterales numerosas gerentes. *Paniculæ* multispiculiferæ, congestæ,  $1\frac{1}{2}$  poll. longæ; axis basi cum bracteis sursum majoribus vaginatus; bracteæ coriaceæ, infimæ ovatæ, tum ovato-oblongæ et breviter mucronatæ, margine ciliolatæ, superiores quæ ramulos subtendunt membranaceæ, lineari-oblongæ, quam inferiores multo longiores, subpollicares, margine densius, dorso sparse hirsutæ, multinerves, cum ligula brevissima et lamina parva triangulari instructæ; spiculæ 3-floræ. *Glumæ* steriles 1 vel 2, inferior bracteis superioribus similis sed paullo brevior, superior inter inferiorem et glumam florentem forma intermedia; glumæ florentes oblongo-lanceolatæ, apice pungentes, dorso rotundatæ et densius hirsutæ, infima  $\frac{2}{3}$  poll. longa; rhachillæ articuli clavati, glabri; palea quam gluma brevior (infima semipollicaris) bidentula, hirsuta præcipue in carinis; lodiculæ 3, ovales vel subobovatae,  $1\frac{1}{2}$  lin. longæ (una autem paullo minor); ovarium et stylus generis.

HUPEH: West (*Wilson*, 10 A!). Herb. Kew.

Near *Phyllostachys nigra*, Munro, but distinguished by the texture of the leaf, which is tougher and has a more rounded base passing abruptly into the petiole; also by the denser panicles.

15. *Phyllostachys viridi-glaucescens*, A. & C. Rivière, in *Bull. Soc. Acclimat. sér. 3*, v. (1878) p. 391.

“Without doubt a native of the north of China” (A. & C. Rivière, loc. cit.).

I have seen no specimen from China; the plant is grown in Kew Gardens, and there are leaf-specimens in the Herbarium.

[*Bambusa agrestis*, Poir. *Encycl.* viii. p. 704; Munro, in *Trans. Linn. Soc.* xxvi. (1868) p. 117.

*Arundo agrestis*, Lour. *Fl. Cochinch.* p. 57.

CHINA? (ex *Loureiro*).

This is insufficiently described; its identity and its occurrence in China are both doubtful.]

1. *Bambusa angulata*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 94; *A. Henry*, in *Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 109; *Hack.* in *Bull. Herb. Boiss.* vii. (1899) p. 725.

FORMOSA (*Oldham*, 650!). Mus. Brit.; Herb. Kew.

2. *Bambusa arundinacea*, *Willd. Sp. Pl.* ii. p. 245; *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 103; *Debeaux*, *Fl. Shanghai*, p. 123; *Gamble*, *Bamb. Brit. Ind.* p. 51, et in *Hook. f. Fl. Brit. Ind.* vii. p. 395.

*Bambusa Arundo*, *Klein, ex Nees*, in *Linnaea*, ix. (1835) p. 471; *Benth. Fl. Hongk.* p. 434.

*Arundo Bambos*, *Linn. Sp. Pl.* p. 81; *Osbeck*, *Voyage, ed. Forster*, ii. p. 343.

Apparently naturalized in many parts of China.

India and Ceylon.

3. *Bambusa Beecheyana*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 108.

*Bambusa verticillata*, *Hook. & Arn. Bot. Beechey's Voy.* p. 254, non *Willd.*

CHINA: without locality (*Staunton! Beechey!*); KWANGTUNG: Canton (*Hance*, 1050 in part!), Ookaisa woods (*Tate!*). Mus. Brit.; Herb. Kew.

4. *Bambusa breviflora*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 96; *A. Henry*, in *Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 109.

FORMOSA: Tamsui (*Oldham*, 651!). Mus. Brit.; Herb. Kew.

5. *Bambusa Cantori*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 111.

HONGKONG: Lantao (*Cantor!*). Herb. Kew.

Leaf-specimens only.

6. *Bambusa flexuosa*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 101; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 137.

KWANGTUNG (*Staunton!*), near Canton (*Hance*, 10000!); HAINAN (*A. Henry*, 7996!). Mus. Brit.; Herb. Kew.

7. *Bambusa nana*, *Roxb. Hort. Bengal.* p. 25, et *Fl. Ind.* (ed. *Carey*) ii. p. 199; *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 89;

*Hack. in Bull. Herb. Boiss.* vii. (1899) p. 720; *Hook. f. Fl. Brit. Ind.* vii. p. 390.

*Bambusa floribunda*, *Zoll. et Morr. ex Steud. Syn. Gram.* p. 330; *Miq. in Ann. Mus. Bot. Lugd.-Bat.* ii. p. 285; *Franch. et Savat. Enum. Pl. Jap.* ii. p. 183.

Japan.

“Keufa, of the Chinese; a native of their country and now plentiful in the Botanic Garden at Calcutta.”—*Roxb. Fl. Ind. (ed. Carey)* ii. p. 199.

I have seen no specimen from China.

8. *Bambusa Oldhami*, *Munro, in Trans. Linn. Soc.* xxvi. (1868) p. 109; *A. Henry, in Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 109; *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 725; *Stapf, in Hook. Ic. Pl.* 2773.

FORMOSA: Tamsui (*Oldham*, 648!), Takow (*A. Henry*, 1955!), South Formosa (*Maries*!). *Mus. Brit.*; *Herb. Kew.*

9. *Bambusa stenostachya*, *Hack. in Bull. Herb. Boiss.* vii. (1899) p. 725.

FORMOSA: Shinchiku (*Makino*, fide *Hackel*).

10. *Bambusa Teba*, *Miq. Fl. Ind. Bat.* iii. p. 418; *Munro, in Trans. Linn. Soc.* xxvi. (1868) p. 122.

*Arundarbor spinosa*, *Rumph. Hort. Amboin.* iv. p. 14, t. 3.

SOUTH CHINA (fide *Rumphius*).

Malaya.

I have not seen this. It is a doubtful species described from leaf-specimens only.

11. *Bambusa tessellata*, *Munro, in Trans. Linn. Soc.* xxvi. (1868) p. 110.

*Bambusa reticulata*, var. *macrophylla*, *Rupr. ex Munro, l. c.* p. 111.

CHINA: without locality or collector's name! *Mus. Brit.*; *Herb. Kew.*

Described from leaf-specimens only.

I do not find the varietal name which *Munro* quotes as of *Ruprecht* (*Bambuseæ*, p. 148), but *Ruprecht's* description of the large leaves which he correlates with his *Bambusa reticulata* agrees exactly with the leaves on which *Munro* based *B. tessellata*.



12. *Bambusa tuldoides*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 93; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 137; *A. Henry*, in *Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 109.

*Bambusa Tulda*, *Benth. Fl. Hongk.* p. 434, non *Roxb.*

FORMOSA: Tamsui (*Oldham*, 652 !, a poor specimen which has been thus determined by *Munro*), South Cape (*Schmürer*, in coll. *A. Henry*, 686 !); KWANGTUNG: Canton (*Millett*!); HONGKONG (*Hance*, 3657, 7486 !), Robinson Road, Victoria (*Lamont*!). Mus. Brit.; Herb. Kew.

13. *Bambusa vulgaris*, *Schrad.* in *Wendl. Collect. Pl.* ii. p. 26, t. 47; var. *striata*, *Gamble*, in *Ann. Bot. Gard. Cale.* vii. p. 44.

*Bambusa striata*, *Lodd. ex Lindl. Penny Cyclop.* iii. (1835) p. 357; *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 121; *Hook. f. in Bot. Mag.* t. 6079.

I have seen no specimen. *Gamble* says (*l. c.*):—"It appears to come from China and Japan and is probably the result of cultivation."

1. *Dendrocalamus affinis*, *Rendle*, n. sp.

*Frutex* elata cujus nihil nisi inflorescentiæ pars exstat. *Inflorescentiæ* axis teres, glaber; rami ad nodos distantes unilateraliter fasciculati, terni, sæpissime ad nodum basale, interdum quoque ad nodum secundum, ramosi, flagelliformes, penduli, subtiliter scabriduli, specimine ad 21 poll. longi; spiculæ ut in *D. flagellifero* paucae ad nodos subdistantes subverticillatæ (glomerulæ rarius plus quam  $\frac{1}{2}$  poll. latæ), ovatæ, ad 5 lin. longæ, fusco-brunnæ, 4-floræ. *Glumæ* coriaceæ, steriles 4, parvæ, sursum majores, 1 ad 2 lin. longæ, ovatæ, obtusæ, multinerves, glabræ, margine autem ciliolatæ; *glumæ* florentes orbiculari-ovatæ, mucronulatæ, multinerves, ut in sterilibus ciliolatæ, infima  $2\frac{1}{2}$  lin. longa; *palea* compressa, oblonga, quam *gluma* brevior (infima  $1\frac{3}{4}$  lin. longa), ad carinas superne ciliolatas implicata, dorso inconspicue binervis, apice obtuse retusa et ciliolata; *lodiculæ* 3, subæquales, obtuse ovatæ, apice ciliatæ,  $\frac{1}{2}$  lin. longæ; *stamina* 3; *ovarium* breviter pedicellatum, albo-pilosum; *stylus* mox in stigmata tria plumosa solutus.

SZECHUEN: Mount Omei (*Faber*, 1118 !). Herb. Kew.

Evidently near the Burmese *Dendrocalamus Brandisii*, *Kurz*,

and the South Indian and Malayan *D. flagellifer*, Munro, especially the latter, from which it differs in the dull polished brown colour of the spikelets, which are also slightly larger, and in the broader lodicules.

2. *Dendrocalamus latiflorus*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 152, t. vi; *Hance*, in *Journ. Linn. Soc., Bot.* xiii. (1873) p. 137; *Gamble*, in *Ann. Bot. Gard. Calc.* vii. p. 131, et in *Hook. f. Fl. Brit. Ind.* vii. p. 407; *A. Henry*, in *Trans. Asiat. Soc. Japan*, xxiv. *Suppl.* (1896) p. 109.

*Bambusa verticillata*, *Benth. Fl. Hongk.* p. 434, non *Willd.*

FORMOSA (*Maries!* *Oldham*, 649), Takow (*A. Henry*, 1803, 1955!), Bankinsing (*A. Henry*, 1760!); HONGKONG (*Harland*, 740 in herb. *Hance*, 1050 in part!), Barracks (*Urquhart!*). *Mus. Brit.*; *Herb. Kew.*

Burma.

? *Pseudostachyum polymorphum*, *Munro*, in *Trans. Linn. Soc.* xxvi. (1868) p. 142, t. 4; *Gamble*, in *Ann. Bot. Gard. Calc.* vii. p. 96, t. 85, et in *Hook. f. Fl. Brit. Ind.* vii. p. 409.

HUPEH: Nanto and mountains to northward (*A. Henry*, 3857!). *Herb. Kew.*

The specimen, which bears no flowers, may belong to this species.

1. *Schizostachyum chinense*, *Rendle*, n. sp.

*Frutex* 10-15-pedalis; internodia glabra teretia. *Foliorum* vaginæ glabræ, ore in specimine non fimbriatæ; ligula brevis truncata,  $\frac{1}{2}$  lin. longa; lamina lanceolata, acuminata, ad 8 poll. longa, vix  $1\frac{1}{2}$  lin. lata, glabra, margine minute scabridula, venæ secundariæ utrinque circa 7 cum nervulis transversis tenuissimis sparsis. *Paniculæ* terminales, aphyllæ, tenues, 14-16 poll. longæ, rhachis tenuis; rami distantes ascendentes, basi cum bractea amplexicauli membranacea lanceolata acuminata vaginati, inferiores ad  $3\frac{1}{2}$  poll. longi, sursum deminuentes; ramuli (in ramis longioribus distantes) similiter vaginati, basi bracteati, bractea infima bicarinata, carinis et apice pilosa; bracteæ superiores dorso rotundatæ spiculæ circumvolutæ; spiculæ ramulos terminantes, unifloræ; *glumæ* steriles binæ florenti similes sed breviores, gl. fertilis ovata, breviter mucronata, glabra, multinervis,

palea glumæ subsimilis, sed paullo longior, magis hyalina et apice obtusiuscula; lodiculæ 3 subæquales, ovatæ vel lanceolatæ, glabræ, 1 lin. paullo excedentes; antheræ 6, inappendiculatæ, 5 lin. longæ; stylus generis.

YUNNAN: Mengtze, S.E. mountains, forests on cliffs 8000 ft. high (*A. Henry*, 10420!). Herb. Kew.

Distinguished from the other species of the genus by its long slender panicle, with a few spikelets clustered on the short ultimate branchlets.

2. *Schizostachyum dumetorum*, *Munro*, in *Seem. Bot. Voy. 'Herald,'* p. 424, et in *Trans. Linn. Soc.* xxvi. (1868) p. 136; *Benth. Fl. Hongk.* p. 435.

*Bambusa dumetorum*, *Hance*, in *Walp. Ann.* iii. p. 781.

CHINA: without locality (*Fortune*, 164!); HONGKONG (*Hance*, 597! *Seemann*!), roadside, Happy Valley (*Hance*, in herb. *Lamont*!), ravine on Mount Davis (*Wilford*, 255!). Mus. Brit.; Herb. Kew.

LIST OF THE GENERA AND SPECIES discovered in China since the publication of the various parts of the "Enumeration," from 1886 to March 1904, alphabetically arranged.

THIS List, compiled by Miss M. Smith, is not offered as an exhaustive one, but as containing references to new species described in publications received at Kew during the period named, with the addition of a number of previously described, mainly Indian, species of later discovery in China.

Finet and Gagnepain's "Contributions à la Flore de l'Asie Orientale d'après l'Herbier du Muséum de Paris," the beginning of which has appeared in the *Bulletin de la Société Botanique de France*, are of the nature of a detailed enumeration of the plants of China and Japan and the adjoining countries, as represented in the Paris herbarium, and have not been drawn upon. Only the new species have been extracted from Gilg and Loesener's "Beiträge zu einer Flora von Kiautschou und einiger angrenzenden Gebiete," in Engler's *Botanische Jahrbücher*, xxxiv. (1904), *Beibl.* n. 75, pp. 1-76.

W. BOTTING HEMSLEY.

- ABELIA ADENOTRICHA, Hance, in Journ. of Bot. ix. (1871) 132.  
 ✓ ANGUSTIFOLIA, Bur. & Franch. in Journ. de Bot. v. (1891) 47.  
 ABUTILON POLYANDRUM, Schlect. ex Franch. Pl. Delav. (1890) 107.  
 ➤ SINENSE, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1750.  
 ✓ ACACIA DELAVAYI, Franch. Pl. Delav. (1890) 194.  
 ✓ PENNATA, Willd. ex Franch. *l. c.* 193.  
 ✓ YUNNANENSIS, Franch. *l. c.* 193.  
 ✓ ACALYPHA GIRALDII, Pax, in Engl. Jahrb. xxix. (1900) 429.  
 ✓ ACANTHOPANAX EVODIÆFOLIUS, Franch. in Journ. de Bot. x. (1896) 306.  
 ✓ HENRYI, Harms, in Engl. & Prantl, Pflanzenf. iii. 8. p. 50.  
 ✓ LEUCORRHIZUS, Harms, *l. c.*  
 ✓ SENTICOSUS, Harms, *l. c.*  
 ✓ SETCHUENENSIS, Harms, in Engl. Jahrb. xxix. (1900) 488.  
 ✓ SETULOSUS, Franch. Pl. David. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 249.  
 ✓ ACER BETULIFOLIUM, Maxim. in Act. Hort. Petrop. xi. (1889) 108.  
 ✓ CAUDATUM, Wall. ex Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 464.

- ✓ ACER CORDATUM, Pax, in Hook. Ic. Pl. xix. (1889) sub t. 1897.
  - ✓ DAVIDI, Franch. Pl. David. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 212.
- ✓ ARIANTHUM, Schwerin, in Engl. Pflanzenreich, Acer. 22.
- ✓ EROSUM, Pax, in Hook. Ic. Pl. xix. (1889) sub t. 1897.
  - ✓ FRANCHETI, Pax, *l. c.*
- ✓ GIRALDII, Pax, in Engl. Pflanzenreich, Acer. 79.
- ✓ GROSSERI, Pax, *l. c.* 80.
- ✓ HENRYI, Pax, in Hook. Ic. Pl. xix. (1889) t. 1896.
- ✓ LÆVIGATUM, Wall. ex Engl. Jahrb. xxix. (1900) 449.
- ✓ LANCEOLATUM, Molliard, in Bull. Soc. Bot. Fr. l. (1903) 134.
- ✓ LAXIFLORUM, Pax, in Engl. Pflanzenreich, Acer. 36.
- ✓ LOBELII, Ten., var.; Pax, in Hook. Ic. Pl. xix. sub t. 1897.
- ✓ MAXIMOWICZII, Pax, *l. c.* sub t. 1897.
- ✓ MULTISERRATUM, Maxim. in Act. Hort. Petrop. xi. (1889) 107.
- ✓ NIKOENSE, Maxim., var.; Franch. in Journ. de Bot. viii. (1894) 294.
- ✓ OLIVERIANUM, Pax, in Hook. Ic. Pl. xix. (1889) sub t. 1897.
- ✓ PAXII, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 464.
- ✓ ROBUSTUM, Pax, in Engl. Pflanzenreich, Acer. 79.
- ✓ SCHENERMARCKIÆ, Pax, *l. c.* 71.
- ✓ SINENSE, Pax, in Hook. Ic. Pl. xix. (1889) sub t. 1897.
- ✓ SUTCHUENENSE, Franch. in Journ. de Bot. viii. (1894) 294.
- ✓ TENELLUM, Pax, in Hook. Ic. Pl. xix. (1889) t. 1897.
- ✓ TETRAMERUM, Pax, *l. c.* sub t. 1897.
- ✓ TRIFLORUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 430.
- ✓ UROPHYLLUM, Maxim. *l. c.* xi. (1889) 105.
  - ✓ VEITCHII, Hort.; Mitth. Deutsch. Dendrol. Ges. 1894, No. 3, 51.
- ✓ ACERIPHYLLUM ROSSII, Engl. & Prantl, Pflanzenf. iii. 2 A, p. 52.
- ✓ ACHLYS JAPONICA, Maxim. & Komarov, in Act. Hort. Petrop. xxii. (1903) 324.
- ✓ ACONITUM ALBO-VIOLACEUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 439.
  - ✓ BODINIERI, Léveillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 45.
- ✓ DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 381.
- ✓ HEMSLEYANUM, E. Pritzel, in Engl. Jahrb. xxix. (1900) 329.
- ✓ HENRYI, E. Pritzel, *l. c.* 329.
  - ✓ LUTEUM, Léveillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 46.
- ✓ PALMATUM, Don; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 381.
- ✓ RACEMULOSUM, Franch. in Journ. de Bot. viii. (1894) 276.
- ✓ ROTUNDIFOLIUM, Kar. & Kir., var.; Maxim. ex Engl. Jahrb. xxix. (1900) 329.
- ✓ SCAPOSUM, Franch. in Journ. de Bot. viii. (1894) 277.
- ✓ UMBROSUM, Komarov, in Act. Hort. Petrop. xxii. (1903) 250.
- ✓ VAGINATUM, E. Pritzel, in Engl. Jahrb. xxix. (1900) 328.
- ✓ VOLUBILE, Pall.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1888) 381.

- ✓ *ACTÆA ERYTHROCARPA*, Fisch. ; Komarov, in Act. Hort. Petrop. xxii. (1903) 237.
- ✓ *SPICATA*, Linn. ex Engl. Jahrb. xxix. (1900) 326.
- ✓ *ACTINIDIA LANATA*, Hemsl. in Ann. Bot. ix. (1895) 146.
- ✓ *MELANANDRA*, Franch. in Journ. de Bot. viii. (1894) 278.
- ✓ *TETRAMERA*, Maxim. in Act. Hort. Petrop. xi. (1889) 35.
- ✓ *TRICHOGYNE*, Franch. in Journ. de Bot. viii. (1894) 278.
- ✓ *ACTINOSTEMMA BIGLANDULOSA*, Hemsl. in Hook. Ic. Pl. xxvii. (1899) tt. 2622, 2645.
- ✓ *MULTILOBUM*, Harms, in Engl. Jahrb. xxix. (1900) 602.
- ✓ *ADENOPHORA AURITA*, Franch. in Journ. de Bot. ix. (1895) 366.
- ✓ *BOCKIANA*, Diels, in Engl. Jahrb. xxix. (1900) 605.
- ✓ *JASIONIFOLIA*, Franch. in Journ. de Bot. ix. (1895) 365.
- ✓ *MARSUPIIFLORA*, Fisch., var.; Diels, in Engl. Jahrb. xxix. (1900) 605.
- ✓ *POTANINI*, Korsh. in Engl. Jahrb. xxix. (1900) 606.
- ✓ *WAWREANA*, Zahlbruckner, in Ann. Hofmus. Wien, x. (1895) 56.
- ✓ *ADINANDRA BOCKIANA*, E. Pritzel, in Engl. Jahrb. xxix. (1900) 474.
- ✓ *ADLUMIA CIRRHOSA*, Rafin. ; Komarov, in Act. Hort. Petrop. xxii. (1903) 343.
- ✓ *ADONIS AMURENSIS*, Regel & Radde ; Komarov, in Act. Hort. Petrop. xxii. (1903) 315.
- ✓ *BREVISTYLA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 372.
- ✓ *DAVIDI*, Franch. Pl. David. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 188.
- ✓ *DELAVAYI*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 91.
- ✓ *SUTCHUENENSIS*, Franch. *l. c.* 89.
- ✓ *ÆGOPodium ALPESTRE*, Ledeb. ; Bull. Herb. Boiss. sér. 2, iii. (1903) 954.
- ✓ *HENRYI*, Diels, in Engl. Jahrb. xxix. (1900) 497.
- ✓ *ÆSCHYNANTHUS BUXIFOLIUS*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 515.
- ✓ *HUMILIS*, Dunn, *l. c.* 516.
- ✓ *ÆSCULUS INDICA*, Colebr. ; Franch. in Journ. de Bot. iv. (1890) 303.
- ✓ *AGANOSMA EDITHÆ*, Hance, in Ann. Sci. Nat. sér. 5, v. (1866) 227.
- ✓ *AGAPETES PARVIFLORA*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 515.
- ✓ *VACCINIOIDEA*, Léveillé, in Bull. Acad. Géog. Bot. xii. (1903) 251.
- ✓ *VACCINOIDES*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 515.
- ✓ *YUNNANENSIS*, Franch. in Journ. de Bot. ix. (1895) 366.
- ✓ *AINSLIÆA ELEGANS*, Hemsl. in Hook. Ic. Pl. xxviii. (1902) t. 2747.
- ✓ *GRACILIS*, Franch. in Journ. de Bot. viii. (1894) 297.
- ✓ *GROSSEDENTATA*, Franch. *l. c.* 297.
- ✓ *HENRYI*, Diels, in Engl. Jahrb. xxix. (1900) 628.
- ✓ *LANCIFOLIA*, Franch. Pl. David. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 41.
- ✓ *NERVOSA*, Franch. in Bull. Mus. Hist. Nat. Par. (1896) no. 2, 64.

- AINSLIÆA OVALIFOLIA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 119.  
     RUBRIFOLIA, Franch. in Journ. de Bot. viii. (1894) 296.  
 ✓ SCABRIDA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 510.  
     SPARSIFLORA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 118.  
     SPICATA, Vaniot, *l. c.* 117.  
     SUTCHUENENSIS, Franch. in Journ. de Bot. viii. (1894) 296.  
     UNDULATA, Diels, in Engl. Jahrb. xxix. (1900) 629.  
 AJUGA OVALIFOLIA, Bur. & Franch. in Journ. de Bot. v. (1891) 150.  
 ✓ AKEBIA LOBATA, Decne., var.; Diels, in Engl. Jahrb. xxix. (1900) 344.  
     LONGERACEMOSA, Matsuma, in Bot. Mag. Tokyo, xiii. (1899) 18.  
 ALANGIUM FABERI, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1774.  
 ALBIZZIA BRACTEATA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 493.  
     ODORATISSIMA, Benth. var.; Franch. Pl. Delav. (1890) 193.  
 ALLIARIA AURICULATA, Komarov, in Act. Hort. Petrop. xviii. (1901) 437.  
 ✓ ALLIUM ZIMMERMANIANUM, Gilg, in Engl. Bot. Jahrb. xxxiv. (1904) Beibl. 75, 23.  
 ALNIPHYLLUM, Matsuma, in Bot. Mag. Tokyo, xv. (1901) 67.  
 ✓ MACRANTHUM, Perk. in Engl. Jahrb. xxxi. (1902) 488.  
     PTEROSPERMUM, Matsuma, in Bot. Mag. Tokyo, xv. (1901) 67.  
 ALNUS FRUTICOSA, Ruprecht; Komarov, in Act. Hort. Petrop. xxii. (1903) 58.  
     HIRSUTA, Turcz.; Komarov, *l. c.* 54.  
     MARITIMA, Nutt., var.; Matsuma, Rev. Alni. sp. jap., ex Bull. Soc. Bot. France, xlix. (1902) 225.  
 ✓ ALPHONSEA MOLLIS, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 485.  
 ALPINIA FORMOSANA, K. Schum. in Engl. Jahrb. xxvii. (1899) 283.  
     SPECIOSA, K. Schum. *l. c.* 284.  
 ALSINE LARICINA, Crantz; Komarov, in Act. Hort. Petrop. xxii. (1903) 186.  
 ✓ ALSOMITRA GRACILIFLORA, Harms, in Engl. Jahrb. xxix. (1900) 602.  
 ✓ AMELANCHIER ASIATICA, Endl.; Engl. Jahrb. xxix. (1900) 389.  
 ANMODENIA PEPLOIDES, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 187.  
 AMPELOCISSUS ARTEMISLÆFOLIA, Planch.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 458.  
     CARDIOSPERMA, Planch.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 460.  
     CARDIOSPERMOIDES, Planch. in DC. Monogr. Phan. v. (1887) 403.  
 AMPELOPSIS DELAVAYANA, Planch.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 460.  
 ✓ MEGALOPHYLLA, Diels & Gilg, in Engl. Jahrb. xxix. (1900) 466.

- ✓ *AMPELOPSIS MIRABILIS*, Diels & Gilg, in Engl. Jahrb. xxix. (1900) 465.  
*TOMENTOSA*, Planch.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 459.  
*AMPELOVITIS*, Carr. in Rev. Hort. (1889) 204.  
*DAVIDI*, Carr. *l. c.* 204.  
*INTERMEDIA*, Carr. *l. c.* (1891) 334.  
*ROMANETI*, Carr. *l. c.* (1892) 94.
- ✓ *AMPHICOME ARGUTA*, Lindl., ex Engl. Jahrb. xxix. (1900) 577.  
*AMYGDALUS TANGUTICA*, Korsh. in Bull. Acad. Pétersb. sér. 5, xiv. (1901) 94.
- ✓ *ANAPHALIS ARANEOSA*, DC.; Engl. Jahrb. xxix. (1900) 613.  
*BODINIERI*, Franch. in Journ. de Bot. iv. (1890) 306.
- ANDROSACE ALCHEMILLOIDES*, Franch. in Journ. de Bot. ix. (1895) 455.  
*AXILLARIS*, Franch. *l. c.* 455.  
*BISULCA*, Bur. & Franch. *l. c.* v. (1891) 103.  
*COCCINEA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 63.  
*CUSCUTIFORMIS*, Franch. in Journ. de Bot. ix. (1895) 454.  
*DELAVAYI*, Franch. *l. c.* 456.  
*DISSECTA*, Franch. *l. c.* 454.  
*HENRYI*, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1973.  
*MARLÆ*, Kanitz, in Pl. Exped. Széchenyi, Asia Centr. (1891) 7.  
*MAXIMA*, Linn.; Engl. Jahrb. xxix. (1900) 522.  
*MIRABILIS*, Bur. & Franch. in Journ. de Bot. ix. (1895) 453.  
*SUTCHUENENSE*, Franch. *l. c.* 454.
- ANEMONE AMURENSIS*, Komarov, in Act. Hort. Petrop. xxii. (1903) 262.  
*BAICALENSIS*, Turcz.; Komarov, *l. c.* 265.  
*BEGONIIFOLIA*, Léveillé & Vaniot, Bull. Acad. Géog. Bot. xi. (1902) 46.  
*BOISSIÆI*, Léveillé & Vaniot, *l. c.* 47.  
*DAVIDI*, Franch. Pl. David. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 185.  
*DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 366.  
*DEMISSA*, Hook. f. & Thoms. var.; Franch. *l. c.* 367.  
*DICHOTOMA*, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 261.  
*FAURIEI*, Léveillé, in Bull. Acad. Géog. Bot. xi. (1902) 300.  
*GELIDA*, Maxim. in Act. Hort. Petrop. xi. (1890) 1. 21.  
*GLAUCIFOLIA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 363.  
*HENRYI*, Oliver, in Hook. Ic. Pl. xvi. (1887) t. 1570.  
*IMBRICATA*, Maxim. Fl. Tangut. (1889) 8.  
*REFLEXA*, Stephan; Komarov, in Act. Hort. Petrop. xxii. (1903) 266.  
*REGELIANA*, Maxim. Fl. Tangut. (1889) 10.
- ✓ *RIVULARIS*, Buch.-Ham.; Engl. Jahrb. xxix. (1900) 331.



- ANEMONE SCABIOSA, Léveillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 47.  
 SILVESTRIS, Linn. ex Léveillé & Vaniot, *l. c.*  
 UMBROSA, C. A. Meyer; Komarov, in Act. Hort. Petrop. xxii. (1903) 263.
- ✓ VITIFOLIA, Buch.-Ham.; Engl. Jahrb. xxix. (1900) 330.
- ANGELICA DIELSII, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 850.
- ✓ DISSOLUTA, Diels, in Engl. Bot. Jahrb. xxix. (1900) 499.  
 FARGESII, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 850.  
 FLACCIDA, Komarov, in Act. Hort. Petrop. xviii. (1901) 431.
- ✓ INVOLUCELLATA, Diels, in Engl. Jahrb. xxix. (1900) 501.  
 KOREANA, Maxim. in Bull. Acad. Pétersb. xxxi. (1887) 51.
- ✓ LAXIFOLIATA, Diels, in Engl. Jahrb. xxix. (1900) 499.  
 MEGAPHYLLA, Diels, *l. c.* 500.  
 PSEUDO-SELINUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1902) 848.  
 SCABERULA, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 141.
- ✓ SETCHUENENSIS, Diels, in Engl. Jahrb. xxix. (1900) 500.
- ✓ POLYMORPHA, Maxim. var.; Oliver, in Hook. Ic. Pl. xx. (1898) t. 1999.  
 UCHIYAMÆ, Yatabe, in Bot. Mag. Tokyo, xvii. (1902) 106.
- ✓ VALIDA, Diels, in Engl. Jahrb. xxix. (1900) 501.
- ANISADENIA PUBESCENS, Oliver, in Hook. Ic. Pl. xxvi. (1901) t. 2593.
- APIOS DELAVAYI, Franch. Pl. Delav. (1890) 180.  
 ✓ GRACILLIMA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 488.  
 MACRANTHA, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1946.
- AQUILEGIA ECALCARATA, Maxim. Fl. Tangut. (1889) 20.  
 OXYSEPALA, Trautv. & Mey.; Komarov, in Act. Hort. Petrop. xxii. (1903) 244.
- ✓ VULGARIS, Linn.; Engl. Jahrb. xxix. (1900) 326.
- ARABIS AXILLARIS, Komarov, in Act. Hort. Petrop. xviii. (1901) 437.  
 HALLERI, Linn.; Komarov, *l. c.* 378.  
 LYRATA, Linn.; Komarov, *l. c.* 377.  
 PANICULATA, Franch. Pl. Delav. (1889) 57.
- ARALIA ATROPURPUREA, Franch. in Journ. de Bot. x. (1896) 301.  
 FARGESII, Franch. *l. c.* 302.
- ✓ HENRYI, Harms, in Engl. Jahrb. xxiii. (1896) 12.  
 PILOSA, Franch. in Journ. de Bot. x. (1896) 302.  
 SEARELLIANA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 498.  
 STIPULATA, Franch. in Journ. de Bot. x. (1896) 304.  
 TOMENTELLA, Franch. *l. c.* 304.  
 YUNNANENSIS, Franch. *l. c.* 303.
- ✓ ARDISIA BREVICAULIS, Diels, in Engl. Jahrb. xxix. (1900) 519.  
 ✓ CORYMBIFERA, Mez, in Engl. Pflanzenreich, Myrsin. 149.  
 ✓ HAINANENSIS, Mez, *l. c.* 138.  
 ✓ HANCEANA, Mez, *l. c.* 149.

## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

The second volume, for 1902, is now in the press.

B. DAYDON JACKSON,  
*General Secretary, Linn. Soc.*

## NOTICE.

---

VOL. XXXV. has been completed by the issue of Part 248.

VOL. XXXVI., of which Parts 249-254 are already published, is reserved for the completion of the Chinese Flora.

Attention to this announcement is specially requested, to prevent application to the Librarian for unpublished Parts.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the "SECRETARIES," but letters on library business only should be addressed to the "LIBRARIAN."

DECEMBER 20.

Price 24s.

THE JOURNAL  
OF  
THE LINNEAN SOCIETY.

---

VOL. XXXVI.

BOTANY.

Nos. 255-56.

---

CONTENTS.

	Page
An Enumeration of all the Plants known from China Proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the Island of Hongkong, together with their Distribution and Synonymy. By FRANCIS BLACKWELL FORBES, F.L.S., Knight Commander of the Swedish Royal Order of Wasa; and WILLIAM BOTTING HEMSLEY, F.R.S., F.L.S., Honorary Member of the Mexican Natural History Society, Keeper of the Herbarium of the Royal Gardens, Kew.— PART XX. ....	457
Index to Names and Synonyms, Vols. 23, 26, and 36.....	531
Titlepage, Contents, Historical Note, etc.	

---

LONDON:

SOLD AT THE SOCIETY'S APARTMENTS, BURLINGTON HOUSE,  
PICCADILLY, W.;

AND BY

LONGMANS, GREEN, AND CO.,

AND

WILLIAMS AND NORGATE.

1905

212

# LINNEAN SOCIETY OF LONDON.

## LIST OF THE OFFICERS AND COUNCIL.

Elected 24th May, 1905.

### PRESIDENT.

Prof. W. A. Herdman, D.Sc., F.R.S.

### VICE-PRESIDENTS.

Charles Baron Clarke, M.A., F.R.S.	Horace W. Monckton, F.G.S.
Frank Crisp, LL.B., B.A.	Dr. A. Smith Woodward, F.R.S.

### TREASURER.

Horace W. Monckton, F.G.S.

### SECRETARIES.

D. H. Scott, M.A., Ph.D., F.R.S.	Rev. T. R. R. Stebbing, M.A., F.R.S.
----------------------------------	--------------------------------------

### GENERAL SECRETARY.

B. Daydon Jackson, Esq.

### COUNCIL.

R. Assheton, M.A.	Prof. F. W. Oliver, D.Sc., F.R.S.
V. H. Blackman, M.A.	Clement Reid, F.R.S.
Gilbert O. Bourne, D.Sc.	Dr. A. B. Rendle, M.A.
Dr. Horace T. Brown, F.R.S.	Dr. W. G. Ridewood, F.Z.S.
Charles Baron Clarke, M.A., F.R.S.	Dukinfield H. Scott, M.A., Ph.D., F.R.S.
Frank Crisp, LL.B., B.A.	David Sharp, F.R.S.
Prof. J. B. Farmer, D.Sc., F.R.S.	Dr. Otto Stapf.
Rev. Canon Fowler, M.A.	Rev. T. R. R. Stebbing, M.A., F.R.S.
Prof. W. A. Herdman, F.R.S.	A. Smith Woodward, F.R.S.
B. Daydon Jackson, Esq.	
Horace W. Monckton, F.G.S.	

### LIBRARIAN.

A. W. Kappel.

### CLERK.

A. R. Hammond.

### LIBRARY COMMITTEE.

This consists of nine Fellows (three of whom retire annually) and of the officers *ex officio*; the former are elected annually by the Council in June, and serve till the succeeding Anniversary. The Committee meets as required during the Session. The Members for 1904-1905, in addition to the officers, are:—

Herbert Druce, F.Z.S.	A. B. Rendle, M.A., D.Sc.
Antony Gepp, M.A.	A. G. Tansley, M.A.
W. Botting Hemsley, F.R.S.	Prof. H. Marshall Ward, F.R.S.
Dr. G. Henderson.	Dr. A. Smith Woodward, F.R.S.
P. Chalmers Mitchell, M.A., F.Z.S.	

- ✓ *ARDISIA MACULOSA*, Mez, in Engl. Pflanzenr., Myrsin. 146.  
 ✓ *OBTUSA*, Mez, *l. c.* 104.  
 ✓ *PATENS*, Mez, *l. c.* 149.  
 ✓ *PENDULIFLORA*, Mez, *l. c.* 150.  
 ✓ *RADIANS*, Hemsl. & Mez, *l. c.* 146.  
 ✓ *TENERA*, Mez, *l. c.* 104.  
 ✓ *UNDULATA*, Mez, *l. c.* 146.  
 ✓ *YUNNANENSIS*, Mez, *l. c.* 107.  
*ARENARIA BARBATA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 430.  
*DELAVAYI*, Franch. *l. c.* 432.  
*LINEARIFOLIA*, Franch. Pl. Delav. (1889) 97.  
*LONGISTYLA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 433.  
*NAPULIGERA*, Franch. *l. c.* 429.  
*NEPALENSIS*, Spreng. ; Williams, in Journ. Linn. Soc., Bot. xxxiii (1898) 429.  
*ORBICULATA*, Royle; Franch. Pl. Delav. (1889) 92.  
*POLYTRICHOIDES*, Edgew.; Williams, in Journ. Linn. Soc., Bot. xxxiii. (1898) 404.  
*PRZEWALSKII*, Maxim. Fl. Tangut. (1889) 88.  
*QUADRIDENTATA*, Williams, in Journ. Linn. Soc., Bot. xxxiii. (1898) 432.  
*ROBOROWSKII*, Maxim. Fl. Tangut. (1889) 87.  
*SZECHUENSIS*, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1900) 437.  
*TRICHOPHORA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 431.  
*YUNNANENSIS*, Franch. *l. c.* 431.  
*ARETHUSANTHA BLETIoidES*, Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 412.  
 ✓ *ARISÆMA DAVIDIANUM*, Engl. Jahrb. xxv. (1898) 27.  
*GIRALDII*, Baroni, in Bull. Soc. Bot. Ital. (1893) No. 10, 515.  
 ✓ *TATARINOWII*, Schott ; Diels, in Engl. Jahrb. xxix. (1900) 236.  
*ARISTOLOCHIA DELAVAYI*, Franch. in Journ. de Bot. xii. (1898) 315.  
*GENTILIS*, Franch. *l. c.* 314.  
*GRACILLIMA*, Hemsl. in Kew Bull. (1900) 143.  
*MANSHURIENSIS*, Komarov, in Act. Hort. Petrop. xxii. (1903) 112.  
*MOUPINENSIS*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 79.  
*SETCHUENENSIS*, Franch. in Journ. de Bot. xii. (1898) 312.  
*SIPHO*, L'Hérit. ; Franch. *l. c.* 311.  
*THIBETICA*, Franch. *l. c.* 313.  
*YUNNANENSIS*, Franch. *l. c.* 313.  
*ARNEBIA SZECHENYII*, Kanitz, Pl. Exped. Szechenyi, Asia Centr. 42.  
*ARRACACHA DELAVAYI*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 115.  
*PEUCEDANIFOLIA*, Franch. *l. c.* 114.

- ARTEMISIA AURATA**, Komarov, in Act. Hort. Petrop. xviii. (1901) 420.  
**DEVERSA**, Diels, in Engl. Jahrb. xxix. (1900) 618.  
**LANCEA**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 500.  
**ASARUM BREVISTYLUM**, Franch. in Journ. de Bot. xii. (1898) 301.  
**CARDIOPHYLLUM**, Franch. in Bull. Mus. Hist. Nat. Par. i. (1895) 65.  
**CAULESCENS**, var., Franch. in Journ. de Bot. xii. (1898) 302.  
**CHINENSE**, Franch. *l. c.* 303.  
**DEBILE**, Franch. *l. c.* 305.  
**DELAVAYI**, Franch. in Bull. Mus. Hist. Nat. Par. i. (1895) 66.  
**FARGESII**, Franch. in Journ. de Bot. xii. (1898) 306.  
**FRANCHETIANUM**, Diels, in Engl. Jahrb. xxix. (1900) 308.  
**SAKAWANUM**, Makino, in Bot. Mag. Tokyo, ix. (1895) 260.  
**ASPARAGUS OLIGOCLONOS**, Maxim. in Act. Hort. Petrop. xix. (1901) 109.  
**ASPIDOPTERYS OBCORDATA**, Hemsl. in Journ. Linn. Soc., Bot. xxxiv. (1900) 475.  
**ASTER AURICULATUS**, Franch. in Journ. de Bot. x. (1896) 379.  
**BATANGENSIS**, Bur. & Franch. *l. c.* v. (1891) 50.  
**BIETII**, Franch. *l. c.* x. (1896) 373.  
**BRACHYTRICHUS**, Franch. *l. c.* 372.  
**BREVISCAPUS**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 495.  
**CANDELABRUM**, Vaniot, *l. c.* 498.  
**CURVATUS**, Vaniot, *l. c.* 499.  
**DELAVAYI**, Franch. in Journ. de Bot. x. (1896) 374.  
**FLABELLUM**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 492.  
**FUSCESCENS**, Bur. & Franch. in Journ. de Bot. v. (1891) 49.  
**INTEGRIFOLIUS**, Franch. in Mém. Soc. Nat. Cherbourg, xxiv. (1884) 222.  
**LATEBRACTEATUS**, Franch. in Journ. de Bot. x. (1896) 371.  
**LATICORYMBUS**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 494.  
**LIKIANGENSIS**, Franch. in Journ. de Bot. x. (1896) 370.  
**LINGULATUS**, Franch. *l. c.* 377.  
**MACILENTUS**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 497.  
**MILLEFOLIUS**, Vaniot, *l. c.* 496.  
**NIGRESCENS**, Vaniot, *l. c.* 493.  
**NIGROMONTANUS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 501.  
**OREOPHILUS**, Franch. in Journ. de Bot. x. (1896) 378.  
**PICCOLII**, Hook. f. Bot. Mag. t. 7669.  
**SENECIOIDES**, Franch. in Journ. de Bot. x. (1896) 381.  
**SETCHUENENSIS**, Franch. *l. c.* 377.  
**SOULMEI**, Franch. *l. c.* 372.  
**STATICEFOLIUS**, Franch. *l. c.* 370.

- ASTER TENUISSIMUS, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 502.  
     TONGOLENSIS, Franch. in Journ. de Bot. x. (1896) 376.  
     TRICAPITATUS, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 493.  
     VELLEREUS, Franch. in Journ. de Bot. x. (1896) 380.  
     VESTITUS, Franch. *l. c.* 378.  
     VILMORINI, Franch. *l. c.* 373.  
     YUNNANENSIS, Franch. *l. c.* 375.  
 ASTILBE PINNATA, Franch. Pl. Delav. (1890) 231.  
     RUBRA, Hook. f. & Thoms. ex Franch. *l. c.* 230.  
 ASTRAGALUS BRACHYCEPHALUS, Franch. *l. c.* (1890) 161.  
     CAMPTODONTUS, Franch. *l. c.* (1889) 160.  
     DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 214.  
     GRAVEOLENS, Buch.-Ham.; Franch. Pl. Delav. (1890) 163.  
 →HENRYI, Oliver, in Hook. Ic. Pl. xx. (1891) 1959.  
     LITANGENSIS, Bur. & Franch. in Journ. de Bot. v. (1891) 24.  
     LOCZII, Kanitz, Pl. Exped. Szechenyi, Asia Centr. 17.  
     MOUPINENSIS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 213.  
     NIGRESCENS, Franch. Pl. Delav. (1890) 162.  
     POLYCLADUS, Bur. & Franch. in Journ. de Bot. v. (1891) 23.  
     SUTCHUENENSIS, Franch. Pl. Delav. (1889) 160.  
     TANGUTICUS, Batalin, in Act. Hort. Petrop. xi. (1891) 485.  
     TATSIENENSIS, Bur. & Franch. in Journ. de Bot. v. (1891) 23.  
     YUNNANENSIS, Franch. Pl. Delav. (1890) 162.  
 ATRAGENE KOREANA, Komarov, in Act. Hort. Petrop. xxii. (1903) 278.  
     OCHOTENSIS, Pallas; Komarov, *l. c.* 276.  
 ATRIPLEX MAXIMOWICZIANA, Makino, in Bot. Mag. Tokyo, x. (1896) No. 107, 2.  
 ATYLOSIA TRICHODON, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 491.  
 AULISCONEMA PERNYI, Hua, in Journ. de Bot. vi. (1892) 472.  
  
 BALANOPHORA KUROIWAI, Makino, in Bot. Mag. Tokyo, xvi. (1902) 213.  
 BARTSIA ODONTITES, Huds.; Franch. in Journ. de Bot. iv. (1890) 317.  
 BAUHINIA BRYONIFLORA, Franch. Pl. Delav. (1890) 191.  
     DELAVAYI, Franch. *l. c.* 192.  
     DENSIFLORA, Franch. *l. c.* 191.  
 →FABERI, Oliver, in Hook. Ic. Pl. xviii. t. 1790.  
     YUNNANENSIS, Franch. Pl. Delav. (1890) 190.  
 →BEAUMONTIA BREVITUBA, Oliver, in Hook. Ic. Pl. xvi. (1887) t. 1582.  
     INDECORA, Baill. in Bull. Soc. Linn. Par. i. (1888) 759.



- BEGONIA AUGUSTINEI, Hemsl. in Gard. Chron. (1900) ii. 286.  
 BOWRINGIANA, Hort. ; Sander, in Gard. Chron. (1903) i. 245.  
 BRETSCHNEIDERIANA, Hemsl. in Hook. Ic. Pl. xxvii. t. 2635.  
 ↪ DIELSIANA, E. Pritzl, in Engl. Jahrb. xxix. (1900) 479.  
 HEMSLEYANA, Hook. f. Bot. Mag. t. 7685.
- BEKETOWIA, Krassn. in Script. Hort. Univ. Petrop. ii. i. (1887-88)  
 12.  
 TIANSCHANICA, Krassn. *l. c.* 13.
- BERBERIS ACUMINATA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
 387.  
 . ANGULOSA, Wall. var., Franch. *l. c.* 388.  
 DICTYOPHYLLA, Franch. Pl. Delav. (1889) 39.  
 GRACILIPES, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1754.  
 ↪ INSIGNIS, Hook. f. ; Engl. Jahrb. xxix. (1900) 341.  
 INTEGERRIMA, Bunge ; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
 386.  
 KOREANA, Palibin, in Act. Hort. Petrop. xvii. i. (1899) 22.  
 LEVIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 386.  
 PALLENS, Franch. Pl. Delav. (1889) 36.  
 POLYANTHA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1892) 302.  
 POTANINI, Maxim. in Act. Hort. Petrop. xi. i. (1890) 41.  
 PRUINOSA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 387.  
 SANGUINEA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
 Par. sér. 2, viii. (1885) 194.  
 SUBTRIPLINERVIS, Franch. in Bull. Mus. Hist. Nat. Par. i. (1895)  
 63.  
 YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 388.
- ↪ BERCHEMIA FLAVESCENS, Wall. ; Engl. Jahrb. xxix. (1900) 458.  
 YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 456.
- BERTEROA POTANINI, Maxim. Bull. Acad. Pétersb. xxvi. (1880) 422.
- BETULA COSTATA, Trautv. ; Komarov, in Act. Hort. Petrop. xxii.  
 (1903) 43.  
 CYLINDROSTACHYS, Wall. var., Diels, in Engl. Jahrb. xxix. (1900)  
 282.  
 ERMANI, Cham. ; Komarov, in Act. Hort. Petrop. xxii. (1903) 49.  
 FRUTICOSA, Pallas ; Komarov, *l. c.* 50.  
 LATIFOLIA, Tausch ; Komarov, *l. c.* 38.  
 SCHMIDTII, Regel ; Komarov, *l. c.* 52.
- BIFARIA DAVIDIANA, Van Tiegh. in Bull. Soc. Bot. Fr. xliii. (1896)  
 173.  
 FASCICULATA, Van Tiegh. *l. c.* 174.
- BLUMEA ADENOPHORA, Franch. in Journ. de Bot. x. (1896) 382.  
 BODINIERI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 23.  
 DUCLOUXII, Vaniot, *l. c.* 25.  
 GLOBATA, Vaniot, *l. c.* 24.

- BLUMEA GRACILIS, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 502.  
 HENRYI, Dunn, *l. c.* 503.  
 HONGKONGENSIS, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 22.  
 MARTINIANA, Vaniot, *l. c.* 26.  
 VERONICÆFOLIA, Franch. in Journ. de Bot. x. (1896) 382.
- BOCCONIA MICROCARPA, Maxim. in Act. Hort. Petrop. xi. i. (1890) 45.
- BODINIERIA, H. Léveillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 48.  
 THALICTRIFOLIA, H. Léveillé & Vaniot, *l. c.* 48.
- BOEHMERIOPSIS, Komarov, in Act. Hort. Petrop. xviii. (1901) 441.  
 PALLIDA, Komarov, *l. c.* 441.
- BOMBAX TENEBROSUM, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 486.
- BOTRYOPLEURON, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2670.  
 AXILLARE, Hemsl. *l. c.*  
 LATIFOLIUM, Hemsl. *l. c.*  
 STENOSTACHYUM, Hemsl. *l. c.*  
 VENOSUM, Hemsl. *l. c.*
- BOURNEA SINENSIS, Oliver, in Hook. Ic. Pl. xxiii. (1893) t. 2254.
- BRACHYACTIS CHINENSIS, Bur. & Franch. in Journ. de Bot. v. (1891) 69.
- BRANDISIA RACEMOSA, Hemsl. in Kew Bull. (1895) 114.
- BRASSAIOPSIS CILIATA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 499.  
 FICIFOLIA, Dunn, *l. c.* 500.
- BRAYA RUBICUNDA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 403.  
 SINENSIS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 303.  
 SINUATA, Maxim. Fl. Tangut. (1889) 69.
- BRETSCHNEIDERA, Hemsl. in Hook. Ic. Pl. xxviii. (1901) t. 2708.  
 SINENSIS, Hemsl. *l. c.*
- ↳ BROUSSONETIA KAZINOKI, Sieb. ; Engl. Jahrb. xxix. (1900) 298.
- BRUNELLA VULGARIS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 70.
- ↳ BUCKLEYA GRAEBNERIANA, Diels, in Engl. Jahrb. xxix. (1900) 306.  
 ↳ HENRYI, Diels, *l. c.* 306.
- BUDDLEIA ACUTIFOLIA, C. H. Wright, in Kew Bull. (1896) 24.  
 DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1888) 65.  
 ↳ GIRALDII, Diels, in Engl. Jahrb. xxix. (1900) 535.  
 HEMSLEYANA, Koehne, in Gartenfl. (1903) 169.  
 OFFICINALIS, Oliver, in Hook. Ic. Pl. xx. (1891) 1972.
- BUPLEURUM CANDOLLEI, Wall. ; Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 805.  
 COMMELINOIDEUM, Boissieu, *l. c.*  
 ↳ GRACILIPES, Diels, in Engl. Jahrb. xxix. (1900) 493.  
 ↳ LONGICAULE, Wall. ; Diels, *l. c.* 493.  
 ↳ LONGIRADIATUM, Turcz. ; Diels, *l. c.* 493.

- ✓ *BUPLEURUM MICROCEPHALUM*, Diels, in Engl. Jahrb. xxix. (1900) 494.  
*PETIOLULATUM*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 119.  
*SACHALINENSE*, F. Schmidt; Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 958.
- ✓ *TENUE*, Buch.-Ham. ex D. Don; Engl. Jahrb. xxix. (1900) 493.  
*YUNNANENSE*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 117.
- CACALIA FIRMA*, Komarov, in Act. Hort. Petrop. xviii. II. (1900) 420.
- CÆSALPINIA MORSEI*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 492.
- CALANTHE ALPINA*, Hook. f.; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 412.  
*BILOBA*, Lindl.; Finet, *l. c.* 412.  
*DAVIDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 85.  
*DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 434.  
*FARGESII*, Franch. *l. c.*  
*FIMBRIATA*, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 86.  
*MEGALOPHA*, Franch. *l. c.* 85.  
*PUSILLA*, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 436.
- ✓ *CALLICARPA VASTIFOLIA*, Diels, in Engl. Jahrb. xxix. (1900) 547.
- CALLITRICHE AQUATICA*, DC.; Vaniot, in Bull. Acad. Géog. Bot. xiii. (1902) 18.
- CALORHABDOS BRUNONIANA*, Benth.; Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 17.  
*FARGESII*, Franch. *l. c.* 16.  
*SUTCHUENENSIS*, Franch. *l. c.* 17.
- CALYPSO JAPONICA*, Maxim. in Act. Hort. Petrop. xx. (1901) 533.
- CAMELLIA GRACILIS*, Hemsl. in Ann. Bot. ix. (1895) 146.
- CAMPANULA CHRYSOSPENIFOLIA*, Franch. in Journ. de Bot. ix. (1895) 364.  
*CRENULATA*, Franch. *l. c.* 365.  
*DELAVAYI*, Franch. *l. c.* 364.
- ✓ *CAMPANUMÆA TRUNCATA*, Wall.; Diels, in Engl. Jahrb. xxix. (1900) 606.
- CAPPARIS FORMOSANA*, Hemsl. in Ann. Bot. ix. (1895) 145.  
*HAINANENSIS*, Oliver, in Hook. Ic. Pl. xvi. (1887) 1588.  
*HENRYI*, Matsum. in Bot. Mag. Tokyo (1899), 33.
- CARAGANA CRASSICAULIS*, Benth.; Franch. Pl. Delav. (1890) 159.  
*GERARDIANA*, Royle; Franch. *l. c.*
- ✓ *CARDAMINE CIRCÆOIDES*, Hook. f. & Thoms.; Engl. Jahrb. xxxii. (1902) 441.  
 ✓ *DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 397.  
*ENGLERIANA*, Schulz, in Engl. Jahrb. xxxii. (1902) 407.

- ✓ *CARDAMINE FLEXUOSA*, Withering; Engl. Jahrb. xxxii. (1902) 474.  
   ✓ Var. *occulta*, Harms; Engl. *l. c.* 479.  
 ✓ *FRAGARIIFOLIA*, Schulz, in Engl. *l. c.* 446.  
 ✓ *INSIGNIS*, Schulz, *l. c.* 439.  
 ✓ *LEUCANTHA*, Tausch; Schulz, in Engl. *l. c.* 403.  
 ✓ *MICROZYGA*, Schulz, *l. c.* 545.  
   *MULTIJUGA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 399.  
   *PROREPENS*, Fisch.; Komarov, in Act. Hort. Petrop. xxii. (1903) 360.  
   *STENOLOBA*, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 303.  
 ✓ *TANGUTORUM*, Schulz, in Engl. Jahrb. xxxii. (1902) 360.  
 ✓ *URBANIANA*, Schulz, *l. c.* 396.  
 ✓ *VIOLIFOLIA*, Schulz, *l. c.* 440.  
   *YUNNANENSIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 398.  
*CARDIANDRA SINENSIS*, Hemsl. in Gard. Chron. (1903) i. 82.  
*CARLESIA*, Dunn, in Hook. Ic. Pl. xxviii. (1902) t. 2739.  
   *SINENSIS*, Dunn, *l. c.* t. 2739.  
*CAROLINELLA*, Hemsl. in Hook. Ic. Pl. xxviii. (1902) t. 2726.  
   *CORDIFOLIA*, Hemsl. *l. c.* (1903) t. 2775.  
   *HENRYI*, Hemsl. *l. c.* (1902) t. 2726.  
*CARPESIUM ATKINSONIANUM*, Hemsl. in Kew Bull. (1893) 79, 157.  
   *EXIMIUM*, Winkl. in Act. Hort. Petrop. xiv. (1895) 58.  
   *FABERI*, Winkl. *l. c.* 65.  
   *HUMILE*, Winkl. *l. c.* 70.  
   *LIPSKYI*, Winkl. *l. c.* 68.  
   *TRISTE*, Maxim., var., Diels, in Engl. Jahrb. xxix. (1900) 615.  
   *VELUTINUM*, Winkl. in Act. Hort. Petrop. xiv. (1895) 73.  
*CARPINUS SEEMENIANA*, Diels, in Engl. Jahrb. xxix. (1900) 279.  
*CARRIERIA*, Franch. in Rev. Hort. (1896) 497.  
   *CALYCINA*, Franch. *l. c.* 497.  
*CARUM CANDOLLEANUM*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 127.  
   *CARDIOCARPUM*, Franch. *l. c.* 120.  
   *CARVI*, Linn.; Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 806.  
   *CAUDATUM*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 126.  
   *CORIACEUM*, Franch. *l. c.* 127.  
   *CRUCIATUM*, Franch. *l. c.* 124.  
   *DELAVAYI*, Franch. *l. c.* 120.  
   *DISSECTUM*, Franch. *l. c.* 123.  
   *FILICINUM*, Franch. *l. c.* 121.  
   *FLACCIDUM*, Franch. *l. c.* 126.  
   *HOOKERI*, Franch. *l. c.* 122.  
   *LOLOENSE*, Franch. *l. c.* 125.  
   *MOLLE*, Franch. *l. c.* 120.  
   *NEUROPHYLLUM*, Maxim.; Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 958.

- CARUM PANICULATUM, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 122.  
 PURPUREUM, Franch. *l. c.* 127.  
 SCABERULUM, Franch. *l. c.* 125.  
 SCHIZOPETALUM, Franch. *l. c.* ii. (1890) 199.  
 SINENSE, Franch. *l. c.* vi. (1894) 119.  
 TANAKÆ, Franch. & Sav.; Franch. *l. c.* 121.  
 TENERUM, Franch. *l. c.* 122.  
 TRICHOMANIFOLIUM, Franch. in Bull. Mus. Hist. Nat. Par. No. 2 (1895) 64.  
 YUNNANENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 128.
- CARYOPTERIS PARVIFOLIA, Batalin, in Act. Hort. Petrop. xiii. (1893) 98.
- CASSINE JAPONICA, O. Kuntze; Engl. & Prantl, Nat. Pflanzenf. iii. 5 (1892) 215.
- CATALPA FARGESII, Bur. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 3, vi. (1894) 195.  
 UMBRACULIFERA, Hort.; Bull. Soc. Tosc. Ort. (1888) 330.  
 ✓ VESTITA, Diels, in Engl. Jahrb. xxix. (1900) 577.
- CATHCARTIA DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 390.  
 LANCIFOLIA, Franch. *l. c.* 391.
- CAULOPHYLLUM ROBUSTUM, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 326.
- CEDRELA TOONA, Roxb.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 449.
- ✓ CELASTRUS FRANCHETIANUS, Loes. in Engl. Jahrb. xxx. (1902) 470.  
 ✓ GEMMATUS, Loes. *l. c.* 468.  
 HYPOGLAUCUS, Hemsl. in Ann. Bot. ix. (1895) 150  
 ✓ HYPOLEUCUS, Oliver; Loes. in Engl. Jahrb. xxix. (1900) 445.  
 ✓ MONOSPERMUS, Roxb.; Loes. *l. c.* xxx. (1902) 467.  
 ✓ ORBICULATUS, var., Loes. *l. c.* 469.  
 ✓ PANICULATUS, Willd.; Loes. *l. c.* 470.  
 RACEMULOSUS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 455.  
 ✓ ROSTHORNIANUS, Loes. in Engl. Jahrb. xxix. (1900) 445.  
 ✓ STYLOSUS, Wall.; Loes. *l. c.* xxx. (1902) 469.
- CENTELLA ASIATICA, Urb.; Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 802.
- CEPHALANTHERA LONGIBRACTEATA, Blume; Act. Hort. Petrop. xix. (1901) 104.
- CEPHALOTAXUS ARGOTÆNIA, Pilg. in Engl. Pflanz., Taxac. (1903) 104.
- CERASTIUM ALPINUM, Linn.; Franch. in Journ. de Bot. iv. (1890) 303.  
 ✓ FIMBRIATUM, E. Pritzell, in Engl. Jahrb. xxix. (1900) 320.  
 PUMILUM, Curt.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 435.

- CERASTIUM SZECHUENSE, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1900) 433.
- ✓ CERATOSTIGMA GRIFFITHII, C. B. Clarke; Engl. Jahrb. xxix. (1900) 525.
- CERCIS RACEMOSA, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1894.
- CHAMÆCYPARIS FORMOSENSIS, Matsum. in Bot. Mag. Tokyo Soc. xv. (1901) 137.
- CHAVICA HAINANA, C. DC., Ann. Conserv. Jard. Bot. Genève, ii. (1898) 275.
- CHEIRANTHUS ROSEUS, Maxim. Fl. Tangut. (1889) 57.
- CHELIDONIUM FRANCHETIANUM, Prain, in Bull. Herb. Boiss. iii. (1895) 586.
- JAPONICUM, Thunb.; Act. Hort. Petrop. xvi. (1899) 23.
- LASIOCARPUM, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1739.
- LEPTOPODUM, Prain, in Bull. Herb. Boiss. iii. (1895) 586.
- SUTCHUENENSE, Franch. in Journ. de Bot. viii. (1894) 293.
- ✓ CHELONOPSIS DEFLEXA, Benth.; Diels, in Engl. Jahrb. xxix. (1900) 554.
- CHENOPODIUM BRYONIAEFOLIUM, Bunge; Komarov, in Act. Hort. Petrop. xxii. (1903) 151.
- RUBRUM, Linn.; Komarov, *l. c.* 154.
- CHIMONANTHUS NITENS, Oliver, in Hook. Ic. Pl. xvi. (1887) t. 1600.
- CHIRITA DEPRESSA, Hook. f. Bot. Mag. t. 7213.
- SPECIOSA, Kurz, in Journ. Bot. xi. (1873) 195.
- CHRYSANTHEMUM TATSIENENSE, Bur. & Franch. in Journ. de Bot. v. (1891) 72.
- CHRYSOSPLENIUM AOMORENSE, Tokubuchi, in Bot. Mag. Tokyo, ix. (1895) 334.
- BAICALENSE, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 423.
- CILIATUM, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 3, ii. (1890) 108.
- GRACILE, Franch. Pl. Dav. *l. c.* sér. 2, viii. (1885) 234.
- ✓ GRAYANUM, Maxim.; Engl. Jahrb. xxix. (1900) 368.
- HENRYI, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 3, ii. (1890) 112.
- MACROPHYLLUM, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1744.
- MICROSPERMUM, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 3, ii. (1890) 109.
- NEPALENSE, Franch. Pl. Delav. i. (1890) 249.
- NODULOSUM, Tokubuchi, Bot. Mag. Tokyo, ix. (1895) 334.
- PILOSUM, Komarov, Act. Hort. Petrop. xxii. (1903) 424.
- PUMILUM, Franch. Pl. Delav. (1889) 249.
- RAMOSUM, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 422.

- CHRYSOSPLENIUM TIANSHANICUM*, Krassn. Script. Hort. Univ. Petrop. ii. 1. (1887-8) 16.  
*VILLOSUM*, Franch. in Journ. de Bot. iv. (1890) 305.
- CIMICIFUGA CALTHÆFOLIA*, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1746.  
 ✓ *FÆTIDA*, Linn.; Engl. Jahrb. xxix. (1900) 326.  
*HERACLEIFOLIA*, Komarov, in Act. Hort. Petrop. xviii. (1901) 438.  
 ✓ *JAPONICA*, Spreng.; Engl. Jahrb. xxix. (1900) 326.  
 ✓ *SIMPLEX*, Wormsk.; Engl. *l. c.* 326.
- ✓ *CIRCÆA QUADRISULCATA*, Maxim.; Engl. Jahrb. xxix. (1900) 485.
- CIRSIUM* = *CNICUS*.  
 ✓ *CHINENSE*, Gardn. & Champ. var., Diels, Engl. Jahrb. xxix. (1900) 628.  
 ✓ *FARGESII*, Franch.; Diels, *l. c.* 627.  
 ✓ *HENRYI*, Franch.; Diels, *l. c.*  
 ✓ *WALLICHII*, DC.; Diels, *l. c.*
- ✓ *CISSUS CARNOSA*, Wall.; Engl. Jahrb. xxix. (1900) 466.  
 ✓ *SETULOSA*, Diels & Gilg, *l. c.* 467.
- ✓ *CITRUS HYSTRIX*, DC.; Engl. Jahrb. xxix. (1900) 424.
- CLADRASTIS SINENSIS*, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 304.
- CLEMATIS ARMANDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 184.  
*ATRAGENOIDES*, Batalin, in Act. Hort. Petrop. xi. (1891) 481.  
*BUCHANANIANA*, DC.; H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. x. (1902) 169.  
*CHRYSOCOMA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 362.  
*CLARKEANA*, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 170.  
*COREANA*, H. Lévillé & Vaniot, *l. c.* 298.  
*DASYANDRA*, Maxim.; Act. Hort. Petrop. xi. (1890) 7.  
*DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 360.  
*DRAKEANA*, H. Lévillé & Vaniot, Bull. Acad. Géog. Bot. xi. (1902) 168.  
*FARGESII*, Franch. in Journ. de Bot. viii. (1894) 273.  
*FASCICULIFLORA*, Franch. Pl. Delav. (1889) 5.  
*FORMOSANA*, O. Kuntze, in Hook. Ic. Pl. xx. (1890) t. 1945.  
*FRUTICOSA*, var., Maxim. in Act. Hort. Petrop. xi. (1890) 5.  
*FUNEBRIS*, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 168.  
*FUSCA*, Turcz.; H. Lévillé & Vaniot, *l. c.* 299.
- ✓ *GOURIEANA*, Roxb.; Engl. Jahrb. xxix. (1900) 332.  
*HEDYSARIFOLIA*, DC. ex H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 169.  
*HENRYI*, Oliver, in Hook. Ic. Pl. xix. (1889) 1819.  
*INTRICATA*, Bunge, var., Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 288.

- CLEMATIS KOREANA, Komarov, in Act. Hort. Petrop. xviii. (1901) 438.
- KUNTZIANA, H. Lévêillé & Vaniot, Bull. Acad. Géog. Bot. xi. (1903) 171.
- LANCIFOLIA, Bur. & Franch. in Journ. de Bot. v. (1891) 18.
- LASIANDRA, Maxim. in Act. Hort. Petrop. xi. (1890) 7.
- LEIOCARPA, Oliver, in Hook. Ic. Pl. xvi. (1888) t. 1533.
- MONTANA, var., Maxim. in Act. Hort. Petrop. xi. (1890) 10.
- OBSCURA, Maxim. *l. c.* 6.
- PHILIPPIANA, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1903) 169.
- POGONANDRA, Maxim. in Act. Hort. Petrop. xi. (1890) 8.
- POTANINI, Maxim. *l. c.* 9.
- PRATTII, Hemsl. in Kew Bull. (1892) 82.
- PTERANTHA, Dunn, in Hook. Ic. Pl. xxviii. (1901) t. 2713.
- RANUNCULOIDES, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 360.
- RUBIFOLIA, C. H. Wright, in Kew Bull. (1896) 21.
- SPECTABILIS, Palibin, in Act. Hort. Petrop. xvii. (1899) 12.
- SPLENDENS, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 171.
- TASHIROI, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 477.
- VITALBA, Linn., var., Franch. Pl. Delav. (1889) 4.
- YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 361.
- CLEMATOCLETHRA ACTINIDIOIDES, Maxim. in Act. Hort. Petrop. xi. (1890) 38.
- CORDIFOLIA, Franch. in Journ. de Bot. viii. (1894) 279.
- FABERI, Franch. *l. c.* 279.
- GIRALDII, Diels, in Engl. Jahrb. xxix. (1900) 472.
- HEMSLEYI, Baill. in Bull. Soc. Linn. Par. ii. (1890) 873.
- INTEGRIFOLIA, Maxim. in Act. Hort. Petrop. xi. (1890) 38.
- LASIOCLADA, Maxim. *l. c.* 38.
- MAXIMOWICZI, Baill. in Bull. Soc. Linn. Par. ii. (1890) 873.
- SCANDENS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 53.
- STRIGILLOSA, Franch. in Journ. de Bot. viii. (1894) 280.
- TOMENTELLA, Franch. *l. c.*
- CLERODENDRON MANDARINORUM, Diels, in Engl. Jahrb. xxix. (1900) 549.
- MOUPINENSE, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 68.
- SUBSCAPOSUM, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2675.
- CLETHRA DELAVAYI, Franch. in Journ. de Bot. ix. (1895) 370.
- FARGESII, Franch. *l. c.* 369.
- CNICUS ARGYRACANTHUS, Hook. f. ; Franch. in Journ. de Bot. xi. (1897) 24.



- CNICUS BODINIERI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 121.  
   CERBERUS, Vaniot, *l. c.* 120.  
   FARGESII, Franch. in Journ. de Bot. xi. (1897) 22.  
   HENRYI, Franch. *l. c.* 21.  
   INVOLUCRATUS, Hook. f.; Franch. *l. c.* 24.  
   LEDUCI, Franch. *l. c.* 23.  
   MONOCEPHALUS, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 122.  
   PROVOSTI, Franch. in Journ. de Bot. xi. (1897) 23.  
   SOULIERI, Franch. *l. c.* 21.  
   SUFFULTUS, Franch. *l. c.* 24.  
 — COCCULUS AFFINIS, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1760.  
 COCHLEARIA SCAPIFLORA, Hook. f. & Thoms.; Franch. Pl. Delav. (1889)  
   61.  
 CODONOPSIS CONVULVULACEA, Hemsl. in Hook. Ic. Pl. xxiv. (1895)  
   t. 2385.  
   HENRYI, Oliver, *l. c.* xx. (1891) t. 1967.  
   SYLVESTRIS, Komarov, in Act. Hort. Petrop. xviii. (1900) 425.  
   TANGSHEN, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1966.  
 CŒLOGYNE BARBATA, Griff.; Finet, in Bull. Soc. Bot. Fr. xlv. (1898)  
   412.  
 —MANDARINORUM, Kränzl.; Engl. Jahrb. xxix. (1900) 269.  
   NITIDA, Lindl.; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 412.  
   OCHRACEA, Lindl.; Finet, *l. c.*  
   PRÆCOX, Lindl.; Finet, *l. c.*  
 COLURIA HENRYI, Batalin, in Act. Hort. Petrop. xiii. (1893) 94.  
 COLUTEA DELAVAYI, Franch. Pl. Delav. (1889) 158.  
 CONYZA PINNATIFIDA, Franch. in Journ. de Bot. x. (1896) 381.  
 COPTIS CHINENSIS, Franch. in Journ. de Bot. xi. (1897) 231.  
 CORALLODISCUS, Batalin, in Act. Hort. Petrop. xii. (1892) 176.  
   CONCHÆFOLIUS, Batalin, *l. c.* 176.  
 ✓ CORCHOROPSIS PSILOCARPA, Harms et Loes. in Engl. Jahrb. xxxiv.  
   (1904) Beibl. 75, p. 51.  
 CORCHORUS OLITORIUS, Linn.; Franch. in Bull. Soc. Bot. Fr. xxxiii.  
   (1886) 441.  
 CORIARIA INTERMEDIA, Matsum. in Bot. Mag. Tokyo, xii. (1898) 62.  
   NEPAULENSIS, Wall.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
   451.  
   TERMINALIS, Hemsl. in Hook. Ic. Pl. xxiii. (1892) t. 2220.  
 CORISPERMUM ELONGATUM, Bunge; Komarov, in Act. Hort. Petrop.  
   xxii. (1903) 158.  
 CORNUS PRETSCHNEIDERI, J. Henry, in Le Jardin (1899) 309.  
   QUINQUENERVIS, Franch. in Journ. de Bot. x. (1896) 307.  
   SCABRIDUS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
   sér. 2, viii. (1885) 250.  
 CORTUSA PEKINENSIS, A. Richt. in Termész. Füzetek (1894) 130.

- CORYDALIS ACUMINATA, Franch. in Journ. de Bot. viii. (1894) 285.
- AEGOPODIOIDES, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 173.
- ALBICAULIS, var., Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 396.
- ANTHRISICIFOLIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 196.
- BALSAMIFLORA, Prain, in Journ. As. Soc. Beng. lxxv. (1896) 41.
- BATALIANA, Prain, ex Bretschn. Hist. Eur. Bot. Disc. China, 1026.
- CHEILANTHIFOLIA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 302.
- CHEIRIFOLIA, Franch. in Journ. de Bot. viii. (1894) 285.
- CHELIDONIIFOLIA, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 174.
- CONSPERSA, Maxim. Fl. Tangut. (1889) 49.
- CRISTA-GALLI, Maxim. *l. c.* 47.
- CRISTATA, Maxim. in Act. Hort. Petrop. xi. (1890) 47.
- CURVIFLORA, Maxim. Fl. Tangut. (1889) 41.
- DAUCIFOLIA, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 172.
- DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 198.
- DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 393.
- DRAKEANA, Prain, in Journ. As. Soc. Beng. lxxv. (1896) 31.
- DUCLOUXII, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 174.
- ECHINOCARPA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 393.
- ELATA, Bur. & Franch. in Journ. de Bot. v. (1891) 20.
- FARGESII, Franch. *l. c.* viii. (1894) 290.
- FLEXUOSA, Franch. Pl. Dav. in Nouv. Arch. Mus. Nat. Hist. Par. sér. 2, viii. (1885) 197.
- FUMARIA, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 172.
- GIGANTEA, Trautv. & Mey.; Komarov, in Act. Hort. Petrop. xxii. (1903) 343.
- GRACILIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 395.
- HAMATA, Franch. in Journ. de Bot. viii. (1894) 192.
- HANNÆ, Kanitz, Pl. Exped. Szechenyi, Asia Centr. (1891) 7.
- HEMSLEYANA, Franch. & Prain, in Journ. As. Soc. Beng. lxxv. (1896) 29.
- LIVIDA, Maxim. Fl. Tangut. (1889) 49.
- LONGICORNU, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 394.
- LOPINENSIS, Franch. in Journ. de Bot. viii. (1894) 284.
- MARTINI, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 173.
- MOUPINENSIS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 198.

- CORYDALIS MUCRONATA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 197.
- OPHIOCARPA, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 355.
- OXYPETALA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 392.
- PACHYCENTRA, Franch. Pl. Delav. (1889) 45.
- PETROPHILA, Franch. *l. c.* 47.
- POTANINI, Maxim. Fl. Tangut. (1889) 48.
- PRATTII, Franch. in Journ. de Bot. viii. (1894) 284.
- PULCHELLA, Franch. Pl. Delav. (1889) 45.
- SCABERULA, Maxim. Fl. Tangut. (1889) 40.
- SCANDENS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 391.
- SOLIDA, Sw.; Palibin, in Act. Hort. Petrop. xvii. (1899) 24.
- SOULIERI, Franch. in Journ. de Bot. viii. (1894) 283.
- STENANTHA, Franch. Pl. Delav. (1889) 49.
- TALIENSIS, Franch. *l. c.* 48.
- TEMULIFOLIA, Franch. in Journ. de Bot. viii. (1894) 291.
- THALICTRIFOLIA, Franch. *l. c.* 291.
- THIBETICA, var., Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 396.
- TOMENTELLA, Franch. in Journ. de Bot. viii. (1894) 292.
- TOMENTOSA, N. E. Br. in Gard. Chron. (1903) ii. 123.
- TONGOLENSIS, Franch. in Journ. de Bot. viii. (1894) 285.
- TRIFOLIATA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 392.
- TRISECTA, Franch. in Journ. de Bot. viii. (1891) 284.
- TRITERNATA, Franch. *l. c.* 290.
- WILSONI, N. E. Br. in Gard. Chron. (1903) ii. 123.
- YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 394.
- CORYLOPSIS, n. sp., Diels, in Engl. Jahrb. xxix. (1900) 380, 381.
- COTONEASTER ACUMINATA, Lindl.; Engl. Jahrb. xxix. (1900) 385.
- ANGUSTIFOLIA, Franch. Pl. Delav. (1890) 221.
- BACILLARIS, Wall.; Franch. *l. c.* 222.
- BUXIFOLIA, Wall.; Franch. *l. c.* 224.
- CORIACEA, Franch. *l. c.* 222.
- DIELSIANA, E. Pritz. in Engl. Jahrb. xxix. (1900) 385.
- FRANCHETI, Bois, in Rev. Hort. (1902) 379.
- FRIGIDA, Wall.; Engl. Jahrb. xxix. (1900) 385.
- GLAUCOPHYLLA, Franch. Pl. Delav. (1890) 222.
- MOUPINENSIS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 224.
- MUCRONATA, Franch. Pl. Delav. (1890) 223.
- NUMMULARIA, Fisch. & Mey.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 225.
- PANNOSA, Franch. Pl. Delav. (1890) 223.
- ROTUNDIFOLIA, Wall.; Engl. Jahrb. xxix. (1900) 385.
- RUGOSA, E. Pritz. in Engl. *l. c.* 385.

- COTONEASTER SALICIFOLIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 225.
- COTYLEDON JAPONICA, Maxim.; Palibin, in Act. Hort. Petrop. xvii. (1899) 93.  
MINUTA, Komarov, *l. c.* xviii. (1900) 436.
- CRASSULA YUNNANENSIS, Franch. in Journ. de Bot. x. (1896) 284.
- CRATÆGUS DSUNGARICA, Zabel; Lange, Revis. Cratæg. 43.  
HENRYI, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 494.  
TENUIFOLIA, Komarov, in Act. Hort. Petrop. xviii. (1901) 435.
- CRAWFURDIA DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 306.  
THIBETICA, Franch. *l. c.* 307.
- CREMANTHODIUM POTANINI, Winkl. in Act. Hort. Petrop. xiv. (1895) 150.
- CREPIS BOCKIANA, Diels, in Engl. Jahrb. xxix. (1900) 633.  
GILLII, S. Moore, in Journ. Bot. xxxvii (1899) 170.
- HENRYI, Diels, in Engl. Jahrb. xxix. (1900) 633.  
HOOKERIANA, C. B. Clarke; Franch. in Journ. de Bot. ix. (1895) 256.  
PHŒNIX, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 511.  
RAPUNCULOIDES, Dunn, *l. c.* 512.
- ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 632.  
TRICHOCARPA, Franch. in Journ. de Bot. ix. (1895) 257.  
UMBRELLA, Franch. *l. c.* 255.
- CROTALARIA CAPITATA, Benth.; Franch. Pl. Delav. (1889) 151.  
FORMOSANA, Matsum.; Ito & Matsum., Journ. Coll. Sci. Tokyo, xii. 395.  
RUFESCENS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 133.  
SIMILIS, Hemsl. in Ann. Bot. ix. (1895) 152.  
YUNNANENSIS, Franch. Pl. Delav. (1889) 151.
- CRYPTOTÆNIOPSIS, Dunn, in Hook. Ic. Pl. xxviii. (1902) 2737.  
ASPLENIOIDES, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 807.  
BOTRYCHIOIDES, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 494.  
CARDIOCARPA, Dunn, *l. c.* 495.  
DELAVAYI, Dunn, *l. c.* 495.  
FILICINA, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 806.  
LEPTOPHYLLA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 495.  
MOLLIS, Dunn, *l. c.*  
TANAKÆ, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 806.
- VULGARIS, Dunn, in Hook. Ic. Pl. xxviii. (1902) t. 2737.
- CUCUBALUS BACCIFER, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 205.
- CYANANTHUS PETIOLATUS, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 147.

- CYCLEA POLYPETALA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 485.  
 — RACEMOSA, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1938.
- CYDONIA CATHAYENSIS, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2657-58.
- CYMBIDIUM GIGANTEUM, Wall.; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 412.
- CYNANCHUM HENRYI, Warb. in Engl. Jahrb. xxix. (1900) 542.  
 NIPPONICUM, Matsum. in Bot. Mag. Tokyo, xii. (1898) 39.  
 VILLOSUM, Matsum. *l. c.*
- CYNOSORCHIS GRACILIS, Blume; Kränzl, Gen. & Sp. i. 488.
- CYPERORCHIS ELEGANS, Blume; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 412.
- CYPRIPEDIUM ARIETINUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 206.  
 — ELEGANS, Reichb. f.; Diels, in Engl. Jahrb. xxix. (1900) 266.
- DALBERGIA DELAVAYI, Franch. Pl. Delav. (1890) 186.  
 — DYERIANA, Prain; Engl. Jahrb. xxix. (1900) 416.  
 MIMUSOIDES, Franch. Pl. Delav. (1890) 187.  
 YUNNANENSIS, Franch. *l. c.*
- DAPHNE GEMMATA, E. Pritz. in Engl. Jahrb. xxix. (1900) 481.  
 — GRACILIS, E. Pritz. *l. c.* 480.  
 RETUSA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 318.  
 TENUIFLORA, Bur. & Franch. in Journ. de Bot. v. (1891) 151.
- DAPHNIDIUM STRYCHNIFOLIUM, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 76.
- DAVIDIA, Baill. Adansonia, x. (1871) 114.  
 INVOLUCRATA, Baill. *l. c.*
- DECAISNEA FARGESII, Franch. in Journ. de Bot. vi. (1892) 234.  
 — INSIGNIS, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 342.
- DECUMARIA SINENSIS, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1741.
- DELAVAYA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 462.  
 TOXOCARPA, Franch. *l. c.*  
 YUNNANENSIS, Franch. Pl. Delav. (1889) 142.
- DELPHINIUM BONVALOTI, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 3, 165.  
 CAMPYLOCENTRUM, Maxim. in Act. Hort. Petrop. xi. i. (1890) 31.  
 CAUCASICUM, var., Maxim. *l. c.* 30.  
 CAVALERIENSE, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 49.  
 CERATOPHORUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 377.  
 CEREFOLIUM, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 49.  
 CHEFOENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, v. 3. (1893) 170.

- DELPHINIUM CÆLESTINUM, Franch. in Journ. de Bot. viii. (1894) 276.  
 DAVIDI, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 179.  
 DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 379.  
 DENSIFLORUM, Duthie; Huth, in Engl. Jahrb. xx. (1895) 393.  
 DENUDATUM, Wall.; var., Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 378.  
 ELATUM, Linn., var., H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. x. (1902) 300.  
 EXIGUUM, E. Pritz. in Engl. Jahrb. xxix. (1900) 327.  
 FARGESII, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 164.  
 FATSIIENSE, E. Pritz. in Engl. Jahrb. xxix. (1900) 328.  
 GILGIANUM, Pilg. *l. c.* xxxiv. (1904), Beibl. 75. 33.  
 HAMATUM, Franch. Pl. Delav. (1889) 28.  
 HENRYI, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 177.  
 HIRTICAULE, Franch. in Journ. de Bot. viii. (1894) 275.  
 ILIENSE, Huth; Engl. Jahrb. xx. (1895) 402.  
 LANKONGENSE, Franch. Pl. Delav. (1889) 26.  
 LIKIANGENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 180.  
 LONGIPES, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 191.  
 MAACKIANUM, Regel; Komarov, in Act. Hort. Petrop. xxii. (1903) 248.  
 MAXIMOWICZII, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 164.  
 MOSOYNENSE, Franch. *l. c.* 168.  
 OLIGANTHUM, Franch. Pl. Delav. (1889) 29.  
 ORTHOCENTRUM, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 178.  
 PACHYCENTRUM, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1892) 301.  
 POTANINI, Huth, in Bull. Herb. Boiss. i. (1893) 332.  
 PYCNOCENTRUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 379.  
 ROBERTIANUM, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 49.  
 SERTIFERUM, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 171.  
 SIWANENSE, Franch. *l. c.* 162.  
 SOULIEI, Franch. *l. c.* 172.  
 SUTCHUENENSE, Franch. *l. c.* 178.  
 TALIENSE, Franch. *l. c.* 174.  
 TANGUTICUM, Maxim. ex Bull. Herb. Boiss. i. (1893) 332.  
 TATSIIENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 169.

- DELPHINIUM TONGOLENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, v. (1893) 166.  
 TRICHOPHORUM, Franch. *l. c.* 166.  
 YUNNANENSE, Franch. *l. c.* 173.
- DENDROBIUM CAPILLIPES, Reichb. f. ; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 411.  
 CHRYSOSTOXUM, Lindl. ; Finet, *l. c.* 411.  
 FARGESII, Finet, *l. c.* 1. (1903) 374.  
 FIMBRIATUM, Hook. ; Finet, *l. c.* xlv. (1898) 412.  
 LONGICORNU, Lindl. ; Finet, *l. c.* 412.  
 ODIOSUM, Finet, *l. c.* 1. (1903) 373.  
 PIERARDI, Roxb. ; Finet, *l. c.* xlv. (1898) 412.  
 YUNNANENSE, Finet, *l. c.* xlv. (1897) 419.
- DERRIS FORDII, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1771.  
 MARGINATA, Benth. ; Franch. Pl. Delav. (1890) 188.
- DESMODIUM AMETHYSTINUM, Dunn, in Gard. Chron. (1902) i. 210.  
 CALLIANTHUM, Franch. Pl. Delav. (1890) 173.  
 CINERASCENS, Franch. *l. c.* 174.  
 GRACILLIMUM, Hemsl. in Ann. Bot. ix. (1895) 152.  
 GRIFFITHIANUM, Benth. ; Franch. Pl. Delav. (1899) 176.  
 HAMULATUM, Franch. *l. c.* 175.  
 HISPIDUM, Franch. *l. c.* 174.  
 RHABDOCLADUM, Franch. *l. c.* 173.  
 — SINUATUM, Blume ; Engl. Jahrb. xxix. (1900) 414.  
 TASHIROI, Matsum. in Journ. Coll. Sci. Tokyo, xii. (1900) 415.  
 YUNNANENSE, Franch. Pl. Delav. (1890) 172.
- DEUTZIA ALBIDA, Batalin, in Act. Hort. Petrop. xiii. vii. (1893) 97.  
 — BARONIANA, Diels, in Engl. Jahrb. xxix. (1900) 372.  
 CORYMBIFLORA, Rev. Hort. (1898) 401.  
 FARGESII, Franch. in Journ. de Bot. x. (1896) 281.  
 — GLABERRIMA, Koehne ; Engl. Jahrb. xxxiv. (1904) Beibl. 75. 38.  
 GLABRATA, Komarov, in Act. Hort. Petrop. xxii. (1903) 433.  
 GLOMERULIFOLIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1886) 236.  
 — HAMATA, Koehne ; Engl. Jahrb. xxxiv. (1904) Beibl. 75. 37.  
 LONGIFOLIA, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1886) 235.  
 SETCHUENENSIS, Franch. in Journ. de Bot. x. (1896) 282.
- DIANELLA STRAMINEA, Yatabe, in Bot. Mag. Tokyo, vii. (1893) 435.
- DIANTHUS SZECHUENSIS, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 428.
- DICENTRA MACRANTHA, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1937.  
 DICHROCEPHALA BODINIERI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 242.

- DICHRORHYZA LEVEILLEI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 241.  
 MINUTIFOLIA, Vaniot, *l. c.* 243.
- DICKINSIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 244.  
 HYDROCOTYLOIDES, Franch. *l. c.* 244.
- DIDISSANDRA DELAVAYI, Franch. in Bull. Mus. Hist. Nat. Par. v. (1899) 249.  
 FARGESII, Franch. in Bull. Soc. Linn. Paris, n. s. i. (1899) 123.  
 GLANDULOSA, Batalin, in Act. Hort. Petrop. xii. (1892) 175.  
 LANCIFOLIA, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 63.  
 LONGIFOLIA, *l. c.*  
 LONGIPES, Hemsl. in Kew Bull. (1895) 115.  
 MORGANI, Franch. in Bull. Soc. Linn. Par. n. s. i. (1899) 124.  
 PPLICATA, Franch. *l. c.*  
 — ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 574.
- DIOSCOREA ACERIFOLIA, Uline, in Engl. Jahrb. xxix. (1900) 261.  
 — BUERGERI, Uline, var.; Engl. Jahrb. xxix. (1900) 260.  
 FARGESII, Franch. Rev. Hort. (1896) 540.  
 — HENRYI, Uline, in Engl. Jahrb. xxix. (1900) 261.  
 PENTAPHYLLA, Linn.; Uline, *l. c.* 260.  
 ROSTHORNII, Diels; Uline, *l. c.* 261.  
 TENUIPES, Franch. & Sav.; Act. Hort. Petrop. xix. (1901) 108.
- DIOSPYROS UTILIS, Hemsl. in Ann. Bot. ix. (1895) 154.
- DIPelta ELEGANS, Batalin, in Act. Hort. Petrop. xiv. 8. (1895) 174.  
 YUNNANENSIS, Franch.; Rev. Hort. (1891) 247.
- DIPHYLLEIA CYMOSA, Michx.; Engl. Jahrb. xxix. (1900) 337.
- DIPLOLEPIS OVATA, Lindl. ex Trans. Hort. Soc. vi. (1826) 268.
- DIPOMA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 404.  
 IBERIDEUM, Franch. *l. c.* 405.
- DIPSACUS CHINENSIS, Batalin, in Act. Hort. Petrop. xiii. 18. (1894) 377.  
 — INERMIS, Wall.; Engl. Jahrb. xxix. (1900) 601.
- DIPTERONIA, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1898.  
 SINENSIS, Oliver, *l. c.*
- DISPOROPSIS ASPERSA, Hua; Engl. Jahrb. xxix. (1900) 249.  
 — PERNYI, Hua; Engl. *l. c.* 249.
- DIURANTHERA, Hemsl. in Hook. Ic. Pl. xxviii. (1902) t. 2734.  
 MAJOR, Hemsl. *l. c.* t. 2734.  
 MINOR, C. H. Wright, *l. c.* sub t. 2734.
- DOBINEA DELAVAYI, Baill. in Bull. Soc. Linn. Par. ii. (1890) 835.
- DOLICHOS LAGOPUS, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 499.



- DONTOSTEMON INTEGRIFOLIUS, C. A. Mey.; Komarov, in Act. Hort. Petrop. xxii. (1903) 383.
- DRABA ALPINA, Linn., var.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 401.
- AMPLEXICAULIS, Franch. *l. c.* 403.
- BRACTEATA, Batalin, in Act. Hort. Petrop. xi. 16. (1891) 482.
- DENTATA, Ledeb.; Maxim. Enum. Pl. Mongol. (1886) 58.
- MOUPINENSIS, Franch. Pl. Delav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 200.
- SURCULOSA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1889) 401.
- YUNNANENSIS, Franch. *l. c.* 402.
- DRACONTOMELUM SINENSE, Stapf, in Hook. Ic. Pl. xxvii. (1900) t. 2641.
- DUMASIA CORDIFOLIA, Hemsl. in Hook. *l. c.* t. 2627.
- DUNBARIA PULCHRA, Benth.; Franch. Pl. Delav. (1889) 184.
- DUPINIA JAPONICA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 439.
- ECHINOCARPUS SINENSIS, Hemsl. in Ann. Bot. ix. (1895) 147.
- EHRETIA CORYLIFOLIA, C. H. Wright, in Kew Bull. (1896) 25.
- ELÆAGNUS BOCKII, Diels, in Engl. Jahrb. xxix. (1900) 483.
- DAVIDII, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 77.
- HENRYI, Warb. in Engl. Jahrb. xxix. (1900) 483.
- LANCEOLATA, Warb. *l. c.* 483.
- ELÆOCARPUS HAINANENSIS, Oliver, in Hook. Ic. Pl. xxv. (1896) t. 2462.
- ELÆODENDRON JAPONICUM, Franch. & Sav.; Engl. Jahrb. xxx. (1902) 473.
- ELSHOLTZIA CALYCOCARPA, Diels, in Engl. Jahrb. xxix. (1900) 560.
- EMBELIA FORDII, Mez, in Engl. Pflanzenr., Myrsin. 326.
- MYRTIFOLIA, Mez, in Notizbl. Bot. Gard. Berlin, iii. (1901) 107.
- PAUCIFLORA, Diels, in Engl. Jahrb. xxix. (1900) 517.
- POLYPODIOIDES, Hemsl. & Mez, in Notizbl. Bot. Gard. Berlin, iii. (1901) 108.
- PROCUMBENS, Hemsl. in Hook. Ic. Pl. xxviii. (1901) t. 2724.
- PRUNIFOLIA, Mez, in Engl. Pflanzenr., Myrsin. 316.
- SAXATILIS, Hemsl. in Hook. Ic. Pl. xxviii. (1901) t. 2724.
- EMMENOPTERYS HENRYI, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1823.
- ENKIANTHUS BRACHYPHYLLUS, Franch. in Journ. de Bot. ix. (1895) 371.
- CHINENSIS, Franch. *l. c.*
- HIMALAICUS, Hook. f. & Thoms., var., Franch.; Engl. Jahrb. xxix. (1900) 509.
- ROSTHORNII, Diels, in Engl. *l. c.*
- EPILOBIUM ANGULATUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 432.

- EPILOBIUM CALYGINUM, Hausskn.; H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 17.  
 CEPHALOSTIGMA, Hausskn.; H. Lévillé, *l. c.*  
 HIMALAYENSE, Hausskn.; Bodin, *l. c.* ii. (1893) 341.  
 LÆTUM, Wall.; Bodin, *l. c.* xii. (1903) 17.  
 LIVIDUM, Hausskn.; H. Lévillé, *l. c.* viii. (1898) 5.  
 MINUTIFLORUM, Hausskn.; H. Lévillé, *l. c.* xii. (1903) 17.  
 MOLLE, Lam.; H. Lévillé, *l. c.* viii. (1898) 4.  
 NEPALENSE, Hausskn.; H. Lévillé, *l. c.* ii. (1893) 341.  
 NERVOSUM, Boiss. & Buhse; H. Lévillé, *l. c.* xii. (1903) 17.  
 NUDICARPUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 432.  
 ROUYANUM, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 17.  
 WALLICHIANUM, Hausskn.; H. Lévillé, *l. c.*  
 EPIMEDIUM ACUMINATUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 109.  
 BREVICORNU, Maxim. in Act. Hort. Petrop. xi. 1. (1889) 42.  
 DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 195.  
 FARGESII, Franch. Journ. de Bot. viii. (1894) 281.  
 SUTCHUENENSE, Franch. *l. c.* 282.  
 EPIPACTIS HELLEBORINE, var., Crantz; Engl. Jahrb. xxix. (1900) 268.  
 LATIFOLIA, Swartz; Bull. Soc. Bot. Fr. xlv. (1898) 413.  
 — MACROSTACHYA, Kränzl.; Engl. Jahrb. xxix. (1900) 268.  
 ERANTHIS ALBIFLORA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 191.  
 VANIOTIANA, H. Lévillé, in Bull. Acad. Géog. Bot. xi. (1902) 299.  
 ERIGERON MOUPINENSIS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 36.  
 ERIBOTRYA DUBIA, Decne.; Franch. Pl. Delav. (1890) 224.  
 GRIFFITHII, Franch. *l. c.*  
 LASIOGYNA, Franch. *l. c.* 225.  
 PRIONOPHYLLA, Franch. *l. c.*  
 ERIOCAULON DECEMFLORUM, Maxim. Diagn. Pl. Nov. As. fasc. viii. (1893) 7.  
 DICLINE, Maxim. *l. c.*  
 KIUSIANUM, Maxim. *l. c.* 22.  
 ERITRICHUM RADICANS, A. DC. in Act. Hort. Petrop. xviii. (1900) 162.  
 ERYNGIUM FÆTIDUM, Linn.; Hemsl. in Journ. Linn. Soc., Bot. xxxiv. (1900) 475.  
 ERYSIMUM AURANTIACUM, Maxim. Enum. Pl. Mong. (1889) 65.  
 YUNNANENSE, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 404.  
 — ERYTHROSPERMUM HYPOLEUCUM, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1899.  
 EUCHRESTA TUBULOSA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 492.

- *EUCOMMIA ULMOIDES*, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1950; xxiv. (1895) t. 2361.
- EUONYMUS ACANTHOCARPA*, Franch. Pl. Delav. (1889) 129.
- ✓ *ACTINOCARPA*, Loes. in Engl. Jahrb. xxx. (1901) 459.
  - ✓ *ACULEATA*, Hemsl. in Kew Bull. (1893) 209.
  - ✓ *AMYGDALIFOLIA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 453.
  - ✓ *BOCKII*, Loes. in Engl. Jahrb. xxix. (1900) 439.
  - ✓ *CHINENSIS*, Lindl. vars., Loes. in Engl. Jahrb. xxix. (1900) 436.
  - ✓ *CORNUTUS*, Hemsl. in Kew Bull. (1893) 209.
  - ✓ *DIELSIANA*, Loes. in Engl. Jahrb. xxix. (1900) 440.
  - ✓ *FIMBRIATA*, Wall. ex Engl. Jahrb. xxx. (1902) 465.
  - ✓ *FLAVESCENS*, Loes. *l. c.* xxix. (1900) 437.
  - ✓ *FORBESIANA*, Loes. *l. c.* xxx. (1902) 457.
  - ✓ *GIRALDII*, Loes. *l. c.* xxix. (1900) 442.
  - ✓ *GRANDIFLORA*, Wall. ex Loes. *l. c.* 439.
  - ✓ *HEMSLEYANA*, Loes. *l. c.* xxx. (1902) 460.
  - ✓ *HUPEHENSIS*, Loes. *l. c.* 454.
  - ✓ *ILICIFOLIA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 452.
  - ✓ *KIAUTSCHOVICA*, Loes. in Engl. Jahrb. xxx. (1902) 453.
  - ✓ *LANCEIFOLIA*, Loes. *l. c.* 462.
  - ✓ *LINEARIFOLIA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 455.
  - ✓ *LUTCHUENSIS*, T. Ito & Matsum. Engl. Jahrb. xxx. (1902) 456.
  - ✓ *MACROPTERA*, Rupr. ex Engl. *l. c.* 465.
  - ✓ *MYRIANTHA*, Hemsl. in Kew Bull. (1893) 210.
  - ✓ *PATENS*, Rehder; Sarg. Trees & Shrubs (1903) 127.
  - ✓ *PHELLOMANA*, Loes. in Engl. Jahrb. xxix. (1900) 444.
  - ✓ *ROSTHORNII*, Loes. *l. c.* 437.
  - ✓ *SALICIFOLIA*, Loes. *l. c.* xxx. (1902) 458.
  - ✓ *SANGUINEA*, Loes. *l. c.* xxix. (1901) 441.
  - ✓ *TASHIROI*, Maxim. in Bull. Acad. Pétersb. xxxi. (1887) 23.
  - ✓ *THEIFOLIA*, Wall. var.; Loes. in Engl. Jahrb. xxx. (1902) 455.
  - ✓ *TONKINENSIS*, Loes. *l. c.* 453.
  - ✓ *VENOSA*, Hemsl. in Kew Bull. (1893) 210.
  - ✓ *VERRUCOSOIDES*, Loes. in Engl. Jahrb. xxx. (1902) 462.
  - ✓ *YUNNANENSIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 454.
- EUPATORIUM CANNABINUM*, Linn.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1888) 34.
- EUPTELEA DAVIDIANA*, Baillon, Adansonia, xi. (1873-76) 305.
- ✓ *DELAVAYI*, van Tiegh. in Journ. de Bot. xiv. (1900) 273.
  - ✓ *FRANCHETII*, van Tiegh. *l. c.* 272.
  - ✓ *PLEIOSPERMA*, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 346.
  - ✓ *POLYANDRA*, Sieb. & Zucc.; Engl. *l. c.*
- EURYA GYNANDRA*, Vesque, in Bull. Soc. Bot. Fr. xlii. (1897) 160.
- ✓ *HENRYI*, Hemsl. in Hook. Ic. Pl. xxviii. (1903) sub t. 2761.
  - ✓ *OBLIQUIFOLIA*, Hemsl. *l. c.* t. 2761.

EUTREMA THIBETICUM, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 201.

YUNNANENSE, Franch. Pl. Delav. (1889) 61.

FAGARA=ZANTHOXYLON.

✓ AVICENNIAE, DC.; Engl. Jahrb. xxix. (1900) 423.

✓ DIMORPHOPHYLLA, Engl.; *l. c.* 422.

✓ DISSITA, Engl.; *l. c.*

✓ ECHINOCARPUS, Engl.; *l. c.*

✓ KHASIANA, Engl.; *l. c.*

✓ MICRANTHA, Engl.; *l. c.*

✓ STENOPHYLLA, Engl.; *l. c.*

✓ VOLUBILIS, E. Pritzl; *l. c.*

✓ FAGUS ENGLERIANA, Seemen, in Engl. Jahrb. xxix. (1900) 285.

✓ SIEBOLDII, Endl.; *l. c.* 284.

✓ FICUS HENRYI, Warb. in Engl. Jahrb. xxix. (1900) 299.

TASHIROI, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 621.

FLEMINGIA VESTITA, Benth.; Franch. Pl. Delav. (1890) 186.

YUNNANENSIS, Franch. *l. c.* 185.

✓ FLUEGGEA CAPILLIPES, Pax, in Engl. Jahrb. xxix. (1900) 427.

FORDIOPHYTON, Stapf, in Ann. Bot. vi. (1892) 314.

CANTONENSE, Stapf, *l. c.*

FABERI, Stapf, *l. c.*

FORDII, Krasser, in Engl. & Prantl, Natürl. Pflanzenf. III. 7. (1893) 175.

FRAXINUS LONGICUSPIS, Sieb. & Zucc.; Palibin, in Act. Hort. Petrop. xviii. 11. (1900) 156.

✓ MALACOPHYLLA, Hemsl. & Hook. Ic. Pl. xxvi. (1899) t. 2598.

PLATYPODA, Oliver, *l. c.* xx. (1890) t. 1929.

RETUSA, var. HENRYANA, Oliver, *l. c.* t. 1930.

FREYCINETIA FORMOSANA, Hemsl. in Kew Bull. (1896) 166.

sp., Makino, in Bot. Mag. Tokyo, x. (1896) no. 113, 222.

FRITILLARIA DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 93.

LOPHOPHORA, Bur. & Franch. in Journ. de Bot. v. (1891) 153.

ROYLEI, Hook.; Franch. *l. c.* xii. (1898) 222.

GALACTIA FORMOSANA, Matsum. in Journ. Coll. Sci. Tokyo, xii. (1898) 424.

✓ GALIUM PARADOXUM, Maxim.; Engl. Jahrb. xxix. (1900) 583.

TRIFLORIFORME, Komarov, in Act. Hort. Petrop. xviii. (1901) 428.

✓ TRIFLORUM, Michx.; Engl. Jahrb. xxix. (1900) 583.

✓ GAULTHERIA FRAGRANTISSIMA, Wall.; Engl. Jahrb. xxix. (1900) 515.

✓ LAXIFLORA, Diels; *l. c.*

✓ NUMMULARIOIDES, D. Don; *l. c.*

- GENTIANA ALGIDA, Pall. ex Engl. Jahrb. xxix. (1900) 536.  
   BELLIDIFOLIA, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 486.  
   BIFLORA, Regel; Kusnezov, in Act. Hort. Petrop. xiii. i. (1893) 62.  
   CHINENSIS, Kusnezov, in Bull. Acad. Petrop. xiii. (1892) 338.  
   CRASSULOIDES, Bur. & Franch. in Journ. de Bot. v. (1891) 104.  
   CRASSULOIDES, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 148.  
   DAHURICA, Fisch.; Act. Hort. Petrop. xv. ii. (1898) 318.  
   DUCLOUXII, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 305.  
   FRIGIDA, Haenke; var., Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1888) 62.  
   GENTILIS, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 491.  
   GRUMII, Kusnezov, in Act. Hort. Petrop. xiii. i. (1893) 63.  
   HAYNALDI, Kanitz, *cf.* Bull. Acad. Pétersb. xxxv. (1894) 352, *et* Pl. Exped. Szech., As. Centr. 39.  
   HEXAPHYLLA, Maxim. in Bull. Acad. Pétersb. xxxv. (1894) 350.  
   JANKÆ, Kanitz, *cf. op. cit.* 352; *et* Pl. Exped. Szechenyi, Asia Centr. 41.  
   KURROO, Royle, var.; Kusnezov, in Bull. Acad. Pétersb. xxxiv. (1892) 508.  
   KUSNEZOWII, Franch. Bull. Soc. Bot. Fr. xliii. (1896) 492.  
   LEUCOMELÆNA, Maxim.; Kusnezov, in Bull. Acad. Pétersb. xxxiv. (1892) 506.  
   MAEULCHANENSIS, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 488.  
   MAXIMOWICZI, Kusnezov, in Bull. Acad. Pétersb. xxxiv. (1892) 505 (*et* xxxv. (1894) 352).  
   MAXIMOWICZII, Kanitz, Pl. Exped. Szechenyi, Asia Centr. 39.  
   MYRIOCLADA, Franch. Bull. Soc. Bot. Fr. xliii. (1896) 487.  
   NAPULIFERA, Franch. *l. c.* 488.  
   ORNATA, Wall.; *l. c.* 493.  
   PHOB, Franch. *l. c.*  
   PRATICOLA, Franch. *l. c.* 489.  
   PRATTII, Kusnezov, in Act. Hort. Petrop. xiii. i. (1893) 63.  
   PSEUDO-AQUATICA, Kusnezov, *l. c.*  
   PURPURATA, Maxim.; Kusnezov, in Bull. Acad. Pétersb. xxxix. (1892) 503.  
   ROSULARIS, Franch. Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 148.  
   SAMOLIFOLIA, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 485.  
   SIPHONANTHA, Maxim.; Kusnezov, in Bull. Acad. Pétersb. xxxiv. (1892) 506.  
   SOULIEI, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 491.  
   SPATHULÆFOLIA, Kusnezov, in Bull. Acad. Pétersb. xxxv. (1894) 351.  
   SZECHENYII, Kanitz, Pl. Exped. Szechenyi, Asia Centr. (1891) 40.  
   TATSIENENSIS, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 489.

- GENTIANA TETRAPHYLLA, Kusnezov, in Bull. Acad. Pétersb. xxxv. (1894) 350.  
 TIZUENSIS, Franch. in Bull. Soc. Bot. Fr. xliii. (1896) 494.  
 TONGOLENSIS, Franch. *l. c.* 490.  
 TRICHOLOBA, Franch. *l. c.*  
 TRICHOTOMA, Kusnezov, in Act. Hort. Petrop. xiii. i. (1893) 60.  
 GERANIUM DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 442.  
 KOREANUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 433.  
 MELANANDRUM, Franch. Pl. Delav. (1889) 112.  
 MOUPINENSE, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 208.  
 NAPULIGERUM, Franch. Pl. Delav. (1889) 115.  
 OCELLATUM, Jacquem.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 444.  
 PALUSTRE, Linn., var., Franch. Pl. Delav. (1889) 109.  
 PLATYPETALUM, Franch. *l. c.* 111.  
 POGONANTHUM, Franch. *l. c.*  
 —ROBERTIANUM, Linn.; Engl. Jahrb. xxix. (1900) 419.  
 SOBOLIFERUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 433.  
 STRIGOSUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 442.  
 UMBELLIFORME, Franch. *l. c.* 443.  
 YUNNANENSE, Franch. Pl. Delav. (1889) 114.  
 GERBERA HENRYI, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 511.  
 TANANTI, Franch. in Journ. de Bot. vii. (1893) 155.  
 —GEUM ELATUM, Wall., var. HUMILE, Royle; Engl. Jahrb. xxix. (1900) 404.  
 —GILBERTIA DENTIGERA, Harms, in Engl. Jahrb. xxix. (1900) 487.  
 PROTEA, Harms, in Engl. & Prantl, Natürl. Pflanzenf. III. 8. (1894) 41.  
 GIRALDIA, Baroni, in Nuov. Giorn. Bot. Ital. n. s. iv. (1897) 431.  
 STAFFII, Baroni, *l. c.*  
 —GIRALDINIA CONDENSATA, Wedd.; Engl. Jahrb. xxix. (1900) 302.  
 VITIFOLIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 80.  
 GLEDITSCHIA DELAVAYI, Franch. Pl. Delav. (1890) 189.  
 OFFICINALIS, Hemsl. in Kew Bull. (1892) 82.  
 GLOCHIDION FORMOSANUM, Hayata, in Bot. Mag. Tokyo, xvii. (1903) 200.  
 LANCEOLATUM, Hayata, *l. c.* 199.  
 GLYCYRRHIZA INFLATA, Batalin, in Act. Hort. Petrop. xi. 16 (1891) 484.  
 GNAPHALIUM BICOLOR, Franch. in Journ. de Bot. x. (1896) 411.  
 CHRYSOCEPHALUM, Franch. *l. c.* 412.  
 CORYMBOSUM, Bur. & Franch. *l. c.* v. (1891) 71.  
 DEDEKENSII, Bur. & Franch. *l. c.* 70.

- GNAPHALIUM DELAVAYI, Franch. in Journ. de Bot. x. (1896) 409.  
     LIKIANGENSE, Franch. *l. c.* 410.  
     NOBILE, Bur. & Franch. *l. c.* v. (1891) 71.  
     PELLUCIDUM, Franch. *l. c.* x. (1896) 411.  
     SERICEO-ALBIDUM, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903)  
         501.  
     SINENSE, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 133.  
     STRACHEYI, Franch. *l. c.*  
     SUBULATUM, Franch. *l. c.* 130.  
     SYLVATICUM, Linn. ; Engl. Jahrb. xxix. (1900) 613.  
     YUNNANENSIS, Franch. Journ. de Bot. x. (1896) 410.  
 GOLDBACHIA LANCIFOLIA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
     408.  
 GORDONIA AXILLARIS, var. ACUMINATA, E. Pritzl, in Engl. Jahrb. xxix.  
     (1900) 473.  
 GULDENSTAEDTIA GIRALDII, Harms, in Engl. Jahrb. xxix. (1900) 413.  
     PAUCIFLORA, Fisch. ex Franch. in Journ. de Bot. iv. (1890) 304.  
     YUNNANENSIS, Franch. Pl. Delav. (1890) 164.  
 GYMNADENIA CRASSINERVIS, Finet, in Rev. Gén. Bot. xiii. (1901) 514.  
     HEMIPILIODES, Finet, *l. c.* 515.  
 GYMNAGATHIS, Stapf, in Ann. Bot. vi. (1892) 315.  
     PEPEROMIIFOLIA, Stapf, *l. c.*  
 GYMNOSPORIA = CELASTRUS.  
     ACUMINATA, Hook. f. & Thoms. ; Engl. Jahrb. xxx. (1902) 472.  
     EMARGINATA, M. Laws. ; Engl. *l. c.* 471.  
     VARIABILIS, Loes. ; Engl. *l. c.* xxix. (1900) 446.  
 GYNOSTEMMA CARDIOSPERMA, Cogn. in Hook. Ic. Pl. xxiii. (1892)  
     t. 2225.  
 GYNURA AURITA, Winkl. in Act. Hort. Petrop. xiv. i. (1895) 151.  
     PINNATIFIDA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 489.  
     SINICA, Diels, in Engl. Jahrb. xxix. (1900) 618.  
 GYPSOPHILA PERFOLIATA, Linn. ; Komarov, in Act. Hort. Petrop. xxii.  
     (1903) 206.  
  
 HABENARIA DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
     Par. sér. 2, x. (1887-88) 86.  
     DELAVAYI, Finet, Rev. Gén. Bot. xiii. (1901) 527.  
     FARGESII, Finet, *l. c.* 528.  
 HARTIA, Dunn, in Hook. Ic. Pl. xxviii. (1902) t. 2727.  
     SINENSIS, Dunn, *l. c.*  
 HEDONA DAVIDI, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899)  
     433.  
 HEDYCHIUM SPICATUM, Buch.-Ham. ; Gagnep. in Bull. Acad. Géog. Bot.  
     xii. (1903) 40.  
 HEDYSARUM SIKKIMENSE, Benth. ; Franch. Pl. Delav. (1890) 171.

- HEDYSARUM TANGUTICUM, B. Fedtsch. in Bot. Centralbl. lxxxiv. (1900) 274.  
 TUBEROSUM, B. Fedtsch. *l. c.*
- HELICIA FORMOSANA, Hemsl. in Ann. Bot. ix. (1895) 156.  
 — GRANDIS, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2631.
- HELLEBORUS CHINENSIS, Maxim. in Act. Hort. Petrop. xi. i. (1890) 27.  
 THIBETANUS, Franch. Pl. Delav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 190.
- HELWINGIA CHINENSIS, Batalin, in Act. Hort. Petrop. xiii. 7. (1893) 97.  
 — HIMALAICA, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 505.
- HEMIBOEA GRACILIS, Franch. in Bull. Soc. Linn. Par. (1899) 124.
- HEMIGRAPHIS SZECHUANICA, Batalin, in Act. Hort. Petrop. xiii. 18. (1894) 384.
- HEMILOPHIA, Franch. Pl. Delav. (1889) 65.  
 PULCHELLA, Franch. *l. c.*
- HEMSLEYA, Cogn.; Hook. Ic. Pl. xix. (1889) t. 1822.  
 CHINENSIS, Cogn. *l. c.*
- HEPTAPLEURUM DELAVAYI, Franch. in Journ. de Bot. x. (1896) 307.  
 FARGESII, Franch. *l. c.* 306.  
 HOI, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 498.  
 MACROPHYLLUM, Dunn, *l. c.* 499.  
 PRODUCTUM, Dunn, *l. c.*
- HERACLEUM ACUMINATUM, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 144.  
 BIVITTATUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 855.  
 CANDICANS, Wall.; Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 144.  
 COREANUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 854.  
 FARGESII, Boissieu, *l. c.* 853.
- HEMSLEYANUM, Diels, Engl. Jahrb. xxix. (1900) 503.  
 RAPULA, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 145.  
 SCABRIDUM, Franch. *l. c.*  
 SOULIEI, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 852.  
 VICINUM, Boissieu, *l. c.* 853.  
 YUNNANENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 143.
- HERMINIUM ALASCHANICUM, Maxim. in Bull. Acad. Pétersb. xxxi. (1887) 105.  
 FALLAX, Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 413.
- HIBISCUS CANCELLATUS, Roxb.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 440.  
 TILIACEUS, Linn., var.; Hochr. in Ann. Conserv. Jard. Bot. Genève, iv. (1900) 64.  
 VENUSTUS, Biume, ex Hochr. *l. c.* 93.
- HIERACIUM SINENSE, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 502.
- HIPTAGE MINOR, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 487.



- HOECKIA, Engl. & Graebn.; Engl. Jahrb. xxix. (1900) 598.
  - ASCHERSONIANA, Engl. & Graebn.; *l. c.*
- HOLBOELLIA ANGUSTIFOLIA, Wall., var.; Diels, in Engl. Jahrb. xxix. (1900) 343.
  - CHINENSIS, Franch.; *l. c.*
  - CORIACEA, Diels, in Engl. *l. c.* 342.
    - CUNEATA, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1817.
  - LATIFOLIA, Wall.; Engl. Jahrb. xxix. (1900) 343.
- HOLOSTEUM SINENSE, Cat. ms. Kew Collect. Henry; ex Diels, in Engl. Jahrb. xxix. (1900) 321.
- HYDRANGEA ALTISSIMA, Wall.; Engl. Jahrb. xxix. (1900) 376.
  - ARBOSTIANA, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 114.
  - GIRALDII, Diels, in Engl. Jahrb. xxix. (1900) 373.
  - HEMSLEYANA, Diels, *l. c.* 376.
    - KAMIENSKII, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 115.
    - LONGIPES, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 227.
  - MANDARINORUM, Diels, in Engl. Jahrb. xxix. (1900) 372.
    - MAXIMOWICZII, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 114.
  - ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 374.
  - XANTHONEURA, Diels, *l. c.* 373.
- HYDROCOTYLE RUBESCENS, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 108.
- HYLOMECON JAPONICUM, Prantl; Engl. Jahrb. xxix. (1900) 353.
  - SUTCHUENSE, Franch.; *l. c.*
    - VERNALE, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 337.
- HYPERICUM ELODEOIDES, Choisy; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 436.
  - GEMINIFLORUM, Hemsl. in Ann. Bot. ix. (1895) 144.
  - LALANDEI, Choisy; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 436.
  - LONGISTYLUM, Oliver, in Hook. Ic. Pl. xvi. (1886) t. 1534.
    - MONANTHEMUM, Hook. f. & Thoms.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 436.
  - NEPAULENSE, Choisy; Engl. Jahrb. xxix. (1900) 476.
    - PETIOLULATUM, Hook. f. & Thoms.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 436.
    - PRATHII, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 303.
    - TRINERVIUM, Hemsl. in Ann. Bot. ix. (1895) 144.
    - YUNNANENSE, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 437.
- IDESIA POLYCARPA, Maxim. var., Diels; Engl. Jahrb. xxix. (1900) 478.

- ILEX AQUIFOLIUM, Linn. ; varr., Loes. in Engl. Bot. Jahrb. xxix. (1900) 435.  
 ARDISIODES, Loes. ; Nov. Act. Nat. Cur. lxxviii. i. (1901) 359.  
 CHAMPIONII, Loes. ; *l. c.* 349.  
 CORALLINA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 452.  
 DELAVAYI, Franch. in Journ. de Bot. xii. (1898) 255.  
 DIPYRENA, Wall., var. LEPTACANTHA, Loes. ; Engl. Jahrb. xxix. (1900) 436.  
 DUBIA, var. HUPEHENSIS, Loes. ; *l. c.*  
 FARGESII, Franch. in Journ. de Bot. xii. (1898) 255.  
 HENRYI, Loes. in Nov. Act. Nat. Cur. lxxviii. i. (1901) 491.  
 INTERMEDIA, Loes. *l. c.* 273.  
 KELUNGENSIS, Loes. *l. c.* 335.  
 LIUKIUENSIS, Loes. *l. c.* 336.  
 MACROCARPA, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1787.  
 MAXIMOWICZIANA, Loes. in Nov. Act. Nat. Cur. lxxviii. i. (1901) 339.  
 METABAPTISTA, Loes. *l. c.* 238.  
 PEDUNCULOSA, Miq. var. ; Loes, in Engl. Jahrb. xxix. (1900) 436.  
 PSEUDO-GODAJAM, Franch. in Journ. de Bot. xii. (1898) 256.  
 PURPUREA, Hassk. var., ex Engl. Jahrb. xxix. (1900) 435.  
 RAMOSA, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1863.  
 SZECHWANENSIS, Loes. in Nov. Act. Nat. Cur. lxxviii. i. (1901) 347.  
 WARBURGII, Loes. *l. c.* 326.  
 YUNNANENSE, Franch. Pl. Delav. (1889) 128.  
 ILLICIUM GRIFFITHII, Hook. f. & Thoms. ; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 383.  
 HENRYI, Diels, in Engl. Jahrb. xxix. (1900) 323.  
 MICRANTHUM, Dunn, in Hook. Ic. Pl. xxviii. (1901) t. 2714.  
 TASHIROI, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 479.  
 VERUM, Hook. f. Bot. Mag. (1888) t. 7005.  
 IMPATIENS ANGUSTIFLORA, Hook. f. ; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 449.  
 BLEPHAROSEPALA, E. Pritz. ; Engl. Jahrb. xxix (1900) 455.  
 CHÆTODONTA, Franch. Pl. Delav. (1889) 155.  
 CORCHORIFOLIA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 448.  
 CORNUCOPIA, Franch. Pl. Delav. (1889) 121.  
 DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 445.  
 DIMORPHOPHYLLA, Franch. *l. c.* 446.  
 DIVARICATA, Franch. *l. c.* 447.  
 DOLICHOCERAS, E. Pritz. in Engl. Jahrb. xxix. (1900) 456.  
 FISSICORNIS, Maxim. in Act. Hort. Petrop. xi. i. (1890) 87.  
 HANCOCKII, C. H. Wright, in Kew Bull. (1896) 21.  
 HENRYI, E. Pritz. in Engl. Jahrb. xxix. (1900) 455.  
 LONGIALATA, E. Pritz. *l. c.* 454.

- IMPATIENS NOTOLOPHA**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 91.  
**ODONTOPETALA**, Maxim. *l. c.* 90.  
**PLATYCERAS**, Maxim. *l. c.* 89.  
**POTANINI**, Maxim. *l. c.* 90.  
**PROCUMBENS**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 447.  
**RADIATA**, Hook. f.; Franch. *l. c.* 449.  
**RECURVICORNIS**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 88.  
**ROSTELLATA**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 209.  
**STENANTHA**, Hook. f.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 449.  
**STENOSEPALA**, E. Pritz.; Engl. Jahrb. xxix. (1900) 453.  
**ULIGINOSA**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 448.  
**VITTATA**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 209.  
**YUNNANENSIS**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 445.  
**INCARVILLEA BERESOVSKII**, Batalin, in Act. Hort. Petrop. xiv. 8. (1895) 181.  
**DELAVAYI**, Bur. & Franch. in Journ. de Bot. v. (1891) 138.  
**GRANDIFLORA**, Bur. & Franch. *l. c.*  
**LUTEA**, Bur. & Franch. *l. c.* 137.  
**VARIABILIS**, Batalin, in Act. Hort. Petrop. xii. 6. (1892) 177.  
**INDIGOFERA CAUDATA**, Dunn, in Gard. Chron. (1902) ii. 210.  
**CHÆTODONTA**, Franch. Pl. Delav. (1889) 155.  
**CINERASCENS**, Franch. *l. c.* 153.  
**DELAVAYI**, Franch. *l. c.* 154.  
**HETERANTHA**, Wall.; Franch. *l. c.* (1889) 154.  
**KIRILOWI**, Maxim. ex Palibin, in Act. Hort. Petrop. xiv. i. (1895) 115.  
**LINIFOLIA**, Retz.; Franch. Pl. Delav. (1889) 153.  
**LIUKIUENSIS**, Makino, in Bot. Mag. Tokyo, x. 115 (1896) 65.  
**MOLLIS**, Franch. Pl. Delav. (1889) 156.  
**PENDULA**, Franch. *l. c.*  
**RETICULATA**, Franch. *l. c.* 153.  
**SCABRIDA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 487.  
**SENSITIVA**, Franch. Pl. Delav. (1889) 157.  
**SERICOPHYLLA**, Franch. *l. c.* 155.  
**SUBULATA**, Vahl; Engl. Jahrb. xxix. (1900) 411.  
**INULA CHRYSANTHA**, Diels, in Engl. Bot. Jahrb. xxix. (1900) 614.  
**CUSPIDATA**, C. B. Clarke; Engl. Jahrb. xxix. (1900) 614.  
**GIRALDII**, Diels; *l. c.* 613.  
**PTEROCAULA**, Franch. in Journ. de Bot. x. (1896) 383.  
**SERICOPHYLLA**, Franch. *l. c.*  
**SERRATA**, Bur. & Franch. *l. c.* v. (1891) 50.

- IPOMŒA TASHIROI, Matsum. in Bot. Mag. Tokyo (1898) 75.  
 — IRIS ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 261.  
 ISOPYRUM AURICULATUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
     376.  
     DELAVAYI, Franch. in Journ. de Bot. xi. (1897) 222.  
     FARGESII, Franch. *l. c.* 220.  
 — HENRYI, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1745.  
     PELTATUM, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
         sér. 2, viii. (1885) 190.  
     SUTCHUENENSE, Franch. in Journ. de Bot. viii. (1894) 274.  
     VAGINATUM, Maxim. Fl. Tangut. (1889) 18.  
 ITEA PARVIFLORA, Hemsl. : Ann. Bot. ix. (1895) 153.  
     YUNNANENSIS, Franch. in Journ. de Bot. x. (1896) 268.  
 ITOA, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2688.  
     ORIENTALIS, Hemsl. *l. c.*
- JASMINUM DISCOLOR, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
     Par. sér. 2, x. (1887-88) 59.  
 — FLORIBUNDUM, Bunge, var., Diels, in Engl. Jahrb. xxix. (1900) 534.  
 — GIRALDII, Diels, *l. c.*  
     POLYANTHUM, Franch. in Rev. Hort. (1891) 207.  
     PRIMULINUM, Hemsl. in Kew Bull. (1895) 109.  
 — JUGLANS SIEBOLDIANA, Maxim. ; Engl. Jahrb. xxix. (1900) 274.  
 JURINEA EDULIS, Franch. in Journ. de Bot. viii. (1894) 337.  
     SOULIEI, Franch. *l. c.*
- JUSSIÆA FAURIEI, H. Lévillé, in Bull. Acad. Géog. Bot. xii.  
     (1903) 18.  
     JAPONICA, H. Lévillé, *l. c.*  
     PARMENTIERI, H. Lévillé, *l. c.*  
     PHILIPPIANA, H. Lévillé, *l. c.*
- KADSURA CHAMPIONI, C. B. Clarke, in Journ. Linn. Soc., Bot. xxv.  
     (1889) 4.  
 KELLOGGIA CHINENSIS, Franch. in Journ. de Bot. vi. (1892) 11.  
 KETELEERIA EVELYNIANA, Masters, Gard. Chron. (1903) i. 194.  
 KOELREUTERIA BIPINNATA, Franch. in Bull. Soc. Bot. xxxiii. (1886)  
     463.  
     — MINOR, Hemsl. in Hook. Ic. Pl. xxvii. t. 2642.  
 — KOLKWITZIA, Graeb. in Engl. Jahrb. xxix. (1900) 593.  
     — AMABILIS, Graeb. *l. c.* 593.  
 KRASCHENINIKOWIA RAPHANORHIZA, Palib. in Act. Hort. Petrop. xvii.  
     i. (1899) 42.  
     SILVATICA, Maxim. ; Komarov, in Act. Hort. Petrop. xxii. (1903)  
     176.

- LACTUCA ATROPURPUREA**, Franch. in Journ. de Bot. ix. (1895) 294.  
**CREPIDIODES**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 244.  
**DIVERSIFOLIA**, Vaniot, *l. c.* 245.  
**ELEGANS**, Franch. in Journ. de Bot. ix. (1895) 262.  
**ERYTHROCARPA**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 31.  
**FABERIA**, Franch. in Journ. de Bot. ix. (1895) 294.  
**GRANDIFLORA**, Franch. *l. c.* 260.  
**HASTATA**, Franch. *l. c.* 261.  
**HEMSLEYI**, Franch. *l. c.* 293.  
**HENRYI**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 512.  
**HIRSUTA**, Franch. in Journ. de Bot. ix. (1895) 258.  
**HUMIFUSA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 512.  
**LIGNEA**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 38.  
**LIKIANGENSIS**, Franch. in Journ. de Bot. ix. (1895) 259.  
**MELANANTHA**, Franch. *l. c.* 291.  
**NAPIFERA**, Franch. *l. c.* 292.  
**OCHROLEUCA**, Franch. *l. c.* 293.  
**POLYPODIIFOLIA**, Franch. *l. c.* 265.  
**PRATTII**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 513.  
**PSEUDOSENECIO**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 320.  
**RADDEANA**, Maxim.; Engl. Jahrb. xxix. (1900) 631.  
**RUBROLUTEA**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 317.  
**SOULIEI**, Franch. in Journ. de Bot. ix. (1895) 257.  
**TALIENSIS**, Franch. *l. c.* 263.  
**TATARINOWII**, Franch. *l. c.* 293.  
**THIBETICA**, Franch. *l. c.*  
**UMBROSA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 513.  
**YUNNANENSIS**, Franch. in Journ. de Bot. ix. (1895) 264.  
**LAGERSTRÆMIA FORDII**, Koehne, in Engl. Pflanzenr., Lythr. 262.  
 — **INTERMEDIA**, Koehne, *l. c.* 260.  
**LAPORTEA BULBIFERA**, Wedd.; Engl. Jahrb. xxix. (1900) 302.  
**EVITTATA**, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887) 80.  
**GIRALDIANA**, E. Pritzl; Engl. Jahrb. xxix. (1900) 301.  
**TERMINALIS**, Wight; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1888) 80.  
**LASIANTHUS FORMOSENSIS**, Matsum. in Bot Mag. Tokyo, xv. (1901) 17.  
**HARTII**, Finet & Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 209.  
**LATHYRUS DIELSIANUS**, Harms, in Engl. Jahrb. xxix. (1900) 417.  
**HUMILIS**, Fisch.; Franch. in Journ. de Bot. iv. (1890) 305.  
**PRATENSIS**, Linn.; Franch. Pl. Delav. (1890) 178.  
**LATOUCHEA**, Finet & Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 212.  
**FOKIENENSIS**, Finet & Franch. *l. c.*  
**LAUNÆA GLABRA**, Franch.; Engl. Jahrb. xxix. (1900) 630.

- LAUNÆA LAMPSANOIDES, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 42.
- LEPTODERMIS DIFFUSA, Batalin, in Act. Hort. Petrop. xiii. 18. (1894) 373.
- POTANINI, Batalin, *l. c.* xiv. II. (1898) 319.
- UMBELLATA, Batalin, *l. c.* xiii. 18. (1894) 374.
- LEPTOPYRUM FUMARIOIDES, Reichb.; Komarov, *l. c.* xxii. (1903) 234.
- LEPYRODICLIS QUADRIDENTATA, Maxim. Fl. Tangut. (1889) 84.
- LESPEDEZA CAPILLIPES, Franch. Pl. Delav. (1890) 165.
- DELAVAYI, Franch. *l. c.*
- DIVERSIFOLIA, Hemsl. in Hook. Ic. Pl. xxvii. (1899) t. 2625.
- ↳ERIOCARPA, Benth.; Engl. Jahrb. xxix. (1900) 414.
- FASCICULIFLORA, Franch. Pl. Delav. (1890) 169.
- GERARDIANA, R. Grah.; Franch. *l. c.*
- HIRTELLA, Franch. *l. c.* (1890) 167.
- LANCEOLATA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 488.
- LATIFOLIA, Dunn, *l. c.*
- TRIGONOCLADA, Franch. Pl. Delav. (1890) 167.
- VARIEGATA, Camb. var.; Franch. *l. c.* (1889) 170.
- ↳VELUTINA, Dunn, in Hook. Ic. Pl. xxvii. (1900) t. 2700.
- YUNNANENSIS, Franch. Pl. Delav. (1890) 166.
- LEVEILLEA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 29.
- CHINENSIS, Vaniot, *l. c.* xiii. (1904) 16.
- MARTINI, Vaniot, *l. c.* xii. (1903) 30.
- LEYCESTERIA SINENSIS, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2633.
- LIGULARIA = SENECIO.
- ↳FARGESII, Franch.; Engl. Jahrb. xxix. (1900) 621.
- JALUENSE, Komarov, in Act. Hort. Petrop. xviii. (1901) 420.
- ↳PRZEWALSKII, Maxim.; Engl. Jahrb. xxix. (1900) 621.
- ↳TENUIPES, Franch. *l. c.*
- ↳VILLIFERA, Franch. *l. c.* 622.
- ↳YESOENSIS, var., Franch. *l. c.*
- LIGUSTICUM ACUMINATUM, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 131.
- ANGELICÆFOLIUM, Franch. *l. c.* 133.
- BRACHIOBUM, Franch. *l. c.* 134.
- DAUCOIDES, Franch. *l. c.* 135.
- DELAVAYI, Franch. *l. c.* 131.
- FERULACEUM, Franch. *l. c.* 135.
- GLAUDESCENS, Franch. *l. c.* 134.
- INVOLUCRATUM, Franch. *l. c.* 136.
- MULTIVITTATUM, Franch. *l. c.* 133.
- PSEUDO-ANGELICA, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 845.
- PTERIDOPHYLLUM, Franch.; Hook. Ic. Pl. xx. (1891) sub t. 1958.

- LIGUSTICUM SINENSE, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1958.  
 TENUIFOLIUM, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 136.  
 TENUISECTUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 843.  
 THOMSONI, C. B. Clarke; Bretsch. Hist. Bot. Disc. China, ii. 976.  
 WALLICHII, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 136.
- LIGUSTRUM DELAVAYANUM, Hariot, in Journ. de Bot. xiv. (1900) 172.  
 ↘ MELLOSUM, Decne.; Engl. Jahrb. xxix. (1900) 532.  
 ↘ MYRIANTHUM, Diels, *l. c.* 533.  
 ↘ PATULUM, Palib. in Act. Hort. Petrop. xviii. ii. (1900) 156.  
 ↘ THIBETICUM, Decne.; Engl. Jahrb. xxix. (1900) 532.  
 ↘ YUNNANENSE, L. Henry, in Rev. Hortie. (1902) 497.
- LILIUM CERNUUM, Komarov, in Act. Hort. Petrop. xx. (1901) 461.  
 CHRISTI, Baroni, in Bull. Soc. Bot. Ital. v. (1896) 213.  
 DAVIDI, Duchartre, Franch. Pl. Dav. (1889) 129.  
 PALIBINIANUM, Yabe, in Bot. Mag. Tokyo, xvii. (1903) 134.  
 ↘ ROSTHORNII, Diels, Engl. Jahrb. xxix. (1900) 243.  
 ↘ SHENSIENSE, Baroni, Bull. Soc. Bot. Ital. v. (1896) 213.  
 ↘ TSINGTANENSE, Gilg, in Engl. Jahrb. xxxiv. (1904) Beibl. 75. 24.
- LIMACIA SAGITTATA, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1749.
- LINARIA THIBETICA, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 11.
- ↘ LINDERA CAUDATA, Diels, in Engl. Jahrb. xxix. (1900) 352.  
 ↘ GLAUCA, Blume, ex Diels, *l. c.* 351.  
 ↘ OBOVATA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 76.  
 ↘ PEDUNCULATA, Diels, in Engl. Jahrb. xxix. (1900) 350.  
 ↘ PUBERULA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 77.  
 ↘ ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 350.  
 ↘ STRYCHNIFOLIA, Villar, var.; Diels, *l. c.* 352.
- LINNÆA = ABELIA.
- ↘ ADENOTRICHIA, Graebn. in Engl. Jahrb. xxix. (1900) 144.  
 ↘ ANGUSTIFOLIA, Bur. & Franch. *l. c.* 135.  
 ↘ ASCHERSONIANA, Graebn. *l. c.* 139 (sphalm. ACHERSONIANA).  
 ↘ BIFLORA, Turcz. *l. c.* 141.  
 ↘ DIELSII, Graebn. *l. c.* 140.  
 ↘ ENGLERIANA, Graebn. *l. c.* 132.  
 ↘ KOEHNEANA, Graebn. *l. c.* 132.  
 ↘ MACROTERA, Graebn. & Buchw. *l. c.* 131.  
 ↘ ONKOCARPA, Graebn. *l. c.* 140.  
 ↘ PARVIFOLIA, Graebn. *l. c.* 129.  
 ↘ RUPESTRIS, A. Br. & Vatke; *l. c.* 138.  
 ↘ SCHUMANNI, Graebn. *l. c.* 130.  
 ↘ SERRATA, Sieb. & Zucc.; Graebn. *l. c.* 133.  
 ↘ TERETICALYX, Graebn. & Buchw. *l. c.* 130.

- ✓ *LINNÆA UMBELLATA*, Graebn. in Engl. Jahrb. xxix. (1900) 143.  
 ✓ *ZANDERI*, Graebn. *l. c.* 142.
- LIPARIS BOOTANENSIS*, Griff.; Finet, in Bull. Soc. Bot. Fr. xlv. (1898) 411.  
*CORNICAULIS*, Makino, in Bot. Mag. Tokyo, x. (1896) no. 114, ii. 56.
- ✓ *LIQUIDAMBAR ROSTHORNII*, Diels, in Engl. Jahrb. xxix. (1900) 380.
- LIRIODENDRON CHINENSE*, Sargent, Trees & Shrubs (1903) 103, t. 52.
- LITHOSPERMUM HANCOCKIANUM*, Oliver, in Hook. Ic. Pl. xxv. (1896) t. 2457.
- LLOYDIA YUNNANENSIS*, Franch. in Journ. de Bot. xii. (1898) 192.
- ✓ *LOBELIA DOLICHOHYRSA*, Diels, in Engl. Jahrb. xxix. (1900) 607.
- ✓ *LONICERA ACUMINATA*, Wall.; Engl. Jahrb. xxix (1900) 594.  
*ADENOPHORA*, Franch. in Journ. de Bot. x. (1896) 311.  
 ✓ *ÆMULANS*, Rehder, Rep. Miss. Bot. Gard. (1903) 47.  
 ✓ *ALSEUOSMOIDES*, Graebn. in Engl. Jahrb. xxix. (1900) 594.  
*BARBINERVIS*, Komarov, in Act. Hort. Petrop. xviii. (1900) 426.  
*CALCARATA*, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2632.  
*CRASSIFOLIA*, Batalin, in Act. Hort. Petrop. xii. (1892) 172.  
*CYANOCARPA*, Franch. in Journ. de Bot. x. (1896) 314.  
*DEFLEXICALYX*, Batalin, in Act. Hort. Petrop. xii. (1892) 173.  
*DELAVAYI*, Franch. in Journ. de Bot. x. (1896) 310.  
*FARGESII*, Franch. *l. c.* 312.  
*FERRUGINEA*, Rehder; Sargent, Trees & Shrubs (1902) t. 22.  
 ✓ *GIRALDII*, Rehder, Rep. Miss. Bot. Gard. (1903) 154.  
 ✓ *GRACILIPES*, Miq.; Engl. Jahrb. xxix. (1900) 596.  
 ✓ *HEMSLEYANA*, Rehder, Rep. Miss. Bot. Gard. (1903) 150.  
*HETEROLOBA*, Batalin, in Act. Hort. Petrop. xii. (1892) 174.  
 ✓ *HILDEBRANDIANA*, Rehder, Rep. Miss. Bot. Gard. (1903) 112.  
*INCONSPICUA*, Batalin, in Act. Hort. Petrop. xiv. (1894) 172.  
*INFUNDIBULUM*, Franch. in Journ. de Bot. x. (1896) 315.  
*KOEHNENANA*, Rehder; Sargent, Trees & Shrubs (1902) t. 21.  
*LIGUSTRINA*, Wall. vars., Franch. in Journ. de Bot. x. (1896) 317.  
 ✓ *LONGA*, Rehder, Rep. Miss. Bot. Gard. (1903) 61.  
*MINUTA*, Batalin, in Act. Hort. Petrop. xii. (1893) 170.  
 ✓ *MUCRONATA*, Rehder, Rep. Miss. Bot. Gard. (1903) 83.  
*ORIENTALIS*, Lam. var.; Franch. in Journ. de Bot. x. (1896) 311.  
*OVALIS*, Batalin, in Act. Hort. Petrop. xiv. (1894) 170.  
 ✓ *PEKINENSIS*, Rehder, Rep. Miss. Bot. Gard. (1903) 95.  
 ✓ *PRÆFLORENS*, Batalin; Rehder, Rep. Miss. Bot. Gard. (1903) 90.  
*RETUSA*, Franch. in Journ. de Bot. x. (1896) 313.  
 ✓ *RHODODENDROIDES*, Graebn. in Engl. Jahrb. xxix. (1900) 595.  
*SACCATA*, Rehder; Sargent, Trees & Shrubs (1902) t. 20.  
*SCABRIDA*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 252.
- ✓ *SERPYLLIFOLIA*, Rehder, Rep. Miss. Bot. Gard. (1903) 58.



- ✓ *LONICERA SETCHUENENSIS*, Rehder, Rep. Miss. Bot. Gard. (1903) 107.  
*SETIFERA*, Franch. in Journ. de Bot. x. (1896) 314.  
*STENOSIPHON*, Franch. *l. c.* 318.  
*STEPHANOCARPA*, Franch. *l. c.* 316.  
 ✓ *SUBÆQUALIS*, Rehder, Rep. Miss. Bot. Gard. (1903) 172.  
*SZECHUANICA*, Batalin, in Act. Hort. Petrop. xiv. (1895) 172.  
*TATSIENENSIS*, Franch. in Journ. de Bot. x. (1896) 313.  
*THIBETICA*, Bur. & Franch. *l. c.* v. (1891) 48.  
*TRICHOPODA*, Franch. *l. c.* x. (1896) 317.  
*TRICHOSANTHA*, Bur. & Franch. *l. c.* v. (1891) 48.  
 ✓ *VEGETA*, Rehder, Rep. Miss. Bot. Gard. (1903) 111.  
*VESICARIA*, Komarov, in Act. Hort. Petrop. xviii. (1900) 427.  
*WEBBIANA*, Sargent, Trees & Shrubs, 137, t. 69.  
*YUNNANENSIS*, Franch. in Journ. de Bot. x. (1896) 310.
- LORANTHUS DELAVAYI*, Engl. in Engl. & Prantl, Natürl. Pflanzenf. Nachtr. (1897) 131.  
*DELAVAYI*, Van Tiegh. in Bull. Soc. Bot. Fr. xli. (1894) 535.
- LOXOSTEMON DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 400.
- LYCHNIS DAVIDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 204.  
*GLANDULOSA*, Maxim. Fl. Tangut. (1889) p. 83.  
*LACINIATA*, Maxim. ex Komarov, in Act. Hort. Petrop. xxii. (1903) 203.  
*YUNNANENSIS*, Baker, in Gard. Chron. i. (1903) 161.
- LYSIMACHIA ALBESCENS*, Franch. in Journ. de Bot. ix. (1895) 460.  
*DELAVAYI*, Franch. *l. c.* 457.  
*DELTOIDEA*, Franch. *l. c.* 461.  
*DRYNARIFOLIA*, Franch. *l. c.* 462.  
*FARGESII*, Franch. *l. c.* 463.  
 ✓ *FORDIANA*, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1983.  
*GLAUCINA*, Franch. in Journ. de Bot. ix. (1895) 457.  
*HEMSLEYANA*, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1980.  
*HEMSLEYI*, Franch. in Journ. de Bot. ix. (1895) 461.  
*HYPERICOIDES*, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 314.  
*INSIGNIS*, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2634.  
*INVOLUCRATA*, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 315.  
*LONGIPES*, Hemsl. *l. c.* 316.  
*MULTANDRA*, Franch. in Journ. de Bot. ix. (1895) 458.  
*NEBELIANA*, Gilg, in Engl. Jahrb. xxxiv. (1904) Beibl. 75. 57.  
*NEMORUM*, Linn., var.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1888) 59.

- LYSIMACHIA NIGROLINEATA**, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 315.  
*OMEIENSIS*, Hemsl. *l. c.* 314.  
**PLATYPETALA**, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 59.  
**PUMILA**, Franch. in Journ. de Bot. ix. (1895) 460.  
**RAMOSA**, Wall. var.; Franch. *l. c.* 463.  
**TRICHOPODA**, Franch. *l. c.* 464.  
✓ **TRIENTALIOIDES**, Hemsl. in Hook. Ic. Pl. xxviii. (1901) t. 2707.  
✓ **UNGUICULATA**, Diels, in Engl. Jahrb. xxix. (1900) 524.  
**VIOLASCENS**, Franch. in Journ. de Bot. ix. (1895) 459.  
**YUNNANENSIS**, Franch. *l. c.* 462.  
**LYSIONOTUS CARNOSA**, Hemsl. in Gard. Chron. (1900) ii. 349.  
**HETEROPHYLLA**, Franch. in Bull. Mus. Hist. Nat. Paris (1899) 249.  
**INVOLUCRATA**, Franch. *l. c.*
- ✓ **MACHILUS PLATYPHYLLA**, Diels, in Engl. Jahrb. xxix. (1900) 348.  
- **THUNBERGII**, Sieb. & Zucc.; Hook. Ic. Pl. xxvi. (1897) t. 2538.  
✓ **MADDENIA HIMALAICA**, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 406.  
✓ **MAESA CASTANEIFOLIA**, Mez, in Engl. Pflanzenr., Myrsin. 44.  
- **DEPAUPERATA**, Diels, in Engl. Jahrb. xxix. (1900) 519.  
- **FORMOSANA**, Mez, in Engl. Pflanzenr., Myrsin. 29.  
- **TENERA**, Mez, *l. c.* 28.  
**MAGNOLIA DELAVAYI**, Franch. Pl. Delav. (1889) 33.  
**HENRYI**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 484.  
✓ **HYPOLEUCA**, Sieb. & Zucc.; Engl. Jahrb. xxix. (1900) 322.  
**PARVIFLORA**, Sieb. & Zucc.; Act. Hort. Petrop. xvii. 1. (1899) 19.  
**MAHONIA = BERBERIS**.  
✓ **EURYBRACTEATA**, Fedde; Engl. Jahrb. xxix. (1900) 338.  
✓ **FORTUNEI**, Fedde; Engl. Jahrb. xxix. (1900) 340.  
✓ **GRACILIPES**, Fedde, in Engl. *l. c.* 338.  
✓ **JAPONICA**, Thunb. var. Fort.; Fedde, *l. c.*  
✓ **POLYODONTA**, Fedde, *l. c.* 339.  
**SUBTRIPLINERVIS**, Franch. in Bull. Mus. Hist. Nat. Paris, i. (1895) 63.  
**TRIFURCA**, Loudon; Engl. Jahrb. xxxi. (1902) 125.  
✓ **MALLOTUS ALBUS**, Muell. Arg.; Engl. Jahrb. xxix. (1900) 429.  
✓ **TENUIFOLIUS**, Pax, in Engl. Jahrb. xxix. (1900) 429.  
✓ **MANGLIETIA FORDIANA**, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1953.  
- **MAPPIA PITTOSPOROIDES**, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1762.  
**MARTINIA**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 31.  
**POLYMORPHA**, Vaniot, *l. c.* 32.  
✓ **MAZUS MACRANTHUS**, Diels, in Engl. Jahrb. xxix. (1900) 566.

- MECONOPSIS BETONICIFOLIA, Franch. Pl. Delav. (1889) 42.  
 CHELIDONIFOLIA, Bur. & Franch. in Journ. de Bot. v. (1891) 19.  
 HENRICI, Bur. & Franch. *l. c.*  
 HORRIDULA, Hook. f. & Thoms., ex Engl. Jahrb. xxix. (1900) 353.  
 INTEGRIFOLIA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 389.  
 PUNICEA, Maxim. Fl. Tangut. (1889) 34.  
 QUINTUPLINERVIA, Regel, Gartenfl. (1876) 291, t. 880.  
 RACEMOSA, Maxim. Fl. Tangut. (1889) 36.
- ✓ MEDICAGO MINIMA, Batalin, ex Engl. Jahrb. xxix. (1900) 411.  
 MEGACARPÆA DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 406.
- MEGADENIA, Maxim. Fl. Tangut. (1889) 76.  
 PYGMÆA, Maxim. *l. c.* 77.
- MELANDRYUM CÆSPITOSUM, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 431.  
 GLANDULOSUM, Williams, *l. c.* 430.  
 KIALENSE, Williams, *l. c.* 432.  
 PLATYPETALUM, Williams, *l. c.* 431.  
 SOULIEI, Williams, *l. c.* 429.
- MELANOSCIADUM, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 803.  
 PIMPINELLOIDEUM, Boissieu, *l. c.* 804.
- ✓ MELIA TOSENDAN, Sieb. & Zucc., ex Engl. Jahrb. xxix. (1900) 426.  
 MELIOSMA CUNEIFOLIA, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 211.  
 ✓ HENRYI, Diels, in Engl. Jahrb. xxix. (1900) 452.  
 YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 465.
- ✓ MELODINUS HEMSLEYANUS, Diels, in Engl. Jahrb. xxix. (1900) 539.  
 MESONA PROCUMBENS, Hemsl. in Ann. Bot. ix. (1895) 155.  
 METAPLEXIS HEMSLEYANA, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1970.
- ✓ MICROGLOSSA SALICIFOLIA, Diels, in Engl. Jahrb. xxix. (1900) 612.  
 MICROMELES ALNIFOLIA, Koehne, ex Palib. in Act. Hort. Petrop. xvii. i. (1899) 75.
- MICROTÆNIA DELAVAYI, Prain, in Bull. Soc. Bot. Fr. xlii. (1895) 424.  
 MOUPINENSIS, Franch. in *l. c.* 426.  
 ✓ PRAINIANA, Diels, in Engl. Jahrb. xxix. (1900) 556.
- ✓ MILLETTIA BOCKII, Harms, in Engl. Jahrb. xxix. (1900) 412.  
 ✓ DIELSIANA, Harms, *l. c.*  
 PACHYCARPA, Benth. Pl. Jungh. 250; Hook. Ic. Pl. xxviii. (1902) t. 2738.  
 PURPUREA, Yatabe, in Bot. Mag. Tokyo, vi. (1892) 379.  
 SCABRICAULIS, Franch. Pl. Delav. (1889) 158.
- MITELLA NUDA, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 419.
- MOHRINGIA LINEARIFOLIA, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 437.

- MOLINIA FAURIEI, Hack. in Bull. Herb. Boiss. sér. 2, iii. (1903) 504.
- ↪ MOSELEYA, Hemsl. in Hook. Ic. Pl. xxvi. (1899) t. 2592.  
 PINNATA, Hemsl. *l. c.*
- MOSLA FORDII, Maxim. in Bull. Acad. Pétersb. xxxi. (1887) 87.  
 GROSSESERRATA, Maxim.; Bot. Mag. Tokyo, ix. (1895) 259.
- MUNRONIA DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 451.  
 ↪ SINICA, Diels, in Engl. Jahrb. xxix. (1900) 425.
- ↪ UNIFOLIATA, Oliver, in Hook. Ic. Pl. xviii. (1887) t. 1705.  
 YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 463.
- MYOSOTIS INTERMEDIA, Link; Palibin, in Act. Hort. Petrop. xviii. (1900) 162.  
 SYLVATICA, Hoffm.; Franch. in Journ. de Bot. iv. (1890) 307.
- ↪ MYRIACTIS NEPALENSIS, Less.; Engl. Jahrb. xxix. (1900) 609.
- MYRICA NANA, Cheval., in Mém. Soc. Nat. Cherb. xxxii. (1901-02) 202.
- MYRICARIA PULCHERRIMA, Batalin, in Act. Hort. Petrop. xi. (1891) 483.
- MYRIPNOIS MAXIMOWICZII, Winkl. *l. c.* xiii. (1893) 12.
- ↪ MYROXYLON CONTROVERSUM, Kuntze; Engl. Jahrb. xxix. (1900) 478.
- ↪ MYRSINE MARGINATA, Mez, in Engl. Pflanzenr., Myrsin. 339.
- NARDOSTACHYS CHINENSIS, Batalin, in Act. Hort. Petrop. xiii. (1894) 376.
- NASTURTIUM BARBARÆFOLIUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 396.  
 HENRYI, Oliver, in Hook. Ic. Pl. xviii. (1887) t. 1719.
- NATSIATUM SINENSE, Oliver, *l. c.* xix. (1889) t. 1900.
- NAUCLEA SINENSIS, Oliver, *l. c.* xx. (1891) t. 1956.
- NEILLIA AFFINIS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1892) 304.  
 GRACILIS, Franch. Pl. Delav. (1890) 202.  
 LONGIRACEMOSA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1892) 304.
- RUBIFLORA, D. Don; Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 217.
- THIBETICA, Bur. & Franch. Journ. de Bot. v. (1891) 45.
- THYRSIFLORA, D. Don; Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 217.
- ↪ NOTHOPANAX BOCKII, Harms, in Engl. Jahrb. xxix. (1900) 488.  
 ↪ DAVIDI, Franch. *l. c.*  
 ↪ ROSTHORNII, Harms, *l. c.* 487.
- NOTOPTERYGIUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 838.  
 FORBESII, Boissieu, *l. c.* 840.  
 FRANCHETI, Boissieu, *l. c.* 839.
- ↪ NYSSA SINENSIS, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1964.
- ŒNANTHE HOOKERI, C. B. Clarke; Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 130.  
 LINEARIS, Wall.; Franch. *l. c.*

- CENANTHE RIVULARIS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 496.  
 ✓ **ROSTHORNII**, Diels, in Engl. Jahrb. xxix. (1900) 498.  
     **SINENSIS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 496.
- OMPHALODES MOUPINENSIS**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 65.
- OMPHALOGRAMMA**, Franch. in Bull. Soc. Bot. Fr. xlv. (1898) 178.  
     **DELAVAYI**, Franch. *l. c.* 179.  
     **SOULIEI**, Franch. *l. c.* 180.  
     **VINCÆFLORA**, Franch. *l. c.*
- ONOSMA EXSERTUM**, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2639.  
     **PANICULATA**, Bur. & Franch. Journ. de Bot. v. (1891) 104.  
 ✓ **SINICA**, Diels, in Engl. Jahrb. xxix. (1900) 547.
- ✓ **OPHIPOGON BOCKIANUS**, Diels, *l. c.* 254.  
     ✓ **JAPONICUS**, Ker-Gawl., var.; Diels, *l. c.*  
     **SPICATUS**, Ker-Gawl., var.; Palib. in Act. Hort. Petrop. xix. (1901) 106.
- OPHIORRHIZA INFLATA**, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 488.  
     **KUROIWAI**, Makino, in Bot. Mag. Tokyo, x. (1896) 68.  
     **TASHIROI**, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 489.
- ORCHIS GENICULATA**, Finet, in Rev. Gén. Bot. xiii. (1901) 505.
- OREOCHARIS FOKKIENENSIS**, Franch. in Bull. Mus. Hist. Nat. Par. v. (1899) 249.  
     **FOKKIENENSIS**, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 212.  
     **HENRYANA**, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1944.  
     **TUBICELLA**, Franch. in Bull. Mus. Hist. Nat. Par. v. (1899) 249.
- OREOPANAX CHINENSIS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 496.
- OREORCHIS FARGESII**, Finet, in Bull. Soc. Bot. Fr. xliii. (1896) 698.
- ✓ **ORIXA JAPONICA**, Thunb.; Engl. Jahrb. xxix. (1900) 423.
- ORMOSIA FORDIANA**, Oliver, in Hook. Ic. Pl. xxv. (1895) t. 2422.  
     **HENRYI**, Prain, in Journ. As. Soc. Beng. lxix. (1900) 180.  
     **STRIATA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 492.  
     **YUNNANENSIS**, Prain, in Journ. As. Soc. Beng. lxix. (1900) 183.
- ORNITHOCHILUS DELAVAYI**, Finet, in Bull. Soc. Bot. Fr. xliii. (1896) 496.
- ✓ **OSMANTHUS ARMATUS**, Diels, in Engl. Jahrb. xxix. (1900) 532.  
     **BRACTEATUS**, Matsum. in Bot. Mag. Tokyo, xii. (1898) 14.  
     **COOPERI**, Hemsl. in Kew Bull. (1896) 18.
- OUROUPARIA = NAUCLEA.**  
     ✓ **SINENSIS**, Diels, in Engl. Jahrb. xxix. (1900) 580.
- ✓ **OXALIS JAPONICA**, Franch. & Sav. ex Engl. Jahrb. xxix. (1900) 419.
- OXYGRAPHIS CYMBALARIA**, Pursh; Komarov, in Act. Hort. Petrop. xxii. (1903) 291.  
     **DELAVAYI**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 374.

- ✓ OXYGRAPHIS GLACIALIS, Bunge, ex Engl. Jahrb. xxix. (1900) 384.  
 ✓ OXYRIA SINENSIS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1892) 317.  
 OXYTROPIS BEKETOWII, Krassn. in Script. Hort. Petrop. ii. 1. (1887-8)  
 15.  
 DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 374.  
 FALCATA, Bunge, in Mém. Acad. Pétersb. sér. 7, xxii. (1874) 38.  
 KANSUENSIS, Bunge, *l. c.* 38.  
 MELANOCALYX, Bunge, *l. c.* 8.  
 OCHROCEPHALA, Bunge, *l. c.* 57.  
 STROBILACEA, Bunge, *l. c.* 103.  
 SYLINCHANENSIS, Franch. in Journ. de Bot. iv. (1890) 304.  
 TRICHOPHORA, Franch. *l. c.* 303.  
 YUNNANENSIS, Franch. Pl. Delav. (1890) 163.
- PACHYSANDRA AXILLARIS, Franch. Pl. Delav. (1889) 135.  
 ✓ PÆONIA ANOMALA, Linn., var. Pall.; Engl. Jahrb. xxix. (1900) 324.  
 DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 382.  
 LUTEA, Franch. *l. c.*  
 PALIURUS HIRSUTUS, Hemsl. in Kew Bull. (1894) 388.  
 ORIENTALIS, Hemsl. *l. c.* 387.  
 PANAX DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, viii. (1885) 248.  
 DELAVAYI, Franch. in Journ. de Bot. x. (1896) 305.  
 PANCOVIA DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii (1886) 461.  
 PANDANUS, sp., Matsum. in Bot. Mag. Tokyo, xi. (1897) 69.  
 PARADISIA MINOR, C. H. Wright, in Kew Bull. (1895) 118.  
 ✓ PARADOMBEYA, Stapf, in Hook. Ic. Pl. xxviii. (1902) t. 2743.  
 SINENSIS, Stapf, *l. c.* t. 2743 B.
- ✓ PARIS BOCKIANA, Diels, in Engl. Jahrb. xxix. (1900) 253.  
 CHRISTII, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 255.  
 FRANCHETIANA, H. Lévillé, *l. c.* 256.  
 ✓ HENRYI, Diels, in Engl. Jahrb. xxix. (1900) 252.  
 MERCIERI, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 255.  
 THIBETICA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, x. (1887-8) 184.
- ✓ PARNASSIA AMENA, Diels, in Engl. Jahrb. xxix. (1900) 370.  
 APPENDICULATA, Batalin, in Bretschn. Hist. Bot. Disc. China, ii.  
 1028.  
 CHINENSIS, Franch. in Bull. Soc. Bot. Fr. xlv. (1897) 252.  
 CRASSIFOLIA, Franch. *l. c.* 253.  
 DAVIDII, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, viii. (1885) 237.  
 DELAVAYI, Franch. in Journ. de Bot. x. (1896) 267.  
 ✓ FABERI, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1778.  
 LAXMANNI, Pall.; Franch. Bull. Soc. Bot. Fr. xlv. (1897) 256.

- PARNASSIA LUTEA**, Batalin, in Act. Hort. Petrop. xiv. 11. (1898) 320.  
**MONOCHORIFOLIA**, Franch. in Bull. Soc. Bot. Fr. xliv. (1897) 260.  
**NOEMIÆ**, Franch. in Journ. de Bot. x. (1896) 281.  
 ✓ **PERCILIATA**, Diels, in Engl. Jahrb. xxix. (1900) 369.  
**PUSILLA**, Hook. f. & Thoms.; Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 237.  
**SETCHUENENSIS**, Franch. in Bull. Soc. Bot. Fr. xliv. (1897) 254.  
**VIRIDIFLORA**, Batalin, in Act. Hort. Petrop. xii. 6. (1892) 168.  
**WIGHTIANA**, Wall., var.; Franch. in Engl. Jahrb. xxix. (1900) 368.  
**YUNNANENSIS**, Franch. in Journ. de Bot. x. (1896) 266.
- PAROCHÆTUS COMMUNIS**, Hamilt.; Franch. Pl. Delav. (1889) 153.
- PARRYA BEKETOWI**, Krassn. in Script. Hort. Petrop. ii. 1. (1887-8) 12.  
**PROLIFERA**, Maxim. Fl. Tangut. (1889) 56.  
**SILIUOSA**, Krassn. in Script. Hort. Petrop. ii. 1. (1887-8) 12.  
**VILLOSA**, Maxim. Fl. Tangut. (1889) 55.
- PARTHENOCISSUS HENRYANA**, Hemsl., var.; Diels, in Engl. Jahrb. xxix. (1900) 464.  
**HIMALAYANA**, Gilg, *l. c.* 463.  
**SINENSIS**, Diels & Gilg, *l. c.*  
**TRICUSPIDATA**, Planch.; *l. c.* 464.
- PARVATIA CHINENSIS**, Franch. in Journ. de Bot. viii. (1894) 281.  
 ✓ **DECORA**, Dunn, in Hook. Ic. Pl. xxviii. (1901) t. 2712.
- PASSIFLORA CUPIFORMIS**, Mast. in Hook. Ic. Pl. xviii. (1888) t. 1768.  
**FRANCHETIANA**, Hemsl. *l. c.* xxvii. (1889) t. 2623.  
**HENRYI**, Hemsl. *l. c.*  
**SEGUINI**, H. Léveillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 174.
- ✓ **PATRINIA DIELSII**, Graebn. in Engl. Jahrb. xxix. (1900) 597.  
**MONANDRA**, var.; Batalin, in Act. Hort. Petrop. xiii. 18. (1894) 377.
- PAULOWNIA FARGESII**, Franch. in Bull. Mus. Hist. Nat. Par. ii. (1896) 280.
- PEDICULARIS BATANGENSIS**, Bur. & Franch. in Journ. de Bot. v. (1891) 106.  
 ✓ **BICOLOR**, Diels, in Engl. Jahrb. xxix. (1900) 570.  
**BIETII**, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 34.  
 ✓ **BIONDIANA**, Diels, in Engl. Jahrb. xxix. (1900) 571.  
**BIROSTRIS**, Bur. & Franch. in Journ. de Bot. v. (1891) 107.  
**BREVILABRIS**, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 33.  
**CINERASCENS**, Franch. *l. c.* 30.  
**CYATHOPHYLLA**, Franch. *l. c.* 25.  
**DAVIDI**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 67.  
**DECORA**, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 28.

- PEDICULARIS ELEGANS, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 32.  
 FARGESII, Franch. *l. c.* 26.  
 FASTIGIATA, Franch. *l. c.* 25.  
 FLACCIDA, Prain, in Kew Bull. (1893) 157.  
 FLORIBUNDA, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 31.  
 GONIANTHA, Bur. & Franch. in Journ. de Bot. v. (1891) 128.  
 HEMSLEYANA, Prain, in Hook. Ic. Pl. xxiii. (1892) t. 2210.  
 IMPERIALIS, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 34.  
 KIALENSIS, Franch. *l. c.* 22.  
 LAXIFLORA, Franch. *l. c.* 27.  
 LONGIFLORA, Rudolph; Franch. in Journ. de Bot. iv. (1890) 317.  
 LYRATA, Prain, var.; Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 33.  
 MACROSIPHON, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
 Par. sér. 2, x. (1887-8) 60.  
 MAXIMOWICZII, Krassn. in Script. Hort. Petrop. ii. i. (1887-8) 18.  
 MICROPHYTON, Bur. & Franch. in Journ. de Bot. v. (1891) 107.  
 MOUPINENSIS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
 Par. sér. 2, x. (1887-8) 67.  
 MUSSOTI, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 24.  
 NASTURTHIFOLIA, Franch. *l. c.* 28.  
 PHACELLÆFOLIA, Franch. *l. c.* 27.  
 PRINCEPS, Franch. in Journ. de Bot. v. (1891) 129.  
 PROVOTI, Franch. *l. c.* iv. (1890) 318.  
 RAMALANA, Britt. ex Gill, River of Golden Sand, ii. 426.  
 RHYNCHODONTA, Franch. *l. c.* v. (1891) 108.  
 SOULIEI, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 23.  
 STENANTHA, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 149.  
 STENOCORYS, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 32.  
 TATSIENENSIS, Franch. in Journ. de Bot. v. (1891) 108.  
 TIBETICA, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 24.  
 TONGOLENSIS, Franch. *l. c.* 29.  
 VERONICIFOLIA, Franch. *l. c.* 30.  
 PENTAPANAX HENRYI, Harms, in Engl. Jahrb. xxiii. (1896) 21.  
 VERTICILLATUM, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903)  
 498.  
 YUNNANENSE, Franch. in Journ. de Bot. x. (1896) 305.  
 PERISTYLIS CÆLOCERAS, Finet, in Rev. Gén. Bot. xiii. (1901) 519.  
 FORCEPS, Finet, *l. c.* 521.  
 MONANTHUS, Finet, *l. c.* 523.  
 TETRALOBUS, Finet, *l. c.* 524.  
 PERROTTETIA RACEMOSA, Oliver, in Engl. Jahrb. xxiv. (1897) 200.  
 PERTYA BODINIERI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 116.  
 SINENSIS, Oliver, in Hook. Ic. Pl. xxiii. (1892) t. 2214.  
 PETROCOSMEA GRANDIFLORA, Hemsl. in Hook. Ic. Pl. xxv. (1895)  
 t. 2410; et in Kew Bull. (1895) 115.



- PETROCOSMEA IODIODES, Hemsl. in Hook. Ic. Pl. xxvi. (1899) t. 2599.  
 MINOR, Hemsl. *l. c.* t. 2600.
- PEUCEDANUM CARTILAGINEO-MARGINATUM, Makino, in Bull. Herb. Boiss. sér. 2, iii. (1903) 958.  
 DELAVAYI, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 143.  
 ELEGANS, Komarov, in Act. Hort. Petrop. xviii. (1901) 430.  
 ✓ GIRALDII, Diels, in Engl. Jahrb. xxix. (1900) 503.  
 GRAVEOLENS, Benth.; Franch. in Bull. Soc. Phil. Par. sér. 2, vi. (1894) 141.  
 HETEROPHYLLUM, Franch. *l. c.*  
 MACILENTUM, Franch. *l. c.* 142.  
 MEDICUM, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 496.  
 PODAGRARIA, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 957.  
 PRÆRUPTORUM, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 497.  
 REPIANS, Diels, in Engl. Jahrb. xxix. (1900) 502.  
 TORILIFOLIUM, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 852.
- PHASEOLUS ACONITIFOLIUS, Jacq.; Franch. Pl. Delav. (1890) 183.  
 GRANDIS, Dalz. & Gibs.; Franch. *l. c.*  
 MUNGO, Linn.; Franch. *l. c.* 182.  
 RADIATUS, Linn.; Franch. *l. c.*  
 ✓ RICCIARDIANUS, Ten.; Engl. Jahrb. xxix. (1900) 418.
- ✓ PHELLODENDRON JAPONICUM, Maxim.; Engl. Jahrb. xxix. (1900) 424.
- PHILADELPHUS INCANUS, Koehne, in Gartenfl. xlv. (1896) 562.  
 SCHRENKII, Rupr.; Komarov, in Act. Hort. Petrop. xxii. (1903) 429.  
 SERICANTHUS, Koehne, var.; Koehne, in Gartenfl. xlv. (1896) 561.
- ✓ PHLOMIS MEDICINALIS, Diels, in Engl. Jahrb. xxix. (1900) 554.  
 SETIFERA, Bur. & Franch. in Journ. de Bot. v. (1891) 149.  
 ✓ STENOCALEYX, Diels, in Engl. Jahrb. xxix. (1900) 555.  
 TATSIENENSIS, Bur. & Franch. in Journ. de Bot. v. (1891) 149.
- PHOTINIA DEFLEXA, Hemsl. in Ann. Bot. ix. (1895) 153.  
 WRIGHTIANA, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 486.
- PHITHEIROSPERMUM TENUISECTUM, Bur. & Franch. in Journ. de Bot. v. (1891) 129.
- PHYLLODESMIS, Van Tiegh. in Bull. Soc. Bot. Fr. xlii. (1895) 255.  
 CORIACEA, Van Tiegh. *l. c.* 256.  
 DELAVAYI, Van Tiegh. *l. c.* 255.  
 PAUCIFOLIA, Van Tiegh. *l. c.* 256.
- PHYSALIS FRANCHETI, Mast. in Gard. Chron. (1894) ii. 434, 441.
- ✓ PHYSOCARPUS ANOMALA, Batalin; Engl. Jahrb. xxix. (1900) 382.
- PHYTOLACCA POLYANDRA, Batalin, in Act. Hort. Petrop. xiii. 7. (1893) 99.
- PICEA NEOVEITCHII, Mast. in Gard. Chron. i. (1903) 116.  
 WILSONI, Mast. *l. c.* 133.
- PICRIS DIVARICATA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 28.  
 ✓ HIERACIOIDES, Linn., var., Thunb.; Engl. Jahrb. xxix. (1900) 630.

- PIERIS BODINIERI, H. Lèveillé, in Bull. Acad. Géog. Bot. xii. (1903) 253.  
 DIVARICATA, H. Lèveillé, *l. c.* 252.  
 DUCLOUXII, H. Lèveillé, *l. c.* 253.  
 GAGNEPAINIANA, H. Lèveillé, *l. c.* 251.  
 KONYANGENSIS, H. Lèveillé, *l. c.* 253.  
 MARTINI, H. Lèveillé, *l. c.* 252.  
 REPENS, H. Lèveillé, *l. c.*
- PILEA FASCIATA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 81.  
 ↙ TRINERVIA, Wight; Engl. Jahrb. xxix. (1900) 302.
- PIMPINELLA, Diels=CARUM.  
 ↗ ARGUTA, Diels, Engl. Jahrb. xxix. (1900) 496.  
 DUNNII, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 841.  
 FARGESII, Boissieu, *l. c.* ii. (1902) 808.  
 FAURIEI, Boissieu, *l. c.* iii. (1903) 955.  
 ↙ FILICINA, Franch.; Engl. Jahrb. xxix. (1900) 494.  
 HELOSCIADOIDEA, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 809.  
 ↙ HENRYI, Diels, in Engl. Jahrb. xxix. (1900) 495.  
 NIKOENSIS, Yabe, in Bot. Mag. Tokyo, xvii. (1903) 106.  
 ↙ RHOMBOIDEA, Diels, in Engl. Jahrb. xxix. (1900) 496.  
 ↙ ROSTHORNII, Diels, *l. c.* 495.  
 SILAIFOLIA, Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 809.  
 SOULIEI, Boissieu, *l. c.* 810.  
 SUTCHUENSIS, Boissieu, *l. c.* 808.  
 ↙ TANAKÆ, Franch. & Sav.; Engl. Jahrb. xxix. (1900) 494.  
 ↙ TRITERNATA, Diels; *l. c.* 496.  
 ↙ TRICHOMANIFOLIA, Franch.; *l. c.* 494.
- ↙ PINGUICULA VULGARIS, Linn.; Engl. Jahrb. xxix. (1900) 573.  
 PINUS FUNEBRIS, Komarov, in Act. Hort. Petrop. xx. (1901) 177.  
 PIPER CHANDOCANUM, C. DC. in Ann. Conserv. Jard. Bot. Genève, ii. (1898) 269.  
 FERRIEI, C. DC. *l. c.* 268.  
 HARMANDI, C. DC. *l. c.*  
 HENRYI, C. DC. *l. c.* 271.  
 LAOSANUM, C. DC. *l. c.* 269.
- PIPTANTHUS NEPALENSIS, D. Don; Franch. Pl. Delav. (1889) 150.  
 TOMENTOSUS, Franch. Pl. Delav. (1889) 150.
- PIRUS, *vide* PYRUS.
- PISTACIA WEINMANNIFOLIA, J. Poisson, ex Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 467.
- ↙ PITTOSPORUM GLABRATUM, Lindl. var., E. Pritz.; Engl. Jahrb. xxix. (1900) 378.  
 HETEROPHYLLUM, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 415.

- ✓ *PITTIOSPORUM NILGHIRENSE*, Wight & Arn. ; Engl. Jahrb. xxix. (1900) 378.  
     ✦ *TRUNCATUM*, E. Pritz. ; Engl. Jahrb. xxix. (1900) 378.  
     *YUNNANENSIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 415.  
*PLAGIOSPERMUM SINENSE*, Oliver, in Hook. Ic. Pl. xvi. (1886) t. 1526.  
 ✓ *PLATANThERA MANUBRIATA*, Kränzl. in Engl. Jahrb. xxix. (1900) 265.  
     ✦ *SEtCHUENICA*, Kränzl. *l. c.*  
*PLECTRANTHUS CALCARATUS*, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2671.  
     ✓ *DICHROMOPHYLLUS*, Diels, in Engl. Jahrb. xxix. (1900) 562.  
     ✦ *LEPTOBOTRYS*, Diels, *l. c.* 561.  
     ✦ *ROSTHORNII*, Diels, *l. c.* 562.  
*PLEIONE BULBOCODIODES*, Rolfe, in Orch. Rev. xi. (1903) 291.  
     *DELAVAYI*, Rolfe, *l. c.*  
     *GRANDIFLORA*, Rolfe, *l. c.*  
     *POGONIOIDES*, Rolfe, *l. c.*  
     *YUNNANENSIS*, Rolfe, *l. c.* 292.  
*PLEUROGYNE CARINTHIACA*, G. Don, var. ; Franch. in Bull. Soc. Bot. Fr. xlv. (1899) 309.  
*PLEUROSPERMUM DAVIDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 247.  
     *DECURRENS*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 138.  
     *FÆTENS*, Franch. *l. c.* 140.  
     *FRANCHETIANUM*, Hemsl. in Hook. Ic. Pl. xxiii. (1892) t. 2244 ; & Journ. Linn. Soc., Bot. xxix. (1893) 307.  
     ✦ *GIRALDII*, Diels, in Engl. Jahrb. xxix. (1900) 493.  
     *GOVANIANUM*, Benth., var. ; Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 138.  
     ✦ *MEOIDES*, Diels, in Engl. Jahrb. xxix. (1900) 493.  
     *NANUM*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 140.  
     *PULSZKYI*, Kanitz, Pl. Exped. Szech., Asia Centr. 25.  
     *SZECHENYI*, Kanitz, *l. c.*  
     *WRIGHTIANUM*, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 847.  
     *YUNNANNENSE*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 137.  
*PODOCHILUS CORNUTUS*, Schlechter, in Mém. Herb. Boiss. i. (1900) 34.  
*PODOON DELAVAYI*, Baill. in Bull. Soc. Linn. Par. (1887) 681.  
     (= *DOBINEA DELAVAYI*.)  
*PODOONACEÆ*, H. Baill. ex Franch. Pl. Delav. (1889) 145.  
*PODOPHYLLUM DELAVAYI*, Franch. in Bull. Mus. Hist. Nat. Par. (1895) 63.  
 ✓ *POGONIA LANCEOLATA*, Kränzl. in Engl. Jahrb. xxix. (1900) 267.  
     ✦ *PLEIONOIDES*, Kränzl. *l. c.*  
*POGOSTEMON FORMOSANUM*, Oliver, in Hook. Ic. Pl. xxiii. (1896) t. 2440.

- POGOSTEMON IANTHINA**, Kanitz, Pl. Exped. Szech., Asia Centr. 46.  
**POLIOTHYRSIS**, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1885.  
     **SINENSIS**, Oliver, *l. c.*  
**POLYGALA CROTALARIOIDES**, Buch.-Ham.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 416.  
     **FLORIBUNDA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 485.  
     **FORBESII**, Chod. in Bull. Herb. Boiss. iv. (1896) 234.  
     **GLOBULIFERA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 483.  
     **LATOCHEI**, Franch. & Finet, in Bull. Soc. Bot. Fr. xlvi. (1899) 206.  
     **PERSICARIOIDES**, DC.; Franch. *l. c.* xxxiii. (1886) 416.  
     **SAXICOLA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 486.  
     **SIBIRICA**, DC.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 417.  
**POLYGONATUM BODINIERI**, H. Lévêillé, in Bull. Soc. Géog. Bot. xii. (1903) 262.  
     **ENSIFOLIUM**, H. Lévêillé, *l. c.* 261.  
     **FALCATUM**, A. Gray; Engl. Jahrb. xxix. (1900) 248.  
     **HENRYI**, Diels, *l. c.* 247.  
     **HUMILE**, Fisch. in Act. Hort. Petrop. xix. (1901) 110.  
     **INFLATUM**, Komarov, *l. c.* xviii. (1900) 442.  
     **MARTINI**, H. Lévêillé, in Bull. Acad. Géog. Bot. xii. (1903) 262.  
**POLYGONUM ACAULE**, Franch. Bull. Mus. Hist. Nat. Par. (1895) 213.  
     **ALPINUM**, All. var., Dammer; Diels, in Engl. Jahrb. xxix. (1900) 314.  
     **AMPHIBIUM**, Linn., varr.; H. Lévêillé, Bull. Acad. Géog. Bot. xi. (1902) 340.  
     **BODINIERI**, H. Lévêillé & Vaniot, *l. c.* 343.  
     **LABORDEI**, H. Lévêillé & Vaniot, *l. c.* 344.  
     **MARTINI**, H. Lévêillé & Vaniot, *l. c.* 340.  
     **MYOSURUS**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88), 73.  
     **NEPALENSE**, Meissn.; Bretsch. Hist. Bot. Disc. China, ii. 1034.  
     **PANDURIFORME**, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 343.  
     **PAUCIFLORUM**, Maxim. in Act. Hort. Petrop. xviii. (1900) 181.  
     **SAGITTIFOLIUM**, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 342.  
     **SINUATUM**, var., Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 74.  
     **SPAETHII**, Dammer, in Notizbl. Bot. Gard. Berl. ii. (1899) 378.  
     **SUFFULTUM**, Maxim.; Franch. in Journ. de Bot. iv. (1890) 318.  
     **UROPHYLLUM**, Bur. & Franch. *l. c.* v. (1891) 150.  
**POPULUS PEKINENSIS**, L. Henry, in Rev. Hort. (1903) 355.  
**PORTULACA VILLOSA**, var.; Matsum. in Bot. Mag. Tokyo, xii. (1898) no. 136, 53.  
**POTENTILLA ALBIFLORA**, Wall.; Franch. Pl. Delav. (1890) 210.

- POTENTILLA AMBIGUA, Cambess. ; Franch. Pl. Delav. (1890) 210.  
 ARTICULATA, Franch. *l. c.*  
 CORIANDRIFOLIA, D. Don, var. ; Franch. *l. c.* 214.  
 DAVIDI, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii.  
 (1885) 222.  
 DELAVAYI, Franch. Pl. Delav. (1890) 215.  
 ✓ ERIOCARPA, Wall. ; Engl. Jahrb. xxix. (1900) 402.  
 FULGENS, Wall. ; Franch. Pl. Delav. (1890) 211.  
 GRIFFITHII, var., Franch. *l. c.* 213.  
 HOLOLEUCA, Boiss. ; Franch. *l. c.* 214.  
 LESCHENAULTIANA, Ser. varr. ; Franch. *l. c.* 212.  
 ✓ LEUCONOTA, D. Don ; Engl. Jahrb. xxix. (1900) 403.  
 MOUPINENSIS, Franch. Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii.  
 (1885) 222.  
 MULTIFIDA, Linn., var. ; Franch. Pl. Delav. (1890) 215.  
 PEDUNCULARIS, D. Don, var. ; Franch. *l. c.* 213.  
 POTERIOIDES, Franch. *l. c.* 212.  
 PURPUREA, Royle ; Franch. *l. c.* 209.  
 SIBBALDI, Hall. f. ; Franch. *l. c.* 210.  
 POTHOS WARBURGII, Engl. Jahrb. xxv. (1898) 2.  
 POUPARTIA FORDII, Hemsl. in Hook. Ic. Pl. xxvi. (1898) t. 2557.  
 POURTHIÆA ARGUTA, Decne., var., Hook. f. & Thoms. ; Engl. Jahrb. xxix.  
 (1900) 388.  
 PARVIFOLIA, E. Pritz, *l. c.*  
 VARIABILIS, Hemsl. ex Palibin, in Act. Hort. Petrop. xvii. i. (1899)  
 76.  
 ✓ PRENANTHES BRUNONIANA, Wall. ; Engl. Jahrb. xxix. (1900) 633.  
 GLANDULOSA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 514.  
 HENRYI, Dunn, *l. c.*  
 MACROPHYLLA, Franch. in Journ. de Bot. iv. (1890) 306.  
 ✓ VITIFOLIA, Diels, in Engl. Jahrb. xxix. (1900) 634.  
 PRIMULA ARGUTIDENS, Franch. in Journ. de Bot. ix. (1895) 451.  
 BARBICALYX, C. H. Wright, in Kew Bull. (1896) 241.  
 BREVISCAPA, Franch. in Bull. Mus. Hist. Nat. Par. (1895) 65.  
 CHARTACEA, Franch. *l. c.* 64.  
 CINERASCENS, Franch. in Journ. de Bot. ix. (1895) 488.  
 COCKBURNIANA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893)  
 313.  
 DAVIDI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 66.  
 DIANTHA, Franch. in Journ. de Bot. v. (1891) 97.  
 FARGESII, Franch. *l. c.* ix. (1895) 452.  
 GEMMIFERA, Batalin, in Act. Hort. Petrop. xi. 16. (1891) 491.  
 GILLII, Britt. ex Gill, River of Golden Sand, ii. 426.  
 HEUCHERIFOLIA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 65.  
 KIALENSIS, Franch. in Journ. de Bot. ix. (1895) 450.

PRIMULA LINEARILOBA, Winkl. ex Bretsch. Hist. Bot. Disc. China, ii. 1033.

LOCZII, Kanitz, Pl. Exped. Szechenyi, Asia Centr. 36.

MOUPINENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 67.

NEUROCALYX, Franch. in Journ. de Bot. ix. (1895) 449.

NIVALIS, Pall., var. ; Franch. *l. c.*

NUTANTIFLORA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 313.

OREODOXA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 66.

✓PAXIANA, Gilg, in Engl. Jahrb. xxxiv. (1904) Beibl. 75. p. 56.

PETIOLARIS, var., Franch. in Journ. de Bot. ix. (1895) 449.

POLYNEURA, Franch. *l. c.* 448.

PRATTII, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 314.

PYCNOLOBA, Bur. & Franch. in Journ. de Bot. v. (1891) 99.

RAMALANA, Bretsch. Hist. Bot. Disc. China, ii. 735.

✓ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 520.

SAXATILIS, Komarov, in Act. Hort. Petrop. xviii. (1901) 429.

SERTULUM, Franch. in Journ. de Bot. ix. (1895) 451.

SINUATA, Franch. in Bull. Mus. Hist. Nat. Par. (1895) no. 2, 65.

SOULIEI, Franch. in Journ. de Bot. ix. (1895) 450.

TONGOLENSIS, Franch. *l. c.* 453.

VIALI, Delav. ex Franch. in Bull. Soc. Phil. Par. sér. 8, viii. (1891) 3. 147.

VITTATA, Bur. & Franch. in Journ. de Bot. v. (1891) 96.

WILSONI, Dunn, in Gard. Chron. i. (1902) 413.

PRINSEPIA UTILIS, Royle ; Franch. Pl. Delav. (1890) 198.

PRUNUS AVIUM, Linn. ; Franch. *l. c.* 197.

BRACHYPODA, Batalin, in Act. Hort. Petrop. xii. (1892) 6. 166.

CAUDATA, Franch. Pl. Delav. (1890) 196.

CINERASCENS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 216.

MAXIMOWICZII, Rupr., var. ; Franch. Pl. Delav. (1890) 195.

MUME, Sieb. & Zucc., var. ; Franch. *l. c.* 197.

NEPALENSIS, var., Batalin, in Act. Hort. Petrop. xiv. (1895) 169.

PADUS, Linn. ; Franch. Pl. Delav. (1890) 198.

PUDDUM, Roxb., var. ; Franch. *l. c.* 168.

SERRULA, Franch. *l. c.* 196.

SETULOSA, Batalin, in Act. Hort. Petrop. xii. (1892) 165.

✓SPINULOSA, Sieb. & Zucc. ; Engl. Jahrb. xxix. (1900) 408.

SZECHUANICA, Batalin, in Act. Hort. Petrop. xiv. (1895) 167.

TATSIENENSIS, Batalin, *l. c.* 322.

THIBETICA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 215.

✓UNDULATA, Wall. ; Engl. Jahrb. xxix. (1900) 408.

VELUTINA, Batalin, in Act. Hort. Petrop. xiv. (1895) 168.

XEROCARPA, Hemsl. in Ann. Bot. ix. (1895) 152.

YUNNANENSIS, Franch. Pl. Delav. (1890) 195.

- PSORALEA CORYLIFOLIA, Franch. Pl. Delav. (1890) 157.  
 PTERNOPETALUM, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, viii. (1885) 246.  
 DAVIDI, Franch. *l. c.*  
 ✓ PTEROCARYA RHOIFOLIA, Sieb. & Zucc.; Engl. Jahrb. xxix. (1900) 274.  
 PTEROCEPHALUS BRETSCHNEIDERI, Batalin, in Act. Hort. Petrop. xiv.  
 (1895) 184.  
 ✓ PTEROLOBIUM ROSTHORNII, Harms, in Engl. Jahrb. xxix. (1900) 410.  
 PTEROSPERMUM FORMOSANUM, Matsum. in Bot. Mag. Tokyo, xv. (1901) 53.  
 PROTEUS, Burkill, in Kew Bull. (1901) 138.  
 PTERYGIELLA DUCLOUXII, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900)  
 22.  
 \* NIGRESCENS, Oliver, in Hook. Ic. Pl. xxv. (1896) t. 2463.  
 PUERARIA CALYCINA, Franch. Pl. Delav. (1890) 181.  
 PEDUNCULARIS, R. Grah., var.; Franch. *l. c.* 182.  
 ✓ TUBEROSA, DC.; Engl. Jahrb. xxix. (1900) 417.  
 YUNNANENSIS, Franch. Pl. Delav. (1890) 181.  
 PULICARIA KOUYANGENSIS, Vaniot, in Bull. Acad. Géog. Bot. sér. 3, xii.  
 (1903) 490.  
 PULSATILLA CERNUA, Spreng.; Komarov, in Act. Hort. Petrop. xxii.  
 (1903) 272.  
 CHINENSIS, Bunge; *l. c.* 271.  
 DAURICA, Spreng.; *l. c.* 273.  
 PYGEUM HENRYI, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 493.  
 PYROLA ATROPURPUREA, Franch. in Journ. de Bot. ix. (1895) 372.  
 CORBIERI, H. Lévêillé, in Bull. Acad. Géog. Bot. xii. (1903) 254.  
 ✓ PYRUS ALNIFOLIA, Decne.; Engl. Jahrb. xxix. (1900) 387.  
 DELAVAYI, Franch. Pl. Delav. (1890) 227.  
 FOLIOLOSA, Wall.; Franch. *l. c.* 230.  
 KANSUENSIS, Batalin, in Act. Hort. Petrop. xiii. 7. (1893) 94.  
 ✓ MICROPHYLLA, Wall.; Engl. Jahrb. xxix. (1900) 387.  
 PASHIA, Buch.-Ham.; Franch. Pl. Delav. (1890) 227.  
 PRATTII, Hemsl. in Kew Bull. (1895) 16.  
 SCABRIFOLIA, Franch. Pl. Delav. (1890) 229.  
 TRANSITORIA, Batalin, in Act. Hort. Petrop. xiii. 7. (1893) 95.  
 VESTITA, Wall.; Franch. Pl. Delav. (1890) 227.  
 WALLICHII, Hook. f.; Franch. *l. c.* 230.  
 YUNNANENSIS, Franch. *l. c.*  
 QUERCUS BLAKEI, Skan, in Hook. Ic. Pl. xxvii. (1900) t. 2662.  
 CAROLINÆ, Skan, in Journ. Linn. Soc., Bot. xxxv. (1903) 518.  
 ✓ DIELSIANA, Seemen, in Engl. Jahrb. xxix. (1900) 291.  
 \* EDITHÆ, Skan, in Hook. Ic. Pl. xxvii. (1900) t. 2662.  
 FORDIANA, Hemsl. in Journ. Linn. Soc., Bot. xxxv. (1903) 478.  
 ✓ GLAUCA, Thunb., var.; Seemen, in Engl. Jahrb. xxix. (1900) 293.

- QUERCUS GROSSESERRATA, Blume; Komarov, in Act. Hort. Petrop. xxii. (1903) 74.  
   ✓ LAMELLOSA, Sm.; Engl. Jahrb. xxix. (1900) 294.  
   ✓ LINEATA, Blume, var.; Engl. *l. c.*  
   REX, Hemsl. in Journ. Linn. Soc., Bot. xxxv. (1903) 477.
- QUINARIA TRICUSPIDATA, Koehne, ex Act. Hort. Petrop. xvii. (1899) 58.
- RADERMACHERA PENTANDRA, Hemsl. in Hook. Ic. Pl. xxviii. (1902) t. 2728.
- ✓ RANDIA HENRYI, E. Pritz. in Engl. Jahrb. xxix. (1900) 581.
- RANUNCULUS AFFINIS, R. Br. var., Maxim.; Engl. Jahrb. xxix. (1900) 334.  
   FLACCIDUS, Hook. f. & Thoms.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 374.  
   HYPERBOREUS, Rottb.; Franch. Pl. Delav. (1889) 21.  
   LABORDEI, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 50.  
   REPTANS, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 294.  
   STENORHYNCHUS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 189.  
   TRICUSPIS, Maxim. Fl. Tangut. (1889) 12.
- ✓ RAPANEA BUXIFOLIA, Mez, in Engl. Pflanzenr., Myrsin. 362.  
   ✓ FABERI, Mez, *l. c.* 358.  
   ✓ PLAYFAIRII, Mez, *l. c.* 361.  
   ✓ YUNNANENSIS, Mez, *l. c.* 359.
- REHMANNIA OLDHAMI, Hemsl. in Ann. Bot. ix. (1895) 154.
- ✓ REINWARDTIA SINENSIS, Hemsl. in Hook. Ic. Pl. xxvi. (1899) t. 2594.
- RHABDOTHAMNOPSIS, Hemsl. in Journ. Linn. Soc., Bot. xxxv. (1903) 517.  
   SINENSIS, Hemsl. *l. c.*
- RHAMNELLA JAPONICA, Miq.; *R. franguloides*, Weberb. in Engl. & Prantl, Natürl. Pflanzenf. III. 5. (1895) 406.
- RHAMNUS FORMOSANA, Matsum. in Bot. Mag. Tokyo, xii. (1898) No. 133. 22.  
   HETEROPHYLLA, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1759.  
   PERSICA, Boiss.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 457.  
   ✓ ROSTHORNII, E. Pritz.; Engl. Jahrb. xxix. (1900) 459.  
   VIRGATA, Roxb. ex Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 457.
- RHAPHIDOPHORA FORMOSANA, Engl. Jahrb. xxv. (1898) 10.
- RHEUM ALEXANDRÆ, Batalin, in Act. Hort. Petrop. xiii. 18. (1894) 384.  
   DELAVAYI, Franch. in Bull. Mus. Hist. Nat. Par. (1895) no. 5, 2.  
   KIALENSE, Franch. *l. c.*  
   STRICTUM, Franch. *l. c.*
- RHODODENDRON ADENOPODUM, Franch. in Journ. de Bot. x. (1895) 391.  
   ANNÆ, Franch. *l. c.* xii. (1898) 258.



- RHODODENDRON ARGYROPHYLLUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 231.
- AUGUSTINII**, Hemsl. (*forma*), in Journ. de Bot. xii. (1898) 261.
- AUREUM**, Franch. in Journ. de Bot. ix. (1895) 394.
- BLEPHAROCALYX**, Franch. *l. c.* 396.
- BODINIERI**, Franch. *l. c.* xii. (1898) 257.
- BONVALOTI**, Franch. *l. c.* v. (1891) 94.
- BREVISTYLUM**, Franch. *l. c.* xii. (1898) 261.
- CALOPHYTON**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 230.
- CHARTOPHYLLUM**, Franch. in Journ. de Bot. ix. (1895) 398.
- ✓ **CHIONOPHYLLUM**, Diels, in Engl. Jahrb. xxix. (1900) 512.
- ✓ **CÆLONEURUM**, Diels, *l. c.* 513.
- CORIACEUM**, Franch. in Journ. de Bot. xii. (1898) 258.
- COSTULATUM**, Franch. *l. c.* ix. (1895) 399.
- CRINIGERUM**, Franch. *l. c.* xii. (1898) 260.
- CYANOCARPUM**, Franch. *l. c.* ix. (1895) 389.
- DAVIDI**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 230.
- DENDROCHARIS**, Franch. *l. c.* 233.
- DETERSILE**, Franch. in Journ. de Bot. xii. (1898) 260.
- DISCOLOR**, Franch. *l. c.* ix. (1895) 391.
- FARGESII**, Franch. *l. c.* 390.
- FLAVIDUM**, Franch. *l. c.* 395.
- FLOCCIGERUM**, Franch. *l. c.* xii. (1898) 259.
- FLORIBUNDUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 233.
- FOKIENSE**, Franch. & Finet, *l. c.* xlvi. (1900) 210.
- FORDII**, Hemsl. in Kew Bull. (1894) 5.
- FORMOSANUM**, Hemsl. *l. c.* (1895) 183.
- GLANDULIFERUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 231.
- GRACILIPES**, Franch. in Journ. de Bot. ix. (1895) 391.
- ✓ **HANCOCKII**, Hemsl. in Hook. Ic. Pl. xxiv. (1895) t. 2381, & Kew Bull. (1895) 107.
- INTRICATUM**, Franch. in Journ. de Bot. ix. (1895) 395.
- KIALENSE**, Franch. *l. c.* 392.
- LATOUCHEÆ**, Franch. & Finet, in Bull. Soc. Bot. Fr. xlvi. (1900) 210.
- LUCIDUM**, Franch. in Journ. de Bot. ix. (1895) 390.
- LUKIANGENSE**, Franch. *l. c.* xii. (1898) 257.
- LUTESCENS**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 235.
- MACULIFERUM**, Franch. in Journ. de Bot. ix. (1895) 393.
- ✓ **MANDARINORUM**, Diels, in Engl. Jahrb. xxix. (1900) 510.
- MEKONGENSE**, Franch. in Journ. de Bot. xii. (1898) 263.
- MOUPINENSE**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 233.
- OREODOXA**, Franch. *l. c.* 230.
- OXYPHYLLUM**, Franch. in Journ. de Bot. xii. (1898) 264.
- PACHYTRICHUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 234.

- ✓ *RHODODENDRON PLATYPODUM*, Diels, in Engl. Jahrb. xxix. (1900) 511.  
   *POLIFOLIUM*, Franch. in Journ. de Bot. ix. (1895) 397.  
   *POLYLEPIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 232.  
   *POTANINI*, Batalin, in Act. Hort. Petrop. xi. (1892) 489.  
   *PRATTII*, Franch. in Journ. de Bot. ix. (1895) 389.  
 ✓ *PRITZELIANUM*, Diels, in Engl. Jahrb. xxix. (1900) 510.   \*  
   *RAMOSISSIMUM*, Bur. & Franch. in Journ. de Bot. xii. (1898) 264.  
 ✓ *ROSTHORNII*, Diels, in Engl. Jahrb. xxix. (1900) 509.  
 ✓ *ROTUNDIFOLIUM*, David; Franch. in Nouv. Arch. Mus. Hist. Nat.  
   Par. sér. 2, x. (1887-88) 47.  
   *RUFESCENS*, Franch. in Journ. de Bot. ix. (1895) 397.  
   *RUFUM*, Batalin, in Act. Hort. Petrop. xi. 16. (1891) 490.  
   *SALUENENSE*, Franch. in Journ. de Bot. xii. (1898) 262.  
   *SANGUINEUM*, Franch. *l. c.* 259.  
   *SELENSE*, Franch. *l. c.* 257.  
   *SIDEROPHYLLUM*, Franch. *l. c.* 262.  
   *SOULIEI*, Franch. *l. c.* ix. (1895) 393.  
   *SPICIFERUM*, Franch. *l. c.* 400.  
   *SPINULIFERUM*, Franch. *l. c.* 399.  
   *STRIGILLOSUM*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 232.  
   *SUTCHUENSE*, Franch. in Journ. de Bot. ix. (1895) 392.  
   *TATSIENSE*, Franch. *l. c.* 394.  
   *THOMPSONI*, Franch. *l. c.* 389.  
   *TRICHOSTOMUM*, Franch. *l. c.* 396.  
   *VERNICOSUM*, Franch. *l. c.* xii. (1898) 258.  
   *VIALI*, Franch. *l. c.* ix. (1895) 398.  
   *YANTHINUM*, Bur. & Franch. *l. c.* v. (1891) 94.  
*RHUS DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 466.  
 ✓ *HENRYI*, Diels, in Engl. Jahrb. xxix. (1900) 432.  
   *POTANINI*, Maxim. in Act. Hort. Petrop. xi. (1890) 110.  
 ✓ *SINICA*, Diels, in Engl. Jahrb. xxix. (1900) 432.  
 ✓ *TOXICODENDRON*, Linn., var.; Diels, *l. c.* 433.  
 ✓ *RHYNCHOSIA DIELSII*, Harms, in Engl. Jahrb. xxix. (1900) 418.  
   *HIMALENSIS*, Benth.; Franch. Pl. Delav. (1890) 185.  
   *LUTEA*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 491.  
   *YUNNANENSIS*, Franch. Pl. Delav. (1890) 185.  
 ✓ *RHYNCHOSPERMUM VERTICILLATUM*, Reinw.; Engl. Jahrb. xxix. (1900)  
   60.  
*RIBES BUREJENSE*, F. Schmidt; Komarov, in Act. Hort. Petrop. xxii.  
   (1903) 435.  
   *DAVIDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
   sér. 2, viii. (1885) 237.  
   *DIACANTHA*, Pall.; Komarov, in Act. Hort. Petrop. xxii. (1903)  
   445.  
   *FARGESII*, Franch. in Bull. Soc. Linn. Par. n. s. (1898) no. 10, 86.

- RIBES GLACIALE**, Wall.; Franch. in *Nouv. Arch. Mus. Hist. Nat. Par.* sér. 2, viii. (1885) 239.  
**HENRYI**, Franch. in *Bull. Soc. Linn. Par. n. s.* (1898) no. 10, 87.  
**LONGERACEMOSUM**, Franch. *Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.* sér. 2, viii. (1885) 238.  
**MANSHURICUM**, Maxim.; Komarov, in *Act. Hort. Petrop.* xxii. (1903) 439.  
**MAXIMOWICZII**, Batalin, in *l. c.* xi. (1891) 16. 487.  
**MOUPINENSE**, Franch. *Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.* sér. 2, viii. (1885) 238.  
**PACHYSANDROIDES**, Hook. f.; *Ic. Pl.* xviii. (1888) t. 1767.  
**PROCUMBENS**, Pall.; Komarov, in *Act. Hort. Petrop.* xxii. (1903) 436.  
 ✓ **ROSTHORNII**, Diels, in *Engl. Jahrb.* xxix. (1900) 378.  
**TRIPARTITUM**, Batalin, in *Act. Hort. Petrop.* xi. 16. (1891) 488.
- RODGERSIA ÆSCULIFOLIA**, Batalin, in *l. c.* xiii. 7. (1893) 96.  
**HENRICI**, Franch. in *Rev. Hort.* (1897) 176.  
**PINNATA**, Franch. *l. c.*  
**TABULARIS**, Hemsl.; Komarov, in *Act. Hort. Petrop.* xxii. (1903) 410.
- RÆTTLERA AUREA**, Franch. in *Bull. Mus. Hist. Nat. Par.* v. (1899) 250.  
**FARGESII**, Franch. *l. c.* 251.  
**MEKONGENSIS**, Franch. *l. c.* 252.  
**THIBETICA**, Franch. *l. c.* 251.  
**UNIFLORA**, Franch. *l. c.* 251.  
**YUNNANENSIS**, Franch. *l. c.* 250.
- ✓ **ROSA BEGGERIANA**, Schrenk; *Engl. Jahrb.* xxix. (1900) 405.  
**BIONDII**, Crép. in *Bull. Soc. Bot. Ital.* (1897) no. 4, 232.  
 ✓ **DAVIDII**, Crép., var. Focke; *Engl. Jahrb.* xxix. (1900) 405.  
**GIRALDII**, Crép. in *Bull. Soc. Bot. Ital.* (1897) no. 4, 233.  
**KOREANA**, Komarov, in *Act. Hort. Petrop.* xviii. 6 (1901) 434.  
**MOSCHATA**, Mill., var. Crép.; Franch. *Pl. Delav.* (1890) 218.  
 ✓ **PRATTII**, Hemsl. in *Journ. Linn. Soc., Bot.* xxix. (1893) 307.  
**SOULIANA**, Crép. in *Compt. Rend. Bull. Soc. Bot. Belg.* xxxv. II. (1896) 21.  
**WEBBIANA**, Wall. var.; Crép. in *Bull. Soc. Bot. Belge*, xiii. (1874) 276.
- ✓ **RUBIA SCHUMANNIANA**, E. Pritz.; *Engl. Jahrb.* xxix. (1900) 583.
- RUBUS ADENANTHUS**, Finet & Franch. in *Bull. Soc. Bot. Fr.* xlvi. (1900) 208.  
**ALLOPHYLLUS**, Hemsl. in *Journ. Linn. Soc., Bot.* xxix. (1893) 305.  
 ✓ **AMPELINUS**, Focke, in *Engl. Jahrb.* xxix. (1900) 396.  
 ✓ **AMPHIDASYS**, Focke, *l. c.*  
 ✓ **BAMBUSARUM**, Focke, in Hook. *Ic. Pl.* xx. (1891) sub t. 1952.  
 ✓ **BIFLORUS**, Buch. ex Focke; *Engl. Jahrb.* xxix. (1900) 397.

- RUBUS BODINIERI, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 92.
- ✓ BUEGGERI, Miq. ex Focke; Engl. Jahrb. xxix. (1900) 394.
- CHAFFANJONI, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 98.
- CHILIADENUS, Focke, in Hook. Ic. Pl. xx. (1891) sub t. 1952.
- CHINENSIS, Franch. Pl. Delav. (1890) 207.
- ✓ CHROOSEPALUS, Focke, in Hook. Ic. Pl. xx. (1891) t. 1952.
- COCHINCHINENSIS, Tratt.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 220.
- COCKBURNIANUS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 305.
- CORCHORIFOLIUS, Linn. f. var., Focke; Engl. Jahrb. xxix. (1900) 391.
- DELAVAYI, Franch. Pl. Delav. (1890) 205.
- ELLIPTICUS, Sm.; Franch. Pl. Delav. 206.
- ✓ EUGENIUS, Focke, in Engl. Jahrb. xxix. (1900) 393.
- ✓ EULEUCUS, Focke; Engl. l. c. 397.
- FARGESII, Franch. in Journ. de Bot. viii. (1894) 294.
- ✓ FLAGELLIFLORUS, Focke, in Engl. Jahrb. xxix. (1900) 393.
- ✓ FLOSCULOSUS, Focke, in Hook. Ic. Pl. xx. (1891) sub t. 1952.
- ✓ FOCKEANUS, Kurz; Engl. Jahrb. xxix. (1900) 397.
- FOLIOLOSUS, D. Don; Franch. Pl. Delav. 208.
- FRAGARIOIDES, Bertol., varr.; Franch. l. c. 202.
- GENTILIANUS, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 99.
- ✓ GIRALDIANUS, Focke, in Engl. Jahrb. xxix. (1900) 401.
- HUPEHENSIS, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1816.
- IDÆUS, Linn. var., Focke; Palibin in Act. Hort. Petrop. xvii. i. (1899) 78.
- ✓ INNOXIUS, Focke, in Engl. Jahrb. xxix. (1900) 395.
- ✓ IRENÆUS, Focke, l. c. 394.
- ✓ JAMINI, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. x. (1902) 102.
- KERRIIFOLIUS, H. Lévillé & Vaniot, l. c. 100.
- LASIOCARPUS, Sm.; Franch. Pl. Delav. (1890) 207.
- LASIOSTYLUS, Focke, in Hook. Ic. Pl. xx. (1891) t. 1951.
- LOROPETALUS, Franch. Pl. Delav. (1890) 203.
- LUTESCENS, Franch. l. c. 206.
- MACILENTUS, Camb.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 221.
- ✓ MALIFOLIUS, Focke, in Hook. Ic. Pl. xx. (1891) t. 1947.
- ✓ MESOGÆUS, Focke, in Engl. Jahrb. xxix. (1900) 399.
- ✓ Var. OXYCOMUS, Focke, l. c.
- ✓ MODICUS, Focke, l. c.

- RUBUS MONGUILLONI, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 101.
- MULTIBRACTEATUS, H. Lévêillé & Vaniot, *l. c.* 99.
- NIVEUS, Thunb. Diss. Rub. 9.
- OTOPHORUS, Franch. Pl. Delav. (1890) 204.
- ✓ PACATUS, Focke, in Engl. Jahrb. xxix. (1900) 395.
- ✓ PELTATUS, Maxim. ; Focke, *l. c.* 391.
- PILEATUS, Focke, in Hook. Ic. Pl. xx. (1891) sub t. 1952.
- PINNATISEPALUS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 305.
- POLYTRICHUS, Franch. Pl. Delav. (1890) 204.
- PYRIFOLIUS, Sm. ; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 219 (*pirifolius*).
- ✓ SEPALANTHUS, Focke, in Engl. Jahrb. xxix. (1900) 391.
- SETCHUENENSIS, Franch. in Journ. de Bot. v. (1891) 46.
- SIMPLEX, Focke, in Hook. Ic. Pl. xx. (1891) t. 1948.
- ✓ SORBIFOLIUS, Maxim. ; Focke, in Engl. Jahrb. xxix. (1900) 391.
- SOZOSTYLUS, Focke, in Hook. Ic. Pl. xx. (1891) sub t. 1952.
- SPINIPES, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 306.
- ✓ TELEDAPOS, Focke, in Engl. Jahrb. xxix. (1900) 398.
- THIBETANUS, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 221.
- VIBURNIFOLIUS, Franch. in Bull. Mus. Hist. Nat. Par. i. (1895) 63.
- XANTHOCARPUS, Bur. & Franch. in Journ. de Bot. v. (1891) 46.
- ✓ XANTHONEURUS, Focke, in Engl. Jahrb. xxix. (1900) 392.
- RUMEX DOMESTICUS, Hartm. ; Komarov, in Act. Hort. Petrop. xxii. (1903) 116.
- STENOPHYLLUS, Ledeb. ; Komarov, *l. c.* 115.
- ✓ SABIA LEPTANDRA, Hook. f. & Thoms. ; Engl. Jahrb. xxix. (1900) 451.
- ✓ PURPUREA, Hook. f. & Thoms. ; Engl. *l. c.*
- ✓ SCHUMANNIANA, Diels. ; *l. c.*
- YUNNANENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 465.
- SACCOLABIUM FARGESII, Kränzl. ; Journ. de Bot. xvii. (1903) 423.
- SAGERETIA FERRUGINEA, Oliver, in Hook. Ic. Pl. xviii. (1887) t. 1710.
- OPPOSITIFOLIA, Brongn., var. ; Franch. Pl. Delav. (1890) 134.
- PAUCICOSTATA, Maxim. in Act. Hort. Petrop. xi. i. (1890) 101.
- SALICORNIA HERBACEA, Linn. ; Act. Hort. Petrop. xviii. (1899) 179.
- ✓ SALIX BOCKII, Seemen, in Engl. Jahrb. xxix. (1900) 278.
- CINEREA, Linn. ; Komarov, in Act. Hort. Petrop. xxii. (1903) 22.
- ✓ DENSIFOLIATA, Seemen, in Engl. Jahrb. xxi. (1896) Beibl. 53. 57.
- ✓ DICTYONEURA, Seemen, *l. c.* xxix. (1900) 275.
- ✓ HETEROCHROMA, Seemen, *l. c.* xxi. (1896) Beibl. 53. 56.
- MAXIMOVICZII, Komarov, in Act. Hort. Petrop. xxii. (1903) 25.

- SALIX MULTINERVIS, Komarov, in Act. Hort. Petrop. xxii. (1903) 25.  
 MYRTILLOIDES, Komarov, *l. c.* 26.
- ✓ ROSTHORNII, Seemen, in Engl. Jahrb. xxix. (1900) 276.  
 VAGANS, Komarov, in Act. Hort. Petrop. xxii. (1903) 31.  
 VIMINALIS, Linn.; Komarov, *l. c.* 32.
- SALVIA BREVILABRA, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891)  
 149.
- ✓ MANDARINORUM, Diels, in Engl. Jahrb. xxix. (1900) 557.  
 PRATTII, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1903) 316.  
 PYGMÆA, Matsum. in Bot. Mag. Tokyo, xi. (1897) 70.  
 TATSIENENSIS, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. (1891) 150.  
 TRICUSPIS, Franch. *l. c.* 150.  
 YUNNANENSIS, C. H. Wright, in Kew Bull. (1896) 164.
- SANICULA CÆRULESCENS, Franch. in Bull. Soc. Phil. Par. sér. 8, vi.  
 (1894) 110.  
 HACQUETIODES, Franch. *l. c.*
- ✓ RUGULOSA, Diels, in Engl. Jahrb. xxix. (1900) 491.  
 YUNNANENSIS, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 108.
- ✓ SARUMA HENRYI, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1895.
- ✓ SATYRIUM SETCHUENICUM, Kränzl. in Engl. Jahrb. xxix. (1900) 266.
- SAURAUJA NAPAULENSIS, DC.; Franch. in Bull. Soc. Bot. Fr. xxxiii.  
 (1886) 438.
- SAUSSUREA ALATIPES, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893)  
 308.  
 AURICULATA, Hemsl. *l. c.* non Franch. = S. OTOPHYLLA, Hemsl.  
*infra.*  
 AURICULATA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, x. (1887-8) 40.
- ✓ BARONIANA, Diels, in Engl. Jahrb. xxix. (1900) 625.  
 BULLOCKII, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 509.  
 CARDUIFORMIS, Franch. in Journ. de Bot. viii. (1894) 343.  
 CAUDATA, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 147.  
 CHETCHOTZE, Franch. in Journ. de Bot. ii. (1888) 359.  
 CILIARIS, Franch. *l. c.* 337.  
 CIRSIODES, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 309.  
 COMPTA, Franch. in Journ. de Bot. x. (1896) 423.  
 CONYZOIDES, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 309.  
 CORDIFOLIA, Hemsl. *l. c.* 310.  
 CRISPA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 21.  
 DECURRENS, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 310.  
 DELAVAYI, Franch. in Journ. de Bot. viii. (1894) 340.  
 DIMORPHÆA, Franch. *l. c.*
- ✓ DOLICHOPODA, Diels, in Engl. Jahrb. xxix. (1900) 623.  
 DUTAILLYANA, Franch. in Journ. de Bot. x. (1896) 421.  
 DZEURENSIS, Franch. *l. c.* viii. (1894) 339.

- SAUSSUREA EDULIS, Franch. in Journ. de Bot. ii. (1888) 337.  
 ERIOCEPHALA, Franch. *l. c.* viii. (1894) 339.  
 FARGESII, Franch. *l. c.* 344.  
 FLEXUOSA, Franch. *l. c.* 341.  
 GOSSYPIPHORA, D. Don; Franch. in Journ. de Bot. ii. (1888) 359.  
 GRAMINEA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 509.  
 GROSSESERRATA, Franch. in Journ. de Bot. ii. (1888) 354.  
 GRUMGRJIMAILO, Winkl.; Bretsch. Hist. Bot. Disc. China, ii. 1039.  
 HENRYI, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 311.  
 KUNTHIANA, C. B. Clarke; Franch. in Journ. de Bot. ii. (1888) 340.  
 LAMPSANIFOLIA, Franch. in Journ. de Bot. ii. (1888) 357.  
 LANUGINOSA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 20.  
 LEONTODON, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 509.  
 LIKIANGENSIS, Franch. in Journ. de Bot. ii. (1888) 356.  
 LINGULATA, Franch. *l. c.* x. (1896) 423.  
 LONGIFOLIA, Franch. *l. c.* ii. 354.  
 MACROTA, Franch. *l. c.* viii. (1894) 343.  
 MANSHURICA, Komarov, in Act. Hort. Petrop. xviii. 2. (1900) 424.  
 ✓ MICROCEPHALA, Diels, in Engl. Jahrb. xxix. (1900) 624.  
 MINUTA, Winkl. in Act. Hort. Petrop. xiii. (1894) 13. 243.  
 MOLLIS, Franch. in Journ. de Bot. ii. (1888) 353.  
 NOBILIS, Franch. *l. c.* 355.  
 ✓ OBVALLATA, Wall. var.; Diels, in Engl. Jahrb. xxix. (1900) 623.  
 OLIGANTHA, Franch. in Journ. de Bot. x. (1896) 421.  
 OTOPHYLLA, Hemsl. n. nov. *Syn. auriculata, Hemsl. non Franch.*  
 PACHYNEURA, Franch. in Journ. de Bot. viii. (1894) 354.  
 PEDUNCULARIS, Franch. *l. c.* ii. (1888) 357.  
 POPULIFOLIA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 311.  
 RADIATA, Franch. in Journ. de Bot. ii. (1888) 356.  
 ROMULEIFOLIA, Franch. *l. c.* 339.  
 ROSTHORNII, Diels, in Engl. Jahrb. xxix. (1900) 625.  
 SALIGNA, Franch. in Journ. de Bot. viii. (1894) 345.  
 SCABRIDA, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 146.  
 SEMILYRATA, Bur. & Franch. in Journ. de Bot. v. (1891) 76.  
 SOULIEI, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 147.  
 SPATULIFOLIA, Franch. in Journ. de Bot. ii. (1888) 338.  
 STRICTA, Franch. *l. c.* viii. (1894) 342.  
 SUGHOO, C. B. Clarke; Franch. *l. c.* ii. (1888) 339.  
 SUTCHUENENSIS, Franch. *l. c.* viii. (1894) 353.  
 TARAXACIFOLIA, Wall.; Franch. *l. c.* ii. (1888) 339.  
 TATSIENENSIS, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891)  
 146.  
 THIBETICA, Franch. in Journ. de Bot. viii. (1894) 338.  
 UMBROSA, Komarov, Act. Hort. Petrop. xviii. 2. (1900) 423.  
 VAGINATA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 510.

- SAUSSUREA VESTITA, Franch. in Journ. de Bot. ii. (1888) 358.  
 VILLOSA, Franch. *l. c.* 353.  
 VIRGATA, Franch. *l. c.* viii. (1894) 340.  
 WOODIANA, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 312.  
 YUNNANENSIS, Franch. in Journ. de Bot. ii. (1888) 340.  
 SAXIFRAGA APLOPHYILLOIDES, Franch. Pl. Delav. (1890) 238.  
 AURANTIACA, Franch. *l. c.* 236.  
 BRACHYPHYLLA, Franch. *l. c.* 237.  
 CANDELABRUM, Franch. *l. c.* 232.  
 CARDIOPHYLLA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat.  
 Par. sér. 2, viii. (1885) 230.  
 CHIONOPHILA, Franch. in Journ. de Bot. x. (1896) 265.  
 CHRYSANTHA, Franch. Pl. Delav. (1890) 236.  
 CRINALIS, Franch. *l. c.* 233.  
 DAVIDII, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par.  
 sér. 2, viii. (1885) 229.  
 DRABÆFORMIS, Franch. Pl. Delav. (1890) 235.  
 FLABELLIFOLIA, Franch. in Journ. de Bot. viii. (1894) 295.  
 ✓ GEMMIGERA, Engl. Jahrb. xxix. (1900) 366.  
 GEMMIPARA, Franch. in Journ. de Bot. x. (1896) 262.  
 GILLII, Trimen, ex Gill, River & Golden Sand, ii. 426.  
 ✓ GIRALDIANA, Engl. Jahrb. xxix. (1900) 365.  
 GLAUCOPHYLLA, Franch. Pl. Delav. (1890) 239.  
 ✓ HISPIDULA, D. Don; Engl. Jahrb. xxix. (1900) 366.  
 HYPERICOIDES, Franch. in Journ. de Bot. x. (1896) 261.  
 ✓ JOSEPHI, Engl. Jahrb. xxix. (1900) 366.  
 LIKIANGENSIS, Franch. in Journ. de Bot. x. (1896) 266.  
 LONGISTYLA, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3. (1891) 144.  
 MACROSTIGMA, Franch. Pl. Delav. (1890) 240.  
 MANSHURIENSIS, Komarov, in Act. Hort. Petrop. xxii. (1903) 415.  
 MELANOCENTRA, Franch. in Journ. de Bot. x. (1896) 263.  
 MICRANTHA, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii.  
 (1885) 229.  
 NUTANS, Franch. *l. c.* 233.  
 OREOPHILA, Franch. in Journ. de Bot. x. (1896) 260.  
 PEPLIDIFOLIA, Franch. Pl. Delav. (1890) 234.  
 PETROPHILA, Franch. *l. c.* 237.  
 PHÆNOPHYLLA, Franch. in Journ. de Bot. x. (1896) 261.  
 PUNCTATA, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 414.  
 PURPURASCENS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2,  
 viii. (1885) 233.  
 RUPICOLA, Franch. in Journ. de Bot. x. (1896) 264.  
 SANGUINEA, Franch. *l. c.* viii. (1894) 294.  
 ✓ SIBIRICA, Linn., var.; Engl. Jahrb. xxix. (1900) 364.



- SAXIFRAGA STELLARLÆFOLIA, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 231.
- TRINERVIA, Franch. *l. c.* 232.
- TSANGCHANENSIS, Franch. Pl. Delav. (1890) 233.
- SCABIOSA BRETSCHNEIDERI, Batalin, in Act. Hort. Petrop. xiv. 8. (1895) 184.
- ✓ SCEPTROCNIDE MACROSTACHYA, Maxim. ; Engl. Jahrb. xxix. (1900) 302.
- ✓ SCHEFFLERA FARGESII, Harms, in Engl. Jahrb. xxix. (1900) 486.
- ✓ MEGALOBOTRYA, Harms, *l. c.*
- ✓ SCHIMA ARGENTEA, E. Pritz. in Engl. Jahrb. xxix. (1900) 473.
- SCHISTOCARYUM, Franch. in Bull. Soc. Linn. Par. ii. (1891) 930.
- CILIARE, Bur. & Franch. in Journ. de Bot. v. (1891) 104.
- MYOSOTIDEUM, Franch. in Bull. Soc. Linn. Par. ii. (1891) 930.
- OVALIFOLIUM, Bur. & Franch. in Journ. de Bot. v. (1891) 104.
- ✓ SCHIZANDRA AXILLARIS, Hook. f. & Thoms. ; Engl. Jahrb. xxix. (1900) 322.
- ✓ ELONGATA, Hook. f. & Thoms. ; Engl. *l. c.* 322.
- ✓ GLAUDESCENS, Diels, *l. c.* 323.
- GRANDIFLORA, Hook. f. & Thoms. ; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 385.
- PROPINQUA, Hook. f. & Thoms. var., Oliver, in Hook. Ic. Pl. xviii. (1889) t. 1715.
- SCHIZOPEPON DIOICUS, Cogn. in Hook. Ic. Pl. xxiii. (1892) t. 2224.
- SCHIZOPHRAGMA HYDRANGEOIDES, Sieb. & Zucc. ; Diels, in Engl. Jahrb. xxix. (1900) 376.
- INTEGRIFOLIA, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1934.
- SCOLOPIA CHINENSIS, Briquet, in Ann. Conserv. Jard. Bot. Genève, ii. (1898) 42.
- SCORZONERA MONGOLICA, var. ; Palibin, in Act. Hort. Petrop. xiv. (1895) 128.
- ✓ SCROPHULARIA ALATA, A. Gray ; Engl. Jahrb. xxix. (1900) 565.
- DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 15.
- DIPLODONTA, Franch. *l. c.* 14.
- FARGESII, Franch. *l. c.* 12.
- KANSUENSIS, Batalin, in Act. Hort. Petrop. xiii. (1894) 381.
- MANDARINORUM, Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 13.
- MICRODONTA, Franch. *l. c.* 11.
- SOULIEI, Franch. *l. c.* 15.
- SPICATA, Franch. *l. c.* 14.
- YUNNANENSIS, Franch. *l. c.* 12.
- SCUTELLARIA AMÆNA, C. H. Wright, in Kew Bull. (1896) 164.
- FORMOSANA, N. E. Br. in Gard. Chron. ii. (1894) 212.
- ✓ SEDUM BRACTEATUM, Diels, in Engl. Jahrb. xxix. (1900) 362.
- ✓ CRASSIPES, Wall. ; Diels, *l. c.* 360.
- DISCOLOR, Franch. in Journ. de Bot. x. (1896) 285.
- GLACIALE, Franch. *l. c.* 290.

- ✓ *SEDUM HENRYI*, Diels, in Engl. Jahrb. xxix. (1900) 361.  
*LEUCOCARPUM*, Franch. in Journ. de Bot. x. (1896) 288.  
*MACROLEPIS*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 240.  
*MIDDENDORFFIANUM*, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 397.  
*MOSOYNENSE*, Franch. in Journ. de Bot. x. (1896) 291.  
*NOBILE*, Franch. *l. c.* 285.  
*OBTUSISEPALUM*, Franch. *l. c.* 289.  
*PLATYSEPALUM*, Franch. *l. c.*  
*PRIMULOIDES*, Franch. *l. c.* 287.  
 ✦ *ROSTHORNIANUM*, Diels, in Engl. Jahrb. xxix. (1900) 361.  
*SCABRIDUM*, Franch. in Journ. de Bot. x. (1896) 284.  
 ✓ *SINICUM*, Diels, in Engl. Jahrb. xxix. (1900) 361.  
*TENUIFOLIUM*, Franch. in Journ. de Bot. x. (1896) 290.  
 ✓ *VALERIANOIDES*, Diels, in Engl. Jahrb. xxix. (1900) 360.  
*VIVIPARUM*, Maxim.; Act. Hort. Petrop. xxii. (1903) 390.  
*YUNNANENSE*, Franch. in Journ. de Bot. x. (1896) 286.  
*SELINUM COREANUM*, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 956.  
*CRYPTOTÆNIUM*, Boissieu, *l. c.* 847.  
*DAUCOIDES*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 135.  
*MELANOTILINGIA*, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 956.  
*OLIVERIANUM*, Boissieu, *l. c.* 846.  
*SENECIO ACERIFOLIUS*, Winkl. in Act. Hort. Petrop. xiii. i. (1893) 9.  
*AINSLIÆFLORUS*, Franch. in Journ. de Bot. viii. (1894) 360.  
*ARACHNANTHUS*, Franch. *l. c.* 355.  
*ASPERIFOLIUS*, Franch. *l. c.* x. (1896) 414.  
*ATROVIOLACEUS*, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 303.  
*BEGONIÆFOLIUS*, Franch. in Journ. de Bot. viii. (1894) 358.  
*BLATTARIÆFOLIUS*, Franch. *l. c.* x. (1896) 415.  
*BODINIERI*, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 348.  
*CAMPANULATUS*, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 284.  
*CAMPTODONTUS*, Franch. in Journ. de Bot. x. (1896) 413.  
*CAROLI*, Winkl. in Act. Hort. Petrop. xiii. i. (1893) 7.  
*CHRYSANTHEMOIDES*, DC., var.; Hook. f., ex Engl. Jahrb. xxix. (1900) 620.  
*CONCINNUS*, Franch. in Journ. de Bot. x. (1896) 418.  
*CYCLAMINIFOLIUS*, Franch. *l. c.* viii. (1894) 362.  
*CYCLOTUS*, Bur. & Franch. *l. c.* v. (1891) 74.  
*DAVIDI*, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 40.  
*DELAVAYI*, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 286.  
 ✓ *DENSIFLORUS*, Wall.; Engl. Jahrb. xxix. (1900) 621.

- SENECIO DIANTHUS**, Franch. in Journ. de Bot. x. (1896) 419.  
**DICTYONEURUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 294.  
**DIDIDYMANTHA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 503.  
**DISCOIDEUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 284.  
**DODRANS**, Winkl. in Act. Hort. Petrop. xiv. (1895) 152.  
**DRYAS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 504.  
**DUCIFORMIS**, Winkl. in Act. Hort. Petrop. xiv. (1895) 155.  
**DUCLouxII**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 504.  
**ERYTHROPAPPUS**, Bur. & Franch. in Journ. de Bot. v. (1891) 73.  
**EURYOPHYLLUS**, Winkl. in Act. Hort. Petrop. xiv. (1895) 156.  
**FARGESII**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 284.  
**FIBRILLOSUS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 505.  
**FILIFERUS**, Franch. in Journ. de Bot. x. (1896) 416.  
**GANPINENSIS**, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 19.  
**GENTILIANUS**, Vaniot, *l. c.* xi. (1902) 350.  
**GLUMACEUS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 505.  
**GYNURA**, Winkl. in Act. Hort. Petrop. xiv. (1895) 157.  
**HELIANTHUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 286.  
**HENRICI**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 351.  
**HOI**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 506.  
**IANTHOPHYLLUS**, Franch. in Journ. de Bot. viii. (1894) 362.  
**KANAITZENSIS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 298.  
**KANSUENSIS**, Franch. *l. c.* 287.  
**KASCHKAROWI**, Winkl. in Act. Hort. Petrop. xiv. i. (1895) 152.  
**KEMATONGENSIS**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 347.  
**KIALENSIS**, Franch. in Journ. de Bot. x. (1896) 413.  
**KOREANUS**, Komarov, in Act. Hort. Petrop. xviii. ii. (1900) 421.  
**KOUALAPENSIS**, Franch. in Journ. de Bot. viii. (1894) 356.  
**LABORDEI**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 345.  
**LANKONGENSIS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 301.  
**LAPATHIFOLIUS**, Franch. *l. c.* 306.  
**LATIPES**, Franch. in Journ. de Bot. viii. (1894) 356.  
**LEUCANTHEMUM**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 507.  
**LEUCOCEPHALUS**, Franch. in Journ. de Bot. viii. (1894) 360.  
**LEVEILLEI**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 346.  
**LIATROIDES**, Winkl. in Act. Hort. Petrop. xiii. i. (1893) 8.  
**LUCORUM**, Franch. in Journ. de Bot. x. (1896) 415.  
**LUTICOLA**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 507.  
**MARTINI**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 346.  
**MAXIMOWICZII**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 287.  
**MELANOCEPHALUS**, Franch. *l. c.* 294.  
**MICRODONTUS**, Bur. & Franch. in Journ. de Bot. v. (1891) 76.  
**MONANTHUS**, Diels, in Engl. Jahrb. xxix. (1900) 621.  
**MOSOYNENSIS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 298.  
**NELUMBIFOLIUS**, Bur. & Franch. in Journ. de Bot. v. (1891) 74.

- SENECIO NIGROCINCTUS**, Franch. in Journ. de Bot. x. (1896) 417.  
**NIMBORUM**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-88) 39.  
**NOBILIS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 287.  
 ✓ **PAUCINERVIS**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 507.  
**PHÆNICOCHÆTUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 295.  
**PHYLLOLEPIS**, Franch. in Journ. de Bot. viii. (1894) 360.  
**PLANTAGINIFOLIUS**, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. III. (1891) 145.  
**PLATYGLOSSUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 293.  
**PLEUROCAULIS**, Franch. in Journ. de Bot. viii. (1894) 365.  
**POTANINI**, Winkl. in Act. Hort. Petrop. xiii. I. (1893) 5.  
**PRATTII**, Hemsl. in Hook. Ic. Pl. xxv. (1896) t. 2491.  
**PRINCIPIS**, Franch. in Journ. de Bot. x. (1896) 412.  
**PRIONOPHYLLUS**, Franch. *l. c.* 420.  
 ✓ **PROFUNDORUM**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 507.  
**PSEUDOSONCHUS**, Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 349.  
**PTERIDOPHYLLUS**, Franch. in Journ. de Bot. viii. (1894) 364.  
**PUTJATÆ**, Winkl. in Act. Hort. Petrop. xiv. I. (1895) 125.  
**QUINQUELOBUS**, Hook. f. & Thoms. ; Engl. Jahrb. xxix. (1900) 620.  
**RENATUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 285.  
**RUFICOMUS**, Franch. *l. c.* 298.  
**RUFIPILIS**, Franch. in Journ. de Bot. viii. (1894) 359.  
**SETCHUENENSIS**, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. III. (1891) 145.  
 ✓ **SOLENOIDES**, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 508.  
**SOULIEI**, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. III. (1891) 145.  
**SPATHIPHYLLUS**, Franch. in Journ. de Bot. x. (1896) 416.  
**STENOGLOSSUS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 304.  
**SUBSPICATUS**, Bur. & Franch. in Journ. de Bot. v. (1891) 75.  
**TALIENSIS**, Franch. *l. c.* viii. (1894) 357.  
**TALONGENSIS**, Franch. *l. c.* x. (1896) 419.  
**TATSIENENSIS**, Bur. & Franch. *l. c.* v. (1891) 75.  
**TENUIPES**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 297.  
**TONGOLENSIS**, Franch. *l. c.* 305.  
**TRICUSPIS**, Franch. in Journ. de Bot. viii. (1894) 357.  
**TSANGCHANENSIS**, Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 299.  
**VELLEREUS**, Franch. *l. c.*  
**VESPERTILIO**, Franch. in Journ. de Bot. viii. (1894) 359.  
**VILLIFERUS**, Franch. *l. c.* 362.  
**YESOENSIS**, var.; Franch. in Bull. Soc. Bot. Fr. xxxix. (1892) 307.  
**YUNNANENSIS**, Franch. *l. c.* 303.  
 ✓ **SERRATULA CORONATA**, Linn. ; Engl. Jahrb. xxix. (1900) 628.

- SERRATULA MARTINI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 21.
- SESELI DELAVAYI, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 130.
- ✓ GIRALDII, Diels, in Engl. Jahrb. xxix. (1900) 497.
- PROVOSTII, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 842.
- YUNNANENSE, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 129.
- SHORTIA ROTUNDIFOLIA, Makino, in Bot. Mag. Tokyo, ix. (1895) 327, & x. (1896) 221.
- SINENSIS, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2624.
- THIBETICA, Franch. Pl. Dav. in Nouv. Arch. Mus. ser. 2, x. (1887–8) 54.
- SHUTERIA LONGIPES, Franch. Pl. Delav. (1896) 179.
- SINENSIS, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2626.
- ✓ SIBBALDIA PROCUMBENS, Linn.; Engl. Jahrb. xxix. (1900) 404.
- SILENE ADENANTHA, Franch. Pl. Delav. (1889) 84.
- ASCLEPIADEA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 422.
- CÆSPITOSA, Bur. & Franch. in Journ. de Bot. v. (1891) 22.
- CAPITATA, Komarov, in Act. Hort. Petrop. xviii. (1901) 440.
- CARDIOPETALA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 419.
- CHINENSIS, Hort. in Bosse, Handl. Blumengartnerei, ed. 3, vii. (1861) 445.
- DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 424.
- FOLIOSA, Maxim.; Franch. Journ. de Bot. iv. (1890) 303.
- GRANDIFOLIA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 427.
- KOREANA, Komarov, in Act. Hort. Petrop. xviii. (1901) 440.
- LANGKONGENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 421.
- LUTEA, Franch. *l. c.* 420.
- MACROSTYLA, Maxim.; Komarov, in Act. Hort. Petrop. xxii. (1903) 193.
- MELANANTHA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 423.
- NAPULIGERA, Franch. Pl. Delav. (1889) 82.
- OTODONTA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 426.
- PACHYRRHIZA, Franch. *l. c.* 427.
- PHÆNICODONTA, Franch. *l. c.* 422.
- PLATYPETALA, Bur. & Franch. Journ. de Bot. v. (1891) 22.
- PLATYPHYLLA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 419.
- POTANINI, Maxim. in Act. Hort. Petrop. xi. i. (1890) 66.
- PTEROSPERMA, Maxim. *l. c.* 67.
- REPENS, Patrin; Franch. in Journ. de Bot. iv. (1890) 302.
- RUBICUNDA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 417.
- SCOPULORUM, Franch. *l. c.* 423.
- SINENSIS, Williams, in Journ. Linn. Soc., Bot. xxxii. (1896) 179.
- SZECHUENSIS, Williams, *l. c.* xxxiv. (1899) 428.
- TANAKÆ, Maxim. in Bull. Acad. Pétersb. xxxii. (1888) 481.

- ✓ *SILENE TENUIS*, Willd.; Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 429.  
*TRACHYPHYLLA*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 418.  
*VISCIDULA*, Franch. *l. c.* 421.  
*YUNNANENSIS*, Franch. *l. c.* 425.
- SISYMBRIUM MAXIMOWICZII*, Palibin, in Act. Hort. Petrop. xvii. (1899) 28.  
*THALIANUM*, J. Gay; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 403.
- SIUM MATSUMURÆ*, Boissieu, in Bull. Herb. Boiss. sér. 2, iii. (1903) 954.  
*NINSI*, Linn.; Boissieu, *l. c.* 958.  
*NIPPONICUM*, Maxim.; Boissieu, *l. c.*
- SKIMMIA FORTUNEI*, Mast. in Gard. Chron. (1889) i. 520.  
 ✓ *JAPONICA*, Thunb.; Engl. Jahrb. xxix. (1900) 424.  
*LAUREOLA*, Franch. in Nouv. Arch. Mus. sér. 2, x. (1888) 171.
- ✓ *SLOANEA HONGKONGENSIS*, Hemsl. in Hook. Ic. Pl. xxvii. (1900) t. 2628.
- SMILACINA = TOVARIA*.  
 ✓ *FARGESII*, Diels, in Engl. Jahrb. xxix. (1900) 246.  
 ✓ *JAPONICA*, A. Gray; Diels, *l. c.*  
 ✓ *STENOLOBA*, Diels, *l. c.* 247.  
 ✓ *TUBIFERA*, Batalin; Diels, *l. c.* 246.
- ✓ *SMILAX BOCKII*, Warb. in Engl. Jahrb. xxix. (1900) 259.  
 ✓ *BREVIPIES*, Warb. *l. c.* 256.  
 ✓ *CINEREA*, Warb. *l. c.* 258.  
 ✓ *COCCULOIDES*, Warb. *l. c.* 257.  
 ✓ *CYCLOPHYLLA*, Warb. *l. c.*  
 ✓ *DISCOTIS*, Warb. *l. c.* 256.  
 ✓ *GLAUCO-CHINA*, Warb. *l. c.* 255.  
 ✓ *LONGIPES*, Warb. *l. c.* 256.  
 ✓ *NEBELII*, Gilg, *l. c.* xxxiv. (1904) Beibl. 75. p. 26.  
 ✓ *POLYCOLEA*, Warb. *l. c.* xxix. (1900) 257.  
 ✓ *TRIGONA*, Warb. *l. c.* 258.
- SMITHIA YUNNANENSIS*, Franch. Pl. Delav. (1890) 170.
- SONERILA PEPEROMIÆFOLIA*, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1814.
- SOPHORA GLAUCA*, Leschen.; Franch. Pl. Delav. (1890) 188.
- SORBUS PEKINENSIS*, Koehne, in Wittm. Gartenfl. (1901) 406.
- SOULIEA*, Franch. in Journ. de Bot. xii. (1898) 69.  
*VAGINATA*, Franch. *l. c.* 70.
- ✓ *SPATHOLOBUS PULCHER*, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 489.  
 ✓ *SUBERECTUS*, Dunn, *l. c.*  
 ✓ *VARIANS*, Dunn, *l. c.* 490.

- ✓ *SPIRÆA ALPINA*, Pall.; Engl. Jahrb. xxix. (1900) 382.  
     *ANOMALA*, Batalin, in Act. Hort. Petrop. xiii. (1893) 92.  
     *ARUNCUS*, Linn.; Franch. Pl. Delav. (1890) 201.  
     *CANESCENS*, D. Don, var., Batalin, in Act. Hort. Petrop. xiv. (1898) 321.  
     *JAPONICA*, Linn.; Franch. Pl. Delav. (1890) 201.  
 ✓ *ROSTHORNII*, E. Pritz. in Engl. Jahrb. xxix. (1900) 383.  
     *VEITCHI*, Hemsl. in Gard. Chron. (1903) i. 258.  
     *VELUTINA*, Franch. Pl. Delav. (1890) 201.  
     *VESTITA*, Wall.; Franch. *l. c.*  
     *VIRGATA*, Franch. *l. c.* 199.  
     *YUNNANENSIS*, Franch. *l. c.* 200.  
*STACHYURUS CHINENSIS*, Franch. in Journ. de Bot. xii. (1898) 254.  
     ✓ *HIMALAICUS*, Hook. f. & Thoms.; Engl. Jahrb. xxix. (1900) 475.  
     *SALICIFOLIUS*, Franch. in Journ. de Bot. xii. (1898) 253.  
     *YUNNANENSIS*, Franch. *l. c.*  
 ✓ *STAPHYLEA BUMALDA*, Sieb. & Zucc.; Engl. Jahrb. xxix. (1900) 447.  
     *HOLOCARPA*, Hemsl. in Kew Bull. (1895) 15.  
*STAURANTHERA CHIRITÆFLORA*, Oliver, in Hook. Ic. Pl. xxv. (1896) t. 2454.  
*STELLARIA DELAVAYI*, Franch. Pl. Delav. (1889) 97.  
     ✓ *DICHASIOIDES*, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 435.  
     *EBRACTEATA*, Komarov, in Act. Hort. Petrop. xviii. (1901) 441.  
     ✓ *FLORIDA*, Fisch.; Engl. Jahrb. xxix. (1900) 320.  
     ✓ *HENRYI*, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 434.  
     *INFRACTA*, Maxim. in Act. Hort. Petrop. xi. 1. (1890) 72.  
     *LONGIFOLIA*, Muhl.; Komarov, in Act. Hort. Petrop. xxii. (1903) 170.  
     ✓ *NUTANS*, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 434.  
     *PILOSA*, Franch. Pl. Delav. (1889) 99.  
     *RADIANS*, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 168.  
 ✓ *SOULIEI*, Williams, in Journ. Linn. Soc., Bot. xxxiv. (1899) 434.  
     ✓ *UDA*, Williams, *l. c.* 435.  
     ✓ *WUSHANENSIS*, Williams, *l. c.* 434.  
     *YUNNANENSIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 433.  
 ✓ *STENOPHRAGMA THALIANUM*, Celak.; Engl. Jahrb. xxix. (1900) 358.  
*STEPHANANDRA INCISA*, Zabel, ex Palib. in Act. Hort. Petrop. xvii. (1899) 73.  
*STEREOSANTHES*, Franch. in Journ. de Bot. x. (1896) 384.  
     *DELAVAYI*, Franch. *l. c.* 385.  
     *SOULIEI*, Franch. *l. c.*  
     *YUNNANENSIS*, Franch. *l. c.*  
*STRANVÆSIA DAVIDIANA*, Decne.; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 179.  
     *GLAUCESCENS*, Lindl., var.; Franch. Pl. Delav. (1890) 226.

- STREPTOCARPUS CHINENSIS, Franch. in Bull. Mus. Hist. Nat. Par. v. (1899) 252.
- STREPTOPUS AJANENSIS, Til., var.; Komarov, in Act. Hort. Petrop. xx. (1901) 476.
- PARVIFLORUS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 189.
- STYRAX CALONEURUM, Perk. in Engl. Jahrb. xxxi. (1902) 484.
- DASYANTHUM, Perk. *l. c.* 485.
- FORMOSANUM, Matsum. in Bot. Mag. Tokyo, xv. 75.
- HEMSLEYANUM, Diels, in Engl. Jahrb. xxix. (1900) 530.
- HYPOGLAUCUM, Perk. *l. c.* xxxi. (1902) 486.
- MACRANTHUM, Perk. *l. c.* 487.
- PRUNIFOLIUM, Perk. *l. c.* 486.
- SWERTIA ASARIFOLIA, Franch. in Bull. Soc. Bot. Fr. xlvi. (1900) 311.
- BIFOLIA, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 378.
- CALYCINA, Franch. in Bull. Soc. Bot. Fr. xlvi. (1900) 311.
- DAVIDI, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x (1887-88) 62.
- DECORA, Franch. in Bull. Soc. Bot. Fr. xlvi. (1900) 317.
- DELAVAYI, Franch. *l. c.* 323.
- DIMORPHA, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 379.
- GENTIANOIDES, Franch. in Bull. Soc. Bot. Fr. xlvi. (1900) 314.
- GRACILIS, Franch. *l. c.* 321.
- KOITCHENSIS, Franch. *l. c.* 320.
- LEDUCI, Franch. *l. c.* 316.
- LONGIPES, Franch. *l. c.* 319.
- MACROSPERMA, C. B. Clarke; Engl. Jahrb. xxix. (1900) 539.
- MEMBRANIFOLIA, Franch. in Bull. Soc. Bot. Fr. xlvi. (1900) 319.
- MUSSOTI, Franch. *l. c.* 316.
- PUBESCENS, Franch. *l. c.* 313.
- STRICTA, Franch. *l. c.* 322.
- TASHIROI, Makino, in Bot. Mag. Tokyo, x. (1896) 57.
- SYCOPSIS SINENSIS, Oliver, in Hook. Ic. Pl. xx. (1890) t. 1931.
- SYMPLOCOS ACUTANGULA, Brand, in Engl. Pflanzenr., Symploc. 65.
- ALATA, Brand, in Engl. Jahrb. xxix. (1900) 529.
- ANOMALA, Brand, *l. c.*
- ARGENTEA, Brand, in Engl. Pflanzenr., Symploc. 67.
- BOTRYANTHA, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 60.
- CUSPIDATA, Brand, in Engl. Pflanzenr., Symploc. 69.
- FORMOSANA, Brand, *l. c.* 67.
- GLANDULIFERA, Brand, *l. c.* 68.
- HENBYI, Brand, *l. c.* 67.
- LEUCOPHYLLA, Brand, *l. c.* 60.
- MODESTA, Brand, *l. c.* 66.



- ✓ SYMPLOCOS SETCHUENSIS, Brand, in Engl. Jahrb. xxix. (1900) 528.  
   ✓ SPECTABILIS, Brand, in Engl. Pflanzenr., Symploc. 69.  
   ✓ STELLARIS, Brand, in Engl. Jahrb. xxix. (1900) 528.  
     TASHIROI, Matsum. in Bot. Mag. Tokyo, xv. 77.  
   ✓ TERMINALIS, Brand, in Engl. Pflanzenr., Symploc. 36.  
   ✓ YUNNANENSIS, Brand, *l. c.*
- ✓ SYRINGA MICROPHYLLA, Diels, in Engl. Jahrb. xxix. (1900) 531.  
   PUBESCENS, var., Batalin, in Act. Hort. Petrop. xiii. [18] (1894)  
     378.  
   TOMENTELLA, Bur. & Franch. in Journ. de Bot. v. (1891) 103.  
   VELUTINA, Bur. & Franch. in Rev. Hort. (1891) 308.  
   VELUTINA, Komarov, in Act. Hort. Petrop. xviii. (1901) 428.  
   YUNNANENSIS, Franch. in Rev. Hort. (1891) 308.
- ✓ TAMARIX PALLASII, Desv.; Engl. Jahrb. xxix. (1900) 476.  
   TANACETUM MYRIANTHUM, Franch. in Bull. Soc. Phil. Par. sér. 8, iii. 3.  
     (1891) 144.  
   ✓ PALLASIANUM, Trautv. & Mey.; Engl. Jahrb. xxix. (1900) 617.
- ✓ TAPISCIA SINENSIS, Oliver, in Hook. Ic. Pl. xx. (1891) t. 1928.  
   TERMINALIA TRIPTERA, Franch. in Journ. de Bot. x. (1896) 291.  
   TETRACENTRON SINENSE, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1892.
- ✓ TETRASTIGMA HEMSLEYANUM, Diels & Gilg, in Engl. Jahrb. xxix. (1900)  
   463.  
   HYPOGLAUCUM, Planch.; Franch. in Bull. Soc. Bot. Fr. xxxiii.  
     (1886) 433.  
   ✓ OBTECTUM, Wall.; Engl. Jahrb. xxix. (1900) 463.
- ✓ TEUCRIUM BIDENTATUM, Hemsl., var.; Diels, in Engl. Jahrb. xxix.  
   (1900) 552.  
   ✓ FRANCHETIANUM, Diels, *l. c.* 551.
- ✓ THALICTRUM AKANENSE, Huth; H. Lévillé, in Bull. Acad. Géog. Bot.  
   xi. (1902) 297.  
   AMPLISSIMUM, H. Lévillé & Vaniot, *l. c.* 51.
- ✓ ANGUSTIFOLIUM, Jacq.; Diels, in Engl. Jahrb. xxix. (1900) 335.  
   AQUILEGIFOLIUM, Linn.; Komarov, in Act. Hort. Petrop. xxii.  
     (1903) 303.  
   CLEMATIDIFOLIUM, Franch. in Journ. de Bot. viii. (1894) 273.  
   CORAIENSE, Matsum. in Bot. Mag. Tokyo, ix. (1895) 276.  
   COREANUM, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi.  
     (1902) 297.  
   CULTRATUM, Wall.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886)  
     369.  
   DELAVAYI, Franch. *l. c.* 367.  
   DIPTEROCARPUM, Franch. *l. c.* 368.  
   GRANDIFLORUM, Maxim. in Act. Hort. Petrop. xi. i. (1890) 11.

- THALICTRUM GRANDISEPALUM**, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 297.  
**HAMATUM**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 16.  
**ACHANGENSE**, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1765.  
**LECOYERI**, Franch. Pl. Delav. (1889) 16.  
**LEUCONOTUM**, Franch. *l. c.* 15.  
**MACRORHYNCHUM**, Franch. in Journ. de Bot. iv. (1890) 302.  
**MICROGYNUM**, Oliver, in Hook. Ic. Pl. xviii. (1888) t. 1766.  
**OLIGANDRUM**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 16.  
**PALLIDUM**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 187.  
**RETICULATUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 371.  
**ROBUSTUM**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 18.  
**SCABRIFOLIUM**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 369.  
**THIBETICUM**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1889) 186.  
**TRICHOPUS**, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 370.  
**TRIPELTATUM**, Maxim. in Act. Hort. Petrop. xi. i. (1890) 13.  
**TUBERIFERUM**, H. Lévillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 298.  
**UNCATUM**, Maxim. in Act. Hort. Petrop. xii. i. (1890) 14.  
**UNCINULATUM**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 187.  
**VIRGATUM**, Hook. f. & Thoms.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 368.
- THEA = CAMELLIA.**  
**ASSIMILIS**, Seem.; Kochs, in Engl. Jahrb. xxvii. (1900) 584.  
**CUSPIDATA**, Kochs, *l. c.* 586.  
**EDITHÆ**, Kuntze; Kochs, *l. c.* 592.  
**EURYOIDES**, Booth; Kochs, *l. c.* 585.  
**GRIJSII**, Kuntze; Kochs, *l. c.* 591.  
**HONGKONGENSIS**, Pierre; Kochs, *l. c.*  
**PUNCTATA**, Kochs, *l. c.* 584.  
**RETICULATA**, Pierre; Kochs, *l. c.* 595.  
**ROSÆFLORA**, Kuntze; Kochs, *l. c.* 585.  
**SPECIOSA**, Kochs, *l. c.* 597.  
**SPECTABILIS**, Kochs, *l. c.* 595.
- THELASIS ELONGATA**, Blume; Mém. Herb. Boiss. (1900) n. 21, 70.  
**THERMOPSIS ALPINA**, Ledeb.; Franch. Pl. Delav. (1889) 150.  
**THESIUM LONGIFOLIUM**, Turcz.; Komarov, in Act. Hort. Petrop. xxii. (1903) 109.  
**THLADIANTHA DAVIDI**, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 243.  
**GLABRA**, Cogn. in Hook. Ic. Pl. xxiii. (1894) sub t. 2223.  
**LONGIFOLIA**, Cogn. *l. c.* t. 2222.

- *THLADIANTHA MACULATA*, Cogn. in Hook. Ic. Pl. xxiii. (1894) sub t. 2223.  
     *OLIVERI*, Cogn. *l. c.* t. 2223.  
     *VERRUCOSA*, Cogn. *l. c.*  
     ✓ *VILLOSULA*, Cogn.; Diels, in Engl. Jahrb. xxix. (1900) 603.  
*THLASPI YUNNANENSIS*, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 407.
- ✓ *TILIA BARONIANA*, Diels, in Engl. Jahrb. xxix. (1900) 468.  
     *CHINENSIS*, Maxim. in Act. Hort. Petrop. xi. 1. (1890) 83.  
     ✓ *CORDATA*, Mill.; Diels, in Engl. Jahrb. xxix. (1900) 468.  
 — *HENRYANA*, Szyszyl.; Hook. Ic. Pl. xx. (1890) t. 1927.  
     *MIQUELIANA*, Maxim., var., Szyszyl. *l. c.* sub t. 1927.  
     *OLIVERI*, Szyszyl. *l. c.* t. 1927.  
     *PAUCICOSTATA*, Maxim. in Act. Hort. Petrop. xi. (1890) 82.  
 — *TUAN*, Szyszyl. in Hook. Ic. Pl. xx. (1890) t. 1926.
- TOFIELDIA MACILENTA*, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887–8) 95.
- TORRICELLIA ANGULATA*, Oliver, in Hook. Ic. Pl. xix. (1889) t. 1893.  
     ✓ *INTERMEDIA*, Harms, in Engl. Jahrb. xxix. (1900) 507.
- ✓ *TRACHELOSPERMUM AXILLARE*, Hook. f.; Diels, in Engl. Jahrb. xxix. (1900) 540.  
     ✓ *DIVARICATUM*, Thunb.; Diels, *l. c.*
- TRACHYDIUM DAUCOIDES*, Franch. Pl. Dav. in Nouv. Arch. Hist. Nat. Mus. Par. sér. 2, viii. (1885) 245.  
     *DELAVAYI*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 110.  
     *HUSPIDUM*, Franch. *l. c.* 113.  
     *NOVEMJUGUM*, C. B. Clarke, var.; Boissieu, in Bull. Herb. Boiss. sér. 2, ii. (1902) 802.  
     *PURPURASCENS*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 112.  
     *ROYLEI*, Lindl.; Boissieu, Bull. Herb. Boiss. sér. 2, ii. (1902) 803.  
     *RUBINERVE*, Franch. in Bull. Soc. Phil. Par. sér. 8, vi. (1894) 112.  
     *VIRIDIFLORUM*, Franch. *l. c.* 111.
- *TRETOCARYA SIKKIMENSIS*, Oliver, in Hook. Ic. Pl. xxiii. (1893) t. 2255.
- ✓ *TRICHOSANTHES ROSTHORNII*, Harms, in Engl. Jahrb. xxix. (1900) 603.
- TRICYRTIS LASIOCARPA*, Matsum. in Bot. Mag. Tokyo, xi. (1897) 79.  
     *STONOLIFERA*, Matsum. *l. c.* 78.
- TRIOSTEUM FARGESII*, Franch. in Journ. de Bot. x. (1896) 318.  
     ✓ *HIMALAYANUM*, Wall., var.; Diels & Graebn. in Engl. Jahrb. xxix. (1900) 590.  
     ✓ *INTERMEDIUM*, Diels & Graebn. *l. c.*  
     ✓ *ROSTHORNII*, Diels & Graebn. *l. c.* 592.
- TRIPLOSTEGIA GRANDIFLORA*, Gagnep. in Bull. Soc. Bot. Fr. xlvii. (1900) 333.
- TRIUMFETTA ANNUA*, Linn.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 441.

- TRIUMFETTA TOMENTOSA, Bojer; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 441.
- TROLLIUS LEDEBOURII, Reichb.; Komarov, in Act. Hort. Petrop. xxii. (1903) 230.
- ✓ PALMATUS, Hook. f. & Thoms.; Diels, in Engl. Jahrb. xxix. (1900) 324.
- PATULUS, Salisb.; Komarov, in Act. Hort. Petrop. xxii. (1903) 232.
- ✓ PUMILUS, D. Don; Diels, in Engl. Jahrb. xxix. (1900) 324.
- ✓ RANUNCULOIDES, Hemsl. in Journ. Linn. Soc., Bot. xxix. (1893) 301.
- ✓ TUPIDANTHUS CALYPTRATUS, Hook. f. & Thoms.; Hemsl. in Journ. Linn. Soc., Bot. xxxiv. (1900) 476.
- TYLOPHORA LIUKIUENSIS, Matsum. in Bot. Mag. Tokyo, xii. (1898) 40.
- TYPHA MARTINI, H. Lévêillé & Vaniot, in Bull. Acad. Géog. Bot. xi. (1902) 166.
- ULMARIA PALMATA, Focke; Palibin, in Act. Hort. Petrop. xvii. 1. (1899) 82.
- ULMUS PUMILA, Linn.; Komarov, in Act. Hort. Petrop. xxii. (1903) 85.
- URARIA LAGOPUS, DC.; Franch. Pl. Delav. (1890) 171.
- PICTA, Desv.; Franch. *l. c.*
- SINENSIS, Franch. *l. c.* 172.
- ✓ URTICA FISSA, E. Pritz. in Engl. Jahrb. xxix. (1900) 301.
- VACCINIUM DELAVAYI, Franch. in Journ. de Bot. ix. (1895) 367.
- DONIANUM, Franch. *l. c.* 369.
- FRAGILE, Franch. *l. c.* 366.
- HIRTUM, Thunb.; Palibin, in Act. Hort. Petrop. xviii. 11. (1900) 149.
- ✓ LÆTUM, Diels, in Engl. Jahrb. xxix. (1900) 516.
- ✓ MANDARINORUM, Diels, *l. c.*
- MOUPINENSE, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 43.
- PUBICALYX, Franch. in Journ. de Bot. ix. (1895) 369.
- SETOSUM, C. H. Wright, in Kew Bull. (1896) 24.
- YUNNANENSE, Franch. in Journ. de Bot. ix. (1895) 368.
- VALERIANA FABERI, Graebn. in Engl. Jahrb. xxiv. (1898) Beibl. n. 59, 32.
- FLACCIDISSIMA, Maxim. *l. c.* xxix. (1900) 598.
- FLAGELLIFERA, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 374.
- HARMSII, Graebn. in Engl. Jahrb. xxiv. (1898) Beibl. n. 59, 32.
- HELICTES, Graebn. *l. c.* xxix. (1900) 600.
- HIEMALIS, Graebn. *l. c.*
- ROSTHORNII, Graebn. *l. c.* 599.
- TANGUTICA, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 375.

- VANDA PARVIFLORA, Lindl.; Franch. in Bull. Soc. Bot. Fr. xlv. (1898) 413.
- VANIOTIA, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 166.  
MARTINI, H. Lévillé, *l. c.*
- VENTILAGO ELEGANS, Hemsl. in Ann. Bot. ix. (1895) 151.
- VERATRILLA BAILLONI, Franch. in Bull. Soc. Bot. Fr. xlvi. (1899) 311.
- VERBASCUM THAPSUS, Linn.; Franch. in Bull. Soc. Bot. Fr. xlvii. (1900) 10.
- VERNONIA AMPLA, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 124.  
✓BOCKIANA, Diels, in Engl. Jahrb. xxix. (1900) 608.  
FARGESII, Franch. in Journ. de Bot. x. (1896) 368.  
✓FLACCIDISSIMA, Maxim.; Engl. Jahrb. xxix. (1900) 608.  
HARDWICKII, Wall.; Engl. *l. c.*  
✓HENRYI, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 500.  
MARTINI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 124.  
MONOSIS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, x. (1887-8) 34.  
PAPILLOSA, Franch. in Journ. de Bot. x. (1896) 369.  
SEGUINI, Vaniot, in Bull. Acad. Géog. Bot. xii. (1903) 241.  
SPELÆICOLA, Vaniot, *l. c.* 123.  
SUBARBOREA, Vaniot, *l. c.* 126.  
✓SYLVATICA, Dunn, in Journ. Linn. Soc., Bot. xxxv. (1903) 501.
- VERONICA FARGESII, Franch. in Bull. Soc. Bot. Fr. xlvii. (1898) 21.  
PIROLÆFORMIS, Franch. *l. c.* 20.  
SUTCHUENENSIS, Franch. *l. c.*  
SZECHUANICA, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 383.
- VIBURNUM ARCUATUM, Komarov, in Act. Hort. Petrop. xviii. (1901) 427.  
BETULIFOLIUM, Batalin, *l. c.* xiii. [18] (1894) 371.  
✓BOCKII, Graebn. in Engl. Jahrb. xxix. (1900) 585.  
BUDDLEIFOLIA, C. H. Wright, in Gard. Chron. (1903) i. 257.  
CEANOTHOIDES, C. H. Wright, in Kew Bull. (1896) 23.  
✓CHINSHANENSE, Graebn. in Engl. Jahrb. xxix. (1900) 585.  
DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 251.  
✓DIELSII, Graebn. in Engl. Jahrb. xxix. (1900) 588.  
✓ERUBESCENS, Wall.; Graebn. *l. c.* 584.  
✓FALLAX, Graebn. *l. c.* 586.  
KANSUENSE, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 372.  
✓LOBOPHYLLUM, Graebn. in Engl. Jahrb. xxix. (1900) 589.  
OLIGANTHUM, Batalin, in Act. Hort. Petrop. xiii. [18] (1894) 372.  
PALLIDUM, Franch. in Journ. de Bot. x. (1896) 308.  
✓PRATTII, Graebn. in Engl. Jahrb. xxix. (1900) 584.  
✓RECTANGULATUM, Graebn. *l. c.* 588.  
✓ROSTHORNII, Graebn. *l. c.* 586.

- VIBURNUM SARGENTI, Koehne, in Wittm. Gartenfl. (1899) 341.  
 STELLULATUM, Wall. ; Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 251.
- ✓ SYMPODIALE, Graebn. in Engl. Jahrb. xxix. (1900) 587.  
 VEITCHI, C. H. Wright, in Gard. Chron. (1903) i. 257.
- ✓ WILLEANUM, Graebn. in Engl. Jahrb. xxix. (1900) 589.
- VICIA CHINENSIS, Franch. Pl. Delav. (1890) 177.  
 MEGALOTROPIS, Ledeb. ; Franch. in Journ. de Bot. iv. (1890) 305.  
 RAMOSISSIMA, Franch. *l. c.*  
 TENERA, R. Grah. ; Franch. Pl. Delav. (1890) 177.
- VILLEBRUNEA PEDUNCULATA, Shirai, in Bot. Mag. Tokyo, ix. (1895) 158.
- VIOLA ALBIDA, Palibin, in Act. Hort. Petrop. xvii. i. (1899) 30.  
 CAMELEO, Boissieu, in Bull. Herb. Boiss. sér. 2, i. (1901) 1074.  
 DACTYLOIDES, Roem. & Schult. ; Palibin, in Act. Hort. Petrop. xvii. i. (1899) 32.  
 DAVIDI, Franch. Pl. Dav. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 203.  
 DELTOIDEA, Makino ; Boissieu, in Bull. Herb. Boiss. sér. 2, i. (1901) 1081.  
 DELAVAYI, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 413.  
 DENTARLÆFOLIA, Boissieu, in Bull. Herb. Boiss. sér. 2, i. (1901) 1076.  
 DISTANS, Wall. ; Boissieu, *l. c.* 1080.  
 FARGESII, Boissieu, *l. c.* ii. (1902) 333.  
 FLAVIDA, Bur. & Franch. in Journ. de Bot. v. (1891) 21.  
 GMELINIANA, Roem. & Schult. ; Boissieu, in Bull. Herb. Boiss. sér. 2, i. (1901) 1079.  
 HENRYI, Boissieu, *l. c.* 1075.  
 LEVEILLEI, Boissieu, in Bull. Acad. Géog. Bot. xi. (1902) 91.  
 MOUPINENSIS, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 412.  
 PALUSTRIS, Franch. in Nouv. Arch. Mus. Hist. Nat. Par. sér. 2, viii. (1885) 202.  
 PHALACROCARPA, Maxim. Diag. Pl. Nov. As. (1876) 726.  
 ✓ ROSTHORNII, E. Pritz. in Engl. Jahrb. xxix. (1900) 477.  
 STRIATELLA, Boissieu, in Bull. Herb. Boiss. sér. 2, i. (1901) 1077.  
 TUBERIFERA, Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 410.  
 UROPHYLLA, Franch. *l. c.* 413.  
 VAGINATA, Maxim. Diagn. Pl. Nov. As. (1876) 733.
- ✓ VITIS ARMATA, Diels & Gilg, in Engl. Jahrb. xxix. (1900) 462.  
 ✓ BETULIFOLIA, Diels & Gilg, *l. c.* 461.  
 ✓ COIGNETIÆ, Pulliat ; Diels, *l. c.*  
 .FORMOSANA, Hemsl. in Ann. Bot. ix. (1895) 151.  
 ✓ PAGNUCCII, Rom. du Caill. ; Diels, in Engl. Jahrb. xxix. (1900).  
 ✓ PENTAGONA, Diels & Gilg, *l. c.* 460.

- VITIS POTANINI, Maxim. in Act. Hort. Petrop. xi. (1890) 104.  
 ✓ ROMANETI, Rom. du Caill.; Diels, in Engl. Jahrb. xxix. (1900) 462.  
 ✓ TRICHOCLADA, Diels & Gilg, *l. c.* 461.  
 VOINIERIANA, Hort.; Gard. and Forest (1897) 293.
- WAHLENBERGIA BREVIPES, Hemsl. in Hook. Ic. Pl. xxviii. (1903) t. 2768.
- WENDLANDIA HENRYI, Oliver, *l. c.* xviii. (1887) t. 1712.
- WICKSTROEMIA RETUSA, A. Gray; Bull. Acad. Pétersb. xxxi. (1887) 98.  
 ✓ STENOPHYLLA, E. Pritz. in Engl. Jahrb. xxix. (1900) 480.
- XANTHOPAPPUS, Winkl. in Act. Hort. Petrop. xiii. i. (1893) 10.  
 SUBACAULIS, Winkl. *l. c.* 11.
- ✓ XYLANCHE HIMALAICA, Beck; Diels, in Engl. Jahrb. xxix. (1900) 573.
- ZANTHOXYLUM ACANTHOPODIUM, DC.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 450.  
 ARNOTTIANUM, Maxim.; Bot. Mag. Tokyo, ix. (1895) 327.  
 DIMORPHOPHYLLUM, Hemsl. in Ann. Bot. ix. (1895) 150.  
 ECHINOCARPUM, Hemsl. *l. c.*  
 EMARGINELLUM, Hemsl. *l. c.* 149.  
 FRAXINOIDES, Hemsl. *l. c.* 150.  
 MICRANTHUM, Hemsl. *l. c.* 147.  
 ▶ MULTIFOLIOLATUM, Hemsl. in Hook. Ic. Pl. xxvi. (1899) t. 2595.  
 MULTIJUGUM, Franch. Pl. Delav. (1889) 124.  
 OXYPHYLLUM, Edgew.; Franch. in Bull. Soc. Bot. Fr. xxxiii. (1886) 450.  
 PIASEZKII, Maxim. in Act. Hort. Petrop. xi. i. (1890) 93.  
 STENOPHYLLUM, Hemsl. in Ann. Bot. ix. (1895) 147.  
 UNDULATIFOLIUM, Hemsl. *l. c.* 148.
- ZENOBIA CERASIFLORA, H. Lévillé, in Bull. Acad. Géog. Bot. xii. (1903) 253.
- ZINGIBER ATRORUBENS, Gagnep. in Bull. Soc. Bot. Fr. xlix. (1902) 252.  
 ROSEUM, Rosc.; Gagnep. in Bull. Acad. Géog. Bot. xii. (1903) 38.  
 ✓ STRIOLATUM, Diels, in Engl. Jahrb. xxix. (1900) 262.
- ZYGOPHYLLUM LOCZII, Kanitz, Pl. Exped. Szechenyi, Asia Centr. 953.

## INDEX.

[Including the names mentioned in Vols. 23 and 26, with corrections, forming a complete index to the Chinese Flora. A star is added to names of genera, species, or varieties which are here first published.]

- Abelia adenotricha*, Hance, 36, 451.  
*angustifolia*, Bur. & Franch., 36, 451.  
*biflora*, Turcz., 23, 358.  
*chinensis*, R. Br., 23, 358.  
*Davidii*, Hance, 23, 358.  
*Hanceana*, Mart., 23, 358.  
*parvifolia*, Hemsl. \*, 23, 358.  
*rupestris*, Lindl., 23, 358.  
— *var. grandiflora*, Carr., 23, 358.  
*serrata*, Sieb. & Zucc., 23, 359.  
*uniflora*, R. Br., 23, 359.
- Abies brachytyla*, Franch., 26, 553.  
*chensiensis*, Van Tiegh., 26, 557.  
*chinensis*, Franch., 26, 556.  
*Davidiana*, Franch., 26, 554.  
*Delavayi*, Franch., 26, 557.  
*dumosa*, Loud., 26, 556.  
*Fargesii*, Franch., 26, 557.  
— *var. sutchuenensis*, Franch., 26, 557.  
*firma*, Sieb. et Zucc., 26, 557.  
*Fortunei*, Hance, 26, 555.  
*Fortunei*, A. Murr., 26, 555.  
*Griffithiana*, Lindl., 26, 558.  
*holophylla*, Maxim., 26, 557.  
*jezoënsis*, Bertrand, 26, 555.  
*Kaempferi*, Lindl., 26, 557.  
*likiangensis*, Franch., 26, 554.  
*Momi*, Sieb., 26, 557.  
*nephrolepis*, Maxim., 26, 557.  
*sacra*, Franch., 26, 555.  
*Schrenkiana*, Lindl., 26, 554.  
*Tsuga*, Sieb. et Zucc., 26, 556.  
*Veitchii*, Lindl., 26, 557.  
*yunnanensis*, Franch., 26, 556.
- Abildgaardia compressa*, Presl, 36, 240.  
*Eragrostis*, Boeck., 36, 240.  
*Eragrostis*, Hance, 36, 235.  
*fusca*, Nees, 36, 236.  
*javanica*, Steud., 36, 241.  
*monostachya*, Vahl, 36, 240.  
*nervosa*, Presl, 36, 243.  
*pauciflora*, Kunth, 36, 236.  
*Rottboelliana*, Nees, 36, 240.
- Abroma fastuosum*, Jacq., 23, 91.  
*Abrus cantoniensis*, Hance, 23, 187.  
*mollis*, Hance, 23, 187.  
*precatorius*, Linn., 23, 187.
- Absinthium maritimum Sinarum*, Pluk., 23, 440.
- Abutilon Avicennæ*, Gaertn., 23, 86.  
*cysticarpum*, Hance, 23, 86.  
*graveolens*, Seem., 23, 86.  
*indicum*, G. Don, 23, 86.  
*polyandrum*, Schlecht., 36, 451.  
*sinense*, Oliver, 36, 451.
- Acacia concinna*, DC., 23, 215.  
*Delavayi*, Franch., 36, 451.  
*Farnesiana*, Willd., 23, 215.  
*Lebbek*, Willd., 23, 216.  
*macrophylla*, Bunge, 23, 216.  
*Nemu*, Willd., 23, 216.  
*pennata*, Willd., 36, 451.  
*Philippinarum*, Benth., 23, 215.  
*Richii*, A. Gray, 23, 215.  
*yunnanensis*, Franch., 36, 451.
- Acalypha acmophylla*, Hemsl. \*, 26, 436.  
*australis*, Linn., 26, 437.  
*boehmerioides*, Miq., 26, 437.



- Acalypha brachystachya*, Hornem., 26, 437.  
*brachystachya*, Maxim., 26, 478.  
*chinensis*, Benth., 26, 438.  
*chinensis*, Roxb., 26, 437.  
*fallax*, Muell. Arg., 26, 437.  
*gemina*, Spreng., 26, 437.  
*Giraldii*, Pax, 36, 451.  
*hispida*, Wall., 26, 437.  
— var. *pubescens*, Hook. & Arn., 26, 437.  
*indica*, Linn., 26, 438.  
*lanceolata*, Wall., 26, 437.  
*pauciflora*, Hornem., 26, 437.  
*sp.*, Houtt., 26, 488.  
*virgata*, Thunb., 26, 437.
- Acampe multiflora*, Lindl., 26, 36.
- ACANTHACEÆ, 26, 237.
- Acanthopanax aculeatum*, Seem., 23, 339.  
*asperatum*, Franch. & Savat., 23, 339.  
*divaricatum*, Seem., 23, 339.  
*diversifolium*, Hemsl. \*, 23, 340.  
*evodiaefolium*, Franch., 36, 451.  
*Henryi*, Harms, 36, 451.  
*leucorrhizum*, Harms, 36, 451.  
*ricinifolium*, Seem., 23, 490.  
*ricinifolium*, Linn., 23, 340.  
*senticosum*, Harms, 36, 451.  
*Sepium*, Seem., 23, 339.  
*sessiliflorum*, Seem., 23, 340.  
*setchuenense*, Harms, 36, 451.  
*setulesum*, Franch., 36, 451.  
*spinatum*, Miq., mentioned, 23, 339.
- Acanthus ilicifolius*, Linn., 26, 242.
- Acarua chinensis*, Bunge, 23, 459.
- Acer betulifolium*, Maxim., 36, 451.  
*caudatum*, Wall., 36, 451.  
*cordatum*, Pax, 36, 452.  
*cultratum*, Walt., 23, 141.  
*Davidi*, Franch., 36, 452.  
*discolor*, Maxim., 23, 140.  
*dissectum*, Thunb., 23, 141.  
*erianthum*, Schwerin, 36, 452.  
*erosum*, Pax, 36, 452.  
*Fabri*, Hance, 23, 140.  
*Francheti*, Pax, 36, 452.  
*Ginnala*, Maxim., 26, 142.  
*Giraldi*, Pax, 36, 452.  
*Grosseri*, Pax, 36, 452.  
*Henryi*, Pax, 36, 452.  
*Hookeri*, Miq., 23, 140.  
*japonicum*, Thunb., 23, 140.  
*lætum*, C. A. Mey., 23, 140.  
*lævigatum*, Wall., 36, 452.  
*lanceolatum*, Molliard, 36, 452.  
*laxiflorum*, Pax, 36, 452.  
*Lobelii*, Ten., 23, 140; 36, 452.  
*Maximowiczii*, Pax, 36, 452.
- Acer Mono*, Maxim., 23, 141.  
*multiserratum*, Maxim., 36, 452.  
*nikoense*, Maxim., 36, 452.  
*oblongum*, Wall., 23, 141.  
*Oliverianum*, Pax, 36, 452.  
*palmatum*, Thunb., 23, 141.  
*Paxii*, Franch., 36, 452.  
*pictum*, Thunb., 23, 141.  
*pilosum*, Maxim., 23, 141.  
*polymorphum*, Sieb. & Zucc., 23, 141.  
*reticulatum*, Champ., 23, 141.  
*robustum*, Pax, 36, 452.  
*rufinerve*, Sieb., 23, 141.  
*Schœnermarkiæ*, Pax, 36, 452.  
*septemlobum*, Thunb., 23, 141.  
*sessilifolium*, Sieb. & Zucc., 23, 141.  
*Sieboldianum*, Miq., var., 23, 140.  
*sinense*, Pax, 36, 452.  
*sterculiaceum*, Wall., 23, 142.  
*sutchuenense*, Franch., 36, 452.  
*tataricum*, Linn., 23, 142.  
*tenellum*, Pax, 36, 452.  
*tetramerum*, Pax, 36, 452.  
*trifidum*, Hook., 23, 142.  
*trifidum*, Thunb., 23, 142.  
*triflorum*, Komarov, 36, 452.  
*truncatum*, Bunge, 23, 142; mentioned, 141.  
*urophyllum*, Maxim., 36, 452.  
*Veitchii*, Hort., 36, 452.  
*villosum*, Wall., 23, 142.
- Aceranthus sagittatus*, Sieb. & Zucc., 23, 33.
- Acerus angustifolia*, Lindl., 36, 50.  
*longicruris*, C. Wright, 36, 51.
- Aceriphyllum Rossii*, Engl. & Prantl, 36, 452.
- Achillea alpina*, Linn., 23, 437.  
*Millefolium*, Linn., 23, 436.  
*mongolica*, Fisch., 23, 437.  
*ptarmicoides*, Maxim., 23, 436.  
*sibirica*, Ledeb., 23, 437.
- Achlys japonica*, Maxim. & Komarov, 36, 452.
- Achudemia japonica*, Maxim., 26, 480.
- Achyranthes aspera*, Linn., 26, 321.  
*bidentata*, Blume, 26, 322.
- Achyrophorus aurantiacus*, DC., 23, 478.  
*grandiflorus*, Ledeb., 23, 478.
- Aclisia sorzogonensis*, E. Meyer, 36, 159.
- Acmena Championi*, Benth., 23, 296.  
*Championi*, Seem., 23, 30?
- Aconitum albo-violaceum*, Komarov, 36, 452.
- Anthora*, Linn., 23, 20.

- Aconitum autumnale*, Lindl., 23, 20.  
*barbatum*, Patr., 23, 21.  
*Bodinieri*, H. Léveillé & Vaniot, 36, 452.  
*Carmichaeli*, Debeaux, 23, 20.  
*chinense* [Paxt.], Bot. Mag., 23, 20.  
*Cynoctonum*, Reichb., 23, 21.  
*Delavayi*, Franch., 36, 452.  
*Fischeri*, Reichb., 23, 20.  
*Fortunei*, Hemsl. \*, 23, 20.  
*gymnandrum*, Maxim., 23, 20.  
*Hemsleyanum*, E. Pritz., 36, 452.  
*Henryi*, E. Pritz., 36, 452.  
*japonicum*, Decne., 23, 20.  
*Kusnezoffii*, Reichb., 23, 20.  
*Lamarckii*, Reichb., 23, 21.  
*luteum*, H. Léveillé & Vaniot, 36, 452.  
*Lycoctonum*, Linn., 23, 20.  
*Napellus*, Linn., 23, 21.  
*ochotense*, Reichb., 23, 21.  
*palmatum*, D. Don, 36, 452.  
*racemosum*, Franch., 36, 452.  
*rotundifolium*, Kar. & Kir., 36, 452.  
*scaposum*, Franch., 36, 452.  
*sinense*, Sieb., 23, 20.  
*umbrosum*, Komarov, 36, 452.  
*vaginatum*, E. Pritz., 36, 452.  
*villosum*, Maxim., 23, 20.  
*volubile*, Pall., 36, 452.  
*Vulparia*, Reichb., 23, 21.  
*Acorus Calamus*, Linn., 36, 187.  
*Calamus*, Lour., 36, 187.  
— var. *terrestris*, Engl., 36, 187.  
*cochinchinensis*, Schott, 36, 187.  
*gramineus*, Ait., 36, 187.  
*Tatarinowii*, Schott, 36, 187.  
*terrestris*, Spreng., 36, 187.  
*Acrocephalus capitatus*, Benth., 26, 269.  
*Acroglochin chenopodioides*, Schrad., 26, 323.  
*persicarioides*, Moq., 26, 323.  
*Acronychia Cyminosma*, F. Muell., 23, 109.  
*laurifolia*, Blume, 23, 108.  
*Acroptilon Picris*, DC., 23, 470.  
*Actæa aspera*, Lour., 23, 22.  
*davurica*, Franch., 23, 21.  
*erythrocarpa*, Fisch., 36, 453.  
*spicata*, Linn., 23, 21; 36, 453.  
*Actinidia arguta*, Planch., 23, 78.  
*callosa*, Lindl., 23, 78.  
*Championi*, Benth., 23, 78.  
*chinensis*, Planch., 23, 78.  
*Davidii*, Franch., 23, 78.  
*eriantha*, Benth., 23, 78.  
*fulvicoma*, Hance, 23, 78.  
*Kolomikta*, Rupr., 23, 78.  
*Actinidia lanata*, Hemsl., 36, 453.  
*melanandra*, Franch., 36, 453.  
*rufa*, Franch. & Sav., mentioned, 23, 78.  
*tetramera*, Maxim., 36, 453.  
*trichogyne*, Franch., 36, 453.  
*Actinodaphne angustifolia*, Benth., 26, 378.  
*angustifolia*, Seem., 26, 378.  
*chinensis*, Nees, 26, 379.  
*lanceifolia*, Meisn., 26, 383.  
*pedunculata*, Meisn., 26, 383.  
*Actinoschænus filiformis*, Benth., 36, 230.  
*sp.*, Benth., 36, 230.  
*Actinospora davurica*, Turcz., 23, 21.  
*Actinostemma biglandulosa*, Hemsl., 36, 453.  
*japonicum*, Miq., 23, 320.  
*lobatum*, Maxim., 23, 320.  
*multilobum*, Harms, 36, 453.  
*paniculatum*, Maxim., 23, 320.  
*racemosum*, Maxim., 23, 320.  
*tenerum*, Griff., 23, 320.  
*tenerum*, Naud., 23, 320.  
*Actinotinus sinensis*, Oliver [a fraudulent imposture of the Collector], 23, 357.  
*Adamia chinensis*, Gard. & Champ., 23, 275.  
*cyanea*, Wall., 23, 275.  
*versicolor*, Fortune, 23, 275.  
*Adenantha pavonina*, Linn., 23, 214.  
*Adenocaulon adhærescens*, Maxim., 23, 432.  
*bicolor*, Hook., 23, 432.  
*himalaicum*, Edgew., 23, 432.  
*Adenophora aurita*, Franch., 36, 453.  
*Boeckiana*, Diels, 36, 453.  
*capillaris*, Hemsl. \*, 26, 10.  
*communis* Fisch., 26, 12.  
*coronopifolia*, Fisch., 26, 11.  
*denticulata*, Fisch., 26, 11.  
*divaricata*, Franch. & Sav., 26, 11.  
*Gmelinii*, Fisch., 26, 11.  
*Isabellæ*, Hemsl., 26, 13.  
*jasionifolia*, Franch., 36, 453.  
*Lamarkii*, Fisch., 26, 11.  
*latifolia*, Fisch., 26, 11.  
*marsupiflora*, Fisch., 26, 11; 36, 453.  
*polymorpha*, Ledeb., 26, 11.  
— var. *latifolia*, Herder, 26, 14.  
*polymorpha*, Ledeb., 26, 12.  
*Potanini*, Korsh., 36, 453.  
*pubescens*, Hemsl. \*, 26, 12.  
*remotidens*, Hemsl. \*, 26, 12.  
*remotiflora*, Miq., 26, 13.  
*rupicola*, Hemsl. \*, 26, 13.

- Adenophora sinensis*, A. DC., 26, 13.  
*stenophylla*, Hemsl. \*, 26, 10.  
*stricta*, Miq., 26, 13.  
*trachelioides*, Maxim., 26, 13.  
— var. *angustifolia*, Debeaux, 26, 13.  
— var. *cordatifolia*, Debeaux, 26, 13.  
*verticillata*, Fisch., 26, 12.  
*verticillata*, Fisch., 26, 14.  
*Wawreana*, Zahlbr., 36, 453.  
*Adenosacme longifolia*, Wall., var. *sinensis*, Hemsl. \*, 23, 379.  
*Adenosma capitata*, Benth., 26, 184.  
*grandiflora*, Benth., 26, 185.  
*microcephala*, Hook., 26, 185.  
*Adenostemma tinctorium*, Cass., 23, 403.  
*viscosum*, Forst., 23, 403.  
*Adhatoda chinensis*, Benth., 26, 244.  
*quadrifaria*, Nees, 26, 246.  
*Vasica*, Nees, 26, 244.  
*Adike pumila*, Rafin., 26, 478.  
*Adina globifera*, Miq., 23, 370.  
*globiflora*, Salisb., 23, 370.  
*peduncularis*, DC., 23, 370.  
*polycephala*, Benth., 23, 371.  
*racemosa*, Miq., 23, 370.  
*rubella*, Hance, 23, 371.  
*sp.*, 23, 371.  
*Adinandra Bockiana*, E. Pritz., 36, 453.  
*Drakeana*, Franch., 23, 76.  
*Millettii*, Benth., 23, 76.  
*Adlumia cirrhosa*, Rafin., 36, 453.  
*Adonis amurensis*, Regel & Radde, 36, 453.  
*apennina*, Linn., 23, 12.  
— var., 23, 12.  
*brevistyla*, Franch., 36, 453.  
*cærulea*, Maxim., 23, 12.  
*Davidi*, Franch., 36, 453.  
*davurica*, Ledeb., 23, 12.  
*Delavayi*, Franch., 36, 453.  
*sutchuenensis*, Franch., 36, 453.  
*Adoxa Moschatellina*, Linn., 23, 347.  
*Æchmandra indica*, Arn., 23, 319.  
*odorata*, Hook. f. & Thoms., 23, 319.  
*Ægiceras fragrans*, König, 26, 67.  
*majus*, Gärtn., 26, 67.  
*Æginetia indica*, Roxb., 26, 220.  
*Ægle sepiaria*, DC., 23, 111.  
*Ægopodium alpestre*, Ledeb., 36, 453.  
*Aërides difforme*, Wall., 36, 38.  
*Hystrix*, Lindl., 36, 38.  
*latifolium*, Thw. Enum., 36, 34.  
*odoratum*, Lour., 36, 35.  
*paniculatum*, Ker-Gawl., 36, 37.  
*Thunbergii*, Miq., 36, 39.  
*Ærua scandens*, Wall., 26, 321.  
*Æschynanthus acuminatus*, Wall., 26, 224.  
*apicidens*, Hance, 26, 224.  
*bracteatus*, Benth., 26, 224.  
*buxifolius*, Dunn, 36, 453.  
*chinensis*, Gardn., 26, 224.  
*humilis*, Dunn, 36, 453.  
*Æschynomene indica*, Linn., 23, 170.  
*Sesban*, Jacq., 23, 162.  
*Æsculus chinensis*, Bunge, 23, 139.  
*indica*, Colebr., 36, 453.  
*turbinata*, Blume, 23, 139.  
*Agalma octophyllum*, Seem., 23, 342.  
*Aganosma Edithæ*, Hance, 26, 100; 36, 453.  
*elegans*, A. DC., 26, 100.  
*lævis*, Champ., 26, 99.  
*Agapetes parviflora*, Dunn, 36, 453.  
*vaccinioides*, H. Léveillé, 35, 453.  
*vaccinoides*, Dunn, 36, 453.  
*yunnanensis*, Franch., 36, 453.  
*Agave americana*, Linn., mentioned, 36, 90.  
*lurida*, Ait., mentioned, 36, 90.  
*Ageratum ciliare*, Lour., 23, 403.  
*conyzoides*, Linn., 23, 403.  
*Aglaia odorata*, Lour., 23, 114.  
*Aglaonema acutispalum*, N. E. Brown, 36, 184, 185.  
*marantifolium*, Engl., 36, 185.  
*modestum*, Schott, 36, 184; mentioned, 185.  
*Agrimonia Eupatoria*, Linn., 23, 246.  
*pilosa*, Ledeb., 23, 246.  
*viscidula*, Bunge, 23, 246.  
*Agriophyllum arenarium*, Bieb., 26, 327.  
*squarrosus*, Moq., 26, 327.  
*Agropyrum caninum*, Beauv., 36, 431.  
*ciliare*, Franch., 36, 432.  
*Pseudo-agropyrum*, Franch., 36, 432.  
*repens*, Beauv., 36, 432.  
*semicostatum*, Nees, 36, 432.  
— var. *ciliare*, 36, 432.  
*strigosum*, Franch., 36, 433.  
*Agrostemma Banksia*, Meerb., 23, 66.  
*Bungeana*, D. Don, 23, 65.  
*fulgens*, Doell, 23, 66.  
*grandiflora*, Doell, 23, 66.  
*Agrostis alba*, Baker & Moore, 36, 390.  
*alba*, Linn., 36, 389.  
*canina*, Linn., 36, 389.  
*diandra*, Retz., 36, 387.  
*Hugoniana*, Rendle \*, 36, 389.  
*inæquiglumis*, Griseb., mentioned, 36, 390.

- Agrostis indica*, Linn., 36, 388.  
*japonica*, Steud., 36, 389.  
*laxiflora*, R. Br., 36, 391.  
*micrantha*, Steud., 36, 390.  
*myriantha*, Hook., 36, 390.  
*perennans*, Tuckerm., 36, 390.  
*scabra*, Willd., 36, 391.  
*virginica*, Linn., 36, 389.  
*vulgaris*, With., var. *ligularis*,  
*Franch.*, 36, 391.  
*Agrostophyllum formosanum*, Rolfe,  
36, 21.  
*Agyneia bacciformis*, A. Juss., 26,  
420.  
*impubes*, Linn., 26, 425.  
*pinnata*, Miq., 26, 425.  
*pubera*, Linn., 26, 425.  
*sinica*, Miq., 26, 425.  
*Aibikkia*, C. B. Clarke, 36, 258.  
sphalm. = *Albikia*, Presl.  
*Ailantus flavescens*, Carr., 23, 114.  
*glandulosa*, Desf., 23, 112.  
*malabarica*, DC., 23, 112.  
*Ainsliæa acerifolia*, Sch.-Bip., 23,  
470.  
*affinis*, Miq., 23, 470.  
*elegans*, Hemsl., 36, 453.  
*fragrans*, Champ., 23, 470; men-  
tioned, 471.  
*glabra*, Hemsl. \*, 23, 471, 492.  
*gracilis*, Franch., 36, 453.  
*grossedentata*, Franch., 36, 453.  
*Henryi*, Diels, 36, 453.  
*lancifolia*, Franch., 36, 453.  
*nervosa*, Franch., 36, 453.  
*ovalifolia*, Vaniot, 36, 454.  
*pertyoides*, Franch., 23, 471.  
*pteropoda*, DC., 23, 471.  
*ramosa*, Hemsl. \*, 23, 471.  
*rubrifolia*, Franch., 36, 454.  
*scabrida*, Dunn, 36, 454.  
*sparsiflora*, Vaniot, 36, 454.  
*spicata*, Vaniot, 36, 454.  
*sutchuenensis*, Franch., 36, 454.  
*undulata*, Diels, 36, 454.  
*Walkeri*, Hook., 23, 472.  
*yunnanensis*, Franch., 23, 472.  
*Aira cæspitosa*, Linn., 36, 399.  
*Ajuga bracteosa*, Wall., 26, 314;  
mentioned, 314.  
*ciliata*, Bunge, 26, 314.  
— var. *glabrescens*, Hemsl. \*, 26,  
315.  
*decumbens*, Thunb., 26, 315.  
*genevensis*, Linn., 26, 315.  
*japonica*, Miq., 26, 295.  
*lupulina*, Maxim., 26, 315.  
*multiflora*, Bunge, 26, 315.  
*ovalifolia*, Bur., 36, 454.  
*remota*, Benth., 26, 315.  
*Ajuga remota*, A. Gray, 26, 315.  
*Akebia lobata*, Decne., 23, 30; 36,  
454.  
*longeracemosa*, Matsum., 36,  
454.  
*quinata*, Decne., 23, 30.  
*Alangium decapetalum*, Lam., 23,  
344.  
*Faberi*, Oliver, 36, 454.  
*hexapetalum*, Lam., 23, 344.  
*Lamarckii*, Thw., 23, 344.  
*Albikia schanoides*, Presl, 36, 258.  
*scirpoides*, Presl, 36, 258.  
*Albizia bracteata*, Dunn, 36, 454.  
*Championi*, Benth., 23, 217.  
*Julibrissin*, Durazz., 23, 216.  
*Lebbek*, Benth., 23, 216.  
*Milletti*, Benth., 23, 216.  
*odoratissima*, Benth., 36, 454.  
*stipulata*, Benth., 23, 215.  
*stipulata*, Boiv., 23, 216.  
*Albucca Gardeni*, Hook., 36, 112.  
*Alcea rosea*, Linn., 23, 83.  
*Alchornea Davidi*, Franch., 26, 438.  
*javensis*, Muell. Arg., 26, 438.  
*rufescens*, Franch., 26, 438.  
*rugosa*, Muell. Arg., 26, 438.  
*trewioides*, Muell. Arg., 26, 438.  
*Alectra indica*, Benth., 26, 201.  
*Aletris chinensis*, Lam., 36, 118.  
*Delavayi*, Franch., 36, 77.  
*foliosa*, var. *sikkimensis*, Franch.,  
36, 75.  
*glabra*, Bur., 36, 75.  
*glandulifera*, Bur., 36, 75.  
*japonica*, Lamb., 36, 76.  
— var. *Fargesii*, C. H. Wright \*,  
36, 76.  
*lactiflora*, Franch., 36, 76.  
*lanuginosa*, Bur., 36, 76.  
*laxiflora*, Bur., 36, 76.  
*nepalensis*, Hook. f., partim, 36, 76.  
*nepalensis*, Hook. f., var. *Delavayi*,  
Franch., 36, 77.  
*revoluta*, Franch., 36, 77.  
*sikkimensis*, Hook. f., 36, 75.  
*spicata*, Franch., 36, 76.  
*stenoloba*, Franch., 36, 77.  
*Aleurites cordata*, Steud., 26, 433.  
*cordifolia*, Steud., 26, 434.  
*japonica*, Blume, 26, 433.  
*moluccana*, Willd., 26, 434.  
*triloba*, Forst., 26, 434.  
*Vernicia*, Hassk., 26, 433.  
*verniciflua*, Baill., 26, 433.  
*Alisma cordifolia*, Thunb., 36, 189.  
*flava*, Thunb., 36, 189.  
*Plantago*, Linn., 36, 189.  
— var. *angustifolium*, Kunth, 36,  
189.

- Alisma Plantago*, var. *latifolium*,  
*Kunth*, **36**, 189.  
 ALISMACEÆ, **36**, 189.  
*Allamanda cathartica*, *Linn.*, **26**,  
 94.  
*Alliaria auriculata*, *Komarov*, **36**,  
 454.  
*Allium angulosum*, *Ker-Gawl.*, **36**,  
 125.  
   *anisopodium*, *Ledeb.*, **36**, 126.  
   *ascalonicum*, *Linn.*, mentioned, **36**,  
   120.  
   *baicalense*, *Willd.*, **36**, 125.  
   *Bakeri*, *Regel*, **36**, 120.  
   *Bouddhæ*, *Debeaux*, **36**, 120.  
   *Cepa*, *Linn.*, **36**, 120.  
   *chinense*, *G. Don*, **36**, 120.  
   *chrysocephalum*, *Regel*, **36**, 121.  
   *condensatum*, *Turcz.*, **36**, 121.  
   *cyaneum*, *Regel*, **36**, 122.  
   — var. *β. brachystemon*, *Regel*,  
   **36**, 123.  
   *cyathophorum*, *Franch.*, **36**, 122.  
   *ellipticum*, *Wall.*, **36**, 126.  
   *exsertum*, *Baker*, **36**, 120.  
   *Fetisowii*, *Regel*, **36**, 122.  
   *fistulosum*, *Linn.*, **36**, 122.  
   *flavovirens*, *Regel*, **36**, 122.  
   *Grayi*, *Regel*, **36**, 121.  
   *Griffithianum*, *Boiss.*, mentioned,  
   **36**, 126.  
   *Henryi*, *C. H. Wright*, **36**, 122;  
   mentioned, 126.  
   *Herderianum*, *Regel*, **36**, 122.  
   *japonicum*, *Regel*, **36**, 122.  
   *jeholense*, *Franch.*, **36**, 123.  
   *kansuense*, *Regel*, **36**, 123.  
   *lineare*, *Linn.*, **36**, 123.  
   *longistylum*, *Baker*, **36**, 124.  
   *macrostemon*, *Bunge*, **36**, 123.  
   *monadelphum*, *Turcz.*, var. *tibeti-*  
   *cum*, *Regel*, **36**, 123.  
   *Nereidum*, *Hance*, **36**, 121.  
   *nerinifolium*, *Baker*, **36**, 120.  
   *odorum*, *Linn.*, **36**, 123.  
   *odorum*, *Lour.*, **36**, 121.  
   *plantagineum*, *Lam.*, **36**, 126.  
   *polyrhizum*, *Turcz.*, **36**, 124.  
   *Pratti*, *C. H. Wright*\*, **36**, 124.  
   *Przewalskianum*, *Regel*, **36**, 124.  
   — var. *planifolium*, *Regel*, **36**,  
   122.  
   *sacculiferum*, *Maxim.*, **36**, 124.  
   *sativum*, *Linn.*, **36**, 124.  
   *Semenovi*, *Regel*, **36**, 125.  
   *senescens*, *Linn.*, **36**, 125.  
   *senescens*, *Miq.*, **36**, 123.  
   *splendens*, *Miq.*, **36**, 120.  
   *sp.*, *Regel*, **36**, 127.  
   *Stevenii*, *Willd.*, **36**, 121.  
   *Allium Stoliczki*, *Regel*, mentioned,  
   **36**, 124.  
   *subangulatum*, *Regel*, **36**, 125.  
   *tanguticum*, *Regel*, **36**, 125; men-  
   tioned, 127.  
   *tehefouense*, *Debeaux*, **36**, 125.  
   *tenuissimum*, *Linn.*, **36**, 125.  
   — var. *anisopodium*, *Regel*, **36**,  
   125.  
   *Thunbergii*, *G. Don*, **36**, 120.  
   *Thunbergii*, *Franch.*, **36**, 120.  
   — var. *Davidii*, *Franch.*, **36**, 120.  
   *triquetrum*, *Lour.*, **36**, 121.  
   *tuberosum*, *Roxb.*, **36**, 123.  
   *uliginosum*, *G. Don*, **36**, 123.  
   *uratense*, *Franch.*, **36**, 126.  
   *venustum*, *C. H. Wright*\*, **36**,  
   126.  
   *Victorialis*, *Linn.*, **36**, 126.  
   — var. *angustifolia*, *Hook. f.*, **36**,  
   126.  
   *weichanicum*, *Palibin*, **36**, 127.  
   *Zimmermannianum*, *Gily*, **36**, 454.  
*Allmania nodiflora*, *R. Br.*, **26**, 321.  
*Allomorpha pauciflora*, *Benth.*, **26**,  
 301.  
*Allophylus Cobbe*, *Blume*, **23**, 138.  
*Alniphyllum*, *Matsum.*, **36**, 454.  
   *macranthum*, *Perk.*, **36**, 454.  
   *pterospermum*, *Matsum.*, **36**,  
   454.  
*Alnus cremastogyne*, *Burkill*\*, **26**,  
 499.  
   *crispa*, *C. Koch*, **26**, 500.  
   *fruticosa*, *Rupr.*, **36**, 454.  
   *hirsuta*, *Turcz.*, **36**, 454.  
   *japonica*, *Sieb. & Zucc.*, **26**, 499.  
   *maritima*, *Henry*, **26**, 500.  
   *maritima*, *Nutt.*, **26**, 499; **36**,  
   454.  
   — var. *arguta*, *Regel*, **26**, 500.  
   — var. *formosana*, *Burkill*\*, **26**,  
   500.  
   — var. *japonica*, *Regel*, **26**, 499.  
   — var. *japonica*, *Regel*, partim,  
   **26**, 499, 500.  
   *nepalensis*, *D. Don*, **26**, 500.  
   *setchuanensis*, *David*, **26**, 500.  
   *sinensis*, *David*, **26**, 500.  
   *sinensis*, *Fauvel*, **26**, 500.  
   *viridis*, *DC.*, **26**, 500.  
   — var. *julacea*, *Franch.*, **26**, 500.  
*Alocasia commutata*, *Schott*, **36**, 184.  
   *cucullata*, *Schott*, **36**, 183.  
   *hainanica*, *N. E. Br.*\*, **36**, 183.  
   *macrorrhiza*, *Schott*, **36**, 184.  
   *odora*, *C. Koch*, **36**, 184.  
*Aloe barbadensis*, var. *chinensis*, *Haw.*,  
**36**, 117.  
   *chinensis*, *Baker*, **36**, 117.

- Alopecurus æqualis*, Sobol., 36, 384.  
*agrestis*, Linn., 36, 385.  
*agrestis*, Pilger, 36, 385.  
*aristulatus*, Michx., 36, 384.  
*fulvus*, Sm., 36, 384.  
*geniculatus*, Linn., 36, 384.  
*hordeiformis*, Lour., 36, 343.  
*hordeiformis*, Osbeck, 36, 338.  
*japonicus*, Steud., 36, 385.  
*malacostachyus*, A. Grav, 36, 385.  
*Alphonsea mollis*, Dunn, 36, 454.  
*Alpinia alba*, Roxb., 36, 70.  
*calcarata*, Rose., 36, 71.  
*cernua*, Sims, 36, 71.  
*chinensis*, Rose., 36, 71.  
*formosana*, K. Schum., 36, 454.  
*Galanga*, Willd., 36, 71; mentioned, 70.  
*globosa*, Horan., 36, 72.  
*japonica*, Miq., 36, 72.  
*nutans*, Rose., 36, 72.  
*officinarum*, Hance, 36, 72.  
*oxyphylla*, Miq., 36, 73.  
*pumila*, Hook., 36, 73.  
*speciosa*, K. Schum., 36, 454.  
*stachyodes*, Hance, 36, 73.  
*viridiflora*, Griff., 36, 72.  
*Alseodaphne breviflora*, Benth., 26, 374.  
*chinensis*, Champ., 26, 374.  
*Alsine laricina*, Crantz, 36, 454.  
*Alsomitra cissoides*, Roem., 23, 321.  
*graciliflora*, Harms, 36, 454.  
*Alstonia macrophylla*, Wall., 26, 96.  
*sp.*, 26, 95.  
*Alternanthera denticulata*, R. Br., 26, 322.  
*nodiflora*, R. Br., 26, 322.  
*sessilis*, R. Br., 26, 322.  
*Althæa chinensis*, Wall., 23, 83.  
*rosea*, Cav., mentioned, 23, 83.  
*sinensis*, Cav., 23, 83.  
*Altingia chinensis*, Benth., 23, 291.  
*Alysicarpus bupleurifolius*, DC., 23, 178.  
*nummulariæfolius*, DC., 23, 179.  
*rugosus*, DC., 23, 179.  
*vaginalis*, DC., 23, 179.  
*Alyssum incanum*, Linn., 23, 41.  
*Alyxia sinensis*, Champ., 26, 95.  
**AMARANTACEÆ**, 26, 318.  
*Amarantus Alopecurus*, Hochst., 26, 319.  
*ascendens*, Hornem., 26, 319.  
*Blitum*, Linn., 26, 319.  
— var. *sylvestris*, Moq., 26, 319.  
*caudatus*, Linn., 26, 319.  
*cruentus*, Linn., 26, 320.  
*gangeticus*, Linn., 26, 319.  
*mangostanus*, Linn., 26, 320.  
*Amarantus melancholicus*, Linn., 26, 319.  
— var. *tricolor*, Moq., 26, 319.  
*oleraceus*, Linn., 26, 319.  
*paniculatus*, Linn., 26, 320.  
*polygamus*, Linn., 26, 320.  
*retroflexus*, Linn., 26, 320.  
*sanguineus*, Linn., 26, 320.  
*spinosus*, Linn., 26, 320.  
*sylvestris*, Desf., 26, 319.  
*tricolor*, Linn., 26, 319.  
*tristis*, Linn., 26, 319.  
*viridis*, Linn., 26, 320.  
**AMARYLLIDEE**, 36, 86.  
*Amaryllis aurea*, L'Hérit., 36, 89.  
*radiata*, L'Hérit., 36, 89.  
*sarniensis*, Thunb., 36, 89.  
*Amberboa moschata*, DC., 23, 470.  
*Ambroma angusta*, Linn., 23, 92.  
*Amelanchier asiatica*, Endl., 36, 454.  
*racemosa*, Lindl., 23, 228.  
*Ameletia indica*, DC., 23, 303.  
*subspicata*, Benth., 23, 303.  
*Amethystea cærulea*, Linn., 26, 310.  
*Ammannia auriculata*, Willd., 23, 304.  
*baccifera*, Linn., 23, 302.  
*indica*, Lam., 23, 303.  
*octandra*, Linn., 23, 303.  
*pentandra*, Roxb., 23, 303.  
*peploides*, Spreng., 23, 303.  
*repens*, Rottl., 23, 303.  
*rotundifolia*, Ham., 23, 303.  
*senegalensis*, Lam., 23, 304.  
*subspicata*, Benth., 23, 303.  
*verticillata*, Lam., 23, 303.  
*vesicatoria*, Roxb., 23, 303.  
*Ammodenia peploides*, Linn., 36, 454.  
*Amomum Curcuma*, Jacq., 36, 69.  
*Galanga*, Lour., 36, 72.  
*globosum*, Lour., 36, 72.  
*medium*, Lour., 36, 70.  
*vittatum*, Hance, 36, 73.  
*Zerumbet*, Linn., 36, 70.  
*Zingiber*, Linn., 36, 70.  
*Amoora Rohituka*, Wight et Arn., 23, 114.  
*Amorphophallus campanulatus*, Decne., 36, 181.  
*Henryi*, N. E. Br. \*, 36, 181.  
*hirtus*, N. E. Br. \*, 36, 181.  
*Konjac*, K. Koch, 36, 182.  
*Rivieri*, Durieu, 36, 182.  
*variabilis*, Blume, 36, 183.  
*virosus*, N. E. Br., 36, 181.  
**AMPELIDEE**, 23, 131.  
*Ampelocissus artemisiæfolia*, Planch., 36, 454.

- Ampelocissus cardiosperma*, *Planch.*, **36**, 454.  
*cardiospermoides*, *Planch.*, **36**, 454.  
*Ampelopsis aconitifolia*, *Bunge*, **23**, 136.  
*Delavayana*, *Planch.*, **36**, 454.  
*heterophylla*, *Sieb. et Zucc.*, **23**, 133.  
*himalayana*, *Royle*, **23**, 133.  
*humulifolia*, *Bunge*, **23**, 133.  
*megalophylla*, *Diels & Gilg*, **36**, 454.  
*mirabilis*, *Diels & Gilg*, **36**, 455.  
*serianæfolia*, *Bunge*, **23**, 136.  
*tomentosa*, *Planch.*, **36**, 455.  
*tricuspidata*, *Sieb. et Zucc.*, **23**, 133.  
*Veitchii*, *Hort.*, **23**, 133.  
*Ampelovitis*, *Carr.*, **36**, 455.  
*Davidi*, *Carr.*, **36**, 455.  
*intermedia*, *Carr.*, **36**, 455.  
*Romaneti*, *Carr.*, **36**, 455.  
*Ampelogyne chinense*, *Lindl.*, **26**, 335.  
*Amphicarpæa Edgeworthii*, *Benth.*, **23**, 188.  
*japonica*, *Oliver*, **23**, 188.  
*monoica*, *Nutt.*, **23**, 188.  
*Amphicome arguta*, *Lindl.*, **36**, 455.  
*Amphidonax bifaria*, *Nees*, **36**, 408.  
*Amphirhapis albescens*, *DC.*, **23**, 419.  
*chinensis*, *Sch.-Bip.*, **23**, 406.  
*leiocarpa*, *Benth.*, **23**, 406.  
*Amygdalopsis Lindleyi*, *Carr.*, **23**, 222.  
*Amygdalus communis*, *Bunge*, **23**, 220.  
*communis*, *Linn.*, **23**, 217.  
*cordifolia*, *Roxb.*, **23**, 221.  
*dasylepis*, *Miq.*, **23**, 220.  
*pedunculata*, *Bunge*, **23**, 222.  
*Persica*, *Linn.*, **23**, 220.  
*pumila*, *Sims*, **23**, 219.  
*tangutica*, *Korsh.*, **36**, 455.  
*Anacampseros spectabile*, *Jord. et Fourr.*, **23**, 287.  
ANACARDIACEÆ, **23**, 146.  
*Anagallis arvensis*, *Linn.*, **26**, 59.  
*phœnicea*, *Thunb.*, **26**, 59.  
*Anagyris fœtida*, *Lour.*, **23**, 204.  
*sinensis*, *Steud.*, **23**, 240.  
*Anandria Bellidiastrum*, *DC.*, **23**, 472.  
*dimorpha*, *Turcz.*, **23**, 472.  
*discoidea*, *Less.*, **23**, 473.  
*radiata*, *Less.*, **23**, 473.  
*Anaphalis adnata*, *DC.*, **23**, 425.  
*alata*, *Maxim.*, **23**, 425.  
*araneosa*, *DC.*, **36**, 455.  
*Anaphalis Bodinieri*, *Franch.*, **36**, 455.  
*cinnamomea*, *C. B. Clarke*, **23**, 426.  
*cuneifolia*, *Hook. f.*, **23**, 425.  
*Hancockii*, *Maxim.*, **23**, 425.  
*lactea*, *Maxim.*, **23**, 425.  
*margaritacea*, *A. Gray*, **23**, 426.  
*margaritacea*, *Benth. & Hook. f.*, **23**, 425.  
*nubigena*, *DC.*, **23**, 426.  
*pteroaulon*, *Maxim.*, **23**, 426; mentioned, 425.  
*sinica*, *Hance*, **23**, 426.  
*sp. n.*, **23**, 426.  
*triplinervis*, *Hance*, **23**, 425.  
*Anchusa officinalis*, *Linn.*, **26**, 155.  
*officinalis*, *Lour.*, **26**, 155.  
*tenella*, *Roem. & Schult.*, **26**, 152.  
*zeylanica*, *Jacq.*, **26**, 152.  
*Ancistrolobus ligustrinus*, *Spach*, **23**, 74.  
*Andrachne chinensis*, *Bunge*, **26**, 420.  
*cordifolia*, *Muell. Arg.*, **26**, 420.  
*fruticosa*, *Linn.*, **26**, 427.  
*Andreoskia dentata*, *Bunge*, **23**, 45.  
*Androglossum reticulatum*, *Benth.*, **23**, 144.  
*Andromeda chinensis*, *Lodd.*, **26**, 14.  
*elliptica*, *Sieb. et Zucc.*, **26**, 17.  
*japonica*, *Thunb.*, **26**, 17.  
*lanceolata*, *Wall.*, **26**, 17.  
*ovalifolia*, *Wall.*, **26**, 17.  
*Andropogon aciculatus*, *Retz.*, **36**, 363.  
*amplexifolius*, *Trin.*, **36**, 360.  
*annulatus*, *Forsk.*, **36**, 369; mentioned, 376.  
— *var. Bladhii*, *Hack.*, **36**, 370.  
— *var. genuinus*, *Hack.*, **36**, 369.  
*apricus*, *Hook. f.*, **36**, 370.  
*apricus*, *Trin.*, *var. indicus*, *Hack.*, **36**, 370.  
— *var. chinensis*, *Trin.*, **36**, 371.  
*assimilis*, *Steud.*, **36**, 370.  
*Bladhii*, *Retz.*, **36**, 370.  
*brevifolius*, *Sw.*, **36**, 370; mentioned, 372.  
*cæsius*, *Nees*, **36**, 377.  
*capilliflorus*, *Steud.*, **36**, 374.  
*caricosus*, *Linn.*, **36**, 371.  
*ciliatus*, *Thunb.*, **36**, 378.  
*contortus*, *Linn.*, **36**, 366.  
*cotulifer*, *Thunb.*, **36**, 351.  
*courtallumensis*, *Steud.*, **36**, 358.  
*crinitus*, *Thunb.*, **36**, 358.  
*Delavayi*, *Hack.*, **36**, 371.  
*fastigiatus*, *Sw.*, **36**, 371.  
*formosanus*, *Rendle* \*, **36**, 371.  
— *var. minor*, *Rendle* \*, **36**, 372.  
*fragilis*, *R. Br.*, **36**, 372.

- Andropogon fragilis*, var. *sinensis*, Rendle \*, 36, 372.  
*glaucopsis*, Steud., 36, 370.  
*Goeringii*, Steud., 36, 376.  
*halepensis*, Brot., 36, 367.  
*hamatulus*, Hack., 36, 376.  
*hamatulus*, Nees, 36, 376.  
*heteroclitus*, Nees, 36, 378.  
*hirtiflorus*, Kunth, 36, 373.  
*intermedius*, R. Br., 36, 373.  
*Ischæmum*, Linn., 36, 374.  
*lasiocoleus*, Steud., 36, 360.  
*Martini*, Benth., 36, 377.  
*micranthus*, Kunth, 36, 374; mentioned, 370.  
— var. *villosulus*, Hack., 36, 375.  
*montanus*, Benth., 36, 374.  
*montanus*, Hack., 36, 376.  
*Nardus*, Linn., 36, 376.  
— var. *genuinus*, Hack., 36, 376.  
*nitidus*, Kunth, 36, 367.  
*Pseudograya*, Steud., 36, 373.  
*punctatus*, Roxb., 36, 373.  
*Schœnanthus*, Linn., 36, 377.  
*Schœnanthus*, Miq., 36, 376.  
*serratus*, Thunb., 36, 367.  
— var. *genuinus*, Hack., 36, 367.  
*sibiricus*, Steud., 36, 353.  
*Sorghum*, Brot., 36, 367.  
— subsp. *halepensis*, Hack., 36, 367.  
— subsp. *sativus*, var. *vulgaris*, Hack., 36, 367.  
*subrepens*, Steud., 36, 370.  
*tropicus*, Spreng., 36, 367.  
*Vachellii*, Nees, 36, 373.  
*vagans*, Rendle \*, 36, 375; mentioned, 372.  
— var. *minor*, Rendle \*, 36, 372.  
— var. *sinensis*, Rendle \*, 36, 372.  
*villosulus*, Steud., 36, 375.  
*yunnanensis*, Hack., 36, 376.  
*zeylanicus*, Arn., 36, 373.
- Androsace Aizoon*, Duby, 26, 43.  
— var. *integra*, Maxim., 26, 44.  
*alchemilloides*, Franch., 36, 455.  
*axillaris*, Franch., 36, 455.  
*bisulca*, Bur., 36, 455.  
*Chamæjasme*, Host, 26, 44.  
*coccinea*, Franch., 36, 455.  
*cuscutiformis*, Franch., 36, 455.  
*Delavayi*, Franch., 36, 455.  
*dissecta*, Franch., 36, 455.  
*elongata*, Linn., 26, 44.  
*filiformis*, Retz., 26, 44.  
*foliosa*, Klatt, 26, 45.  
*Gmelini*, Gaertn., 26, 44.  
*Henryi*, Oliver, 36, 455.
- Androsace Jacquemontii*, Duby, 26, 46.  
Marlæ, Kanitz, 36, 455.  
*maxima*, Linn., 36, 455.  
*mirabilis*, Bur. & Franch., 36, 455.  
*mutica*, Franch., 26, 45.  
*patens*, Wright, 26, 45.  
*rotundifolia*, Hardw., 26, 45; mentioned, 46.  
— var. *axillaris*, Franch., 26, 45.  
— var. *dissecta*, Franch., 26, 45.  
*sarmentosa*, var. *grandifolia*, Hook. f., 26, 45.  
*saxifragæfolia*, Bunge, 26, 45; mentioned, 46.  
*sempervivoides*, Jacquem., 26, 45.  
*strigillosa*, Franch., 26, 45.  
— var. *spinulifera*, Franch., 26, 45.  
*sutchuenense*, Franch., 36, 455.  
*tapete*, Maxim., 26, 46.  
*villosa*, Jacq., 26, 44.  
*villosa*, Linn., 26, 46.
- Androscepiæ gigantea*, var. *armata*, Anderss., 36, 377.
- Aneilema angustifolium*, N. E. Br. \*, 36, 151.  
*didymum*, Seem., 36, 159.  
*divergens*, C. B. Clarke, 36, 151.  
*esculentum*, Wall., mentioned, 36, 155.  
*formosanum*, N. E. Br. \*, 36, 152.  
*herbaceum*, Wall., 36, 151.  
— var. *divergens*, C. B. Clarke, 36, 151.  
*japonicum*, Maxim., 36, 152.  
*Keisak*, Hassk., 36, 152; mentioned, 154.  
*Loureirii*, Hance, 36, 152.  
*melanostictum*, Hance, 36, 154.  
*medicum*, R. Br., 36, 153.  
*nanum*, Kunth, 36, 154.  
*nudiflorum*, R. Br., 36, 153.  
— var. *rigidior*, Benth., 36, 151.  
*oliganthum*, Franch., 36, 152.  
*paucifolium*, N. E. Br. \*, 36, 153.  
*scapiflorum*, Wight, mentioned, 36, 152.  
— var. *latifolium*, N. E. Br. \*, 36, 151.  
*secunda*, Wight, 36, 154.  
*sinicum*, Lindl., 36, 154; mentioned, 151.  
*spectabile*, Kurz, 36, 152.  
*spiratum*, R. Br., 36, 154; mentioned, 155.



- Aneilema vaginatum*, R. Br., 36, 155.  
*Anemarrhena asphodeloides*, Bunge, 36, 118.  
*Anemone altaica*, Fisch., 23, 10.  
   *amurensis*, Komarov, 36, 455.  
   *baicalensis*, Turcz., 36, 455.  
   *barbata*, Turcz., 23, 10.  
   *barbatula*, Turcz., 23, 10.  
   *begoniifolia*, H. Léveillé & Vaniot, 36, 455.  
   *Boissiaei*, H. Léveillé & Vaniot, 36, 455.  
   *cernua*, Thunb., 23, 10.  
   *chinensis*, Bunge, 36, 10.  
   *cœlestina*, Franch., 23, 11.  
   *Davidi*, Franch., 36, 455.  
   *Delavayi*, Franch., 36, 455.  
   *demissa*, Hook. f., 36, 455.  
   *dichotoma*, Linn., 36, 455.  
   *exigua*, Maxim., 23, 11.  
   *Fauriei*, H. Léveillé, 36, 455.  
   *flaccida*, F. Schmidt, 23, 11.  
   *gelida*, Maxim., 36, 455.  
   *glaucifolia*, Franch., 36, 455.  
   *Henryi*, Oliver, 36, 455.  
   *Hepatica*, Linn., 23, 11.  
   *imbricata*, Maxim., 36, 455.  
   *japonica*, Sieb., 23, 11.  
   *narcissiflora*, Linn., 23, 11.  
   *nikoënsis*, Maxim., 23, 11.  
   *obtusiloba*, D. Don, 23, 11.  
   *polyanthes*, D. Don, 23, 12.  
   *Raddeana*, Regel, 23, 12.  
   *reflexa*, Steph., 36, 455.  
   *Regeliana*, Maxim., 36, 455.  
   *rivularis*, Buch.-Ham., 23, 10; 36, 455.  
   *Rossii*, S. Moore, 23, 12.  
   *rupestris*, Wall., 23, 12.  
   *scabiosa*, H. Léveillé & Vaniot, 36, 456.  
   *silvestris*, Linn., 36, 456.  
   *umbrosa*, C. A. Mey., 36, 456.  
   *vitifolia*, Buch.-Ham., 36, 456.  
*Anethum graveolens*, Linn., 23, 335.  
*Angelica anomala*, Lallem., 23, 333.  
   *citriodora*, Hance, 23, 334.  
   *decursiva*, Franch., 23, 335.  
   *Dielsii*, Boissieu, 36, 456.  
   *dissoluta*, Diels, 36, 456.  
   *Fargesii*, Boissieu, 36, 456.  
   *flaccida*, Komarov, 36, 456.  
   *glauca*, Edgew., mentioned, 23, 334.  
   *gracilis*, Franch., 23, 334.  
   *grosseserrata*, Maxim., 23, 334.  
   *involutellata*, Diels, 36, 456.  
   *kiusiana*, Maxim., 23, 334.  
   *koreana*, Maxim., 36, 456.  
   *lævigata*, Franch., 23, 334.  
*Angelica laxifoliata*, Diels, 36, 456.  
   *megaphylla*, Diels, 36, 456.  
   *Miqueliana*, Maxim., 23, 334.  
   *mongolica*, Franch., 23, 333.  
   *montana*, Schleich., 23, 333.  
   *polymorpha*, Maxim., 36, 456.  
   *Pseudo-selinum*, Boissieu, 36, 456.  
   *scaberula*, Franch., 36, 456.  
   *setchuenensis*, Diels, 36, 456.  
   *sp.*, 23, 334.  
   *Uchiyamæ*, Yatabe, 36, 456.  
   *valida*, Diels, 36, 456.  
   *viridiflora*, [Hemsl.], mentioned, 23, 334.  
*Angræcum falcatum*, Lindl., 36, 38.  
*Anguina sinensis*, Mich., 23, 313.  
*Ania angustifolia*, Benth., 36, 20.  
*Anisadenia pubescens*, Oliver, 36, 456.  
*Aniseia biflora*, Choisy, 26, 157.  
   *calycina*, Choisy, 26, 157, 160.  
*Anisocalyx limnanthiflorus*, Hance, 26, 187.  
*Anisomeles ovata*, R. Br., 26, 299.  
*Anisopappus chinensis*, Hook. & Arn., 23, 431.  
*Anodendron Benthamianum*, Hemsl.\*, 26, 98.  
   *læve*, Maxim., 26, 99.  
*Anæctochilus Roxburghii*, Lindl., 36, 42.  
   *sp.*, Hance, 36, 42.  
   *Tashiroi*, Maxim., 36, 42.  
*Anona uncinata*, Lam., 23, 26.  
*Anosporum monocephalum*, Nees, 36, 209.  
*Anotis boerhaavioides*, Maxim., 23, 377.  
   *Thwaitesiana*, Maxim., 23, 378.  
*Anplectrum parviflorum*, Benth., 23, 301.  
*Antennaria cinnamomea*, DC., 23, 426.  
   *leontopodina*, DC., 23, 424.  
   *margaritacea*, R. Br., 23, 425.  
*Anthericum Gmelinianum*, Schultes, 36, 148.  
   *parviflorum*, Benth., 36, 119.  
   *serotinum*, Linn., 36, 139.  
*Anthistiria arguens*, Franch., 36, 378.  
   *arguens*, Willd., var. *japonica*, Anderss., 36, 378.  
   *caudata*, Hance, 36, 379.  
   *caudata*, Nees, 36, 377.  
   *ciliata*, A. Henry, 36, 377.  
   *ciliata*, Retz., 36, 377.  
   *Forskali*, Kunth, 36, 378.  
   *gigantea*, Cav., 36, 377.  
   *heteroclita*, Roxb., 36, 378.  
   *imberbis*, Retz., 36, 378.

- Anthistiria japonica*, Willd., 36, 378.  
*villosa*, Lam., 36, 377.  
*Anthocephalus chinensis*, Hassk., 23, 370.  
*Anthogonium gracile*, Lindl., 36, 21.  
*Griffithii*, Reichb. f., 36, 21.  
*sp.*, Griff., 36, 21.  
*Anthoxanthum Hookeri*, Rendle\*, 36, 380.  
*Anthriscus nemorosa*, Baker & S. Moore, 23, 330.  
*nemorosa*, Spreng., 23, 330.  
*sylvestris*, Hoffm., 23, 330.  
*Anticholerica*, Rumph., 23, 201.  
*Anticlea sibirica*, Kunth, 36, 148.  
*Antidesma apiculatum*, Hemsl.\*, 26, 430; mentioned, 432.  
*Bunius*, Spreng., 26, 430.  
*Fordii*, Hemsl. \*, 26, 430.  
*Ghaesembilla*, Gaertn., 26, 431.  
*gracile*, Hemsl. \*, 26, 431.  
*Henryi*, Hemsl. \*, 26, 431.  
*japonicum*, Sieb. & Zucc., 26, 432; mentioned, 431.  
*microphyllum*, Hemsl. \*, 26, 432.  
*paniculatum*, Roxb., 26, 432.  
*velutinosum*, Blume, mentioned, 26, 430.  
*Antirrhinum Linaria*, Linn., 26, 178.  
*Antirrhœa chinensis*, Benth., 23, 384.  
*Aopla reniformis*, Lindl., 36, 61.  
*Aparisthium javense*, Hassk., 26, 438.  
*Apaturia chinensis*, Lindl., 36, 17.  
*Aperula citriodora*, Blume, 26, 387.  
*Apetalon minutum*, Wight, 36, 47.  
*Aphananthe aspera*, Planch., 26, 452.  
*Aphanochilis polystachyus*, Benth., 26, 278.  
*Apios Delavayi*, Franch., 36, 456.  
*Fortunei*, Maxim., 23, 189.  
*gracillima*, Dunn, 36, 456.  
*macrantha*, Oliver, 36, 456.  
*Apium cicutæfolium*, Benth., 23, 328.  
*graveolens*, Linn., 23, 328.  
*ternatum*, Sieb., 23, 332.  
*Aplotaxis Bungei*, DC., 23, 463.  
*carthamoides*, DC., 23, 463.  
*multicaulis*, DC., 23, 463.  
*Apluda aristata*, Linn., 36, 379.  
*mutica*, Linn., 36, 379.  
— var. *aristata*, Hack., 36, 379.  
*varia*, subsp. *mutica*, Hack., 36, 379.  
*Apochoris pentapetala*, Duby, 26, 56.  
*Apocopsis Wightii*, Nees, 36, 358.  
*Apocopsis Wightii*, subsp. *genuinus*, var. *Wrightii*, Hack., 36, 358.  
*Wrightii*, Munro, 36, 358.  
APOCYNACEÆ, 26, 93.  
*Apocynum alternifolium*, Lour., 26, 98.  
*sibiricum*, Pall., 26, 98.  
*tiliæfolium*, Linn., 26, 115.  
*venetum*, Linn., 26, 98.  
*Aponogeton monostachyon*, Linn. f., 36, 192.  
*monostachys*, Edgew., 36, 192.  
*sp.*, 36, 193.  
*Aporosa frutescens*, Benth., 26, 429.  
*lanceolata*, Hance, 26, 428.  
*leptostachys*, Benth., 26, 429.  
*microcalyx*, var. *chinensis*, Hassk., 26, 429.  
*sinensis*, Baill., 26, 429.  
*Aporum concinnum*, Lindl., 36, 9.  
*Appendicula bifaria*, Lindl., 36, 39.  
*sp.*, Hance, 36, 38.  
*Aquilaria chinensis*, Spreng., 23, 402.  
*grandiflora*, Benth., 26, 402.  
*malaccensis*, Benth., 26, 402.  
*Aquilegia atropurpurea*, Willd., 23, 18.  
*ecalcarata*, Maxim., 36, 456.  
*oxypetala* [Franch.], 23, 18.  
*oxysepala*, Trautv. & Mey., 36, 456.  
*viridiflora*, Pall., 23, 18.  
*vulgaris*, Linn., 23, 18; 36, 456.  
— var. *oxypetala*, Franch., 23, 18.  
*Arabis alpina*, Linn., 23, 42.  
*axillaris*, Komarov, 36, 456.  
*Halleri*, Linn., 36, 456.  
*hirsuta*, Scop., 23, 42.  
*hypoglauca*, Franch., 23, 42.  
*japonica*, A. Gray, 23, 42.  
*lyrata*, Linn., 36, 456.  
*paniculata*, Franch., 36, 456.  
*pendula*, Linn., 23, 42.  
*perfoliata*, Lam., 23, 42.  
*Piasezkii*, Maxim., 23, 42.  
*serrata*, Franch., 23, 42.  
*spp.*, 23, 42, 43.  
*Stelleri*, DC., 23, 42.  
*Arachis asiatica*, Lour., 23, 171.  
*hypogæa*, Linn., mentioned, 23, 171.  
*Aralia atropurpurea*, Franch., 36, 456.  
*canescens*, Sieb. & Zucc., 23, 338.  
*chinensis*, Blume, 23, 337.  
*chinensis*, Linn., 23, 338.  
*dasyphylla*, Miq., 23, 337.  
*Decaisneana*, Hance, 23, 338.  
*elata*, Seem., 23, 338.  
*Fargesii*, Franch., 36, 456.

- Aralia Henryi*, Harms, **36**, 456.  
*hypoleuca*, Presl, **23**, 338.  
*mandshurica*, Seem., **23**, 338.  
*Maximowiczii*, Van Houtte, **23**, 340.  
*octophylla*, Lour., **23**, 342.  
*palmata*, Lour., **23**, 338.  
*papyrifera*, Hook., **23**, 341.  
*pentaphylla*, Thunb., **23**, 341.  
*pilosa*, Franch., **36**, 456.  
*Planchoniana*, Hance, **23**, 338.  
*quinquefolia*, A. Gray, **23**, 338.  
*quinquefolia*, Planch., **23**, 338.  
*Searelliana*, Dunn, **36**, 456.  
*sp. ?*, **23**, 339.  
*spinosa*, Linn., **23**, 338.  
*stipulata*, Franch., **36**, 456.  
*tomentella*, Franch., **36**, 456.  
*trifoliata*, Meyen, **23**, 339.  
*yunnanensis*, Franch., **36**, 456.
- ARALIACEÆ**, **23**, 337.  
*Archangelica*, Benth. & Hook. f., **23**, 334.
- Arctium Lappa*, Linn., **23**, 460.  
*majus*, Schkuhr, **23**, 460.
- Ardisia affinis*, Hemsl. \*, **26**, 63; mentioned, 65, 67.  
*brevicaulis*, Diels, **36**, 456.  
*caudata*, Hemsl. \*, **26**, 63.  
*chinensis*, Benth., **26**, 63; mentioned, 65, 67.  
*corymbifera*, Mez, **36**, 456.  
*crenata*, Sims, **26**, 63.  
*crenulata*, Lodd., **26**, 64.  
*crispa*, A. DC., **26**, 63.  
 — var. *elegans*, A. DC., **26**, 63, 64.  
*elegans*, Andrews, **26**, 64.  
*Faberii*, Hemsl. \*, **26**, 64.  
*Fordii*, Hemsl. \*, **26**, 64; mentioned, 67.  
*formosana*, Rolfe, **26**, 67.  
*hainanensis*, Mez, **36**, 456.  
*Hanceana*, Mez, **36**, 456.  
*Henryi*, Hemsl. \*, **26**, 65.  
*humilis*, Vahl, **26**, 65.  
*japonica*, Benth., **26**, 63.  
*japonica*, Blume, **26**, 65.  
*lentiginosa*, Lindl., **26**, 64.  
*littoralis*, Andrews, **26**, 65.  
*maculosa*, Mez, **36**, 457.  
*mamillata*, Hance, **26**, 66.  
*obtusata*, Mez, **36**, 457.  
*patens*, Mez, **26**, 457.  
*pauciflora*, Benth., **26**, 66.  
*pauciflora*, Heyne, **26**, 66.  
*pedunculosa*, Wall., **26**, 65.  
*penduliflora*, Mez, **36**, 457.  
*prinuulæfolia*, Gardn. & Champ., **26**, 66.  
*punctata*, Lindl., **26**, 64.
- Ardisia pusilla*, A. DC., **26**, 66.  
*quinquegona*, Blume, **26**, 66.  
*radians*, Hemsl., **36**, 457.  
*rigida*, Kurz, **26**, 66.  
*Sieboldii*, Miq., **26**, 67.  
*spp. n.*, **26**, 67.  
*tenera*, Mez, **36**, 457.  
*triflora*, Hemsl. \*, **26**, 67.  
*undulata*, Mez, **36**, 457.  
*yunnanensis*, Mez, **36**, 457.
- Areca Catechu*, Willd., **36**, 167.  
*Catechu*, Linn., **36**, 167.  
*hortensis*, Lour., **36**, 167.
- Arenaria barbata*, Franch., **36**, 457.  
*Delavayi*, Franch., **36**, 457.  
*holosteoides*, Edgew., **23**, 70.  
*juncea*, Bieb., **23**, 70.  
*kansuensis*, Maxim., **23**, 70.  
*lateriflora*, Linn., **23**, 70.  
*linearifolia*, Franch., **36**, 457.  
*longistyla*, Franch., **36**, 457.  
*napuligera*, Franch., **36**, 457.  
*nepalensis*, Spreng., **36**, 457.  
*orbiculata*, Royle, **36**, 457.  
*polytrichoides*, Edgew., **36**, 457.  
*Przewalskii*, Maxim., **23**, 70; **36**, 457.  
*quadridentata*, Williams, **36**, 457.  
*Roborowskii*, Maxim., **36**, 457.  
*serpyllifolia*, Linn., **23**, 70.  
*szechuensis*, Williams, **36**, 457.  
*trichophora*, Franch., **36**, 457.  
*yunnanensis*, Franch., **36**, 457.
- Arenga Engleri*, Becc., **36**, 167.  
*saccharifera*, Labill., **36**, 167.
- Arethusa ecristata*, Griff., **36**, 47.  
*sinensis*, Rolfe \*, **36**, 46.
- Arethusantha bletioides*, Finet, **36**, 457.
- Argemone mexicana*, Linn., mentioned, **23**, 34.
- Argyreia acuta*, Benth., **26**, 156.  
*acuta*, Lour., **26**, 156.  
*Championi*, Benth., **26**, 156.  
*festiva*, Wall., **26**, 156.  
*mollis*, Choisy, mentioned, **26**, 156.  
*obtectata*, C. B. Clarke, mentioned, **26**, 156.  
*speciosa*, Sweet, **26**, 156.
- Argyrothamnia cantonensis*, Hance, **26**, 435.  
*tuberculatus*, Muell. Arg., **26**, 436.
- Aria alnifolia*, Decne., **23**, 255.
- Arisæma amurense*, Maxim., **36**, 175.  
 — var. *magnidens*, N. E. Br. \*, **36**, 176.  
 — var. *robustum*, Engl., **36**, 175.  
 — var. *Sazensoo*, Engl., **36**, 175; mentioned, 176.

- Arisæma asperatum*, *N. E. Br.* \*, **36**, 176.  
*Bockii*, *Engl.*, **36**, 176.  
*consanguineum*, *Schott*, **36**, 176.  
*cordatum*, *N. E. Br.* \*, **36**, 177.  
*curvatum*, *Hance*, **36**, 178.  
*Davidianum*, *Engl.*, **36**, 457.  
*erubescens*, *Schott*, var. *consanguineum*, *Engl.*, **36**, 177.  
*Franchetianum*, *Engl.*, **36**, 177.  
*Giraldii*, *Baroni*, **36**, 177, 457.  
*heterophyllum*, *Blume*, **36**, 178.  
*japonicum*, *Blume*, **36**, 178.  
— var. *Sazensoo*, *Blume*, **36**, 175.  
*Konjac*, *Sieb.*, **36**, 182.  
*laminatum*, *Benth.*, **36**, 179.  
*latisectum*, *Blume*, **36**, 178.  
*lobatum*, *Engl.*, **36**, 178.  
— var. *Rosthornianum*, *Engl.*, **36**, 178.  
*nepenthoides*, *Mart.*, mentioned, **36**, 177.  
*parvum*, *N. E. Br.*, **36**, 178.  
*penicillatum*, *N. E. Br.*, **36**, 179.  
*pentaphyllum*, *Schott*, **36**, 180.  
*pictum*, *N. E. Br.*, **36**, 179.  
*præcox*, *De Vriese*, **36**, 179.  
*ringens*, *Schott*, **36**, 179.  
— var. *Sieboldii*, *Engl.*, **36**, 179.  
— var. *præcox*, *Engl.*, **36**, 179.  
*serratum*, *Schott*, var., *Baker & S. Moore*, **36**, 175.  
*Sieboldii*, *De Vriese*, **36**, 179.  
*Tatarinowii*, *Schott*, **36**, 177, 457.  
*Thunbergii*, *Blume*, **36**, 180.  
— var. *heterophyllum*, *Engl.*, **36**, 178.  
*tripartitum*, *Engl.*, **36**, 174.  
*verrucosum*, *Schott*, mentioned, **36**, 176.
- Aristida Adscensionis*, *Linn.*, **36**, 381.  
*capillacea*, *Cav.*, **36**, 381.  
*chinensis*, *Munro*, **36**, 381.  
*cærulescens*, *Desf.*, **36**, 381.  
*Cumingiana*, *Trin.*, **36**, 381.  
*vulpioides*, *Hance*, **36**, 381.
- Aristolochia acuminata*, *Roxb.*, **26**, 363.  
*contorta*, *Bunge*, **26**, 361.  
*debilis*, *Sieb. & Zucc.*, **26**, 361; mentioned, 363.  
*Delavayi*, *Franch.*, **36**, 457.  
*Fordiana*, *Hemsl.*, **26**, 361.  
*gentilis*, *Franch.*, **36**, 457.  
*gracillima*, *Hemsl.*, **36**, 457.  
*heterophylla*, *Hemsl.* \*, **26**, 361.  
*Kæmpferi*, *Willd.*, **26**, 362.  
*longa*, *Thunb.*, **26**, 361.  
*longifolia*, *Champ.*, **26**, 362.
- Aristolochia manshuriensis*, *Komarov*, **36**, 457.  
*mollissima*, *Hance*, **26**, 362.  
*moupinensis*, *Franch.*, **36**, 457.  
*recurvilabra*, *Hance*, **26**, 361.  
*Roxburghiana*, *Klotzsch*, **26**, 363.  
*setchuenensis*, *Franch.*, **36**, 457.  
*Sinarum*, *Lindl.*, **26**, 363.  
*Sipho*, *L'Hérit.*, **26**, 362; **36**, 457.  
*Tagala*, *Cham. & Schlecht.*, **26**, 363.  
*thibetica*, *Franch.*, **36**, 457.  
*Westlandi*, *Hemsl.*, **26**, 363.  
*yunnanensis*, *Franch.*, **36**, 457.
- ARISTOLOCHIACEÆ, **26**, 358.  
*Aristotelea spiralis*, *Lour.*, **36**, 41.  
*Armeniaca vulgaris*, *Lam.*, **23**, 217.  
*Arnebia saxatilis*, *Benth. & Hook. f.*, **26**, 155.  
*Szechenyii*, *Kanitz*, **36**, 457.  
*Arnica japonica*, *Thunb.*, **23**, 453.  
*palmata*, *Thunb.*, **23**, 458.
- AROIDEÆ, **36**, 173.  
*Arracacha Delavayi*, *Franch.*, **36**, 457.  
*peucedanifolia*, *Franch.*, **36**, 457.  
*Arsis rugosa*, *Lour.*, **23**, 92.  
*Artabotrys Blumei*, *Hook.*, **23**, 26.  
*hongkongensis*, *Hance*, **23**, 26.  
*odoratissimus*, *R. Br.*, **23**, 26.
- Artemisia annua*, *Linn.*, **23**, 441.  
*anomala*, *S. Moore*, **23**, 441.  
*apiacea*, *Hance*, **23**, 441.  
*aquatica*, *Lour.*, **23**, 441.  
*aurata*, *Komarov*, **36**, 458.  
*Besseriana*, *Ledeb.*, **23**, 441.  
*brachyloba*, *Franch.*, **23**, 442.  
*campestris*, *Linn.*, **23**, 442.  
*capillaris*, *Thunb.*, **23**, 442; mentioned, 445.  
— var. *arbuscula*, *Miq.*, **23**, 442.  
*chinensis*, *Bess.*, **23**, 441.  
*chinensis*, *Linn.*, **23**, 440.  
*cuneifolia*, *DC.*, **23**, 443.  
*Desertorum*, *Spreng.*, mentioned, **23**, 443.  
*deversa*, *Diels*, **36**, 458.  
*eriopoda*, *Bunge*, **23**, 442.  
*frigida*, *Willd.*, **23**, 443.  
*glabrata*, *DC.*, **23**, 443.  
*glaucæ*, *Pall.*, **23**, 443.  
*igniaria*, *Maxim.*, **23**, 446.  
*indica*, *Willd.*, **23**, 446.  
*integrifolia*, *Linn.*, **23**, 446.  
*japonica*, *Thunb.*, **23**, 442.  
— var. *rotundifolia*, *Debeaux*, **23**, 443.  
*judaica*, *Lour.*, **23**, 443.  
*Keiskeana*, *Miq.*, **23**, 444.  
*Kruhsiana*, *Bess.*, mentioned, **23**, 442.

- Artemisia laciniata*, Willd., 23, 444.  
*lactiflora*, Wall., 23, 444.  
*lagocephala*, Fisch., mentioned, 23, 442.  
*lancea*, Vaniot, 36, 458.  
*latifolia*, Ledeb., 23, 444.  
*lavandulæfolia*, DC., 23, 446.  
*lithophila*, Turcz., mentioned, 23, 442.  
*macrantha*, Ledeb., 23, 444.  
*Messerschmidiana*, Bess., 23, 444.  
*Moxa*, DC., 23, 444.  
*parviflora*, Buch.-Ham., 23, 443.  
*Sacrorum*, Ledeb., 23, 444; mentioned, 445.  
*scoparia*, Waldst. & Kit, 23, 445; mentioned, 442.  
*sibirica*, Maxim., 23, 440.  
*Sieversiana*, Willd., 23, 445.  
*sp. nov.?*, 23, 446.  
*Thunbergiana*, Maxim., 23, 441.  
*vestita*, Wall., 23, 445.  
*vulgaris*, Linn., 23, 446.
- Arthraxon breviaristatus*, Hack., 36, 359.  
*ciliaris*, Beauv., 36, 359.  
 — *subsp. Langsdorffii*, var. *genuinus*, Hack., 36, 360.  
 — *subsp. submuticus*, Hack., 36, 360.  
*ciliaris*, Beauv., partim, 36, 360.  
*cuspidatus*, Hochst., 36, 360.  
 — *var. micans*, Hance, 36, 360.  
*japonicus*, Miq., 36, 360.  
*lanceolatus*, Hochst., 36, 361.  
*Langsdorffianus*, Hochst., 36, 360.  
*micans*, Hochst., 36, 360.  
*nodosus*, Komarov, 36, 361.
- Arthrocnemum indicum*, Moq., 26, 328.
- Arthrosprion stipulatum*, Hassk., 23, 215.
- Arthrostylis chinensis*, Benth., 36, 230.  
*filiformis*, Thw., 36, 230.
- Artocarpus hypargyræa*, Hance, 26, 470.  
*integrifolia*, Linn. f., 26, 470.
- Arum campanulatum*, Roxb., 36, 181.  
*cucullatum*, Lour., 36, 183.  
*divaricatum*, Linn., 36, 180.  
*diversifolium*, Blume, 36, 180.  
*Dracontium*, Thunb., 36, 180.  
*Dracunculus*, Thunb., 36, 178.  
*esculentum*, Linn., 36, 183.  
*macrorrhizon*, Linn., 36, 184.  
*macrourum*, Bunge, 36, 174.  
*odorum*, Roxb., 36, 184.  
*pentaphyllum*, Linn., 36, 180.  
*sagittifolium*, Lour., 36, 184.
- Arum ternatum*, Thunb., 36, 174.  
*trilobatum*, Curt., 36, 180.
- Aruncus sylvester*, Kostel., 23, 223.
- Arundarbor spinosa*, Rumph., 36, 446.
- Arundina chinensis*, Blume, 36, 27.  
*Meyenii*, Reichb. f., 36, 27.  
*Philippii*, Reichb. f., 36, 27.  
*pulchella*, Teijsm. & Binn., 36, 27.  
*pulchra*, Miq., 36, 27.
- Arundinaria densiflora*, Rendle\*, 36, 434.  
*elegans*, Kurz, 36, 342.  
*Faberi*, Rendle\*, 36, 435.  
*flexuosa*, Hance, 36, 436.  
*Hindsii*, Munro, 36, 436.  
*khasiana*, Bean, 36, 436.  
*linearis*, Hack., 36, 446.  
*longiramea*, Munro, 36, 436.  
*nitida*, Mitford, 36, 436.  
*racemosa*, Munro, mentioned, 36, 435.  
*sinica*, Hance, 36, 436.  
*sparsiflora*, Rendle\*, 36, 436.  
*vaginata*, Hack., 36, 437.  
*Wightii*, Benth., 36, 436.  
*Wilsoni*, Rendle\*, 36, 437.
- Arundinella anomala*, Steud., 36, 341.  
*brasiliensis*, Raddi, mentioned, 36, 342.  
 — *var. depauperata*, Rendle\*, 36, 341.  
*glabra*, Nees, 36, 341.  
*miliacea*, Nees, 36, 341.  
*nepalensis*, Trin., 36, 341.  
*setosa*, Trin., 36, 342.  
*sinensis*, Rendle\*, 36, 342.  
*stricta*, Nees, 36, 342.
- Arundo agrestis*, Lour., 36, 444.  
*Bambos*, Linn., 36, 445.  
*bengalensis*, Retz., 36, 408.  
*bifaria*, Retz., 36, 408.  
*Donax*, Linn., 36, 408.  
 — *var. coleotricha*, Hack., 36, 408.  
*epigejos*, Linn., 36, 391.  
*formosana*, Hack., 36, 408.  
 — *var. gracilis*, Hack., 36, 408.  
*Henslowiana*, Hook. & Arn., 36, 409.  
*madagascariensis*, Kunth, 36, 409.  
*Phragmites*, Linn., 36, 409.  
*Reynaudiana*, Kunth, 36, 409.  
*Roxburghii*, Kunth, 36, 410.
- Asarum Blumei*, Hance, 26, 358.  
*brevistylum*, Franch., 36, 458.  
*cardiophyllum*, Franch., 36, 458.  
*caulescens*, var., Franch., 36, 458.  
*chinense*, Franch., 36, 458.

- Asarum debile*, Franch., 36, 458.  
*Delavayi*, Franch., 36, 458.  
*Fargesii*, Franch., 36, 458.  
*Forbesii*, Maxim., 26, 358.  
*Franchetianum*, Diels, 36, 458.  
*geophilum*, Hemsl., 26, 358.  
*heterotropoides*, F. Schmidt, 26, 360.  
*himalaicum*, Hook. f. & T. Thoms., 26, 359.  
*macranthum*, Hook. f., 26, 359.  
*maximum*, Hemsl., 26, 359.  
*pulchellum*, Hemsl., 26, 360.  
*sakawanum*, Makino, 36, 458.  
*Sieboldi*, Miq., 26, 360.  
*Thunbergii*, A. Br., 26, 359.  
*Ascarina serrata*, Blume, 26, 367.  
*Asclepiadea dubia*, Oliver, 26, 109.  
**ASCLEPIADEÆ**, 26, 100.  
*Asclepias carnosu*, Linn., 26, 116.  
*curassavica*, Linn., mentioned, 26, 102.  
*hastata*, Bunge, 26, 105.  
*paniculata*, Bunge, 26, 102.  
*purpurea*, Pall., 26, 108.  
*sibirica*, Linn., 26, 108.  
*tingens*, Buch., 26, 112.  
*volubilis*, Linn. f., 26, 115.  
*Ascolepis kyllingioides*, Steud., 36, 257.  
*tenuior*, Steud., 36, 253.  
*Asparagopsis sinica*, Miq., 36, 103.  
*Asparagus brachyphyllus*, Turcz., 36, 101.  
*davuricus*, Fisch., 36, 101.  
*falcatus*, Benth., 36, 102.  
*filicinus*, Buch.-Ham., 36, 102.  
*gibbus*, Bunge, 36, 102.  
*insularis*, Hance, 36, 102.  
*longiflorus*, Franch., 36, 102.  
*lucidus*, Lindl., 36, 102; mentioned, 146.  
*maritimus*, Pall., 36, 101.  
*micranthus*, Sieb., 36, 103.  
*officinalis*, Linn., 36, 103.  
*oligoclonus*, Maxim., 36, 458.  
*parviflorus*, Turcz., 36, 103.  
*schoberioides*, Kunth, 36, 103.  
— var. *subsetaceus*, Franch., 36, 103.  
*Sieboldi*, Debeaux, 36, 102.  
*Sieboldi*, Maxim., 36, 103.  
*sinicus*, C. H. Wright\*, 36, 103.  
*trichophyllus*, Bunge, 36, 103.  
— var. *trachyphyllus*, Kunth, 36, 101.  
*Wrightii*, A. Gray, 36, 103.  
*Asperula Platygaliu*, Maxim., 23, 395.  
*Aspidistra elatior*, Blume, 36, 113.  
*lurida*, Ker-Gawl., 36, 113.  
*Aspidistra minutiflora*, Stapf\*, 36, 131.  
*punctata*, Carrière, 36, 113.  
— var. *albo-maculata*, Hook., 36, 113.  
*punctata*, Lindl., 36, 114.  
*sp.*, Baill., 36, 113.  
*variegata*, Hort., 36, 113.  
*Aspidopterys obcordata*, Hemsl., 36, 458.  
*Asprelia Duthiei*, Stapf, 36, 434.  
*Aster ageratoides*, Turcz., 23, 416.  
— var. *heterophyllus*, Debeaux, 23, 416.  
*alatipes*, Hemsl. \*, 23, 407.  
*alpinus*, Linn., 23, 408.  
*altaicus*, Willd., 23, 408; mentioned, 409, 414, 416.  
*alyssoides*, Turcz., 23, 409.  
*auriculatus*, Franch., 36, 458.  
*baccharoides*, Benth., 23, 409.  
*baccharoides*, Steetz, 23, 409.  
*batangensis*, Bur. & Franch., 36, 458.  
*Benthami*, Steetz, 23, 416.  
*biennis*, Ledeb., 23, 412.  
*Bietii*, Franch., 36, 458.  
*brachytrichus*, Franch., 36, 458.  
*brevipes*, Benth., 23, 409.  
*breviscapus*, Vaniot, 36, 458.  
*Candelabrum*, Vaniot, 36, 458.  
*chinensis*, Linn., 23, 407.  
*ciliatus*, A. Gray, 23, 414.  
*curvatus*, Vaniot, 36, 458.  
*Delavayi*, Franch., 36, 458.  
*dimorphophyllus*, Franch., 23, 410.  
*fastigiatus*, Fisch., 23, 410.  
*Flabellum*, Vaniot, 36, 458.  
*Fordii*, Hemsl. \*, 23, 410.  
*fuscescens*, Bur. & Franch., 36, 458.  
*Gerlachii*, Hance, 23, 411, 490.  
*Glehnii*, F. Schmidt, 23, 411.  
*Henryi*, Hemsl. \*, 23, 411.  
*hispidus*, Thunb., 23, 411.  
*holophyllus*, Hemsl. \*, 23, 412.  
*incisus*, Fisch., 23, 412.  
*indicus*, Linn., 23, 413.  
*integrifolius*, Franch., 23, 412; 36, 458.  
*japonicus*, Franch., mentioned, 23, 410.  
*latebracteatus*, Franch., 36, 458.  
*laticorymbus*, Vaniot, 36, 458.  
*Latureanus*, Franch., 23, 413.  
*likiangensis*, Franch., 36, 458.  
*limosus*, Hemsl. \*, 23, 413.  
*lingulatus*, Franch., 36, 458.  
*macilentus*, Vaniot, 36, 458.  
*millefolius*, Vaniot, 36, 458.

- Aster mongolicus*, Franch., 23, 414.  
*nigrescens*, Vaniot, 36, 458.  
*nigromontana*, Dunn, 36, 458.  
*Oldhami*, Hemsl. \*, 23, 414.  
*oreophilus*, Franch., 36, 458.  
*panduratus*, Nees, 23, 415.  
*Piccolii*, Hook., 36, 458.  
*procerus*, Hemsl. \*, 23, 415.  
*Sampsoni*, Hemsl. \*, 23, 415.  
*scaber*, Thunb., 23, 415; mentioned, 408.  
*scaberulus*, Miq., 23, 416.  
*senecioides*, Franch., 36, 458.  
*setchuensis*, Franch., 36, 458.  
*Souliei*, Franch., 36, 458.  
*staticifolius*, Franch., 36, 458.  
*striatus*, Camp., 23, 416.  
*striatus*, Franch., 23, 409.  
*tataricus*, Linn., 23, 416; mentioned, 415.  
*tenuissimus*, Dunn, 36, 459.  
*tongolensis*, Franch., 36, 459.  
*tricapitatus*, Vaniot, 36, 459.  
*trinervius*, Roxb., 23, 416; mentioned, 409.  
*Tripolium*, Linn., 23, 417.  
*turbinatus*, S. Moore, 23, 417.  
*vellereus*, Franch., 36, 459.  
*vestitus*, Franch., 36, 459.  
*Vilmorini*, Franch., 36, 459.  
*Walpersianus*, Nees, 23, 417.  
*yunnanensis*, Franch., 36, 459.  
*Asteromæa indica*, Blume, 23, 413.  
*pekinensis*, Hance, 23, 412.  
*Asterostoma repens*, Blume, 23, 300.  
*Astilbe chinensis*, Franch. & Sav., 23, 265; mentioned, 266.  
 — *var. Davidi*, Franch., 23, 265.  
 — *var. japonica*, Franch., 23, 265.  
*Davidii* [Forbes & Hemsl.], 23, 495.  
*japonica*, A. Gray, mentioned, 23, 265, 266.  
*odontophylla*, Miq., 23, 265.  
*pinnata*, Franch., 36, 459.  
*polyandra*, Hemsl. \*, 23, 265, 490.  
*rubra*, Hook. f. & Thoms., 36, 459.  
*Thunbergi*, Miq., 23, 266.  
*Astragalus adsurgens*, Pall., 23, 164.  
*brachycephalus*, Franch., 36, 459.  
*camptodontus*, Franch., 36, 459.  
*capillipes*, Fisch., 23, 165.  
*chinensis*, Linn., 23, 165.  
*chrysopterus*, Bunge, 23, 165.  
*complanatus*, R. Br., 23, 165.  
*daluricus*, DC., 23, 165.  
*Davidi*, Franch., 36, 459.  
*dependens*, Bunge, 23, 165.  
*glanduliferus*, Debeaux, 23, 165.  
*graveolens*, Buch.-Ham., 36, 459.  
*Hancockii*, Bunge, 23, 165.  
*Astragalus Henryi*, Oliver, 36, 459.  
*Laxmanni*, Jacq., 23, 165.  
*litangensis*, Bur. & Franch., 36, 459.  
*Loczii*, Kanitz, 36, 459.  
*lotoides*, Lam., 23, 166.  
*melilotoides*, Pall., 23, 166.  
*Mœllendorffi*, Bunge, 23, 166.  
*monadelphus*, Bunge, 23, 166.  
*moupinensis*, Franch., 36, 459.  
*nigrescens*, Franch., 36, 459.  
*polycladus*, Bur. & Franch., 36, 459.  
*Przewalskii*, Bunge, 23, 166.  
*scaberrimus*, Bunge, 23, 166.  
*sciadophorus*, Franch., 23, 166.  
*sinicus*, Linn., 23, 166.  
*skythropos*, Bunge, 23, 166.  
*sutchuenensis*, Franch., 36, 459.  
*tanguticus*, Batalin, 36, 459.  
*tartaricus*, Franch., 23, 167.  
*tatsiensis*, Bur. & Franch., 36, 459.  
*tenuis*, Turcz., 23, 167.  
*variabilis*, Bunge, 23, 167.  
*yunnanensis*, Franch., 36, 459.  
*Astranthus cochinchinensis*, Lour., 23, 312.  
*Asystasia chelonioides*, Nees, mentioned, 26, 243.  
*chinensis*, S. Moore, 26, 243.  
*violacea*, Dalz., 26, 243.  
*Atalantia buxifolia*, Oliver, 23, 110.  
*Hindsii*, Oliver, 23, 110.  
*monophylla*, Benth., 23, 110.  
*monophylla*, Hook. & Arn., 23, 110.  
*Ataxia Hookeri*, Griseb., 36, 380.  
*Athamanta chinensis*, Linn., 23, 333.  
*Atherurus ternatus*, Blume, 36, 175.  
*tripartitus*, Blume, 36, 174.  
*Atractylis chinensis*, DC., 23, 459.  
*lancea*, DC., 23, 459.  
*lancea*, Thunb., 23, 459; mentioned, 460.  
*ovata*, Thunb., 23, 459; mentioned, 460.  
*Atractylodes lancea*, DC., 23, 459.  
*lyrata*, Sieb. & Zucc., 23, 459.  
*ovata*, DC., 23, 459.  
*Atragene alpina*, Linn., 23, 1.  
*koreana*, Komarov, 36, 459.  
*macropetala*, Ledeb., 23, 5.  
*Atriplex arenaria*, Nutt., 26, 325.  
*littoralis*, Linn., 26, 325.  
*Maximowicziana*, Makino, 36, 459.  
*nummularia*, Lindl., 26, 326.  
*sibirica*, Linn., 26, 326.  
*Atropis distans*, Griseb., 36, 428.  
*Atylosia scarabæoides*, Benth., 23, 195.  
*subrhombica*, Miq., 23, 195.  
*trichodon*, Dunn, 36, 459.

- Aubletia ramosissima*, Lour., 23, 126.
- Aucuba chinensis*, Benth., 23, 346.  
*japonica*, Thunb., 23, 346.
- Augia sinensis*, Lour., 23, 149.
- Auliconema aspera*, Hua, 36, 110.  
 Pernyi, Hua, 36, 459.
- Avena Delavayi*, Hack., 36, 401.  
*fatua*, Linn., 36, 401.  
 — var. *glabrata*, Stapf \*, 36, 401.  
*flavescens*, Linn., 36, 399.  
*nuda*, Linn., 36, 401.  
 — var. *chinensis* [Fisch.], Cat. Gor., 36, 401.  
*pratensis*, Linn., 36, 401.  
*sativa*, Linn., 36, 402.  
*strigosa*, Schreb., mentioned, 36, 401.  
*subspicata*, Clairv., 36, 400.
- Averrhoa Bilimbi*, Linn., mentioned, 23, 100.  
*Carambola*, Linn., 23, 100.  
*sinica*, Hance, 23, 149.
- Avicennia officinalis*, Linn., 26, 265.  
*tomentosa*, Jacq., 26, 266.
- Axonopus cimicinus*, Beauv., 36, 334.  
*semialatus*, Hook. f., 36, 334.
- Axyris amaranthoides*, Linn., 26, 326.
- Azalea amœna*, Lindl., 26, 26.  
*crispiflora*, Hook., 26, 26.  
*Danielsiana*, Paxt., 26, 26.  
*decumbens*, DC., 26, 26.  
*fragrans*, Adams, 26, 23.  
*indica*, Linn., 26, 25.  
 — var. *alba*, Lindl., 26, 27.  
 — var. *lateritia*, Lindl., 26, 25.  
 — var. *variegata*, Lindl., 26, 25.  
 — var. *vittata*, Fl. des Serres, 26, 31.  
*japonica*, A. Gray, 26, 30.  
*ledifolia*, Hook., 26, 27.  
 — var. *phœnicea*, Hook., 26, 28.  
*macrantha*, Bunge, 26, 25.  
*mollis*, Blume, 26, 30.  
*mucronata*, Blume, 26, 27.  
*myrtifolia*, Champ., 26, 28.  
*narcissiflora*, Fortune, 26, 27.  
*obtusata*, Lindl., 26, 26.  
*ovata*, Lindl., 26, 28.  
*pontica*, var. *sinensis*, Lindl., 26, 30.  
*punicea*, Sweet, 26, 28.  
*ramentacea*, Lindl., 26, 26.  
*rosmarinifolia*, Burm. f., 26, 27.  
*sinensis*, Lodd., 26, 30.  
*squamata*, Lindl., 26, 23.
- Baccharis chinensis*, Lour., 23, 420.  
*Dioscoridis*, Linn., 23, 420.  
*Salvia*, Lour., 23, 422.
- Baea*, see Boea.
- Bæckeia chinensis*, Gaertn., 23, 295.  
*frutescens*, Linn., 23, 295.
- Balanophora Harlandi*, Hook. f., 26, 410.  
 Henryi, Hemsl. \*, 26, 410.  
 Hookeriana, Hemsl., mentioned, 26, 410.  
*involuta*, Hook. f., 26, 410.  
 Kuroiwai, Makino, 36, 459.  
*laxiflora*, Hemsl. \*, 26, 410.  
*minor*, Hemsl. \*, 26, 410.  
*polyandra*, Griff., 26, 411.  
 sp.?, 26, 411.
- BALANOPHOREÆ, 26, 410.
- Baliospermum*, Blume, sp.?, 26, 444.
- Balsamina chinensis*, DC., 23, 101.  
*cochleata*, DC., 23, 101.  
*hortensis*, Desf., 23, 100.  
*mutila*, DC., 23, 101.
- Bambusa agrestis*, Poir., 36, 444.  
*angulata*, Munro, 36, 445.  
*arundinacea*, Willd., 36, 445.  
*Arundo*, Klein, 36, 445.  
*Beecheyana*, Munro, 36, 445.  
*breviflora*, Munro, 36, 445.  
*Cantori*, Munro, 36, 445.  
*dumetorum*, Hance, 36, 449.  
*flexuosa*, Munro, 36, 445.  
*floribunda*, Zoll. & Mor., 36, 446.  
*nana*, Roxb., 36, 445.  
*Oldhami*, Munro, 36, 446.  
*quadrangularis*, Fenzi, 36, 443.  
*reticulata*, var. *macrophylla*, Rupr., 36, 446.  
*stenostachya*, Hack., 36, 446.  
*striata*, Lindl., 36, 447.  
 Teba, Miq., 36, 446.  
*tessellata*, Munro, 36, 446.  
*Tulda*, Benth., 36, 447.  
*tuldoides*, Munro, 36, 447.  
*verticillata*, Benth., 36, 448.  
*verticillata*, Hook. & Arn., 36, 445.  
*vulgaris*, Schrad., var. *striata*, Gamble, 36, 447.
- Banksia speciosa*, Koen., 36, 71.
- Barbarea vulgaris*, R. Br., 23, 41.
- Barbula sinensis*, Lour., 26, 263.
- Barkhausia repens*, DC., 23, 474.  
*tenella*, Benth., 23, 485.
- Barleria cristata*, Linn., 26, 242.  
*nepalensis*, Nees, 26, 242.  
*procumbens*, Lour., 26, 243.
- Barnardia scilloides*, Lindl., 36, 127.
- Barthea chinensis*, Hook., 23, 300.



- Bartsia Odontites*, *Huds.*, **36**, 459.  
*Basella alba*, *Linn.*, **26**, 331.  
     *nigra*, *Lour.*, **26**, 331.  
     *rubra*, *Linn.*, **26**, 331.  
*Batatas acetosæfolia*, *Choisy*, **26**, 158.  
     *edulis*, *Choisy*, **26**, 157.  
     *littoralis*, *Choisy*, **26**, 158.  
     *paniculata*, *Choisy*, **26**, 159.  
*Bauhinia acuminata*, *Linn.*, **23**, 212.  
     *bryoniflora*, *Franch.*, **36**, 459.  
     *Championi*, *Benth.*, **23**, 212.  
     *chinensis*, *Vogel*, **23**, 213.  
     *corymbosa*, *Roxb.*, **23**, 212.  
     *Delavayi*, *Franch.*, **36**, 459.  
     *densiflora*, *Franch.*, **36**, 459.  
     *Faberi*, *Oliver*, **36**, 459.  
     *glauca*, *Wall.*, **23**, 212.  
     *parviflora*, *Vahl*, **23**, 212.  
     *parvifolia*, *Teysm. et Binn.*, **23**,  
     212.  
     *racemosa*, *Lam.*, **23**, 212.  
     *retusa*, *Roxb.*, mentioned, **23**, 213.  
     *scandens*, *Linn.*, mentioned, **23**,  
     213.  
     *sp.*, **23**, 213.  
     *tomentosa*, *Linn.*, **23**, 213.  
     *variegata*, *Linn.*, **23**, 213.  
     *yunnanensis*, *Franch.*, **36**, 459.  
*Baumea Brownei*, *Boeck.*, **36**,  
     262.  
     *rubiginosa*, *Boeck.*, **36**, 262.  
*Beaumontia breviflora*, *Oliver*, **36**,  
     459.  
     *indecora*, *Baill.*, **36**, 459.  
*Beckmannia erucæformis*, *Host*, **36**,  
     405.  
*Begonia Augustinei*, *Hemsl.*, **36**, 460.  
     *Bowringiana*, *Champ.*, **23**, 323.  
     *Bowringiana*, *Hort.*, **36**, 460.  
     *Bretschneideriana*, *Hemsl.*, **36**,  
     460.  
     *circumlobata*, *Hance*, **23**, 321.  
     *cyclophylla*, *Hook.*, **23**, 321.  
     *Dielsiana*, *E. Pritz.*, **36**, 460.  
     *discolor*, *Ait.*, **23**, 321.  
     *Evansiana*, *Andrews*, **23**, 321.  
     *fimbristipula*, *Hance*, **23**, 321.  
     *Fordii*, *Hemsl.* \*, **23**, 322; corrected  
     to *fimbristipula*, 490.  
     *grandis*, *Dryand.*, **23**, 321.  
     *Hemsleyana*, *Hook. f.*, **36**, 460.  
     *Henryi*, *Hemsl.* \*, **23**, 322, 490.  
     *laciniata*, *Roxb.*, **23**, 322.  
     *leprosa*, *Hance*, **23**, 322.  
     *Roxburghii*, *A. DC.*, **23**, 322.  
     *sinensis*, *A. DC.*, **23**, 323.  
     *sp.*, **23**, 323.  
 BEGONIACEÆ, **23**, 321.  
*Beilschmiedia assamica*, *Meism.*, **26**,  
     370.  
*Beilschmiedia chinensis*, *Hance*, **26**,  
     370.  
     *fagifolia*, *Nees*, **26**, 370; men-  
     tioned, 378.  
*Beketowia*, *Krassn.*, **36**, 460.  
     *tianschanica*, *Krassn.*, **36**, 460.  
*Belamcanda chinensis* [*DC.*], **36**, 86.  
     *punctata*, *Moench*, **36**, 86.  
*Belis jaculifolia*, *Salisb.*, **26**, 548.  
*Benincasa cerifera*, *Savi*, **23**, 315.  
     *hispida*, *Cogn.*, **23**, 315.  
*Benthania fragifera*, *Lindl.*, **23**, 345.  
     *japonica*, *Benth.*, **23**, 345.  
     — var. *sinensis*, *Benth.*, **23**, 345.  
*Benzoin citrodorum*, *Sieb. & Zucc.*,  
     **26**, 387.  
     *glaucum*, *Sieb. & Zucc.*, **26**, 388.  
     *præcox*, *Sieb. & Zucc.*, **26**, 391.  
     *sericeum*, *Sieb. & Zucc.*, **26**, 391.  
     *Thunbergii*, *Sieb. & Zucc.*, **26**, 393.  
     *trilobum*, *Sieb. & Zucc.*, **26**, 392.  
 BERBERIDEÆ, **23**, 30.  
*Berberis acuminata*, *Franch.*, **36**, 460.  
     *amurensis*, *Franch.*, **23**, 32.  
     *angulosa*, *Wall.*, **36**, 460.  
     *Bealei*, *Fortune*, **23**, 31.  
     — var. *planifolia* [*Hook.*], *Bot.*  
     *Mag.*, **23**, 31.  
     *brachypoda*, *Maxim.*, **23**, 31.  
     *caraganeæfolia*, *DC.*, **23**, 163.  
     *dasystachya*, *Maxim.*, **23**, 31.  
     *diaphana*, *Maxim.*, **23**, 31.  
     *dictyophylla*, *Franch.*, **36**, 460.  
     *Fortunei*, *Lindl.*, **23**, 31.  
     *gracilipes*, *Oliver*, **36**, 460.  
     *insignis*, *Hook. f.*, **36**, 460.  
     *integerrima*, *Bunge*, **36**, 460.  
     *koreana*, *Palibin*, **36**, 460.  
     *levis*, *Franch.*, **36**, 460.  
     *Maximowiczii*, *Regel*, **23**, 32.  
     *nepalensis*, *Spreng.*, **23**, 31.  
     *pallens*, *Franch.*, **36**, 460.  
     *polyantha*, *Hemsl.*, **36**, 460.  
     *Potanini*, *Maxim.*, **36**, 460.  
     *pruinosa*, *Franch.*, **36**, 460.  
     *sanguinea*, *Franch.*, **36**, 460.  
     *sinensis*, *Desf.*, **23**, 31.  
     *stenophylla*, *Hance*, **23**, 31.  
     *subtriplinervis*, *Franch.*, **36**, 460.  
     *Thunbergii*, *DC.*, **23**, 32.  
     *vulgaris*, *Linn.*, **23**, 32.  
     *Wallichiana*, *DC.*, **23**, 32.  
     *yunnanensis*, *Franch.*, **36**, 460.  
*Berchemia congesta*, *S. Moore*, **23**,  
     127.  
     *flavescens*, *Wall.*, **36**, 460.  
     *lineata*, *Benth.*, **23**, 127.  
     *lineata*, *DC.*, **23**, 127.  
     *Loureiriana*, *DC.*, mentioned, **23**,  
     127.

- Berchemia Poiretiana*, DC., 23, 127.  
*racemosa*, Sieb., 23, 127.  
*sessiliflora*, Benth., 23, 128.  
*yunnanensis*, Franch., 36, 460.  
*Berghausia mutica*, Munro, 36, 387.  
*patula*, Munro, 36, 306.  
*Bergia ammannioides*, Roxb., 23, 72.  
*Berteroa incana*, DC., 23, 41.  
*Potanini*, Maxim., 36, 460.  
*Beta vulgaris*, Linn., mentioned, 26, 325.  
*Betonica officinalis*, Lour., 26, 288.  
*Betula acuminata*, Wall., 26, 497.  
*alba*, Linn., 26, 496; mentioned, 497.  
— *var. mandschurica*, Regel, 26, 496.  
— *var. Tauschii*, Regel, 26, 497.  
— *var. verrucosa*, Franch., 26, 497.  
— *var. vulgaris*, Regel, 26, 497.  
*albo-sinensis*, Burkill\*, 26, 497.  
*alnoides*, Buch.-Ham., 26, 497.  
— *var. pyrifolia*, Franch., 26, 497.  
*Bhojpattra*, Wall., 26, 497, 499.  
— *var. sinensis*, Franch., 26, 497.  
*chinensis*, Maxim., 26, 498.  
*costata*, Trautv., 36, 460.  
*cylindrostachys*, Wall., 36, 460.  
*dahurica*, Pall., 26, 498.  
*Delavayi*, Franch., 26, 498.  
*Ermani*, Cham., 36, 460.  
*exalata*, S. Moore, 26, 498.  
*Fargesii*, Franch., 26, 498.  
*fruticosa*, Pall., 36, 460.  
*insignis*, Franch., 26, 498.  
*latifolia*, Tausch, 36, 460.  
*Potanini*, Batalin, 26, 498.  
*rosea*, David, 26, 499.  
*Schmidtii*, Regel, 36, 460.  
*utilis*, D. Don, 26, 429; mentioned, 497.  
— *var. Prattii*, Burkill\*, 26, 499.  
*verrucosa*, Ehrh., 26, 497.  
*Bidaria tingens*, Decne., 26, 112.  
*Bidens bipinnata*, Linn., 23, 434; mentioned, 435.  
*cernua*, Linn., 23, 435.  
*chinensis*, Wall., 23, 435, 436.  
*leucantha*, Willd., 23, 436.  
*leucorhiza*, DC., 23, 435.  
*Meyeniana*, Walp., 23, 435.  
*parviflora*, Willd., 23, 435.  
*pilosa*, Linn., 23, 435.  
— *var. bipinnata*, Hook. f., 23, 435.  
*tenuifolia*, Labill., 23, 436.  
*tripartita*, Linn., 23, 437.  
*Wallichii*, DC., 23, 435.  
*Biebersteinia heterostemon*, Maxim., 23, 97.  
*Bifaria Davidiana*, Van Tiegh., 36, 460.  
*fasciculata*, Van Tiegh., 36, 460.  
*Bigelovia lasiocarpa*, Wight et Arn., 23, 392.  
*Bignonia chinensis*, Lam., 23, 235.  
*grandiflora*, Thunb., 26, 235.  
*indica*, Linn., 26, 234.  
*pentandra*, Lour., 26, 234.  
BIGNONIACEÆ, 26, 234.  
*Biophytum Reinwardtii*, Edgew., 23, 99.  
*sensitivum*, DC., 23, 100.  
*Biota chinensis*, Hort., 26, 540.  
*orientalis*, Endl., 26, 540.  
*Biotia discolor*, Maxim., 23, 415.  
*Bischofia javanica*, Blume, 26, 428.  
*oblongifolia*, Decne., 26, 428.  
*trifoliata*, Hook., 26, 428.  
BIXINEÆ, 23, 57.  
*Blachia Pentzii*, Benth., 26, 435.  
*Blackwellia fagifolia*, Lindl., 23, 312.  
*Loureiri*, Benth., 23, 312.  
*padiflora*, Lindl., 23, 312.  
*Bludhia*, Thunb., 26, 63 note.  
*crispa*, Thunb., 26, 64.  
*glabra*, Thunb., 26, 65.  
*japonica*, Thunb., 26, 65.  
*villosa*, Thunb., 26, 66.  
*Blastus cochinchinensis*, Lour., 23, 301.  
*Hindsii*, Hance, 23, 301.  
*parviflorus*, Triana, 23, 301.  
*Bletia Gebina*, Lindl., 36, 19.  
*hyacinthina*, R. Br., 36, 19.  
— *var. Gebina*, Blume, 36, 19.  
*Tankervilliae*, R. Br., 36, 18.  
*Bletilla striata*, Reichb. f., 36, 19.  
— *var. Gebinæ*, Reichb. f., 36, 19.  
*Gebinæ*, Reichb. f., 36, 19.  
*Blumea adenophora*, Franch., 36, 460.  
*alata*, DC., 23, 422.  
*amethystina*, Hance, 23, 420.  
*balsamifera*, DC., 23, 420.  
*barbata*, var. *sericans*, Kurz, 23, 422.  
*Bodinieri*, Vaniot, 36, 460.  
*chinensis*, Hook. et Arn., 23, 420, 429.  
*chinensis*, DC., 23, 420.  
*crinita*, Steetz, 23, 421.  
*Duclouxii*, Vaniot, 36, 460.  
*glandulosa*, Benth., 23, 421.  
*globata*, Vaniot, 36, 460.  
*glomerata*, DC., 23, 420.  
*gracilis*, Dunn, 36, 461.  
*Henryi*, Dunn, 36, 461.  
*hieracifolia*, Benth., 23, 422.  
*hieracifolia*, DC., 23, 421; cf. 420.

- Blumea hongkongensis*, Vaniot, **36**, 461.  
*lacera*, DC., **23**, 421.  
 — *var. glandulosa*, Hook. f., **23**, 421.  
*laciniata*, DC., **23**, 421.  
*Martiniana*, Vaniot, **36**, 461.  
*myriocephala*, DC., **23**, 421.  
*runcinata*, DC., **23**, 421.  
*sericans*, Hook. f., **23**, 422.  
*sonchifolia*, DC., **23**, 421.  
*veronicæfolia*, Franch., **36**, 461.  
*Wightiana*, DC., **23**, 422.  
*Blysmus compressus*, Link, **36**, 248.  
*Blyxa octandra*, Planch., **36**, 2.  
*Bobua japonica*, Miers, **26**, 73.  
*lanceifolia*, Miers, **26**, 73.  
*myrtacea*, Miers, **26**, 73.  
*neriifolia*, Miers, **26**, 74.  
*prunifolia*, Miers, **26**, 74.  
*Bocconia cordata*, Willd., **23**, 35.  
*microcarpa*, Maxim., **36**, 461.  
*Bodiniera*, H. Léveillé, **36**, 461.  
*thalictrifolia*, H. Léveillé, **36**, 461.  
*Boea Clarkeana*, Hemsl. \*, **23**, 232.  
*crassifolia*, Hemsl. \*, **26**, 233.  
*dictyoneura*, Hance, **26**, 233.  
*Hancei*, C. B. Clarke, **26**, 233.  
*hygrometrica*, R. Br., **26**, 234; mentioned, 233.  
*rufescens*, Franch., **26**, 234.  
*Swinhooi*, Hance, **26**, 234.  
*Boehmeria alienata*, Willd., **26**, 490.  
*clidemioides*, Miq., **26**, 487.  
*cochinchinensis*, Spreng., **26**, 490.  
*comosa*, Wedd., **26**, 484.  
*densiflora*, Hook. et Arn., **26**, 484; mentioned, 485.  
*diffusa*, Wedd., **26**, 484.  
*diversiflora*, Miq., **26**, 488.  
*frutescens*, Thunb., **26**, 491.  
*gracilis*, C. H. Wright \*, **26**, 485.  
*grandiflora*, Wedd., **26**, 485.  
*japonica*, *var. platanifolia*, Maxim., **26**, 486.  
 — *var. tricuspis*, Hance, **26**, 488.  
*longispica*, Franch. & Sav., **26**, 485.  
*longispica*, Steud., mentioned, **26**, 488.  
*macrophylla*, D. Don, **26**, 485; mentioned, 484.  
*macrophylla*, Sieb. & Zucc., **26**, 485.  
*nivea*, Gaudich., **26**, 486.  
 — *var. crassifolia*, C. H. Wright \*, **26**, 486.  
 — *var. tenacissima*, Kew Bull., **26**, 486.  
*ourantha*, Miq., **26**, 487.  
*penduliflora*, Wedd., **26**, 485.  
*Boehmeria pilosiuscula*, Hassk., **26**, 487.  
*platanifolia*, Franch. & Savat., **26**, 486.  
*platyphylla*, D. Don, **26**, 487.  
 — *var. clidemioides*, Wedd., **26**, 487.  
 — *var. japonica*, Wedd., **26**, 488.  
 — *var. loochooensis*, Wedd., **26**, 488.  
 — *var. macrostachya*, Wedd., **26**, 487; cf. 485.  
 — *var. scabrella*, Wedd., **26**, 487.  
 — *var. stricta*, C. H. Wright \*, **26**, 487.  
*polystachya*, Wedd., **26**, 488.  
*sidæfolia*, Hook. f., **26**, 484.  
*sidæfolia*, Wedd., **26**, 485.  
*spicata*, Thunb., **26**, 488; mentioned, 485.  
 — *var. duploserrata*, C. H. Wright \*, **26**, 488.  
*squamigera*, Wedd., **26**, 489.  
*tenacissima*, Gaudich., **26**, 486.  
*urantha*, Blume, **26**, 487.  
*Weddeliana*, Vidal, **26**, 484.  
*Zollingeriana*, Wedd., **26**, 488.  
*Boehmeriopsis*, Komarov, **36**, 461.  
*pallida*, Komarov, **36**, 461.  
*Bœninghausenia albiflora*, Reichb., **23**, 102; mentioned, 103.  
*Boerhaavia diffusa*, Choisy, **26**, 317.  
*repanda*, Willd., **26**, 317.  
*repens*, Linn., **26**, 317.  
*Boltonia incisa*, Benth., **23**, 412.  
*indica*, Benth., **23**, 413.  
*japonica*, Miq., mentioned, **23**, 410.  
*Lautureana*, Debeaux, **23**, 413.  
*pekinensis*, Hance, **23**, 412.  
*Bombax malabaricum*, DC., **23**, 89.  
*pentandrum*, Linn., **23**, 89.  
*tenebrosum*, Dunn, **36**, 461.  
*Bonnaya brachiata*, Link & Otto, **26**, 191.  
*grandiflora*, Spreng., **26**, 192.  
*hyssopioides*, Benth., **26**, 191.  
*reptans*, Spreng., **26**, 192.  
*tenuifolia*, Spreng., **26**, 192.  
*verbenæfolia*, Spreng., **26**, 192.  
*veronicæfolia*, Benth., **26**, 192.  
*veronicæfolia*, Spreng., **26**, 192.  
**BORAGINÆ**, **26**, 141.  
*Borreria discolor*, Benth., **23**, 392.  
*Boschniakia himalaica*, Hook. f. & Thoms., **26**, 221.  
*Bostrychanthera deflexa*, Benth., **26**, 298.  
*Bothriospermum asperugoides*, Sieb. & Zucc., **26**, 152.  
*bicarınatum*, Bunge, **26**, 151.

- Bothriospermum bicarunculatum*,  
 Fisch. & Mey., 26, 151.  
*chinense*, Bunge, 26, 151.  
*chinense*, Fisch. & Mey., 26, 151.  
*Kusnezowii*, Bunge, 26, 151.  
*secundum*, Maxim., 26, 151.  
*tenellum*, Fisch. & Mey., 26, 152.
- Botryopleuron*, Hemsl., 36, 461.  
*axillare*, Hemsl., 36, 461.  
*latifolium*, Hemsl., 36, 461.  
*stenostachyum*, Hemsl., 36, 461.  
*venosum*, Hemsl., 36, 461.
- Bournea sinensis*, Oliver, 36, 461.
- Bowringia callicarpa*, Champ., 23, 201.
- Boymia glabrifolia*, Champ., 23, 104.  
*rutæcarpa*, A. Juss., 23, 104.
- Brachyactis chinensis*, Bur. & Franch.,  
 36, 461.  
*ciliata*, Ledeb., 23, 417.
- Brachybotrys paridiformis*, Maxim.,  
 26, 152.
- Brachypodium chinense*, S. Moore, 36,  
 432.  
*sylvaticum*, Beauv., 36, 431.
- Brachyramphus ramosissimus*, Benth.,  
 23, 481.  
*sinicus*, Miq., 23, 481.
- Brachyspatha Konjac*, K. Koch, 36,  
 182.  
*variabilis*, Schott, 36, 183.
- Bradleia mollis*, Steud., 26, 424.  
*pinnata* Roxb., 26, 425.  
*sinensis*, Sieb., 26, 425.  
*sinica*, Gaertn., 26, 425.
- Brandisia discolor*, Hance, 26, 179.  
*Hancei*, Hook. f., 26, 179.  
*racemosa*, Hemsl., 36, 461.
- Brassaiopsis ciliata*, Dunn, 36, 461.  
*ficifolia*, Dunn, 36, 461.  
*Hainla*, Seem., mentioned, 23, 338.  
*ricinifolia*, Seem., 23, 340.
- Brassica campestris*, Linn., 23, 46 ;  
 mentioned, 47.  
*cernua*, Thunb., 23, 47.  
*chinensis*, Linn., 23, 46.  
*juncea*, Hook. f. & Thoms., 23, 47.  
*oleracea*, Lour., 23, 46.  
*violacea*, Linn., 23, 47.
- Brathys japonica*, Blume, 23, 73.  
*laxa*, Blume, 23, 73.
- Braya rubicunda*, Franch., 36, 461.  
*sinensis*, Hemsl., 36, 461.  
*sinuata*, Maxim., 36, 461.
- Bredia Oldhami*, Hook. f., 23, 301.
- Bretschneidera*, Hemsl., 36, 461.  
*sinensis*, Hemsl., 36, 461.
- Breynia fruticosa*, Hook. f., 26, 427.  
*officinalis*, Hemsl. \*, 26, 427.  
*patens*, Rolfe, 26, 428.  
*rhamnoides*, Muell. Arg., 26, 428.
- Bridelia Fordii*, Forbes & Hemsl. \*,  
 26, 419.  
*Lourieri*, Hook. & Arn., 26, 419.  
*retusa*, Spreng., mentioned, 26,  
 419.  
*tomentosa*, Blume, 26, 419.
- Briza elegans*, Osbeck, 36, 421.  
*minor*, Linn., 36, 422.
- Bromus asper*, Murr., 36, 430.  
*japonicus*, Thunb., 36, 430.  
*patulus*, Mert. & Koch, 36, 430.  
*pauciflorus*, Hack., 36, 430.  
*squarrosus*, Linn., 36, 431.  
*tectorum*, Linn., 36, 431.
- Broussonetia Kämpferi*, Sieb., 26,  
 455.  
*Kazinoki*, Sieb., 36, 461.  
*monoica*, Hance, 26, 455.  
*papyrifera*, Vent., 26, 455.
- Bucea sumatrana*, Roxb., 23, 112.
- Bruguiera cylindrica*, Blume, 23, 293.  
*gymnorrhiza*, Engl., 23, 293.  
*gymnorrhiza*, Lam., 23, 294.  
*Rheedii*, Miq., 23, 294.
- Brunella vulgaris*, Franch., 36, 461.
- Bryonia cucumeroides*, Ser., 23, 313.  
*hastata*, Lour., 23, 318.  
*scabrella*, Linn., 23, 319.
- Bryophyllum calycinum*, Salisb., 23,  
 280.
- Buchnera coccinea*, Benth., 26, 201.  
*cruciata*, Buch.-Ham., 26, 201.  
*densiflora*, Hook. & Arn., 26, 201.  
*phænicea*, Benth., 26, 201.  
*stricta*, Benth., 26, 201.
- Buckleya Graebneriana*, Diels, 36,  
 461.  
*Henryi*, Diels, 36, 461.  
*lanceolata*, Miq., 26, 409.  
*Quadriala*, Benth. & Hook. f., 26,  
 409.
- Buddleia acutifolia*, C. H. Wright,  
 36, 461.  
*albiflora*, Hemsl. \*, 26, 118.  
*alternifolia*, Maxim., 26, 119.  
*asiatica*, Lour., 26, 119.  
*crispa*, Benth., 26, 120.  
*curviflora*, Hook. & Arn., 26, 119.  
*curviflora*, hort., 26, 119.  
*Davidi*, Franch., 36, 461.  
*discolor*, Buch.-Ham., 26, 119.  
*Giraldii*, Diels, 36, 461.  
*Hemsleyana*, Koehne, 36, 461.  
*intermedia*, Carr., 26, 119.  
*japonica*, Hemsl. \*, 26, 119 in note.  
*Lindleyana*, Fortune, 26, 119.  
 — var. *sinuatodentata*, Hemsl. \*,  
 26, 120.  
 — var. *sinuato-denticulata*,  
 Hemsl. \*, 26, 567 = præc.

- Buddleia madagascariensis*, Hance, 26, 120.  
*Neemda*, Buch.-Ham., 26, 119.  
*officinalis*, Maxim., 26, 120; 36, 461.  
*paniculata*, Wall., 26, 120.  
*variabilis*, Hemsl. \*, 26, 120.  
*Buergeria floribunda*, Miq., 23, 201.  
*obovata*, Sieb. & Zucc., 23, 23.  
*Buettneria aspera*, Colebr., 23, 92.  
*Bulbophyllum affine*, Lindl., mentioned, 36, 14.  
*bicolor*, Lindl., 36, 14.  
*chinense*, Reichb. f., 36, 14.  
*delitescens*, Hance, 36, 15.  
*odoratissimum*, Lindl., 36, 13.  
*radiatum*, Lindl., 36, 13.  
*retusiusculum*, Reichb. f., 36, 15.  
*tigrinum*, Hance, 36, 15.  
*Watsonianum*, Reichb. f., 36, 14.  
*yunnanense*, Rolfe \*, 36, 14.  
*Bulbostylis barbata*, Kunth, 36, 247.  
*capillaris*, Kunth, 36, 248.  
— var. *trifida*, C. B. Clarke, 36, 248.  
*trifida*, Kunth, 36, 248.  
*Bungea Sheareri*, S. Moore, 26, 203.  
*Bunias tcheliensis*, Debeaux, 23, 49.  
*Buphthalmum oleraceum*, Lour., 23, 431.  
*Bupleurum Candollei*, Wall., 36, 461.  
*chinensis*, DC., 23, 327.  
*chinensis*, Franch., 23, 327.  
*commelinoideum*, Boissieu, 36, 461.  
*falcatum*, Linn., 23, 327; mentioned, 328.  
*gracilipes*, Diels, 36, 461.  
*longicaule*, Wall., 36, 461.  
*longiradiatum*, Turcz., 36, 461.  
*microcephalum*, Diels, 36, 462.  
*octoradiatum*, Bunge, 23, 327.  
*pekinense*, Franch. \*, 23, 327.  
*petiolulatum*, Franch., 36, 462.  
*sachalinense*, F. Schmidt, 36, 462.  
*scorzoneræfolium*, Willd., 23, 327.  
*tenue*, Buch.-Ham., 36, 462.  
*Vanheurckii*, Muell. Arg., 23, 327.  
*yunnanense*, Franch., 36, 462.  
*Burmannia azurea*, Griff., 36, 4.  
*cœlestis*, D. Don, 36, 4.  
*Dalzieli*, Rendle, 36, 4.  
*distachya*, R. Br., 36, 4.  
*disticha*, Linn., 36, 4.  
*nepalensis*, Hook. f., 36, 4.  
*sp.*, Griff., 36, 5.  
*spica gemina Linnæi*, Burm., 36, 4.  
*Wallichii*, Miers, 36, 5.  
**BURMANNIACEÆ**, 36, 4.  
**BURSERACEÆ**, 23, 113.  
*Butomus junceus*, Turcz., 36, 191.  
*umbellatus*, Linn., 36, 191.  
— var. *junceus*, Turcz., 36, 191.  
— var. *minor*, Ledeb., 36, 191.  
*Buxoides aculeata*, Osbeck, 23, 110.  
*Buxus chinensis*, Link, 26, 419.  
*Fortunei*, Carr., 26, 418.  
*Harlandi*, Hance, 26, 418.  
*japonica*, Muell. Arg., 26, 418.  
*microphylla*, Sieb. & Zucc., 26, 418; mentioned, 419.  
*sempervirens*, Linn., 26, 418; mentioned, 419.  
*sinensis*, Link, 26, 418.  
*stenophylla*, Hance, 26, 418.  
*virens*, Thunb., 26, 411.  
*Wallichiana*, Baill., mentioned, 26, 418, 419.  
*Cacalia aconitifolia*, Bunge, 23, 449.  
*aconitifolia*, Miq., 23, 458.  
*auriculata*, DC., 23, 451.  
*bicolor*, Roxb., 23, 447.  
*bulbosa*, Lour., 23, 448.  
*coccinea*, Sims, 23, 449.  
*firma*, Komarov, 36, 462.  
*hastata*, Linn., 23, 456.  
*ovalis*, Ker-Gawl., 23, 448.  
*pinnatifida*, Lour., 23, 448.  
*procumbens*, Lour., 23, 448.  
*sagittata*, Vahl, 23, 449.  
*Cacoucia chinensis*, A. Juss., 23, 294.  
**CACTACEÆ**, 23, 323.  
*Cæsalpinia Bonducella*, Fleming, 23, 205.  
*chinensis*, Roxb., 23, 206.  
*lebbekkoides*, DC., 23, 216.  
*Millettii*, Hook., 23, 205.  
*minax*, Hance, 23, 205.  
*Morsei*, Dunn, 36, 462.  
*Nuga*, Ait., 23, 206.  
*paniculata*, Desf., 23, 206.  
*pulcherrima*, Sw., 23, 206.  
*Sappan*, Linn., 23, 206.  
*sepiaria*, Roxb., 23, 206.  
*vernalis*, Champ., 23, 206.  
**CÆSALPINIÆÆ**, 23, 205.  
*Cajanus bicolor*, DC., 23, 195.  
*flavus*, DC., 23, 195.  
*indicus*, Spreng., 23, 195.  
*Caladium odoratum*, Roxb., 36, 184.  
*odorum*, Lindl., 36, 184.  
*Calamagrostis arundinacea*, Roth, 36, 395.  
*brachytricha*, Steud., 36, 396.  
*collina*, Franch., 36, 396.  
*epigejos*, Roth, 36, 391.  
— var. *densiflora*, Ledeb., 36, 391.

- Calamagrostis hakonensis*, Franch. & Sav., mentioned, 36, 394.  
*Langsdorffii*, Trin., 36, 391.  
*moupinensis*, Franch., 36, 395.  
*Onoei*, Franch., 36, 392.  
*pulchella*, Griseb., 36, 395.  
*robusta*, Franch. & Sav., mentioned, 36, 397.  
*sachalinensis*, Fr. Schmidt, 36, 395.  
*scabrescens*, Griseb., 36, 395.  
 — var. *elatio*r, Griseb., 36, 393.  
*sciuroides*, Franch. & Sav., 36, 398; mentioned, 396.  
*sylvatica*, DC., 36, 395.  
*villosa*, Mutel, 36, 392.  
 — var. *Langsdorffii*, Hack., 36, 391.
- Calamintha chinensis*, Benth., 26, 283.  
*Clinopodium*, Benth., mentioned, 26, 283.  
 — var. *chinensis*, Miq., 26, 283.  
 — var. *urticifolia*, Hance, 26, 283.  
*confinis*, Hance, 26, 283.  
*gracilis*, Benth., 26, 283.  
 — var. *densiflora*, Franch. & Sav., 26, 283.  
*repens*, Benth., 26, 284.  
*sp. nov.?*, 26, 284.  
*umbrosa*, Benth., 26, 284; mentioned, 283.  
 — var. *japonica*, Franch. & Sav., 26, 284.
- Calamus formosanus*, Becc., 36, 170.  
*Henryanus*, Becc., 36, 170.  
*Margaritæ*, Hance, 36, 170.  
*tetradactylus*, Hance, 36, 170.  
*thysanolepis*, Hance, 36, 170.  
*Walkerii*, Hance, 36, 170.
- Calanthe alpina*, Hook., 36, 462.  
*angusta*, Lindl., 36, 24.  
*arcuata*, Rolfe, 36, 25.  
*biloba*, Lindl., 36, 462.  
*brevicornu*, Lindl., mentioned, 36, 25, 26, 27.  
*buccinifera*, Rolfe, 36, 25.  
*Davidi*, Franch., 36, 25, 462.  
*Delavayi*, Franch., 36, 462.  
*ecarinata*, Rolfe, 36, 25.  
*ensifolia*, Rolfe, 36, 25.  
*Fargesii*, Franch., 36, 462.  
*fimbriata*, Franch., 36, 462.  
*formosana*, Rolfe, 36, 25.  
*gracilis*, Lindl., 36, 25.  
*Hancockii*, Rolfe, 36, 25.  
*Henryi*, Rolfe, 36, 25.  
*herbacea*, Lindl., 36, 26.  
*lamellosa*, Rolfe, 36, 26.  
*megalopha*, Franch., 36, 462.
- Calanthe plantaginea*, Lindl., mentioned, 36, 26.  
*puberula*, Lindl., 36, 26.  
*pusilla*, Franch., 36, 462.  
*sp.*, 36, 27.  
*speciosa*, Vieill., 36, 18.  
*striata*, R. Br., mentioned, 36, 25.  
*tricarinata*, Lindl., 36, 26.  
*veratrifolia*, R. Br., 36, 26.  
*yunnanensis*, Rolfe\*, 36, 27.
- Calimeris altaica*, Nees, 23, 408.  
*alyssoides*, DC., 23, 409.  
*biennis*, Ledeb., 23, 412.  
*chinensis*, Sch.-Bip., 23, 408.  
*ciliata*, A. Gray, 23, 409, 416.  
*ciliosa*, Turcz., 23, 408, 416.  
*coronata*, Sch.-Bip., 23, 409.  
*hispida*, Nees, 23, 412.  
*incisa*, DC., 23, 412.  
*integrifolia*, Turcz., 23, 412.  
*tatarica*, Lindl., 23, 412.
- Callerya tomentosa*, Endl., 23, 159.
- Callianthemum cachemirianum*, Camb., 23, 12.
- Callicarpa americana*, Lour., 26, 252.  
*breviceps*, Hance, 26, 252.  
*brevipes*, Hance, 26, 252.  
*cana*, Linn., 26, 252.  
*formosana*, Rolfe, 26, 252, 254.  
*gracilis*, Sieb. & Zucc., 26, 254.  
*incana*, Roxb., 26, 255.  
*integrifolia*, Champ., 26, 253.  
*japonica*, Thunb., 26, 253; mentioned, 252.  
*longifolia*, Hance, 26, 253.  
*longifolia*, Hook., 26, 252.  
*longifolia*, Lam., 26, 253; mentioned, 252, 254.  
 — var. *brevipes*, Benth., 26, 252.  
 — var. *japonica*, DC., 26, 253.  
 — var. *longissima* Hemsl.\* , 26, 253.  
 — var. *subglabra*, Schau., 26, 253.  
*macrophylla*, Vahl, 26, 254.  
*mollis*, Sieb. & Zucc., 26, 254.  
*nudiflora*, Hook. & Arn., 26, 255.  
*pilosissima*, Maxim., 26, 254.  
*purpurea*, Juss., 26, 254; mentioned, 252.  
*Reevesii*, Wall., 26, 254.  
*Roxburghii*, Schau., 26, 255.  
*rubella*, Lindl., 26, 255.  
*tenuifolia*, Champ., 26, 255.  
*tomentosa*, Willd., 26, 255.  
*triloba*, Lour., 26, 255.  
*vastifolia*, Diels, 36, 462.  
*Zollingeriana*, Schau., 26, 254.
- Calligonum asperum*, Lour., 23, 22.
- Callistemma hortense*, Cass., 23, 407.

- Callistephus biennis*, Lindl., 23, 412.  
*chinensis*, Nees, 23, 407.  
*hortensis*, Cass., 23, 407.  
*Callitriche aquatica*, DC., 36, 462.  
*verna*, Linn., 23, 293.  
*Calocedrus macrolepis*, Kurz, 26, 540.  
*Calodium cochinchinensis*, Lour., 26, 393.  
*Calogyne chinensis*, Benth., 26, 1.  
*Calonyction speciosum*, var. *muricatum*, Choisy, 26, 161.  
— var. *vulgare*, Choisy, 26, 158.  
*Calophyllum Inophyllum*, Linn., 23, 74.  
*membranaceum*, Gardn., 23, 74.  
*spectabile*, Hook., 23, 75.  
*Calorhabdos axillaris*, Benth. & Hook. f., 26, 195.  
*Brunoniana*, Benth., 36, 462.  
*cauloptera*, Hance, 26, 195.  
*Fargesii*, Franch., 36, 462.  
*latifolia*, Hemsl. \*, 26, 196.  
*stenostachya*, Hemsl. \*, 26, 196.  
*sutehuenensis*, Franch., 36, 462.  
*venosa*, Hemsl. \*, 26, 197.  
*Calosanthus indica*, Blume, 26, 234.  
*Caloscordum exsertum*, Baker, 36, 120.  
*nerinifolium*, Baker, 36, 120.  
*Calotropis gigantea*, R. Br., 26, 102.  
CALYCANTHACEÆ, 23, 22.  
*Calycanthus præcox*, Linn., 23, 22.  
*Calypso japonica*, Maxim., 36, 462.  
*Calyptranthes mangiferifolia*, Hance, 23, 297.  
*Calysaccion chinense*, Walp., 23, 75.  
*Calysphyrum floridum*, Bunge, 23, 368.  
*pauciflorum*, Bunge, 23, 368.  
*Calystegia acetosæfolia*, Turcz., 26, 164.  
*dahurica*, Choisy, 26, 163.  
*hederacea*, Wall., 26, 164; mentioned, 166.  
*pellita*, Ledeb., 26, 164.  
*pubescens*, Lindl., 26, 164.  
*Sepium*, R. Br., 26, 164.  
*Soldanella*, R. Br., 26, 165.  
*subrotundifolia*, Ledeb., 26, 164.  
*Camellia anemoniflora*, Seem., 23, 82.  
*assimilis*, Champ., 23, 80.  
*axillaris*, [Roxb.] Bot. Reg., 23, 80.  
*Bohea*, Griff., 23, 82.  
*caudata*, Wall., 23, 80.  
*drupifera*, Lour., 23, 80.  
*Edithæ*, Hance, 23, 81.  
*euryoides*, Lindl., 23, 81.  
*Camellia fraterna*, Hance, 23, 81.  
*gracilis*, Hemsl., 36, 462.  
*Grijsii*, Hance, 23, 81.  
*hongkongensis*, Seem., 23, 81.  
*integrifolia*, Choisy, 26, 379.  
*japonica*, Champ., 23, 81.  
*japonica*, Linn., 23, 81.  
*maliflora*, Lindl., 23, 82.  
*oleifera*, Abel, 23, 82.  
*oleosa*, Lour., 23, 82.  
*reticulata*, Lindl., 23, 82.  
*rosæflora*, Hook., 23, 82; mentioned, 83.  
*sulcifolia*, Champ., 23, 82.  
*Sasanqua*, Thunb., 23, 82; mentioned, 81.  
— var. *anemoniflora*, Seem., 23, 82.  
— var. *stricta*, Edwards, 23, 82.  
*spectabilis*, Champ., 23, 82.  
*Thea*, Link, 23, 82.  
*theifera*, Griff., 23, 82.  
*theiformis*, Hance, 23, 81.  
*viridis*, Link, 23, 83.  
*Camirium cordifolium*, Gaertn., 26, 434.  
*Campanula canescens*, Wall., 26, 8.  
*chryso-splenifolia*, Franch., 36, 462.  
*circæoides*, F. Schmidt, 26, 9.  
*colorata*, Wall., 26, 9; mentioned, 10.  
*crenulata*, Franch., 36, 462.  
*Delavayi*, Franch., 36, 462.  
*fulgens*, Wall., 26, 9.  
*gentianoides*, Lam., 26, 5.  
*glomerata*, Linn., 26, 9.  
*grandiflora*, Jacq., 26, 5.  
*nobilis*, Lindl., 26, 10.  
*remotiflora*, Sieb. & Zucc., 26, 13.  
*veronicifolia*, Hance, 26, 10; mentioned, 9.  
*verticillata*, Pall., 26, 14.  
CAMPANULACEÆ, 26, 2.  
*Campanula axillaris*, Oliver, 26, 7.  
*celebica*, C. B. Clarke, 26, 7, 8.  
*japonica*, Maxim., 26, 8.  
*javanica*, Blume, 26, 8.  
*lanceolata*, Sieb. & Zucc., 26, 5.  
*pilosula*, Franch., 26, 8.  
*truncata*, Wall., 36, 462.  
*villosula*, Franch., 26, 8.  
*Camphora chinensis*, Nees 26, 372.  
*Parthenoxylon*, Nees, 26, 372.  
*Campsis adrepens*, Lour., 26, 235.  
*Fortunei*, Seem., 26, 180.  
*Camptotheca acuminata*, Decne., 23, 346.  
*Campuleia coccinea*, Hook., 26, 201.  
*Campylandra Wattii*, Hook. f., 36, 114.

- Campylotropis chinensis*, Bunge, **23**, 182.  
*Cananga odorata*, Hook. f. & Thoms., **23**, 27.  
*Canarium album*, *Raensch.*, **23**, 113.  
*Pimela*, *Kon.*, **23**, 113.  
*sinense*, *Rumph.*, **23**, 113.  
*Canavalia ensiformis*, *DC.*, **23**, 192.  
— *var. turgida*, *Baker*, **23**, 192.  
*gladiata*, *DC.*, **23**, 192.  
*obtusifolia*, *DC.*, **23**, 192.  
*Canna chinensis*, *Willd.*, **36**, 74.  
*indica*, *var. orientalis*, *Hook.*, **36**, 73.  
*indica*, *Linn.*, *partim*, **36**, 73.  
*juncea*, *Retz.*, **36**, 74.  
*orientalis*, *Rosc.*, **36**, 73.  
*Cannabis chinensis*, *Del.*, **26**, 453.  
*sativa*, *Linn.*, **26**, 453.  
— *var. chinensis*, *Franch.*, **26**, 453.  
*Cansjera lanceolata*, *Benth.*, **23**, 115.  
*Rheedii*, *Gmel.*, **23**, 115.  
*Canthium chinense*, *Pers.*, **23**, 381.  
*didymum*, *Gaertn.*, **23**, 384.  
*dubium*, *Lindl.*, **23**, 384.  
*horridum*, *Benth.*, **23**, 385.  
*parvifolium*, *Roxb.*, **23**, 385.  
*undulatum*, *Champ.*, **23**, 385.  
*Capnorchis spectabilis*, *Borck.*, **23**, 36.  
CAPPARIDÆ, **23**, 50.  
*Capparis cantoniensis*, *Lour.*, **23**, 51.  
*falcata*, *Lour.*, **23**, 51.  
*flexicaulis*, *Hance*, **23**, 50.  
*formosana*, *Hemsl.*, **36**, 462.  
*hainanensis*, *Oliver*, **36**, 462.  
*hastigera*, *Hance*, **23**, 51.  
*Henryi*, *Matsum.*, **36**, 462.  
*magna*, *Lour.*, **23**, 51.  
*membranacea*, *Gardn.*, **23**, 50.  
*pumila*, *Champ.*, **23**, 51.  
*sciaphila*, *Hance*, **23**, 51.  
*sepiaria*, *Linn.*, **23**, 51.  
*Swinhooii*, *Hance*, **23**, 51.  
CAPRIFOLIACEÆ, **23**, 347.  
*Caprifolium longiflorum*, *Lindl.*, **23**, 367.  
*Capsella Bursa-pastoris*, *Moench*, **23**, 48.  
*Capsicum annuum*, *Linn.*, **26**, 169.  
*baccatum*, *Linn.*, **26**, 169.  
*frutescens*, *Linn.*, **26**, 169.  
*sinense*, *Jacq.*, **26**, 169.  
*Caragana Altagana*, *Poir.*, **23**, 163.  
*Chamlagu*, *Lam.*, **23**, 163.  
*chinensis*, *Turcz.*, **23**, 163.  
*crassicaulis*, *Benth.*, **36**, 462.  
*frutescens*, *DC.*, **23**, 163.  
*Gerardiana*, *Royle*, **36**, 462.  
*Caragana microphylla*, *Lam.*, **23**, 163.  
*rosea*, *Turcz.*, **23**, 163.  
*Carallia integerrima*, *DC.*, **23**, 294.  
*sinensis*, *Arn.*, **23**, 294.  
*zeylanica*, *Arn.*, **23**, 294.  
*Cardamine Chelidonia*, *Linn.*, **23**, 44.  
*circæoides*, *Hook. f. & Thoms.*, **36**, 462.  
*Delavayi*, *Franch.*, **36**, 462.  
*Engleriana*, *Schulz*, **36**, 462.  
*flexuosa*, *Engl.*, **36**, 463.  
*fragarifolia*, *Schulz*, **36**, 463.  
*Griffithii*, *Hook. f. & Thoms.*, **23**, 43.  
*hirsuta*, *Linn.*, **23**, 43.  
— *subsp. flexuosa*, *Withering*, **23**, 43.  
*Impatiens*, *Linn.*, **23**, 43.  
*insignis*, *Schulz*, **36**, 463.  
*Lamontii*, *Hance*, **23**, 43.  
*leucantha*, *Tausch*, **36**, 463.  
*lyrata*, *Bunge*, **23**, 43.  
*macrophylla*, *Willd.*, **23**, 43.  
*microzyga*, *Schulz*, **36**, 463.  
*multicaulis*, *Hoppe*, **23**, 43.  
*multijuga*, *Franch.*, **36**, 463.  
*paradoxa*, *Hance*, **23**, 44.  
*parviflora*, *Linn.*, **23**, 44.  
*pratensis*, *Linn.*, **23**, 44.  
— *var. ?*, **23**, 43.  
*prorepens*, *Fisch.*, **36**, 463.  
*scaposa*, *Franch.*, **23**, 44.  
*stenoloba*, *Hemsl.*, **36**, 463.  
*sylvatica*, *Link.*, **23**, 43.  
*Tangutorum*, *Schulz*, **36**, 463.  
*tenuifolia*, *Turcz.*, **23**, 44.  
*umbrosa*, *Andrz.*, **23**, 43.  
*Urbaniana*, *Schulz*, **36**, 463.  
*violifolia*, *Schulz*, **36**, 463.  
*yunnanensis*, *Franch.*, **36**, 463.  
*Cardiandra alternifolia*, *Sieb. & Zucc.*, **23**, 278.  
*sinensis*, *Hemsl.*, **36**, 463.  
*Cardiospermum canescens*, *Wall.*, **23**, 138.  
*Corindum*, *Linn.*, **23**, 138.  
*Halicacabum*, *Linn.*, **23**, 138.  
*microcarpum*, *H. B. K.*, **23**, 138.  
*Carduus acaulis*, *Thunb.*, **23**, 461.  
*chinensis*, *DC.*, **23**, 461.  
*crispus*, *Linn.*, **23**, 460.  
*japonicus*, *Franch.*, **23**, 461.  
*lanceolatus*, *Lour.*, **23**, 461.  
*linearis*, *Thunb.*, **23**, 461.  
*multiflorus*, *DC.*, **23**, 461.  
*Segetum*, *Franch.*, **23**, 462.  
*tuberosus*, *Linn.*, **23**, 462.  
*Carex abacta*, *L. H. Bailey*, **36**, 298.



- Carex acutiformis*, Ehrh., 36, 269.  
*agglomerata*, C. B. Clarke\*, 36, 269.  
*albata*, Miq., 36, 301.  
*albomas*, C. B. Clarke\*, 36, 270.  
*alliiformis*, C. B. Clarke\*, 36, 270.  
*alopeuroides*, D. Don, 36, 271.  
— var. *calorostachya*, C. B. Clarke\*, 36, 271.  
*alta*, Boott, 36, 272.  
— var. *Rochebruni*, Franch., 36, 272.  
*amurensis*, Kuek., 36, 273.  
*angusticruris*, Franch., 36, 313.  
*anomala*, Boott, 36, 287.  
*anthericoides*, Presl, 36, 296.  
*aphanolepis*, Franch., 36, 292.  
*arenicola*, F. Schmidt, 36, 272.  
*argyrolepis*, Maxim., 36, 301.  
*aristata*, Boott, 36, 272.  
*aristata*, R. Br., 36, 272.  
— var. *orthostachys*, C. B. Clarke\*, 36, 272.  
*Arnellii*, Christ, 36, 273.  
*arridens*, C. B. Clarke, mentioned, 36, 306.  
*ascotreta*, Franch., 36, 282.  
*atrata*, Linn., 36, 274.  
— var., 36, 284.  
*atrofusca*, Schkuhr, 36, 317.  
*baccans*, Nees, 36, 274.  
*basiflora*, C. B. Clarke\*, 36, 274.  
*bella*, C. B. Clarke, 36, 314.  
*bengalensis*, Boeck., 36, 280.  
*benghalensis*, Roxb., 36, 281.  
*bidentula*, Franch., 36, 275.  
*biwensis*, Franch., 36, 275.  
*Bodinieri*, Franch., 36, 296.  
*Bongardi*, Boott, 36, 275, 276.  
— var., 36, 318.  
*Boottiana*, Hook. & Arn., 36, 275.  
— var. *bracteosa*, C. B. Clarke\*, 36, 275.  
*breviculmis*, R. Br., 36, 276; mentioned, 277.  
*brevicuspis*, C. B. Clarke\*, 36, 277; mentioned, 275.  
*Brizopyrum*, Kunze, 36, 272.  
*Brownii*, Tuckerm., 36, 277.  
*brunnea*, Thunb., 36, 278; mentioned, 299, 311 (brunca).  
*bulbosa*, Boott, 36, 277.  
*Bungeana*, Debeaux, 36, 289.  
*Burbaumii*, Franch., 36, 288.  
*cæspitosa*, Linn., 36, 278.  
*cæspitosa*, Linn., partim, 36, 287.  
*cæspitosa*, Palmst., mentioned, 36, 288.
- Carex capillacea*, Boott, 36, 278; mentioned, 275, 295.  
— var. *yunnanensis*, Franch., 36, 278.  
*capilliformis*, Franch., 36, 279; mentioned, 313.  
*capitellata*, Boiss., 36, 278.  
*cardiolepis*, Nees, 36, 279.  
*celsa*, Boott, 36, 307.  
*cercidaseus*, C. B. Clarke\*, 36, 279.  
*cercostachys*, Franch., 36, 267.  
*cernua*, Boott, 36, 279.  
*cernua*, Franch., 36, 280, 306.  
*chaetorrhiza*, Franch., 36, 272.  
*chinensis*, Retz., 36, 280.  
*chlorostachya*, D. Don, 36, 271.  
*clandestina*, Good., 36, 290.  
*Commersoniana*, Kunth, 36, 278.  
*condensata*, Franch., 36, 281.  
*condensata*, Nees, 36, 280.  
*conica*, Boott, 36, 280.  
*consocialis*, Steud., 36, 271.  
*continua*, C. B. Clarke, 36, 281.  
*cremostachys*, Franch., 36, 281.  
*cruciata*, Franch., 36, 280, 281.  
*cruciata*, Wahlenb., 36, 281.  
*cryptostachys*, Bronq., 36, 281.  
*curaica*, Franch. & Sav., 36, 272.  
*cylindrostachys*, Franch., 36, 282.  
*cyrtostachys*, C. B. Clarke, 36, 282.  
*Davidi*, Franch., 36, 282; mentioned, 286.  
*decidua*, Boott, 36, 287.  
*Delavayi*, Franch., 36, 282.  
*dichroa*, Franch., 36, 314.  
*Dickinsii*, Franch., 36, 282.  
*digitata*, Linn., 36, 283.  
*dimorpholepis*, Steud., 36, 306.  
*dineuros*, C. B. Clarke\*, 36, 283.  
*discoidea*, Boott, 36, 276.  
*dispalata*, Boott, 36, 283.  
*disticha*, DC., 36, 291.  
*divulsa*, Good., 36, 311.  
*Doniana*, Spreng., 36, 271.  
— var., 36, 292.  
*drepanorrhyncha*, Franch., 36, 284.  
*drymophila*, Turcz., 36, 272.  
*Duthiei*, C. B. Clarke, 36, 284.  
*Duvaliana*, Franch. & Sav., mentioned, 36, 318.  
*echinata*, Boeck., 36, 311.  
*echinata*, Murr., 36, 311.  
*Emodorum*, Spreng., 36, 271.  
*excurva*, Boott, 36, 300.  
*Fabri*, Hance, 36, 278.  
*falcata*, Turcz., 36, 317.  
*fallax*, Steud., 36, 301.

- Carex Fargesii*, Franch., **36**, 306.  
*fastigiata*, Franch., **36**, 284.  
*Fedia*, Nees, **36**, 318.  
*ferruginea*, Scop., **36**, 284.  
— var. *tatsiensis*, Franch., **36**, 284.  
*filicina*, Nees, **36**, 285.  
*filipes*, Franch. & Sav., **36**, 285; mentioned, 309.  
*finitima*, Boott, **36**, 285.  
*fluvialilis*, C. B. Clarke, partim, **36**, 316.  
*folliculata*, Rudge, **36**, 298.  
*foraminata*, C. B. Clarke\*, **36**, 285.  
*Forbesii*, C. B. Clarke\*, **36**, 286.  
*Forficula*, Franch. & Sav., **36**, 286; mentioned, 311.  
*fragilis*, Boott, mentioned, **36**, 296.  
*gentilis*, Franch., **36**, 286; mentioned, 299.  
*gibba*, Boott, **36**, 287.  
*gibba*, Wahlenb., **36**, 287.  
*Godenowii*, err. typ., **36**, 287 = seq.  
*Goodenovii*, J. Gay, **36**, 287; mentioned, 278.  
*gracilirostris*, Franch., **36**, 313; mentioned, 300.  
*gracilis*, R. Br., **36**, 278.  
*Hancei*, C. B. Clarke\*, **36**, 288.  
*Hancockiana*, Maxim., **36**, 288.  
*Harlandi*, Boott, **36**, 289.  
*hebecarpa*, C. B. Clarke, **36**, 297.  
— var., **36**, 297.  
*Henryi*, C. B. Clarke\*, **36**, 295.  
*heterolepis*, Bunge, **36**, 289; mentioned, 315.  
*heterostachya*, Bunge, **36**, 289.  
*hirta*, Boiss., **36**, 318.  
*hongkongensis*, Franch., **36**, 294.  
*humilis*, Leysser, **36**, 290.  
*ichangensis*, C. B. Clarke\*, **36**, 290; mentioned, 279.  
*immanis*, C. B. Clarke\*, **36**, 290.  
*incurva*, Lightf., **36**, 291.  
*indica*, Munro, **36**, 281.  
*indica*, Nees, **36**, 280.  
*instabilis*, Boott\*, **36**, 292.  
*intermedia*, Good., **36**, 291.  
*ischnostachya*, Steud., **36**, 292.  
*Jackiana*, Boott, **36**, 292.  
— var. *minor*, C. B. Clarke, **36**, 292.  
*japonica*, Boott, **36**, 271, 299.  
— varr., **36**, 271.  
*japonica*, Thunb., **36**, 292; mentioned, 299.
- Carex lamprosandra*, Franch., **36**, 293.  
*lanceolata*, Boott, **36**, 303.  
— var. *macrosandra*, Franch., **36**, 303.  
*lanceifolia*, C. B. Clarke\*, **36**, 293.  
*Langsdorffii*, Boott, **36**, 276.  
*laticeps*, Franch., **36**, 293.  
*Lehmanni*, Drej., **36**, 293.  
*leiocarpa*, Franch., **36**, 294.  
*leiocarpa*, C. A. Mey., mentioned, **36**, 294.  
*leiorrhyncha*, C. A. Mey., **36**, 294.  
*lenticularis*, D. Don, **36**, 304.  
*leucochlora*, Bunge, **36**, 276.  
*ligata*, Boott, **36**, 294; mentioned, 300, 311.  
*ligulata*, Nees, **36**, 294.  
— var. *angustifolia*, Boott\*, **36**, 292.  
— var. *minor*, Boott\*, **36**, 297.  
*lithosperma*, Forst., **36**, 266.  
*litorrhyncha*, Franch., **36**, 295.  
*littorea*, Labill., **36**, 307.  
*longerostrata*, C. A. Mey., mentioned, **36**, 288, 308.  
*longieruris*, Diels, **36**, 295; mentioned, 287.  
*longieruris*, Nees, **36**, 295.  
— var. *Henryi*, C. B. Clarke\*, **36**, 295.  
*longipes*, D. Don, **36**, 295.  
*longistolon*, Franch., **36**, 296.  
*longkiensis*, Franch., **36**, 296.  
*luctuosa*, Franch., **36**, 295.  
*macrocephala*, Spreng., **36**, 296.  
*macrorrhyncha*, Kar. & Kir., **36**, 303.  
*maculata*, Boott, **36**, 297.  
*manca*, Boott, **36**, 297; mentioned, 309, 310, 311.  
— var. *Harlandi*, Boott, **36**, 289.  
*manciformis*, Franch., **36**, 297.  
*marginaria*, Franch., **36**, 283, 284.  
*Maubertiana*, Boott, **36**, 297.  
*Maximowiczii*, Boeck., **36**, 273.  
*Maximowiczii*, Miq., **36**, 306.  
*melinacra*, Franch., **36**, 298.  
*micans*, Boott, **36**, 297.  
*Michauxiana*, Boeck., **36**, 298.  
*microglochin*, Wahlenb., **36**, 298.  
*microstoma*, Franch., **36**, 298, 299.  
*Mielichhoferi*, Schkuhr, **36**, 284.  
*Mielichhoferi*, Sm., **36**, 317.  
*mollicula*, Boott, **36**, 277.  
*monadelphica*, Boott, **36**, 315.  
*Moorcroftii*, Franch., **36**, 317.  
*mosoynensis*, Franch., **36**, 299.  
*Motoskei*, Miq., **36**, 292.

- Carex moupinensis*, Franch., **36**, 299.  
*multinervia*, Kuek., **36**, 298.  
*muricata*, Boott, **36**, 272.  
*mutans*, Franch., **36**, 267.  
*nana*, Boott, mentioned, **36**, 278.  
*nemostachys*, Steud., **36**, 300.  
*neurocarpa*, Maxim., **36**, 300.  
*nexa*, Boott, **36**, 300.  
*notha*, Baker, **36**, 286.  
*notha*, Kunth, **36**, 286, 298.  
— var., **36**, 286.  
*nubigena*, D. Don, **36**, 300.  
— var. *fallax*, C. B. Clarke\*, **36**, 301.  
*nuptialis*, Boott, **36**, 275.  
*nutans*, Host, **36**, 301; mentioned, 289.  
— var. *pumila*, Boeck., **36**, 307.  
*obscura*, Nees, **36**, 302.  
*ontakensis*, Franch. & Sav., mentioned, **36**, 278.  
*oxyphylla*, Franch., **36**, 292.  
*pachyrrhiza*, Franch., **36**, 302.  
*pallescens*, Linn., forma, **36**, 299.  
*paludosa*, Good., **36**, 269.  
*panicea*, Bunge, **36**, 313.  
*panicea*, Linn., **36**, 302; mentioned, 313.  
*parva*, Nees, **36**, 302.  
*patens*, Franch., **36**, 271.  
*pediformis*, C. A. Mey., **36**, 303.  
— var. *macrosandra*, C. B. Clarke\*, **36**, 303.  
— var. *rostrata*, Maxim., **36**, 303.  
*Peiktusani*, Komarov, **36**, 288.  
*petræa*, Schkuhr, **36**, 309.  
*phacodes*, Spreng., **36**, 303.  
*phacota*, Franch., **36**, 279.  
*phacota*, Spreng., err. typ., **36**, 303.  
*phænocarpa*, Franch., **36**, 304.  
*phæostachya*, Sm., **36**, 317.  
*picta*, Boott, **36**, 306.  
*Pierotii*, Miq., **36**, 304.  
*pisiformis*, Boott, **36**, 304; mentioned, 277.  
*platycarpa*, Steud., **36**, 304.  
*pocilliformis*, Boott, **36**, 315.  
*pollens*, C. B. Clarke\*, **36**, 305.  
— var. *angustior*, C. B. Clarke\*, **36**, 305.  
*præcox*, Franch., **36**, 270.  
*præcox*, Jacq., **36**, 270.  
*Prainii*, C. B. Clarke\*, **36**, 305.  
*Prattii*, Franch., **36**, 317.  
*prava*, Boott, **36**, 311.  
*Prescottiana*, Boott, **36**, 306; mentioned, 280.  
*pruinosa*, Boott, **36**, 306.  
— var. *picta*, Boott, **36**, 306.
- Carex psychophila*, Nees, **36**, 306.  
*pterolepta*, Franch., **36**, 301.  
*pteroloma*, Kunze, **36**, 287.  
*puberula*, Boott, **36**, 276.  
*pumila*, Thunb., **36**, 307; mentioned, 286.  
*rara*, Palibin, **36**, 278.  
*remota*, Linn., **36**, 307.  
— var. *reptans*, Franch., **36**, 308.  
— var. *Rochebruni*, C. B. Clarke\*, **36**, 307.  
*remota*, Thunb., **36**, 287.  
*repanda*, C. B. Clarke, **36**, 308.  
*retrorsa*, Palibin, **36**, 282.  
— var. *minor*, Boott\*, **36**, 282.  
*retrorsa*, Schwein., **36**, 283.  
*Retzii*, Nees, **36**, 280.  
*rhynchophora*, Franch., **36**, 308.  
*rigens*, L. Bail., **36**, 315.  
*rigida*, Good., **36**, 308; mentioned, 296.  
*rigidula*, Franch. \*, **36**, 275.  
*ringens*, Boott, **36**, 277.  
*Ringoldiana*, Boott, **36**, 292.  
*Rochebruni*, Franch., **36**, 272.  
*rostrata*, Mich., **36**, 298.  
*Rouyana*, Franch., **36**, 309.  
*Royleana*, Nees, **36**, 276.  
*rubrobrunea*, C. B. Clarke, **36**, 309.  
*rubrobrunea*, Franch., **36**, 304.  
*rupestris*, All., **36**, 309.  
*Sampsoni*, Hance, **36**, 309; mentioned, 288.  
*saxatilis*, Oeder, **36**, 308.  
*scaposa*, C. B. Clarke, **36**, 309.  
*sclerocarpa*, Franch., **36**, 310.  
*Scopoliana*, Willd., **36**, 284.  
*scotica*, Spreng., **36**, 317.  
*setchuensis*, Franch., **36**, 312.  
*setosa*, Boott, mentioned, **36**, 310.  
*shimidzensis*, Franch., **36**, 280.  
*siccata*, Franch., **36**, 291.  
*siderosticta*, Hance, **36**, 310.  
*simplicissima*, F. Muell., **36**, 278.  
*simulans*, C. B. Clarke\*, **36**, 310.  
*sociata*, Boott, **36**, 311.  
*songarica*, Franch., **36**, 289.  
*songorica*, Kar. & Kir., **36**, 289.  
*Souliei*, Franch., **36**, 302.  
*Spachiana*, Boott, **36**, 311.  
*sparsinux*, Franch., **36**, 285.  
*stachydesma*, Franch., **36**, 313.  
*stellulata*, Good., **36**, 311.  
*stenantha*, Franch. & Sav., **36**, 311.  
*stenolepis*, Franch., **36**, 311.  
*stenophylla*, Wahlenb., **36**, 312.  
*stipitinux*, Franch., **36**, 312.

- Carex stramentitia*, Franch., 36, 281.  
*striata*, R. Br., 36, 277.  
*suberea*, Boott, 36, 304.  
*sutchuensis*, Franch., 36, 312.  
*sylvatica*, Huds., mentioned, 36, 274.  
*sylvatica*, Maxim., 36, 273.  
*sylvatica*, Turcz., 36, 273.  
*taliensis*, Franch., 36, 313.  
*tapintzensis*, Franch., 36, 313.  
*tenebrosa*, Boott, 36, 313; mentioned, 275.  
*tenuissima*, Boott, 36, 313.  
*thibetica*, Franch., 36, 314; mentioned, 284.  
*Thompsoni*, Franch., 36, 314.  
*Thomsoni*, Boott, 36, 314.  
*Tonnerrei*, C. B. Clarke \*, 36, 314; mentioned, 299.  
*transversa*, Boott, 36, 314.  
*Trappistarum*, Franch., 36, 315.  
*trichocarpa*, Muhl., 36, 272; mentioned, 273.  
— var. *orthostachys*, Kuek., 36, 272.  
*trichostyles*, Franch., 36, 292.  
*tristachya*, Thunb., 36, 315.  
*truncatigluma*, C. B. Clarke \*, 36, 315.  
*tsangensis*, Franch., 36, 316.  
*Turczaninowiana*, Meinsh., 36, 273.  
*tumida*, Boott, 36, 316.  
*uncinoides*, Boott, 36, 269.  
*unisexualis*, C. B. Clarke \*, 36, 316.  
*Urvillei*, Brongn., 36, 307.  
*ustulata*, Wahlenb., 36, 307; mentioned, 284.  
*vaginata*, Tausch, 36, 317.  
*valida*, Nees, 36, 281.  
*vesicaria*, Linn., 36, 317.  
*villosa*, Boott, 36, 318.  
*vulgaris*, Fries, 36, 287; mentioned, 275.  
— var. *alpina*, Boott, 36, 308.  
*wahuensis*, Boott, 36, 275, 318; mentioned, 276.  
*Wallichiana*, Prescott, 36, 318.  
*Wightiana*, Boott, 36, 308.  
*Wilfordii*, C. B. Clarke \*, 36, 318.  
*xanthophysa*, varr. *nana* & *minor*, Dewey, 36, 298.  
*yedoensis*, Boeck., 36, 301.  
*yunnanensis*, Franch., 36, 319.  
*Zollingeri*, Kunze, 36, 300.  
*Zollingeri*, Steud., 36, 271.  
*Carissa Carandas*, Linn., 26, 94.  
*Carlesia*, Dunn, 36, 463.  
*sinensis*, Dunn, 36, 463.  
*Carmona heterophylla*, Cav., 26, 144.  
*Carolinella*, Hemsl., 36, 463.  
*cordifolia*, Hemsl., 36, 463.  
*Henryi*, Hemsl., 36, 463.  
*Caroxylon indicum*, Wight, 26, 331.  
*recurvum*, Moq., 26, 331.  
*Carpesium abrotanoides*, Linn., 23, 430.  
*Atkinsoniana*, Hemsl., 36, 463.  
*cernuum*, Linn., 23, 430.  
— var. *lanatum*, C. B. Clarke, 23, 431.  
— var. *trachelifolium*, C. B. Clarke, 23, 431.  
*eximium*, Winkl., 36, 463.  
*Faberi*, Winkl., 36, 463.  
*humile*, Winkl., 36, 463.  
*Lipskyi*, Winkl., 36, 463.  
*minus*, Hemsl. \*, 23, 431; mentioned, 493.  
*trachelifolium*, Less., 23, 431.  
*triste*, Maxim., 36, 463.  
*velutinum*, Winkl., 36, 463.  
*Carpinus cordata*, Blume, 26, 501.  
— var. *chinensis*, Franch., 26, 501.  
*faginea*, Lindl., 26, 501.  
*Fargesii*, Franch., 26, 501.  
*laxiflora*, Blume, 26, 501.  
— var. *Davidi*, Franch., 26, 501.  
— var. *Fargesii*, Burkill \*, 26, 501.  
— var. *macrostachya*, Oliver, 26, 501.  
*laxiflora*, Franch., 26, 501.  
*polyneura*, Franch., 26, 501; mentioned, 502.  
*pubescens*, Burkill \*, 26, 502.  
*Seemeniana*, Diels, 36, 463.  
*Turczaninowii*, Hance, 26, 502; mentioned, 501.  
*viminea*, Sargent, 26, 501.  
*yedoensis*, Maxim., 26, 502.  
— var. *cinerascens*, Franch., 26, 503.  
*Carrieria*, Franch., 26, 503.  
*calycina*, Franch., 36, 463.  
*Carthamus tinctorius*, Linn., mentioned, 23, 470.  
*Carum buriaticum*, Turcz., 23, 328.  
*Candolleianum*, Franch., 36, 463.  
*cardiocarpum*, Franch., 36, 463.  
*Carvi*, Linn., 36, 463.  
*caudatum*, Franch., 36, 463.  
*coriaceum*, Franch., 36, 463.  
*cruciatum*, Franch., 36, 463.  
*Delavayi*, Franch., 36, 463.  
*dissectum*, Franch., 36, 463.  
*flicinum*, Franch., 36, 463.  
*flaccidum*, Franch., 36, 463.  
*Hookeri*, Franch., 36, 463.

- Carum loloense*, Franch., 36, 463.  
*molle*, Franch., 36, 463.  
*neurophyllum*, Maxim., 36, 463.  
*paniculatum*, Franch., 36, 464.  
*purpureum*, Franch., 36, 464.  
*scaberulum*, Franch., 36, 464.  
*schizopetalum*, Franch., 36, 464.  
*sinense*, Franch., 36, 464.  
*Tanakæ*, Franch., 36, 464.  
*tenerum*, Franch., 36, 464.  
*trichomanifolium*, Franch., 36, 464.  
*yunnanense*, Franch., 36, 464.
- CARYOPHYLLEÆ**, 23, 63.
- Caryopteris divaricata*, Maxim., 26, 263.  
*incana*, Miq., 26, 263.  
*Mastacanthus*, Schau., 26, 263.  
*mongholica*, Bunge, 26, 264.  
*nepetaefolia*, Maxim., 26, 264; mentioned, 265.  
*ningpoënsis*, Hemsl. \*, 26, 264.  
*ovata*, Miq., 26, 265.  
*parvifolia*, Batalin, 36, 464.  
*tangutica*, Maxim., 26, 265.  
*terniflora*, Maxim., 26, 265.
- Caryota mitis*, Lour., 36, 167.  
*ochlandra*, Hance, 36, 167.  
*urens*, Jacq., 36, 167.
- Casearia glomerata*, Roxb., 23, 311.  
*membranacea*, Hance, 23, 311.  
*sp.*, 23, 311.  
*subrhombea*, Hance, 23, 311.
- Cassia bicapsularis*, Linn., 23, 210.  
*chinensis*, Jacq., 23, 211.  
*chinensis*, Lam., 23, 210.  
*glauca*, Lam., 23, 210.  
*Horsfieldi*, Miq., 23, 210.  
*mimosoides*, Linn., 23, 210.  
*nictitans*, Linn., 23, 211.  
*obtusifolia*, Linn., 23, 211.  
*occidentalis*, Linn., 23, 211.  
*procumbens*, Lour., 23, 211.  
*Sophera*, Linn., 23, 211; mentioned, 210.  
*suffruticosa*, Koen., 23, 210.  
*Tora*, Linn., 23, 211.  
*torosa*, Cav., 23, 211.
- Cassine japonica*, O. Kuntze, 36, 464.
- Cassiope selaginoides*, Hook. f. & Thoms., 26, 16.
- Cassytha filiformis*, Linn., 26, 393.
- Castanea chinensis*, Hassk., 26, 525.  
*chinensis*, Spreng., 26, 523.  
*concinna*, Champ., 26, 523.  
*Hystrix*, Hook., 26, 524.  
*japonica*, Blume, 26, 525.  
*mollissima*, Blume, 26, 525.  
*regia*, Hance, 26, 512.  
*rufescens*, Hook. f., 26, 524.  
*sativa*, Mill., 26, 525.
- Castanea vesca*, Gaertn., 26, 525.  
*vulgaris*, Lam., 26, 525.  
 — var. *japonica*, DC., 26, 525.  
 — var. *yunnanensis*, Franch., 26, 525.
- Castanopsis caudata*, Franch., 26, 522.  
*chinensis*, Hance, 26, 523.  
*concinna*, A. DC., 26, 523.  
*Delavayi*, Franch., 26, 523.  
*Fabri*, Hance, 26, 523.  
*Fargesii*, Franch., 26, 523.  
*Fordii*, Hance, 26, 523.  
*Henryi*, Skan \*, 26, 523.  
*Hystrix*, A. DC., 26, 524.  
*jucunda*, Hance, 26, 524.  
*Lamontii*, Hance, 26, 524.  
*orthacantha*, Franch., 26, 524.  
*tibetana*, Hance, 26, 524.  
*tribuloides*, A. DC., 26, 524.  
 — var. *formosana*, Skan \*, 26, 524.
- Catalpa Bungei*, C. A. Mey., 26, 234.  
*Fargesii*, Bur., 36, 464.  
*Kämpferi*, Sieb. & Zucc., 26, 235.  
*syringæfolia*, Bunge, 26, 235.  
*umbraculifera*, Hort., 36, 464.  
*vestita*, Diels, 36, 464.
- Catapodium unilaterale*, Hance, 36, 404.
- Catenaria laburnifolia*, Benth., 23, 173.
- Catha Benthami*, Gardn., 23, 122.  
*monosperma*, Benth., 23, 123.
- Cathcartia Delavayi*, Franch., 36, 464.  
*integrifolia*, Maxim., 23, 34.  
*lanceifolia*, Franch., 36, 464.
- Cathetus fasciculata*, Lour., 26, 421.
- Caucalis Anthriscus*, Scop., 23, 337.  
*japonica*, Houtt., 23, 337.  
*orientalis*, Lour., 23, 337.
- Caulinia ovalis*, R. Br., 36, 3.
- Caulophyllum robustum*, Maxim., 36, 464.
- Cedrela Rosmarinus*, Lour., 23, 278.  
*sinensis*, A. Juss., 23, 114.  
*Toona*, Roxb., 36, 464.
- Cedronella cordata*, Benth., mentioned, 26, 293.  
*urticifolia*, Maxim., 26, 293.
- Celastrus alatus*, Thunb., 23, 121.  
*angulatus*, Maxim., 23, 122.  
*articulatus*, Thunb., 23, 122.  
 — var. *punctatus*, Franch., 23, 122.  
*cantonensis*, Hance, 23, 122.  
*Championi*, Benth., 23, 122.  
*ciliidens*, Miq., 23, 123.  
*crispulus*, Regel, 23, 123.  
*diversifolius*, Hemsl. \*, 23, 123.  
*flagellaris*, Rupr., 23, 123.  
*Franchetianus*, Loes., 36, 464.

- Celastrus gemmatus*, *Loes.*, **36**, 464.  
*Hindsii*, *Benth.*, **23**, 123.  
*hypoglaucus*, *Hemsl.*, **36**, 464.  
*hypoleucus*, *Oliver*, **36**, 464.  
*latifolius*, *Hemsl.* \*, **23**, 123.  
*monospermus*, *Roxb.*, **36**, 464.  
*monospermus*, *Benth.*, **23**, 123.  
*orbiculatus*, var., *Loes.*, **36**, 464.  
*paniculatus*, *Willd.*, **36**, 464.  
*punctatus*, *Thunb.*, **23**, 122.  
*racemosus*, *Franch.*, **36**, 464.  
*Rosthornianus*, *Loes.*, **36**, 464.  
*Royleanus*, *Wall.*, **23**, 124.  
*striatus*, *Thunb.*, **23**, 121.  
*stylosus*, *Wall.*, **36**, 464.  
*Tatarinotvii*, *Rupr.*, **23**, 122.  
*variabilis*, *Hemsl.* \*, **23**, 124.  
*Wallichianus*, *Hance*, **23**, 123.  
*Wallichianus*, *Wight & Arn.*, mentioned, **23**, 123.
- Celosia argentea*, *Linn.*, **26**, 318.  
*castrensis*, *Linn.*, **26**, 318.  
*cristata*, *Linn.*, **26**, 318.  
*margaritacea*, *Linn.*, **26**, 318.  
*Swinhoëi*, *Hemsl.* \*, **26**, 318.
- Celsia coromandeliana*, *Vahl*, **26**, 177.
- Celtis amboinensis*, *Willd.*, **26**, 451.  
*Bungeana*, *Blume*, **26**, 449.  
*caudata*, *Hance*, **23**, 221; **26**, 450.  
*chinensis*, *Bunge*, **26**, 449.  
*Davidiana*, *Carr.*, **26**, 450; mentioned, 449.  
*japonica*, *Planch.*, **26**, 449, 450.  
*Muku*, *Sieb. & Zucc.*, **26**, 452.  
*nervosa*, *Hemsl.* \*, **26**, 450.  
*philippinensis*, *Blanco*, **26**, 450.  
*sinensis*, *Pers.*, **26**, 450; mentioned, 449.  
*sinensis*, *Planch.*, **26**, 449.  
*Willdenowiana*, *Roem. & Schult.*, **26**, 450.
- Cenchrus racemosus*, *Linn.*, **36**, 343.
- Centaurea grandiflora*, *Pall.*, **23**, 470.  
*membranacea*, *Lam.*, **23**, 470.  
*monantha*, *Georgi*, **23**, 470.  
*monanthos*, [*Ledeb.*], **23**, 470 = præc.  
*moschata*, *Linn.*, mentioned, **23**, 470.  
*Picris*, *Pall.*, **23**, 470.
- Centella asiatica*, *Urb.*, **36**, 464.  
*Centipeda orbicularis*, *Lour.*, **23**, 440.  
*Centotheca lappacea*, *Desv.*, **36**, 419.  
 — var. *inermis*, *Rendle* \*, **36**, 420.
- Centranthera Brunoniana*, *Wall.*, **26**, 201; mentioned, 202.  
*hispida*, *Benth.*, **26**, 202.
- Centranthera hispida*, *R. Br.*, **26**, 202.  
*humifusa*, *Wall.*, **26**, 202.
- Centratherum chinense*, *Less.*, **23**, 401.  
*Centrophorum chinense*, *Trin.*, **36**, 368.
- Cepa Victorialis*, *Moench*, **36**, 126.  
*Cephalandra indica*, *Naud.*, **23**, 318.
- Cephalanthera erecta*, *Lindl.*, **36**, 48.  
*falcata*, *Lindl.*, **36**, 48.  
*japonica*, *A. Gray*, **36**, 48.  
*longibracteata*, *Blume*, **36**, 464.  
*platycheila*, *Reichb. f.*, **36**, 48.  
*Royleana*, *Regel*, **36**, 49.
- Cephalanthus chinensis*, *Lam.*, **23**, 370.  
*montanus*, *Lour.*, **23**, 370.  
*naucleoides*, *DC.*, **23**, 370.  
*occidentalis*, *Linn.*, **23**, 369.  
*orientalis*, *Roem. & Schult.*, **23**, 370.  
*pilulifera*, *Lam.*, **23**, 370.
- Cephaloschænus articulatus*, *Nees*, **36**, 259.  
*parvus*, *Nees*, **36**, 261.
- Cephalotaxus argotænia*, *Pilg.*, **36**, 464.  
*drupacea*, *Sieb. & Zucc.*, **26**, 544.  
*Fortunei*, *sphalm.*, **26**, 545 = seq.  
*Fortunei*, *Hook.*, **26**, 545.  
 — var. *concolor*, *Franch.*, **26**, 545.  
*Griffithii*, *Oliver*, **26**, 545.  
*koraiana*, *Hort.*, **26**, 546.  
*Mannii*, *Hook. f.*, **26**, 545.  
*Oliveri*, *Mast.*, **26**, 545.  
*pedunculata*, *Sieb. & Zucc.*, **26**, 545.  
 — var. *fastigiata*, *Carr.*, **26**, 546.
- Ceraia simplicissima*, *Lour.*, **36**, 10.
- Cerastium alpinum*, *Linn.*, **36**, 464.  
*aquaticum*, *Linn.*, **23**, 67.  
*arvense*, *Linn.*, **23**, 66.  
*fimbriatum*, *E. Pritz.*, **26**, 464.  
*melanandrum*, *Maxim.*, **23**, 66.  
*petiolare*, *Hance*, **23**, 67.  
*pilosum*, *Ledeb.*, **23**, 67.  
*pumilum*, *Curt.*, **36**, 464.  
*repens*, *Linn.*, **23**, 66.  
*szechuense*, *Williams*, **36**, 465.  
*triviale*, *Link*, **23**, 67.  
*viscosum* et *C. vulgatum*, *auctor.*, **23**, 67.
- Cerasus glandulosa*, *Loisel.*, **23**, 219.  
*japonica*, *Loisel.*, **23**, 219.  
*Padus*, *DC.*, **23**, 220.  
*serrulata*, *G. Don*, **23**, 221.  
*Sieboldii*, *Carr.*, **23**, 221.
- CERATOPHYLLACEÆ, **26**, 538.
- Ceratophyllum demersum*, *Linn.*, **26**, 538.  
*verticillatum*, *Roxb.*, **26**, 538.
- Ceratostigma Griffithii*, *C. B. Clarke*, **36**, 465.

- Ceratostigma plumbaginoides*, Bunge, **26**, 36.  
*Cerbera manghas*, Linn., **26**, 95.  
 Odollam, Gaertn., **26**, 95.  
*Cercis canadensis*, Linn., mentioned, **23**, 213.  
*chinensis*, Bunge, **23**, 213.  
*Griffithii*, Boiss., mentioned, **23**, 213.  
*occidentalis*, Torr., mentioned, **23**, 213.  
*racemosa*, Oliver, **36**, 465.  
*Siliquastrum*, Linn., mentioned, **23**, 213.  
*Ceropegia trichantha*, Hemsl., **26**, 117.  
*Chenomeles japonica*, Lindl., **23**, 257.  
*Cheradoplectron spiranthes*, Schau., **36**, 53.  
*Cherophyllum scabrum*, Thunb., **23**, 337.  
*Chaetaria trichodes*, Nees, **36**, 381.  
*Chaetocyperus costulatus*, Nees, **36**, 225.  
*Limnocharis*, Nees, **36**, 225.  
*Chaetospora imberbis*, R. Br., **36**, 261.  
*japonica*, Franch., **36**, 261.  
*Chailletia bainanensis*, Hance, **23**, 114.  
 CHAILLETIACEÆ, **23**, 114.  
*Chalcas japonensis*, Lour., **23**, 109.  
*paniculata*, Lour., **23**, 109.  
*Chalynochlamis anomala*, Franch., **36**, 341.  
*Chamabainia cuspidata*, Wight, **26**, 489.  
*squamigera*, Wedd., **26**, 489.  
*Chamaecyparis formosensis*, Matsum., **36**, 465.  
*Chamaefistula chinensis*, G. Don, **23**, 211.  
*Chamaemelum limosum*, Maxim., **23**, 439.  
*Chamaeraphis setosa*, var. *Schultzii*, Benth., **36**, 334.  
*spinescens*, Poir., **36**, 339.  
*Chamaerhodos erecta*, Bunge, **23**, 246.  
*Chamaerops excelsa*, Thunb., **36**, 168.  
*Fortunei*, Hook., **36**, 168.  
*khasiana*, Griff., **36**, 169.  
*Chamaesarache heterophylla*, Hemsl.\*, **26**, 174.  
*japonica*, Franch. & Sav., mentioned, **26**, 174.  
*sinensis*, Hemsl. \*, **26**, 174.  
*Chamissoa dichotoma*, Mart., **26**, 321.  
*nodiflora*, Mart., **26**, 321.  
*Champereia Griffithiana*, Planch., **26**, 409.  
*Chapelliera glomerata*, Nees, **36**, 262.  
*Chaptalia lyrata*, Spreng., **23**, 473.  
*Chasalia* sp., **23**, 386.  
*Chavica Benthamiana*, Miq., **26**, 366.  
*Belle*, Seem., **26**, 366.  
*hainana*, C. DC., **36**, 465.  
*leptostachya*, Hance, **26**, 355.  
*officinarum*, Miq., **26**, 364.  
*puberula*, Benth., **26**, 365.  
*sarmentosa*, Miq., **26**, 366.  
*sinensis*, Champ., **26**, 366.  
*Wallichii*, Miq., **26**, 364.  
*Cheiranthus apricus*, Stev., **23**, 44.  
 — var. *trichosepala*, Franch., **23**, 44.  
*aurantiacus*, Bunge, **23**, 39.  
*roseus*, Maxim., **36**, 465.  
*Cheirostylis chinensis*, Rolfe, **36**, 43.  
*flabellata*, Wight, **36**, 43.  
*yunnanensis*, Rolfe \*, **36**, 43.  
*Chelidonium Franchetianum*, Prain, **36**, 465.  
*japonicum*, Thunb., **36**, 465.  
*lasiocarpum*, Oliver, **36**, 465.  
*leptopodium*, Prain, **36**, 465.  
*majus*, Linn., **23**, 35.  
*majus*, Lour., **23**, 35.  
*sinense*, DC., **23**, 35.  
*sutchuenense*, Franch., **36**, 465.  
*Chelonopsis Benthamiana*, Hemsl. \*, **26**, 298.  
*deflexa*, Benth., **36**, 465.  
*moschatus*, Miq., **26**, 298.  
*Chenolea divaricata*, Hook. f., **26**, 327.  
 CHENOPODIACEÆ, **26**, 323.  
*Chenopodina australis*, Moq., **26**, 328.  
*glaucæ*, Moq., **26**, 329.  
*maritima*, Moq., **26**, 328.  
*Salsa*, Moq., **26**, 328.  
*Chenopodium acuminatum*, Willd., **26**, 323.  
*album*, Linn., **26**, 323.  
 — var. *rhombofolium*, Franch., **26**, 323.  
*ambrosioides*, Hance, **26**, 323.  
*ambrosioides*, Linn., **26**, 324.  
*aristatum*, Linn., **26**, 324.  
*australe*, R. Br., **26**, 328.  
*Botrys*, Linn., **26**, 324.  
*bryoniæfolium*, Bunge, **36**, 465.  
*ficifolium*, Sm., **26**, 324.  
*glaucum*, Linn., **26**, 324.  
*hybridum*, Linn., **26**, 325; mentioned, 324.  
*rubrum*, Linn., **36**, 465.  
*Scoparia*, Linn., **26**, 328.  
*urbicum*, Linn., **26**, 325.  
*Vachelli*, Hook. & Arn., **26**, 323.  
*Chimaphila japonica*, Miq., **26**, 33.

- Chimaphila umbellata*, Nutt., mentioned, **26**, 33.  
*Chimonanthus fragrans*, Lindl., **23**, 22.  
     *nitens*, Oliver, **36**, 465.  
*Chionanthus chinensis*, Maxim., **26**, 88.  
     *retusus*, Lindl. & Paxt., **26**, 88.  
     *terminalis*, Roxb., **26**, 92.  
*Chirita anachoreta*, Hance, **26**, 231.  
     *bifolia*, D. Don, mentioned, **26**, 228.  
     *cortusifolia*, Hance, **26**, 231.  
     *depressa*, Hook. f., **36**, 465.  
     *eburnea*, Hance, **26**, 231.  
     *Fauriei*, Franch., **26**, 231.  
     *macrosiphon*, Hance, **26**, 231.  
     *monophylla*, C. B. Clarke, mentioned, **26**, 228.  
     *obtusata*, C. B. Clarke, **26**, 232.  
     *sinensis*, Lindl., **26**, 232.  
     *speciosa*, Kurz, **36**, 465.  
*Chloopsis acaulis*, Blume, **36**, 78.  
**CHLORANTHACEÆ**, **26**, 367.  
*Chloranthus angustifolius*, Oliver, **26**, 367.  
     *brachystachys*, Blume, **26**, 367; mentioned, 369.  
     *denticulatus*, Cordem., **26**, 367.  
     *elatiior*, R. Br., **26**, 369.  
     *Fortunei*, Solms, **26**, 367.  
     *Henryi*, Hemsl. \*, **26**, 367.  
     *inconspicuus*, Sw., **26**, 368.  
     *indicus*, Wight, **26**, 368.  
     *japonicus*, Sieb., **26**, 368.  
     *macrostachys*, R. Br., **26**, 369.  
     *mandshuricus*, Rupr., **26**, 368.  
     *monander*, R. Br., **26**, 369.  
     *monostachys*, R. Br., **26**, 369.  
     *officinalis*, Blume, **26**, 369.  
     *Oldhami*, Solms, **26**, 369.  
     *Pernyanus*, Solms, **26**, 369.  
     *serratus*, Roem. & Schult., **26**, 369.  
*Chloripetalum Benthalianum*, Hance, **26**, 62.  
     *obovatum*, Benth., **26**, 62.  
*Chloris barbata*, Sw., **36**, 403.  
     *caudata*, Trin., **36**, 404.  
     *digitata*, Steud., **36**, 404.  
     *incompleta*, Roth, **36**, 404.  
     *virgata*, Sw., **36**, 404.  
*Chlorophytum chinense*, Franch., **36**, 118.  
     *falcatum*, Baker, **36**, 119.  
     *laxiflorum*, Baker, **36**, 119.  
     *laxum*, R. Br. **36**, 119.  
*Chondrachne articulata*, R. Br., **36**, 258.  
*Choricarpa aphylla*, Boeck., **36**, 258.  
*Chorisis repens*, DC., **23**, 483.  
*Chorispora tenella*, DC., **23**, 50.  
*Chrysanthemum coronarium*, Linn., **23**, 437; mentioned, 438.  
     *indicum*, Linn., **23**, 437.  
     *monspeliense*, Schkuhr, **23**, 438.  
     *oreastrum*, Hance, **23**, 438.  
     *procumbens*, Lour., **23**, 437.  
     *Segetum*, Linn., **23**, 438; mentioned, 437.  
     *sibiricum*, Fisch., **23**, 438.  
     *sinense*, Sabine, **23**, 438.  
     — *var. vestitum*, Hemsl. \*, **23**, 439.  
     *tatsiense*, Bur. & Franch., **36**, 465.  
     *tripartitum*, Sweet, **23**, 437.  
*Chrysophyllum pentagonum*, Hance, **26**, 68.  
     *Roxburghii*, G. Don, **26**, 68.  
*Chrysopogon aciculatus*, Trin., **36**, 368.  
     *pictus*, Hance, **36**, 370.  
     *sinensis*, Rendle \*, **36**, 368.  
*Chrysosplenium alternifolium*, Linn., **23**, 270.  
     *aomorense*, Tokubuchi, **36**, 465.  
     *axillare*, Maxim., **23**, 270.  
     *baicalense*, Maxim., **36**, 465.  
     *ciliatum*, Franch., **36**, 465.  
     *Davidianum*, Decne., **23**, 270.  
     *Delavayi*, Franch., **23**, 270.  
     *flageliferum*, F. Schmidt, **23**, 270.  
     *gracile*, Franch., **36**, 465.  
     *Grayanum*, Maxim., **36**, 465.  
     *Griffithii*, Hook. f. & Thoms., **23**, 270.  
     *Henryi*, Franch., **36**, 465.  
     *macrophyllum*, Oliver, **36**, 465.  
     *microspernum*, Franch., **36**, 465.  
     *nepalense*, Franch., **36**, 465.  
     *nodulosum*, Tokubuchi, **36**, 465.  
     *nudicaule*, Bunge, **23**, 271.  
     *pilosum*, Komarov, **36**, 465.  
     *pumilum*, Franch., **36**, 465.  
     *ramosum*, Maxim., **36**, 465.  
     *sinicum*, Maxim., **23**, 271.  
     *sphærospermum*, Maxim., **23**, 271.  
     *tianschanicum*, Krassn., **36**, 466.  
     *trachyspermum*, Maxim., **23**, 271.  
     *uniflorum*, Maxim., **23**, 271.  
     *villosum*, Franch., **36**, 466.  
     *yunnanense*, Franch., **23**, 271.  
*Chylocalyx perfoliatus*, Hassk., **26**, 344.  
     *senticosus*, Meisn., **26**, 349.  
*Cicca flexuosa*, Sieb. & Zucc., **26**, 421.  
     *microcarpa*, Benth., **26**, 421.  
*Cichorium Endivia*, Linn., mentioned, **23**, 474.  
     *Intybus*, Linn., **23**, 474.  
*Cicutaria aquatica*, Lam., **23**, 328.



- Cicuta sinensis*, [Zuccagni] Roem., **23**, 333.  
*virosa*, Linn., **23**, 328.
- Cimicifuga calthæfolia*, Oliver, **36**, 466.  
*davurica*, Maxim., **23**, 21.  
*fœtida*, Linn., **36**, 466.  
*heracleifolia*, Komarov, **36**, 466.  
*japonica*, Spreng., **36**, 466.  
*simplex*, Wormsk., **36**, 466.
- Cineraria californica*, Lam., **23**, 454.  
*campestris*, Retz., **23**, 451.  
*chinensis*, Spreng., **23**, 457.  
*repanda*, Lour., **23**, 457.  
*sibirica*, Linn., **23**, 454.  
*speciosa*, Schrad., **23**, 454.  
*subdentata*, Bunge, **23**, 451.
- Cinnamomum aromaticum*, Nees, **26**, 371.  
*Burmanni*, Blume, **26**, 371.  
*Camphora*, Nees & Eberm., **26**, 371.  
*Cassia*, Blume, **26**, 371.  
*chinense*, Blume, **26**, 371.  
*Doederleinii*, Engl., **26**, 372.  
*dulce*, Nees, **26**, 371.  
*Neesianum*, Meisn., **26**, 372.  
— var. *petiolare*, Meisn., **26**, 372.  
*Parthenoxylon*, Meisn., **26**, 372.  
*pedunculatum*, Nees, **26**, 372; mentioned, 373.  
— var. *angustifolium*, Hemsl. \*, **26**, 372.  
*Tamala*, Fr. Nees, **26**, 373.  
— var. *albiflorum*, Meisn., **26**, 371.  
*validinerve*, Hance, **26**, 373.  
*zeylanicum*, var. *commune*, Nees, **26**, 373.
- Circæa alpina*, Linn., **23**, 310.  
*cordata*, Royle, **23**, 310.  
*lutetiana*, Linn., **23**, 310.  
*mollis*, Sieb. & Zucc., **23**, 310.  
*quadrisulcata*, Maxim., **36**, 466.
- Circæaster agrestis*, Maxim., **26**, 370.
- Cirrhopetalum bicolor*, Rolfe \*, **36**, 14.  
*chinense*, Lindl., **36**, 14.  
*delitescens*, Rolfe, **36**, 15.  
*Fordii*, Rolfe, **36**, 15.  
*gamosepalum*, Griff., mentioned, **36**, 15.  
*Henryi*, Rolfe \*, **36**, 15.  
*retusiusculum*, Hemsl., **36**, 15.  
*tigrinum*, Rolfe \*, **36**, 15.  
*Wallichii*, Lindl., **36**, 15.
- Cirsium* = *Cnicus*.  
*arvense*, Scop., var. *setosum*, Ledeb., **23**, 460.  
*brevicaule*, A. Gray, **23**, 461.  
*chinense*, Gardn. & Champ., **23**, 461; **36**, 466.  
*Fargesii*, Franch., **36**, 466.  
*Henryi*, Franch., **36**, 466.  
*japonicum*, DC., **23**, 461.  
*lineare*, Sch.-Bip., **23**, 461.  
*littorale*, Maxim., **23**, 461.  
*lyratum*, Bunge, **23**, 463.  
*Moackii*, Maxim., **23**, 461.  
*oreithales*, Hance, **23**, 461.  
*Segetum*, Bunge, **23**, 462.  
*tehefouense*, Debeaux, **23**, 462.  
*Vlassovianum*, Fisch., **23**, 462.  
*Wallichii*, DC., **36**, 466.
- Cissus brevipedunculata*, Maxim., **23**, 133.  
*bryonifolia*, Regel, **23**, 133.  
*cantoniensis*, Hook. & Arn., **23**, 131.  
*carnosa*, Wall., **36**, 466.  
*cordata*, Roxb., **23**, 135.  
*diversifolia*, Walp., **23**, 131.  
*himalayana*, Walp., **23**, 133.  
*japonica*, Willd., **23**, 134.  
*setulosa*, Diels & Gilg, **36**, 466.  
*Thunbergii*, Sieb. & Zucc., **23**, 133.  
*viticifolia*, Sieb. & Zucc., **23**, 136.
- Cistanche salsa*, Benth. & Hook. f., **26**, 222.
- Citrullus vulgaris*, Schrad., mentioned, **23**, 318.
- Citrus Aurantium*, Linn., **23**, 110.  
— var. *japonica*, Hook. f., **23**, 111.  
— var. *sinensis*, Linn., **23**, 111.  
— *sinense*, Gall., **23**, 110.  
*bracteata*, Kurz, **23**, 5.  
*buxifolia*, Poir., **23**, 111.  
*Decumana*, Lour., **23**, 111.  
*delicosa*, Tenore, **23**, 111.  
*digitata* seu *chirocarpus*, Lour., **23**, 111.  
*fusca*, Lour., **23**, 111.  
*Hystrix*, DC., **36**, 466.  
*inermis*, Roxb., **23**, 111.  
*japonica*, Thunb., **23**, 111.  
*madurensis*, Lour., **23**, 111.  
*Medica*, Linn., **23**, 111.  
*microcarpa*, Bunge, **23**, 111.  
*nobilis*, Lour., **23**, 111.  
*sinensis*, Pers., **23**, 110.  
*trifolia*, Thunb., **23**, 111.  
*trifoliata*, Linn., **23**, 111.  
*triptera*, Desf., **23**, 111.  
*vulgaris*, Rasso, **23**, 111.
- Cladium chinense*, Nees, **36**, 262.  
*ensigerum*, Hance, **36**, 262.  
*germanicum*, Schrad., **36**, 262.  
*glomeratum*, R. Br., **36**, 262.  
*jamaicense*, Crantz, **36**, 262.

- Cladium leptostachyum*, Nees, **36**, 262.  
*Mariscus*, R. Br., **36**, 262.  
*Cladrastis amurensis*, Maxim., mentioned, **23**, 201.  
*sinensis*, Hemsl., **36**, 466.  
 sp., **23**, 201.  
*Clandestinaria indica*, Spach, **23**, 40.  
*Claoxylum indicum*, Hassk., **26**, 436.  
*muricatus*, Wight, mentioned, **26**, 440.  
*parviflorum*, Hook. & Arn., **26**, 436.  
*Clausena Wampi*, Oliver, **23**, 110.  
*Claytonia* sp., **23**, 71.  
*Cleidion ulmifolium*, Muell. Arg., **26**, 442.  
*Cleisostoma virginale*, Hance, **36**, 38.  
*Clematis acerifolia*, Maxim., **23**, 1.  
*æthusæfolia*, Turcz., **23**, 1.  
*alpina*, Mill., **23**, 1.  
*angustifolia*, Jacq., **23**, 2.  
*apiifolia*, DC., **23**, 2.  
*Armandi*, Franch., **36**, 466.  
*atragenoides*, Batalin, **36**, 466.  
*Benthamiana*, Hemsl. \*, **23**, 2; mentioned, 3.  
*brachyura*, Maxim., **23**, 2.  
*bracteata*, Kurz, **23**, 5.  
*brevicaudata*, DC., **23**, 3.  
*Buchaniana*, DC., **36**, 466.  
*cadmia*, Buch.-Ham., **23**, 5.  
*cærulea*, Lindl., **23**, 6.  
*casariata*, Hance, **23**, 3.  
*chinensis*, Retz., **23**, 3; mentioned, 2.  
*chrysocoma*, Franch., **36**, 466.  
*chusanensis*, Pluk., **26**, 195.  
*Clarkeana*, H. Léveillé & Vaniot, **36**, 466.  
*coreana*, H. Léveillé & Vaniot, **36**, 466.  
*crassifolia*, Benth., **23**, 3.  
*dasyandra*, Maxim., **36**, 466.  
*Davidiana*, Decne., **23**, 4.  
*Delavayi*, Franch., **36**, 466.  
*Drakeana*, H. Léveillé & Vaniot, **36**, 466.  
*Fargesii*, Franch., **36**, 466.  
*fasciculiflora*, Franch., **36**, 466.  
*Flammula*, Linn., var., **23**, 7.  
*florida*, var. *lanuginosa*, Kuntze, **23**, 5.  
*formosana*, O. Kuntze, **36**, 466.  
*fruticosa*, var., Maxim., **36**, 466.  
*funebri*, H. Léveillé & Vaniot, **36**, 466.  
*fusca*, Turcz., **36**, 466.  
*Gourieana*, Roxb., **36**, 466.  
*grata*, Wall., **23**, 3.  
*Hancockiana*, Maxim., **23**, 4.  
*Clematis hedysarifolia*, DC., **23**, 5; **36**, 466.  
*Henryi*, Oliver, **36**, 466.  
*heracleæfolia*, DC., **23**, 4.  
 — var. *Davidiana*, Franch., **23**, 4.  
*Hookeri*, Decne., **23**, 4.  
*intricata*, Bunge, **36**, 466.  
*Kirilowi*, Maxim., **23**, 5.  
*koreana*, Komarov, **36**, 467.  
*Kuntziana*, H. Léveillé & Vaniot, **36**, 467.  
*lancifolia*, Bur. & Franch., **36**, 467.  
*lanuginosa*, Lindl., **23**, 5.  
*lasiandra*, Maxim., **36**, 467.  
*leiocarpa*, Oliver, **36**, 467.  
*leptomera*, Hance, **23**, 6.  
*longiloba*, DC., **23**, 5.  
*macropetala*, Ledeb., **23**, 5.  
*mandshurica*, Rupr., **23**, 7.  
*Massoniana*, Maxim., **23**, 5.  
*Meyeniana*, Walp., **23**, 5; mentioned, 6.  
*minor*, DC., **23**, 3.  
*montana*, var., Maxim., **36**, 467.  
*nannophylla*, Maxim., **23**, 6.  
*nutans*, var. *æthusifolia*, O. Kuntze, **23**, 1.  
*obscura*, Maxim., **36**, 467.  
*oreophila*, Hance, **23**, 5, 6.  
*orientalis*, Linn., **23**, 6.  
 — var. *intricata*, Maxim., **23**, 6.  
*paniculata*, Thunb., **23**, 6; mentioned, 2, 5.  
*parviloba*, Gardn. & Champ., **23**, 6.  
*patens*, Morr., **23**, 6.  
*pauciflora*, Miq., **23**, 2.  
*Philippiana*, H. Léveillé & Vaniot, **36**, 467.  
*pinnata*, Maxim., **23**, 7.  
*pogonandra*, Maxim., **36**, 467.  
*Potanini*, Maxim., **36**, 467.  
*Prattii*, Hemsl., **36**, 467.  
*pterantha*, Dunn, **36**, 467.  
*ranunculoides*, Franch., **36**, 467.  
*recta*, Linn., **23**, 7; mentioned, 2.  
 — var. *angustifolia*, O. Kuntze, **23**, 1.  
*rubifolia*, C. H. Wright, **36**, 467.  
*sinensis*, Lour., **23**, 3.  
*spectabilis*, Palibin, **36**, 467.  
*splendens*, H. Léveillé & Vaniot, **36**, 467.  
*stans*, Sieb., **23**, 4.  
*Strouachii*, Hance, **23**, 7; mentioned, 5.  
*Tashiroyi*, Maxim., **36**, 467.  
*Tatarinowii*, Maxim., **23**, 7.  
*tchefouensis*, Franch., **23**, 2.  
*tenuiflora*, auct., **23**, 2.

- Clematis tenuiflora*, DC., 23, 2, 7.  
*terniflora*, Benth., 23, 2.  
*terniflora*, DC., 23, 2, 3, 7.  
*ternifolia*, Benth., 23, 2, 3.  
*tubulosa*, Turcz., 23, 4.  
— var. *Hookeri*, Hook., 23, 4.  
*uncinata*, Champ., 23, 7.  
*urophylla*, Franch., 23, 7.  
*virginiana*, Lour., 23, 2.  
*virginica*, Maxim., 23, 2.  
*Vitalba*, var., Franch., 36, 467.  
*Viticella*, var. *flore pleno*, Hook. & Arn., 23, 7.  
*yunnanensis*, Franch., 36, 467.  
*zeylanica*, Poir., 23, 8.  
*Clematoclethra actinidioides*, Maxim., 36, 467.  
*cordifolia*, Franch., 36, 467.  
*Faberi*, Franch., 36, 467.  
*Giraldii*, Diels, 36, 467.  
*Hemsleyi*, Baill., 36, 467.  
*integrifolia*, Maxim., 36, 467.  
*lasioclada*, Maxim., 36, 467.  
*Maximowiczii*, Baill., 36, 467.  
*scandens*, Franch., 36, 467.  
*strigillosa*, Franch., 36, 467.  
*tomentella*, Franch., 36, 467.  
*Cleome icosandra*, Linn., 23, 50.  
*viscosa*, Linn., 23, 50.  
*Clerodendron amplius*, Hance, 26, 259.  
*Bungei*, Steud., 26, 259.  
*canescens*, Wall., 26, 259.  
*castaneifolium*, Hook. & Arn., 26, 261.  
*cyrtophyllum*, Turcz., 26, 259.  
*divaricatum*, Sieb. & Zucc., 26, 263.  
*foetidum*, Bunge, 26, 259.  
*formosanum*, Maxim., 26, 259.  
*Fortunei*, Hemsl. \*, 26, 259.  
*fortunatum*, Linn., 26, 260.  
*fragrans*, Vent., 26, 260.  
*Griffithianum*, C. B. Clarke, 26, 261.  
*hæmatocalyx*, Hance, 26, 259.  
*inerme*, Gaertn., 26, 261.  
*infortunatum*, Gaertn., mentioned, 26, 261.  
*Lindleyi*, Decne., 26, 260.  
*lividum*, Lindl., 26, 261.  
*Mandarinorum*, Diels, 36, 467.  
*moupinense*, Franch., 36, 467.  
*neriifolium*, Wall., 26, 261.  
*oxysepalum*, Miq., 26, 261.  
*paniculatum*, Linn., 26, 261.  
*pentagonum*, Hance, 26, 261.  
*pubescens*, Wall., 26, 259.  
*pumilum*, Spreng., 26, 262.  
*pyramidale*, Andr., 26, 262.  
*Clerodendron sericeum*, "Wall.," 26, 263.  
*serotinum*, Carr., 26, 262.  
*Siphonanthus*, R. Br., 26, 262.  
*squamatum*, Vahl, 26, 262.  
*subscaposum*, Hemsl., 36, 467.  
*trichotomum*, Thunb., 26, 262.  
*Clethra barbinervis*, Sieb. & Zucc., 26, 33.  
*canescens*, Reinw., 26, 33.  
*Delavayi*, Franch., 36, 467.  
*Fabri*, Hance, 26, 34.  
*Fargesii*, Franch., 36, 467.  
*lancifolia*, Turcz., 26, 34.  
*scandens*, Franch., mentioned, 26, 34.  
sp. ?, 26, 34.  
*Cleyera dubia*, Champ., 23, 75.  
*fragrans*, Champ., 23, 75.  
*japonica*, Sieb. & Zucc., 23, 76.  
*japonica*, Thunb., 23, 75.  
*Millettii*, Hook., 23, 76.  
*ochracea*, DC., 23, 76.  
*Clinacanthus Burmanni*, Nees, 26, 244.  
*Clintonia udensis*, Trautv., 36, 144.  
*Clitoria cajanifolia*, Benth., mentioned, 23, 188.  
*Hanceana*, Hemsl. \*, 23, 187.  
*macrophylla*, Hance, 23, 188.  
*macrophylla*, Wall., 23, 188.  
*Ternatea*, Linn., 23, 188.  
*Cnemidia angulosa*, Lindl., 36, 39.  
*semilibera*, Lindl., 36, 40.  
*Cnicus argyranthus*, Hook., 36, 467.  
*arvensis*, Hoffm., 23, 460; mentioned, 462.  
— var. *setosus*, Maxim., 23, 462.  
*Bodinieri*, Vaniot, 36, 468.  
*Cerberus*, Vaniot, 36, 468.  
*chinensis*, Maxim., 23, 461.  
*Fargesii*, Franch., 36, 468.  
*Henryi*, Franch., 36, 468.  
*involutus*, Hook., 36, 468.  
*japonicus*, Maxim., 23, 461.  
*Leduci*, Franch., 36, 468.  
*linearis*, Maxim., 23, 461.  
*monocephalus*, Vaniot, 36, 468.  
*Provosti*, Franch., 36, 468.  
*Segetum*, Maxim., 23, 462.  
*sinensis*, C. B. Clarke, 23, 461.  
*Souliei*, Franch., 36, 468.  
*suffultus*, Franch., 36, 468.  
*tchefouensis*, Franch., 23, 462.  
*Vlassovianus*, Maxim., 23, 462.  
*Cnidium Monnieri*, Cuss., 23, 333.  
*Coccinia cordifolia*, Cogn., 23, 318.  
*indica*, Wight & Arn., 23, 318.  
*Coccoloba platyclada*, [F. Muell.] Bot. Mag., 26, 358.

- Cocculus affinis*, *Oliver*, **36**, 468.  
*cuneatus*, *Benth.*, **23**, 28.  
*diantherus*, *Hook. & Arn.*, **23**, 28.  
*incanus*, *Colebr.*, **23**, 29.  
*macrocarpus*, *Wight & Arn.*, **23**, 28.  
*ovalifolius*, *DC.*, **23**, 28.  
*Thunbergii*, *DC.*, **23**, 28.  
*Cochlearia scapiflora*, *Hook.*, **36**, 468.  
*Cocos nucifera*, *Linn.*, **36**, 170.  
*Codiaeum Pentzii*, *Muell. Arg.*, **26**, 435.  
*umbellatum*, *Hance*, **26**, 435.  
*Codonacanthus pauciflorum*, *Nees*, **26**, 244.  
*Codonopsis convolvulacea*, *Kurz*, **26**, 5; **36**, 468.  
*cordata*, *Hook.*, **26**, 8.  
*Henryi*, *Oliver*, **36**, 468.  
*javanica*, *Hook. f.*, **26**, 8.  
*lanceolata*, *Benth. & Hook. f.*, **26**, 5.  
— var. *ussuriensis*, *Trautv.*, **26**, 6.  
*sylvestris*, *Komarov*, **36**, 468.  
*Tangshen*, *Oliver*, **36**, 468.  
*truncata*, *Wall.*, **26**, 7.  
*ussuriensis*, *Hemsl. \**, **26**, 6.  
*viridiflora*, *Maxim.*, **26**, 6.  
*Cœlachne pulchella*, *R. Br.*, **36**, 399.  
*pulchella*, *Benth.*, **36**, 399.  
*Cœlodiscus lappaceus*, *Kurz*, **26**, 438.  
*Cœloglossum acuminatum*, *Lindl.*, **36**, 53.  
*densum*, *Lindl.*, **36**, 57.  
*lacertiferum*, *Lindl.*, **36**, 53.  
*peristyloides*, *Reichb.*, **36**, 57.  
*Cœlogyne barbata*, *Griff.*, **36**, 468.  
*bulbocodioides*, *Franch.*, **36**, 21; mentioned, 23.  
*chinensis*, *Lindl.*, **36**, 24.  
*corymbosa*, *Lindl.*, **36**, 22.  
*Delavayi*, *Rolfe*, **36**, 21; mentioned, 23.  
*elata*, *Lindl.*, **36**, 22.  
*fimbriata*, *Lindl.*, **36**, 22.  
*grandiflora*, *Rolfe \**, **36**, 22.  
*Henryi*, *Rolfe*, **36**, 22.  
*Mandarinorum*, *Kränzl.*, **36**, 468.  
*nitida*, *Lindl.*, **36**, 468.  
*ochracea*, *Lindl.*, **36**, 468.  
*Pholas*, *Reichb. f.*, **36**, 24.  
*pogonoides*, *Reichb. f.*, **36**, 23.  
*præcox*, *Lindl.*, **36**, 468; mentioned, 22.  
*yunnanensis*, *Rolfe \**, **36**, 23.  
*Cœlonema draboides*, *Maxim.*, **23**, 49.  
*Coix agrestis*, *Lour.*, **36**, 345.  
*chinensis*, *Tod.*, **36**, 345.  
*Lachryma*, *Linn.*, **36**, 345.  
*Lachryma-Jobi*, *Linn.*, **36**, 345.  
*Coldenia procumbens*, *Linn.*, **26**, 146.  
*Colebrookia oppositifolia*, *Lodd.*, **26**, 278.  
*Colocasia Antiquorum*, *Schott*, **36**, 183.  
*Antiquorum*, *Wight*, **36**, 183.  
*indica*, *Benth.*, **36**, 184.  
*nymphæfolia*, *Wight*, **36**, 183.  
*odora*, *Brongn.*, **36**, 184.  
*prunipes*, *C. Koch*, **36**, 183.  
*Colquhounia coccinea*, *Wall.*, **26**, 299.  
*Coluria Henryi*, *Batalin*, **36**, 468.  
*longifolia*, *Maxim.*, **23**, 239.  
*Colutea Delavayi*, *Franch.*, **36**, 468.  
COMBRETACEÆ, **23**, 294.  
*Combretum Alfredi*, *Hance*, **23**, 294.  
*pilosum*, *Roxb.*, **23**, 294.  
*Commelina auriculata*, *Blume*, **36**, 155.  
*benghalensis*, *Linn.*, **36**, 155.  
*communis*, *C. B. Clarke*, **36**, 156.  
*communis*, *Linn.*, **36**, 155.  
— var. *ludens*, *C. B. Clarke*, **36**, 156.  
*delicatula*, *Schlecht.*, **36**, 155.  
*japonica*, *Thunb.*, mentioned, **36**, 152.  
*longicaulis*, *Jacq.*, **36**, 156.  
*Loureirii*, *Kunth*, **36**, 156.  
*ludens*, *Miq.*, **36**, 156.  
*medica*, *Lour.*, **36**, 153.  
*nudiflora*, *Linn.*, **36**, 156; mentioned, 153.  
*obliqua*, *Buch.-Ham.*, **36**, 157.  
*ochreatea*, *Schau.*, **36**, 156.  
*polygama*, *Roth*, **36**, 156.  
*polyspatha*, *Wight*, **36**, 157.  
*procurrens*, *Schlecht.*, **36**, 155.  
*salicifolia*, *Benth.*, **36**, 156.  
*sinica*, *Roem. & Schult.*, **36**, 154.  
*tuberosa*, *Lour.*, **36**, 152.  
*undulata*, *R. Br.*, **36**, 157.  
*vulgaris*, *Schmidel*, **36**, 156.  
COMMELINACEÆ, **36**, 151.  
COMPOSITÆ, **23**, 400.  
*Conchidium sinicum*, *Lindl.*, **26**, 17.  
CONIFERÆ, **26**, 540.  
*Conioselinum Czernævia*, *Fisch. & Mey.*, **23**, 147.  
CONNARACEÆ, **23**, 149.  
*Connarus juglandifolius*, *Hook. & Arn.*, **23**, 147.  
*microphylla*, *Hook. & Arn.*, **23**, 149.  
*Roxburghii*, *Hook. & Arn.*, **23**, 150.  
*Conocephalus amœnus*, *King*, mentioned, **26**, 471.

- Conocephalus sinensis*, *C. H. Wright*,  
**26**, 471.  
*Conophallus Konjac*, *Schott*, **36**, 182.  
*Conopodium Cyminum*, *Benth.*, **23**,  
 330.  
*Convallaria cirrhifolia*, *Wall.*, **36**,  
 104.  
     *commuta*, *Schult.*, **36**, 106.  
     *Govaniana*, *Wall.*, **36**, 106.  
     *japonica*, *Linn.*, **36**, 78.  
     — var. *major*, *Thunb.*, **36**, 77.  
     — var. *minor*, *Thunb.*, **36**, 78.  
     *majalis*, *Linn.*, **36**, 112.  
     *multiflora*, *Linn.*, **36**, 106.  
     *Polygonatum*, *Linn.*, **36**, 107.  
     *rosea*, *Ledeb.*, **36**, 108.  
     *spicata*, *Thunb.*, **36**, 79.  
     *umbellata*, *Bunge*, **36**, 106.  
     *verticillata*, *Linn.*, **36**, 109.  
 CONVULVULACEÆ, **26**, 156.  
*Convolvulus acetosæfolius*, *Turcz.*, **26**,  
 164.  
     *althæoides*, *Linn.*, **26**, 165.  
     *Ammannii*, *Desr.*, **26**, 165.  
     *arvensis*, *Linn.*, **26**, 165.  
     *Batatas*, *Linn.*, **26**, 157.  
     *biflorus*, *Linn.*, **26**, 157.  
     *bryoniæfolius*, *Willd.*, mentioned,  
     **26**, 165.  
     *calycinus*, *Roxb.*, **26**, 160.  
     *calystegoides*, *Choisy*, **26**, 165.  
     *chinensis*, *Ker-Gawl.*, **26**, 165.  
     *dahuricus*, *Sims*, **26**, 164.  
     *fastigiatus*, *Roxb.*, **26**, 159.  
     *Hardwickii*, *Spreng.*, **26**, 160.  
     *ianthinus*, *Hance*, **26**, 166.  
     *involutus*, *Ker-Gawl.*, **26**, 163.  
     *Nil*, *Linn.*, **26**, 160.  
     *obscurus*, *Linn.*, **26**, 161.  
     *parviflorus*, *Vahl*, **26**, 166.  
     *pellita*, *Ledeb.*, **26**, 164.  
     *repens*, *Vahl*, **26**, 157.  
     *reptans*, *Linn.*, **26**, 157.  
     *Sepium*, *Linn.*, **26**, 164.  
     — var. *americanus*, [*Sims*] *Bot.*  
     *Mag.*, **26**, 164.  
     — var. *brevipes*, *Franch.*, **26**, 164.  
     *sinensis*, *Desr.*, **26**, 162.  
     *Soldanella*, *Linn.*, **26**, 165.  
     *spinosus*, *Bunge*, **26**, 166.  
     *subvolubilis*, *Ledeb.*, **26**, 164.  
     *tomentosus*, *Lour.*, **26**, 163.  
     *tragacanthoides*, *Turcz.*, **26**, 166.  
     *Turpethum*, *Linn.*, **26**, 163.  
*Conyza ægyptiaca*, *Ait.*, **23**, 419.  
     *altaica*, *DC.*, **23**, 417.  
     *ambigua*, *DC.*, **23**, 418.  
     *balsamifera*, *Linn.*, **23**, 420.  
     *chinensis*, *Lam.*, **23**, 401.  
     *chinensis*, *Linn.*, **23**, 420.  
*Conyza hirsuta*, *Linn.*, **23**, 422.  
     *indica*, *Miq.*, **23**, 422.  
     *japonica*, *Less.*, **23**, 419.  
     *lanceolaria*, *Roxb.*, **23**, 421.  
     *leucodasys*, *Miq.*, **23**, 419.  
     *patula*, *Ait.*, **23**, 401.  
     *pinnatifida*, *Franch.*, **36**, 468.  
     *pyrifolia*, *Lam.*, **23**, 419.  
     *syringæfolia*, *Meyen & Walp.*, **23**,  
     420.  
     *veronicæfolia*, *Wall.*, **23**, 419.  
*Cookia punctata*, *Sonnerat*, **23**, 110.  
     *Wampi*, *Blanco*, **23**, 110.  
*Coprosma*, *Forst.*, mentioned, **23**,  
 391.  
*Coprosmanthus japonicus*, *Kunth*, **36**,  
 96.  
*Coptis chinensis*, *Franch.*, **36**, 468.  
     *Teeta*, *Wall.*, **23**, 17.  
*Corallodiscus*, *Batalin*, **36**, 468.  
     *conchæfolius*, *Batalin*, **36**, 468.  
*Corallorhiza innata*, *R. Br.*, **36**, 9.  
*Corchoropsis crenata*, *Sieb. & Zucc.*,  
**23**, 94.  
     *psilocarpa*, *Harms & Loes.*, **36**, 468.  
*Corchorus acutangulus*, *Lam.*, **23**,  
 93.  
     *capsularis*, *Linn.*, **23**, 93.  
     *hirtus*, *Thunb.*, **26**, 449.  
     *japonicus*, *Thunb.*, **23**, 229.  
     *olitorius*, *Linn.*, **36**, 468.  
     *serratus*, *Thunb.*, **26**, 449.  
*Cordia Loureiri*, *Roem. & Schult.*, **26**,  
 143.  
     *Myxa*, *Linn.*, **26**, 143.  
     *sinensis*, *Lam.*, **26**, 143.  
     *thyrsiflora*, *Sieb. & Zucc.*, **26**, 144.  
     *venosa*, *Hemsl. \**, **26**, 143.  
*Cordyline Jacquini*, *Kunth*, **36**, 118.  
     *terminalis*, *Kunth*, **36**, 118.  
     — var. *ferrea*, *Baker*, **36**, 118.  
*Cordyloblaste Henschelii*, *Mor.*, **26**,  
 73.  
*Coreopsis biternata*, *Lour.*, **23**, 434.  
     *leucorhiza*, *Lour.*, **23**, 435.  
*Coriandrum sativum*, *Linn.*, **23**, 336.  
*Coriaria intermedia*, *Matsum.*, **36**, 468.  
     *nepalensis*, *Wall.*, **23**, 149; **36**,  
     468.  
     *sinica*, *Maxim.*, **23**, 149.  
     *terminalis*, *Hemsl.*, **36**, 468.  
 CORIARIÆ, **23**, 149.  
*Corispermum elongatum*, *Bunge*, **36**,  
 468.  
     *hyssopifolium*, *Linn.*, **26**, 326;  
     mentioned, 327.  
     *orientale*, *Lam.*, **26**, 327.  
     *Redowskii*, *Fisch.*, **26**, 327.  
     *Stauntoni*, *Moq.*, **26**, 327.  
     *tylocarpum*, *Hance*, **26**, 327.

## CORNACEÆ, 23, 344.

- Cornus alba*, Linn., 23, 344.  
*brachypoda*, O. A. Mey., 23, 345.  
*Bretschneideri*, J. Henry, 36, 468.  
*canadensis*, Linn., 23, 344.  
*capitata*, Wall., 23, 345.  
*crispula*, Hance, 23, 345.  
— var. *foliis latioribus*, 23, 346.  
*hongkongensis*, Hemsl. \*, 23, 345.  
*Kousa*, Buerger, 23, 345.  
*macrophylla*, Wall., 23, 345.  
*Mas*, Linn., mentioned, 23, 345.  
*officinalis*, Sieb. & Zucc., 23, 345.  
*paucinervis*, Hance, 23, 346.  
*quinquenervis*, Franch., 36, 468.  
*scabridus*, Franch., 36, 468.  
*tartarica*, Mill., 23, 344.  
*Cornutia quinata*, Lour., 26, 257.  
*Cortusa Matthioli*, Linn., 26, 46.  
*pekinensis*, A. Richt., 36, 468.  
*Corydalis acuminata*, Franch., 36, 469.  
*adunca*, Maxim., 23, 36.  
*ægopodioides*, H. Léveillé & Vaniot, 36, 469.  
*albicaulis*, var., Franch., 36, 469.  
*anthriscifolia*, Franch., 36, 469.  
*aurea*, Willd., mentioned, 23, 38.  
— var. *speciosa*, Baker & S. Moore, 23, 38.  
*balsamiflora*, Prain, 36, 469.  
*Bataliana*, Prain, 36, 469.  
*bulbosa*, DC., 23, 36; mentioned, 39.  
*Bungeana*, Turcz., 23, 36.  
— var. *odontopetala*, Hemsl. \*, 23, 36.  
*cheilanthifolia*, Hemsl., 36, 469.  
*cheirifolia*, Franch., 36, 469.  
*chelonifolia*, H. Léveillé & Vaniot, 36, 469.  
*chinensis*, Franch., 23, 37.  
*conspersa*, Maxim., 36, 469.  
*Crista-galli*, Maxim., 36, 469.  
*cristata*, Maxim., 36, 469.  
*curviflora*, Maxim., 23, 37; 36, 469.  
*dasyptera*, Maxim., 23, 37.  
*daucifolia*, H. Léveillé & Vaniot, 36, 469.  
*Davidi*, Franch., 36, 469.  
*decumbens*, Pers., 23, 37.  
*Delavayi*, Franch., 36, 469.  
*Drakeana*, Prain, 36, 469.  
*Duelouxii*, H. Léveillé & Vaniot, 36, 469.  
*echinocarpa*, Franch., 36, 469.  
*edulis*, Maxim., 23, 37.  
*elata*, Bur. & Franch., 36, 469.  
*Fargesii*, Franch., 36, 469.  
*flexuosa*, Franch., 36, 469.

- Corydalis Fumaria*, H. Léveillé & Vaniot, 36, 469.  
*gamosepala*, Maxim., 23, 36.  
*gigantea*, Trautv. & Mey., 36, 469.  
*gracilipes*, S. Moore, 23, 37.  
*gracilis*, Franch., 36, 469.  
*hamata*, Franch., 36, 469.  
*Hannæ*, Kunitz, 36, 469.  
*Hemsleyana*, Franch. & Prain, 36, 469.  
*heterocarpa*, Sieb. & Zucc., 23, 38.  
*incisa*, Pers., 23, 37.  
*linarioides*, Maxim., 23, 37.  
*livida*, Maxim., 36, 469.  
*longicornu*, Franch., 36, 469.  
*lopinensis*, Franch., 36, 469.  
*Martini*, H. Léveillé & Vaniot, 36, 469.  
*melanochlora*, Maxim., 23, 37.  
*micropoda*, Franch., 23, 37.  
*moupinensis*, Franch., 36, 469.  
*mucronata*, Franch., 36, 470.  
*ochotensis*, Turcz., 23, 37.  
*ophiocarpa*, Hook. f. & Thoms., 36, 470.  
*oxypetala*, Franch., 36, 470.  
*pachycentra*, Franch., 36, 470.  
*pallida*, Pers., 23, 37; mentioned, 38.  
*petrophila*, Franch., 36, 470.  
*Potanini*, Maxim., 36, 470.  
*Prattii*, Franch., 36, 470.  
*pulchella*, Franch., 36, 470.  
*racemosa*, Bunge, 23, 36.  
*racemosa*, Pers., 23, 38.  
*Raddeana*, Regel, 23, 37.  
*remota*, Baker & S. Moore, 23, 36.  
*remota*, Maxim., 23, 36.  
*rosea*, Maxim., 23, 38.  
*scaberula*, Maxim., 36, 470.  
*scandens*, Franch., 36, 470.  
*Sheareri*, S. Moore, 23, 38.  
*solida*, Sw., 23, 36; 36, 470.  
*Souliei*, Franch., 36, 470.  
*speciosa*, Maxim., 23, 38.  
*spectabilis*, Willd., 23, 36.  
*stenantha*, Franch., 36, 470.  
*straminea*, Maxim., 23, 38.  
*streptocarpa*, Maxim., 23, 38.  
*suaveoleus*, Hance, 23, 39.  
*taliensis*, Franch., 36, 470.  
*temulifolia*, Franch., 36, 470.  
*thalictrifolia*, Franch., 36, 470.  
*thibetica*, var., Franch., 36, 470.  
*tomentella*, Franch., 36, 470.  
*tomentosa*, N. E. Br., 36, 470.  
*tongolensis*, Franch., 36, 470.  
*trachycarpa*, Maxim., 23, 39; mentioned, 38.  
*trifoliata*, Franch., 36, 470.

- Corydalis trisecta*, Franch., 36, 470.  
*triternata*, Franch., 36, 470.  
*Wilfordi*, Regel, 23, 38.  
*Wilsoni*, N. E. Br., 36, 470.  
*yunnanensis*, Franch., 36, 470.  
*Corylopsis multiflora*, Hance, 23, 290.  
*sp.*, Diels, 36, 470.  
*Corylus Avellana*, Baker & S. Moore, 26, 504.  
*chinensis*, Franch., 26, 503.  
*Columna*, Linn., 26, 503; mentioned, 504.  
— *var. chinensis*, Burkill\*, 26, 503.  
*Davidiana*, Baill., 26, 503.  
*ferox*, Wall., 26, 503.  
— *var. thibetica*, Franch., 26, 503.  
*heterophylla*, Fisch., 26, 504.  
— *var. Crista-galli*, Burkill\*, 26, 504.  
— *var. sutchuenensis*, Franch., 26, 504.  
— *var. yunnanensis*, Franch., 26, 504.  
*mandshurica*, Maxim., 26, 505.  
— *var. Fargesii*, Franch., 26, 505.  
*rostrata*, *var. mandshurica*, Regel, 26, 505.  
*thibetica*, Batalin, 26, 503.  
*Costus arabicus*, Jacq., 36, 71.  
*speciosus*, Sm., 36, 71.  
*Cotinus coggyria*, Scop., 23, 146.  
*Cotoneaster acuminata*, Lindl., 36, 470.  
*acutifolia*, Maxim., 23, 260.  
*angustifolia*, Franch., 36, 470.  
*bacillaris*, Wall., 36, 470.  
*buxifolia*, Wall., 36, 470.  
*coriacea*, Franch., 36, 470.  
*Dielsiana*, E. Pritz., 36, 470.  
*Fontanesii*, Spach, mentioned, 23, 261.  
*Fortuncana*, Wenzig, 23, 262.  
*Francheti*, Bois, 36, 470.  
*frigida*, Wall., 36, 470.  
*glaucophylla*, Franch., 36, 470.  
*integerrima*, Medic., 23, 260; mentioned, 261.  
*microphylla*, Wall., 23, 261.  
*moupinensis*, Franch., 36, 470.  
*mucronata*, Franch., 36, 470.  
*multiflora*, Bunge, mentioned, 23, 261.  
*nummularia*, Fisch. & Mey., 36, 470.  
*pannosa*, Franch., 36, 470.  
*Pyracantha*, Spach, 23, 260.  
*Cotoneaster reflexa*, Carr., 23, 261.  
*rotundifolia*, Wall., 36, 470.  
*rugosa*, E. Pritz., 36, 470.  
*salicifolia*, Franch., 36, 471.  
*vulgaris*, Lindl., 23, 26.  
*Cottonia Championi*, Lindl., 36, 34.  
*Cotula anthemoides*, Linn., 23, 439.  
*Cotyledon fimbriata*, Turcz., 23, 281.  
*japonica*, Maxim., 36, 471.  
*laciniata*, Linn., 23, 281.  
*leucantha*, Ledeb., 23, 281.  
*malacophylla*, Pall., 23, 282.  
*minuta*, Komarov, 36, 471.  
*spinosa*, C. B. Clarke, 23, 281.  
*spinosa*, Linn., 23, 282.  
*thyrsiflora*, Maxim., 23, 281.  
*Covellia demonum*, Miq., 26, 462.  
*dasycarpa*, Miq., 26, 462.  
*hispida*, Miq., 26, 462.  
*oppositifolia*, Miq., 26, 462.  
*radiata*, Miq., 26, 464.  
*rapiformis*, Miq., 26, 464.  
*venosa*, Miq., 26, 464.  
*Crantzia lineata*, Linn., mentioned, 23, 281.  
*Crassula mongolica*, Franch., 23, 280.  
*pinnata*, Linn., 23, 280.  
*yunnanensis*, Franch., 36, 471.  
*Cratægus Bibas*, Lour., 23, 261.  
*crenulata*, Roxb., 23, 260.  
*cuneata*, Sieh. & Zucc., 23, 259.  
*dsungarica*, Zabel, 36, 471.  
*glabra*, Hort., 23, 263.  
*glabra*, Thunb., 23, 262; mentioned, 263.  
*Henryi*, Dunn, 36, 471.  
*indica*, Linn., 23, 264.  
*lævis*, Thunb., 23, 263.  
*Oxyacantha*, *var. pinnatifida*, Regel, 23, 259.  
*pinnatifida*, Bunge, 23, 259.  
*Pyracantha*, Pers., 23, 260; mentioned, 262.  
*rubra*, Lour., 23, 264.  
*sanguinea*, Pall., 23, 260.  
*serratifolia*, Desf., 23, 263.  
*spiralis*, Steud., 23, 264.  
*tenuifolia*, Komarov, 36, 471.  
*villosa*, Thunb., 23, 263.  
*Cratæva falcata*, DC., 23, 51.  
*magna*, DC., 23, 51.  
*Nurvala*, Buch.-Ham., 23, 51.  
*religiosa*, Forst., 23, 51.  
*Cratoxylon cochinchinense*, Blume, 23, 74.  
*ligustrinum*, Blume, 23, 74.  
*petiolatum*, Blume, 23, 74.  
*polyanthum*, Korth., 23, 74.  
*Crawfordia affinis*, Wall., 26, 123.  
*Delavayi*, Franch., 36, 471.

- Crawfordia fasciculata*, Wall., 26, 122.  
*japonica*, Sieb. & Zucc., 26, 122.  
*luteo-viridis*, C. B. Clarke, 26, 123.  
*Pterygocalyx*, Hemsl.\*, 26, 123.  
 sp., 26, 123.  
*thibetica*, Franch., 36, 471.
- Cremanthodium discoideum*, Maxim., 23, 447.  
*humile*, Maxim., 23, 447.  
*lineare*, Maxim., 23, 447.  
*plantagineum*, Maxim., 23, 447.  
*Potanini*, Winkl., 36, 471.
- Creodus odorifer*, Lour., 26, 368.
- Crepis Bockiana*, Diels, 36, 471.  
*Gillii*, S. Moore, 36, 471.  
*Henryi*, Diels, 36, 471.  
*heterophylla*, Hemsl.\*, 23, 475.  
*Hookerana*, C. B. Clarke, 36, 471.  
*integra*, Miq., 23, 475.  
*japonica*, Benth., 23, 475; mentioned, 476, 477.  
*longipes*, Hemsl.\*, 23, 476; mentioned, 475.  
*lyrata*, Benth., 23, 475.  
*nana*, Sch.-Bip., 23, 475.  
*Phoenix*, Dunn, 36, 471.  
*preanthoides*, Hemsl.\*, 23, 477.  
*rapunculoides*, Dunn, 36, 471.  
*Rosthornii*, Diels, 36, 471.  
*tangeana*, Miq., 23, 475.  
*trichocarpa*, Franch., 36, 471.  
*Turczaninowii*, C. A. Mey., 23, 477.  
*Umbrella*, Franch., 36, 471.
- Crinum anomalum*, Herb., 36, 88.  
*asiaticum*, Benth., 36, 88.  
*asiaticum*, Linn., 36, 88.  
 — *var. anomalum*, Herb., 36, 88.  
 — *var. declinatum*, Herb., 36, 68.  
 — *var. sinicum*, Baker, 36, 88.  
*declinatum*, Herb., 36, 88.  
*latifolium*, Thunb., 36, 88.  
*Loureirii*, M. Roem., 36, 88.  
*plicatum*, Livingst., 36, 88.  
*sinicum*, Roxb., 36, 88.  
*zeylanicum*, Lour., 36, 88.
- Crithmum*, mentioned, 23, 332.
- Crossandra infundibuliformis*, Nees, 26, 243.  
*undulæfolia*, Salisb., 26, 243.
- Crossostephium artemisioides*, Less., 23, 440.
- Crotalaria albida*, Heyne, 23, 150.  
*brevipes*, Champ., 23, 152.  
*calycina*, Schrank, 23, 151.  
*capitata*, Benth., 36, 471.  
*chinensis*, Lam., 23, 152.  
*chinensis*, Linn., 23, 151.  
*elliptica*, Roxb., 23, 151.  
*eriantha*, Sieb. & Zucc., 23, 152.
- Crotalaria ferruginea*, R. Grah., 23, 151.  
*formosana*, Matsum., 36, 471.  
*juncea*, Linn., 23, 151.  
*leiocarpus*, Vogel, 23, 151.  
*linifolia*, Linn., 23, 151.  
*medicaginea*, Lam., 23, 152.  
*mollis*, Weinm., 23, 152.  
*Oldhami*, Miq., 23, 152.  
*paniculata*, Willd., 23, 152.  
*retusa*, Linn., 23, 152.  
*rufescens*, Franch., 23, 151; 36, 471.  
*sessiliflora*, Linn., 23, 152.  
*similis*, Hemsl., 36, 471.  
*splendens*, Vogel, 23, 151.  
*striata*, DC., 23, 153.  
*Vachellii*, Hook. & Arn., 23, 150.  
*verrucosa*, Linn., 23, 153.  
*yunnanensis*, Franch., 36, 471.
- Croton aromaticus*, Linn., 26, 434.  
*aromaticus*, Lour., 26, 434.  
*chinensis*, Benth., 26, 434.  
*chinensis*, Geisel., 26, 439.  
*congestus*, Lour., 26, 434.  
*crassifolius*, Geisel., 26, 434.  
*Cumingii*, Muell. Arg., 26, 434.  
*Hancei*, Benth., 26, 434.  
*Kroneanus*, Miq., 26, 435.  
*lachnocarpus*, Benth., 26, 435.  
*laevigatus*, Vahl, 26, 435.  
*longifolius*, Seem., 26, 434.  
*mollissimus*, Geisel., 26, 442.  
*polystachyus*, Hook. & Arn., 26, 434.  
*sebiferus*, Linn., 26, 445.  
*Tiglium*, Linn., 26, 435.  
*tomentosus*, Muell. Arg., 26, 434.  
*tuberculatus*, Bunge, 26, 436.
- Crozophora mollissima*, Spreng., 26, 442.
- Crucianella stylosa*, Linn., 23, 395.
- CRUCIFERÆ, 23, 39.
- Crypsis aculeata*, Ait., 36, 384.
- Cryptocarya chinensis*, Hemsl.\*, 26, 370.  
*concinna*, Hance, 26, 370.
- Cryptolepis Edithæ*, Maxim., 26, 100.  
*elegans*, Wall., 26, 100.  
*laxiflora*, Blume, 26, 100.  
*pauciflora*, Wight, 26, 100.
- Cryptomeria Fortunei*, Otto & Dietr., 26, 544.  
*japonica*, D. Don, 26, 544.
- Cryptonema malaccensis*, Turcz., 36, 4.
- Cryptotænia canadensis*, DC., 23, 329.  
*japonica*, Hassk., 23, 329.
- Cryptotæniopsis*, Dunn, 36, 471.  
*asplenioides*, Boissieu, 36, 471.  
*botrychioides*, Dunn, 36, 471.



- Cryptotæniopsis cardiocarpa*, *Dunn*, 36, 471.  
*Delavayi*, *Dunn*, 36, 471.  
*flicina*, *Boissieu*, 36, 471.  
*leptophylla*, *Dunn*, 36, 471.  
*mollis*, *Dunn*, 36, 471.  
*Tanakæ*, *Boissieu*, 36, 471.  
*vulgaris*, *Dunn*, 36, 471.  
*Cubospermum palustre*, *Lour.*, 23, 309.  
*Cucubalus baccifer*, *Linn.*, 36, 471.  
*Cucumis cantonians*, *Debeaux*, 23, 317.  
*maderaspatanus*, *Linn.*, 23, 319.  
*Melo*, *Linn.*, 23, 317.  
*sativus*, *Linn.*, 23, 317.  
*trigonus*, *Roxb.*, 23, 317.  
*Cucurbita maxima*, *Duch.*, 23, 317.  
*moschata*, *Duch.*, 23, 318.  
*Pepo*, *Linn.*, 23, 317.  
*sinensis*, *Til.*, 23, 313.  
 CUCURBITACEÆ, 23, 312.  
*Cudrania crenata*, *C. H. Wright*\*, 26, 469.  
*javanensis*, *Trécul*, 26, 469.  
*obovata*, *Trécul*, 26, 469.  
*rectispina*, *Hance*, 26, 470.  
*tricuspidata*, *Bur.*, 26, 470.  
*triloba*, *Hance*, 26, 470.  
*Cudranus*, *Rumph.*, 26, 469.  
*Cunninghamia lanceolata*, *Lamb.*, 26, 548.  
*sinensis*, *R. Br.*, 26, 548.  
*Cypia corymbosa*, *Hook. & Arn.*, 23, 380.  
*mollissima*, *Hook. & Arn.*, 23, 381.  
*Cupressus amœna*, *C. Koch*, 26, 541.  
*funeris*, *Endl.*, 26, 540.  
*horizontalis*, *Mill.*, 26, 541.  
*japonica*, *Linn. f.*, 26, 544.  
*patula*, *Spad.*, 26, 541.  
*pendula*, *Abel*, 26, 541.  
*sempervirens*, *Linn.*, 26, 541.  
*Tourneforti*, *Audib.*, 26, 541.  
 CUPULIFERÆ, 26, 496.  
*Curanga amara*, *Juss.*, 26, 187.  
*Curculigo ensifolia*, *R. Br.*, 36, 87.  
*graminifolia*, *Nimmo*, 36, 86.  
*orchioides*, *Gaertn.*, 36, 87.  
 — var. *minor*, *Benth.*, 36, 87.  
*recurvata*, *Dryand.*, 36, 87.  
*stans*, *Labill.*, 36, 87.  
*Curcuma longa*, *Linn.*, 36, 69.  
*pallida*, *Lour.*, 36, 70.  
*rotunda*, *Linn.*, 36, 69.  
*Cuscuta chinensis*, *Lam.*, 26, 167.  
*colorans*, *Maxim.*, 26, 168.  
*fimbriata*, *Bunge*, 26, 168.  
*Grasii*, *Delpon*, 26, 168.  
*hyalina*, *Wight*, 26, 168.  
*japonica*, *Choisy*, 26, 168.  
*Cuscuta lupuliformis*, *Krock.*, 26, 168.  
*Millettii*, *Hook. & Arn.*, 26, 168.  
*monogyna*, *Vahl*, 26, 168.  
*obtusiflora*, *H. B. K.*, 26, 168.  
*pedicellata*, *Ledeb.*, 26, 169.  
*Cyananthus barbatus*, *Franch.*, 26, 6.  
*Delavayi*, *Franch.*, 26, 6.  
*Hookeri*, *C. B. Clarke*, 26, 6.  
 — var. *hispidus*, *Franch.*, 26, 6.  
 — var. *levicaulis*, *Franch.*, 26, 6.  
*incanus*, *Hook. f. & Thoms.*, 26, 6.  
 — var. *leiocalyx*, *Franch.*, 26, 6.  
*inflatus*, *Hook. f. & Thoms.*, 26, 7.  
 — var. *rufus*, *Franch.*, 26, 7.  
 — var. *tenuis*, *Franch.*, 26, 7.  
*longiflorus*, *Franch.*, 26, 7.  
*petiolatus*, *Franch.*, 36, 471.  
*Cyanitis sylvatica*, *Reinw.*, 23, 275.  
*Cyanopsis pubescens*, *DC.*, 23, 401.  
*villosa*, *DC.*, 23, 401.  
*Cyanosis arachnoidea*, *C. B. Clarke*, 36, 157.  
*axillaris*, *Roem. & Schult.*, 36, 157.  
*barbata*, *D. Don*, 36, 157.  
*eristata*, *Roem. & Schult.*, 36, 158.  
 — var. *Griffithii*, *C. B. Clarke*, 36, 158.  
*fasciculata*, *Benth.*, 36, 157.  
*nepalensis*, *Miers*, 36, 4.  
*nobilis*, *Hassk.*, 36, 157.  
*pilosa*, *Wight*, 36, 157.  
*vaga*, *C. B. Clarke*, 36, 160.  
*vaga*, *Roem. & Schult.*, 36, 160.  
*Cyanthillium pubescens*, *Blume*, 23, 401.  
*villosum*, *Blume*, 23, 401.  
*Cyathocline lyrata*, *Cass.*, 23, 407.  
*Cyathula prostrata*, *Blume*, 26, 321.  
 CYCADACEÆ, 26, 559.  
*Cycas Bellefonti*, *Lind. & Rod.*, 26, 560.  
*Boddami*, *Hort.*, 26, 560.  
*circinalis*, *Linn.*, mentioned, 26, 560.  
*inermis*, *Lour.*, 26, 559.  
*pectinata*, *Griff.*, mentioned, 26, 560.  
*revoluta*, *Thunb.*, 26, 559.  
*Rumphii*, *Miq.*, 26, 559, 560.  
 — var. *bifida*, *Dyer*\*, 26, 560.  
*siamensis*, *Miq.*, 26, 560.  
*taiwaniana*, *Carruth.*, 26, 560.  
*tonkinensis*, *Dyer*, 26, 560.  
*Cyclea deltoidea*, *Miers*, 23, 29.  
*polypetala*, *Dunn*, 36, 472.  
*racemosa*, *Oliver*, 36, 472.  
*Cyclodon adnatus*, *Griff.*, 26, 7.  
*distans*, *Griff.*, 26, 7.

- Cyclodon truncatus*, Hook. f. & Thoms., 26, 7, 8.
- Cydonia cathayensis*, Hemsl., 36, 472.
- chinensis*, DC., 23, 256.
- japonica*, Pers., 23, 257.
- sinensis*, Thouin, 23, 256.
- Cymbaria daurica*, Linn., 26, 203.
- Cymbidium aloifolium*, Lodd., 36, 31.
- aloifolium*, Sw., 36, 31.
- Corallorhiza*, Sw., 36, 9.
- cyperifolium*, Wall., mentioned, 36, 30.
- eburneum*, Lindl., not Chinese, 36, 29.
- ensifolium*, Hook., 36, 31.
- ensifolium*, Sw., 36, 29; mentioned, 31.
- erythræum*, Lindl., 36, 31.
- Faberi*, Rolfe, 36, 30.
- floribundum*, Lindl., 36, 30.
- fragrans*, Salisb., 36, 31.
- giganteum*, Lindl., 36, 30.
- giganteum*, Wall., 36, 472.
- grandiflorum*, Griff., 36, 30.
- Hookerianum*, Reichb., 36, 30.
- hyacinthinum*, Sm., 36, 19.
- Leachianum*, Reichb. f., 36, 30.
- longifolium*, D. Don, 36, 31.
- Meyenii*, Schau., 36, 27.
- micans*, Schau., 36, 29.
- pendulum*, Sw., 36, 31; mentioned, 32.
- sinense*, Willd., 36, 31; mentioned, 30.
- sp., 36, 32.
- sp., Griff., 36, 30.
- striatum*, Sw., 36, 19.
- triste*, Lindl., 36, 33.
- xiphiifolium*, Lindl., 36, 29.
- Cymbopogon Nardus*, Rendle, 36, 376.
- *subsp. hamatulus*, Hack., 36, 376.
- *subsp. marginatus*, var. *Goeringii*, Hack., 36, 376.
- Schoenanthus*, Spreng., 36, 377.
- *var. cæsius*, Hack., 36, 377.
- Cyminosma pedunculata*, DC., 23, 108.
- resinosa*, DC., 23, 108.
- Cymopterus littoralis*, A. Gray, 23, 331.
- Cynanchum*, a species of *Polygonum* resembling, 26, 338.
- acuminatifolium*, Hemsl. \*, 26, 104.
- affine*, Hemsl. \*, 26, 104.
- amplexicaule*, Hemsl. \*, 26, 104.
- atratum*, Bunge, 26, 104.
- auriculatum*, Royle, 26, 105; mentioned, 109.
- Bungei*, Decne., 26, 105; mentioned, 110.
- caudatum*, Maxim., 26, 105.
- caudatum*, Hemsl. \*, 26, 110; sphalm. vide *Endotropis caudata*, Miq.
- chinense*, R. Br., 26, 105.
- deltoideum*, Hance, 26, 105.
- Fordii*, Hemsl. \*, 26, 106.
- formosanum*, Hemsl. \*, 26, 106.
- grandifolium*, Hemsl. \*, 26, 106.
- Henryi*, Warb., 36, 472.
- inodorum*, Lour., 26, 107.
- insulanum*, Hemsl. \*, 26, 107.
- japonicum*, Hemsl. \*, 26, 107.
- linearifolium*, Hemsl. \*, 26, 107.
- mandshuricum*, Hemsl. \*, 26, 107.
- mongolicum*, Hemsl. \*, 26, 107.
- Mooreanum*, Hemsl. \*, 26, 108.
- nipponicum*, Matsum., 36, 472.
- odoratissimum*, Lour., 26, 114.
- ovalifolium*, Wight, mentioned, 26, 106.
- pubescens*, Bunge, 26, 105.
- roseum*, R. Br., 26, 108.
- sibiricum*, R. Br., 26, 108.
- stenophyllum*, Hemsl. \*, 26, 108.
- versicolor*, Bunge, 26, 109.
- verticillatum*, Hemsl. \*, 26, 109.
- villosum*, Matsum., 36, 472.
- viridiflorum*, Sims, 26, 113.
- volubile*, Hemsl. \*, 26, 109; mentioned, 104.
- Wilfordi*, Franch. & Sav., 26, 109; mentioned, 110.
- Cynoctoum formosanum*, Maxim., 26, 106.
- insulanum*, Hance, 26, 107.
- ovalifolium*, Wight, mentioned, 26, 106.
- roseum*, Decne., 26, 108.
- Wilfordi*, Maxim., 26, 109.
- Cynodon Dactylon*, Pers., 36, 402.
- Cynoglossum divaricatum*, Steph., 26, 149.
- furcatum*, Wall., 26, 149.
- javanicum*, Thunb., 26, 149.
- micranthum*, Desf., 26, 149, 150.
- nervosum*, Benth., 26, 150.
- sp. n. ?, 26, 150.
- Cynosorchis gracilis*, Blume, 36, 472.
- Cynosurus ægyptius*, Linn., 36, 406.
- coracanus*, Linn., 36, 405.
- indicus*, Linn., 36, 406.
- CYPERACEÆ, 36, 202.
- Cyperorchis elegans*, Blume, 36, 472.
- Cyperus albidus*, Retz., 36, 208.
- alopecuroides*, Roxb., 36, 212.

- Cyperus amoyensis*, Hance, **36**, 215, 216.  
*amuricus*, Maxim., **36**, 208; mentioned, 209.  
*angulatus*, Nees, **36**, 202.  
*arenarius*, Hance, **36**, 217.  
*aristatus*, Rottb., **36**, 209.  
— var. *Maingayi*, C. B. Clarke, **36**, 208.  
*atratus*, Steud., **36**, 206.  
*auricomus*, Benth., **36**, 211.  
*auriculatus*, Nees, **36**, 222.  
*Baccha*, Kunth, **36**, 203.  
*bracteatus*, Steud., **36**, 219.  
*bromoides*, Vahl, **36**, 220.  
*brunneus*, Hook. & Arn., **36**, 206.  
*bulboso-stoloniferus*, Steud., **36**, 217.  
*canescens*, Vahl, **36**, 220.  
*capillaris*, Roxb., **36**, 203.  
*castaneus*, Miq., **36**, 219.  
*castaneus*, Willd., **36**, 209; mentioned, 219.  
*cephalotes*, Vahl, **36**, 209.  
*compactus*, Retz., **36**, 221.  
*complanatus*, Presl, **36**, 204.  
*compressus*, Linn., **36**, 210.  
*conjunctus*, Steud., **36**, 217.  
*cuspidatus*, H. B. K., **36**, 219.  
*cylindrostachys*, Boeck., **36**, 221.  
*cylindrostachyus*, Boeck., **36**, 221.  
*densiflorus*, G. F. W. Mey., **36**, 222.  
*dichroostachyus*, Hochst., **36**, 210.  
*difformis*, Linn., **36**, 210.  
*diffusus*, Vahl, **36**, 211.  
*digitatus*, Roxb., **36**, 211; mentioned, 207.  
*dilutus*, Vahl, **36**, 221.  
*diluvialis*, Schultes, **36**, 207.  
*distans*, Linn., **36**, 211.  
*dulcis*, Rumph., **36**, 229.  
*elegans*, Sw., **36**, 211.  
*eleusinoides*, Kunth, **36**, 212.  
*erectus*, Roxb., **36**, 205.  
*eriophorus*, Steud., **36**, 212.  
*exaltatus*, Retz., **36**, 212.  
*ferax*, Rich., **36**, 222, 223.  
*ferox*, Vahl, **36**, 222, 223.  
*fimbriatus*, Franch. & Sav., **36**, 218.  
*flavescens*, var. *rubromarginata*, Schrenk, **36**, 206.  
*flexuosus*, Vahl, **36**, 222.  
*Fortunei*, Steud., **36**, 214.  
*fuscus*, Linn., **36**, 212.  
*fuscus*, Maxim., **36**, 210.  
— var. *chinensis*, Debeaux, **36**, 218.  
*giganteus*, Boeck., **36**, 221.  
*globosus*, Allioni, **36**, 203.
- Cyperus glomeratus*, Linn., **36**, 213.  
*Griffithii*, Steud., **36**, 216.  
*gymnoleptus*, Steud., **36**, 206.  
*Haspan*, Linn., **36**, 213.  
*hexastachys*, Rottb., **36**, 217.  
*hexastachyus*, Nees, **36**, 219.  
*Heynei*, Boeck., **36**, 215.  
*infra-apicalis*, Nees, **36**, 212.  
*inundatus*, Nees, **36**, 203.  
*inundatus*, Roxb., **36**, 207.  
*Iria*, Linn., partim, **36**, 207.  
*Iria*, Linn., **36**, 213.  
— forma *japonica*, Boeck., **36**, 201.  
— var. (*amabilis*), C. B. Clarke, **36**, 208.  
— var. *paniciformis*, C. B. Clarke, **36**, 214.  
*Iria*, Thwaites, **36**, 214.  
*Jacquini*, Schrad., **36**, 215.  
*japonicus*, Miq., **36**, 208.  
*Kegelianus*, Steud., **36**, 259.  
*lævissimus*, Steud., **36**, 217.  
*latespicatus*, Boeck., **36**, 205.  
*leucocephalus*, Hassk., **36**, 224.  
*Linnæanus*, Kunth, **36**, 205.  
*littoralis*, R. Br., **36**, 217.  
*lomentaceus*, Nees, **36**, 222.  
*lucidulus*, C. B. Clarke, **36**, 220.  
*macropus*, Miq., **36**, 216.  
*macrosciadion*, Steud., **36**, 216.  
*malaccensis*, Lam., **36**, 214; mentioned, 223.  
— var. *brevifolia*, Boeck., **36**, 218.  
*marginellus*, Nees, **36**, 215.  
*Meyenii*, Nees, **36**, 210.  
*mæstus*, Kunth, **36**, 211.  
*monogynus*, Boeck., **36**, 209.  
*monostachyus*, Linn., **36**, 240.  
*Monti*, Linn., **36**, 208.  
*multiceps*, Hook., **36**, 222.  
*Neesii*, Kunth, **36**, 211.  
*nigro-viridis*, Thwaites, **36**, 211.  
*nitens*, Retz., **36**, 206.  
*niveus*, Retz., **36**, 215.  
*nutans*, Vahl, **36**, 215.  
*odoratus*, Linn., **36**, 205, 214, 222.  
*odoratus*, Osbeck, **36**, 217.  
*orthostachyus*, Franch. & Sav., **36**, 218.  
*ovularis*, Boeck., **36**, 221.  
*paniciformis*, Franch., **36**, 214.  
*parviflorus*, Nees, **36**, 214.  
*pauciflorus*, Steud., **36**, 215.  
*pectiniformis*, Nees, **36**, 210.  
*pennatus*, Lam., **36**, 220.  
*Pes-avium*, Bertol., **36**, 213.  
*pilosus*, Vahl, **36**, 215; mentioned, 203.

- Cyperus piptolepis*, Steud., 36, 215.  
*polystachyus*, R. Br., 36, 205.  
*Prescottianus*, Hook., 36, 222.  
*procerus*, Rottb., 36, 215.  
*pulvinatus*, Nees, 36, 206.  
*pumilus*, Linn., 36, 206.  
*puncticulatus*, Trimen, 36, 215.  
*puncticulatus*, Vahl, 36, 203, 216.  
*pygmæus*, Rottb., 36, 207, 208.  
— var.  $\beta$ , Boeck., 36, 252.  
*racemosus*, Boeck., 36, 212.  
*radians*, Nees, 36, 216.  
*radicans*, Kunth, 36, 216.  
*radiatus*, Vahl, 36, 216.  
*Retzii*, Nees, 36, 219.  
*Retzii*, Poir., 36, 208.  
*rotundus*, Benth., 36, 219.  
*rotundus*, Kunth, 36, 217.  
*rotundus*, Linn., 36, 216.  
*sanguinolentus*, Vahl, 36, 206.  
*Santonici*, Rottb., 36, 214.  
*septatus*, Steud., 36, 221.  
*serotinus*, Rottb., 36, 208.  
*Sieberianus*, Diels, 36, 221.  
*sinensis*, Debeaux, 36, 216.  
*solutus*, Steud., 36, 219.  
*sp.*, Rottb., 36, 207.  
*spaniophyllus*, Steud., 36, 214.  
*squarrosus*, Linn., 36, 209.  
*stolonifer*, Retz., 36, 217.  
*strictus*, Lam., 36, 205.  
*strigosus*, Hook., 36, 222.  
*stuppeus*, Forst., 36, 220.  
*subalatus*, Steud., 36, 215.  
*subfuscus*, Debeaux, 36, 218.  
*tegetiformis*, Roxb., 36, 218.  
*tenuiculmis*, Boeck., 36, 219.  
*tenuiflorus*, Roxb., 36, 219.  
*teretifructus*, Steud., 36, 206.  
*Teysmanni*, Boeck., 36, 207.  
*tortuosus*, Roxb., 36, 205.  
*truncatus*, Turcz., 36, 218.  
— var. *orthostachya*, C. B. Clarke\*, 36, 218.  
*tuberosus*, Bojer, 36, 217.  
*tuberosus*, Rottb., 36, 219.  
*umbellatus*, Benth., 36, 220.  
*umbellatus*, Miq., 36, 221.  
*uncinatus*, Poir., 36, 219.  
*unioloides*, R. Br., 36, 202.  
*venustus*, Nees, 36, 211.  
*venustus*, R. Br., 36, 212.  
*verticillatus*, Nees, 36, 211.  
*verticillatus*, Roxb., 36, 216.  
*vulgaris*, Kunth, 36, 204.  
*Wightii*, Hance, 36, 220.  
*Zollingeri*, Steud., 36, 219.  
*Cypripedium arietinum*, R. Br., 36, 63, 472.  
*Calceolus*, Linn., 36, 64.  
*Cypripedium chinense*, Franch., 36, 65.  
*corrugatum*, Franch., 36, 64.  
— var. *obesa*, Franch., 36, 64.  
*ebracteatum*, Rolfe, 36, 64.  
*elegans*, Reichb. f., 36, 472.  
*Fargesii*, Franch., 36, 64.  
*fasciolatum*, Franch., 36, 64.  
*guttatum*, Sw., 36, 64.  
*Henryi*, Rolfe, 36, 65.  
*himalaicum*, Rolfe, 36, 65.  
*japonicum*, Thunb., 36, 65; mentioned, 64.  
*luteum*, Franch., 36, 65.  
*macranthum*, Hook., 36, 65, 66.  
*macranthum*, Sw., 36, 66.  
*margaritaceum*, Franch., 36, 66; mentioned, 64.  
*micranthum*, Franch., 36, 66; mentioned, 64.  
*Parishii*, Reichb. f., 36, 67.  
*plectrochilon*, Franch., 36, 63.  
*purpuratum*, Lindl., 36, 67.  
*sinicum*, Hance, 36, 67.  
*tibeticum*, King, 36, 66.  
*venustum*, Wall., mentioned, 36, 67.  
*yunnanense*, Franch., 36, 66.  
*Cyrtia agrestis*, Miers, 26, 77.  
*japonica*, Miers, 26, 76.  
*serrulata*, Miers, 26, 77.  
*suberifolia*, Miers, 26, 77.  
*virgata*, Miers, 26, 77.  
*Cyrtandra Fortunei*, C. B. Clarke, 26, 224.  
*Cyrtopera bicarinata*, Lindl., mentioned, 36, 28.  
*flava*, Lindl., 36, 28.  
*formosana*, Rolfe, 36, 28.  
*Cyrtophyllum fragrans*, DC., 26, 121.  
*peregrinum*, Blume, 26, 121.  
*Cystorchis nebulorum*, Hance, 36, 45.  
*Cytisus Cajan*, Linn., 23, 195.  
*Czernævia lævigata*, Turcz., 23, 334.  
*Dactylæa schizopetalum*, Franch., 36, 472.  
*Dactylis glomerata*, Linn., 36, 421.  
*Dactyloctenium ægyptiacum*, Willd., 36, 406.  
*mucronatum*, Willd., 36, 406.  
*Dædalacanthus nervosus*, T. Anders., 26, 238.  
*Dalbergia Delavayi*, Franch., 36, 472.  
*Dyeriana*, Prain, 36, 472.  
*frondosa*, Roxb., 23, 198.

- Dalbergia Hancei*, Benth., 23, 198.  
*hupeana*, Hance, 23, 198.  
*lanceolaria*, Linn., 23, 198.  
*Millettii*, Benth., 23, 198.  
*mimusoides*, Franch., 36, 472.  
*monosperma*, Dalz., 23, 198.  
*polyphylla*, Benth., 23, 198.  
*rubiginosa*, Roxb., 23, 198.  
*scandens*, Roxb., 23, 199.  
*yunnanensis*, Franch., 36, 472.  
*Damasonium indicum*, Willd., 36, 3.  
*Damnacanthus indicus*, Gaertn., 23, 386.  
*macrophyllus*, Sieb., 23, 387.  
*major*, Sieb. & Zucc., 23, 387.  
*Danthonia neuroelytrum*, Steud., 36, 342.  
*Daphne cannabina*, Lour., 26, 398.  
*cannabina*, Wall., 26, 395.  
*Championi*, Benth., 26, 395.  
*Fortunei*, Benth., 26, 395.  
*Fortunei*, Lindl., 26, 395.  
*gemmata*, E. Pritz., 36, 472.  
*Genkwa*, Sieb. & Zucc., 26, 395.  
*gracilis*, E. Pritz., 36, 472.  
*indica*, [Loisel.] Herb. Gén., 26, 395.  
*indica*, Linn., 26, 398.  
*japonica*, Sieb. & Zucc., 26, 395.  
*odora*, Thunb., 26, 395.  
*retusa*, Hemsl., 26, 401; 36, 472.  
*Roumea*, Meisn., 26, 395.  
*sinensis*, Lam., 26, 395.  
*tangutica*, Maxim., 26, 396.  
*tenuiflora*, Bur. & Franch., 26, 401; 36, 472.  
*triflora*, Lour., 26, 396.  
*Daphnidium bifarium*, Nees, 26, 386.  
*Cubeba*, Nees, 26, 380.  
*elongatum*, Nees, 26, 381.  
*lanceifolium*, Sieb. & Zucc., 26, 383.  
*Myrrah*, Sieb. & Zucc., 26, 392.  
*sinense*, Blume, 26, 392.  
*strychnifolium*, Sieb. & Zucc., 26, 392; 36, 472.  
*Daphniphyllum Calleryanum*, Baill., 26, 429.  
*calycinum*, Benth., 26, 429.  
*Gaudichaudianum*, Baill., 26, 429.  
*glaucescens*, Blume, 26, 429.  
— var. *Oldhami*, Hemsl. \*, 26, 429.  
*macropodium*, Miq., 26, 423.  
*Roxburghii*, Baill., 26, 429.  
*Dasyloma benghalense*, DC., 23, 331.  
*glaucum*, DC., 23, 331.  
*japonicum*, Miq., 23, 331.  
*subpinnatum*, Miq., 23, 331.  
*Datura alba*, Nees, 26, 175.  
*fastuosa*, var. *alba*, C. B. Clarke, 26, 175.  
*ferox*, Linn., 26, 176.  
*Metel*, Linn., 26, 176.  
*Stramonium*, Linn., 26, 176.  
*Daucus Carota*, Linn., 23, 336.  
*Davidia*, Baill., 36, 472.  
*involuta*, Baill., 36, 472.  
*Debregeasia edulis*, Wedd., 26, 492.  
*obovata*, C. H. Wright \*, 26, 492.  
*Decaisnea angulosa*, Lindl., 36, 39.  
*Fargesii*, Franch., 36, 472.  
*insignis*, Hook. f. & Thoms., 36, 472.  
*Decumaria sinensis*, Oliver, 36, 472.  
*Deeringia baccata*, Moq., 26, 318.  
*celosioides*, R. Br., 26, 318.  
*Delavaya*, Franch., 36, 472.  
*toxocarpa*, Franch., 36, 472.  
*yunnanensis*, Franch., 36, 472.  
*Delina sarmentosa*, Linn., 23, 22.  
*Delphinium albocaruleum*, Maxim., 23, 19.  
*anthriscifolium*, Hance, 23, 19.  
*Bouvaloti*, Franch., 36, 472.  
*Callerii*, Franch., 23, 19.  
*campylocentrum*, Maxim., 36, 472.  
*caucasicum*, var., Maxim., 36, 472.  
*cavaleriense*, H. Lévillé & Vaniot, 36, 472.  
*ceratophorum*, Franch., 36, 472.  
*cerefolium*, H. Lévillé & Vaniot, 36, 472.  
*chefoense*, Franch., 36, 472.  
*cheilanthum*, Fisch., 23, 19.  
*chinense*, Lodd., 23, 19.  
*cœlestinum*, Franch., 36, 473.  
*Davidi*, Franch., 36, 473.  
*Delavayi*, Franch., 36, 473.  
*densiflorum*, Duthie, 36, 473.  
*denudatum*, Wall., 36, 473.  
*elatum*, Linn., 36, 473.  
*exiguum*, E. Pritz., 36, 473.  
*Fargesii*, Franch., 36, 473.  
*fatsienense*, E. Pritz., 36, 473.  
*Gilgianum*, Pilg., 36, 473.  
*grandiflorum*, Linn., 23, 19.  
— var. *chinense*, Fisch., 23, 19.  
*hamatum*, Franch., 36, 473.  
*Henryi*, Franch., 36, 473.  
*hirticaule*, Franch., 36, 473.  
*iliense*, Huth, 36, 473.  
*lankongense*, Franch., 36, 473.  
*likiangense*, Franch., 36, 473.  
*longipes*, Franch., 36, 473.  
*Maackianum*, Regel, 36, 473.  
*Maximowiczii*, Franch., 36, 473.  
*mosoynense*, Franch., 36, 473.  
*oliganthum*, Franch., 36, 473.

- Delphinium orthocentrum*, *Franch.*, 36, 473.  
*pachycentrum*, *Hemsl.*, 36, 473.  
*Potanini*, *Huth*, 36, 473.  
*pycnocentrum*, *Franch.*, 36, 473.  
*Robertianum*, *H. Léveillé & Vaniot*, 36, 473.  
*sertifera*, *Franch.*, 36, 473.  
*siwanense*, *Franch.*, 36, 473.  
*Souliei*, *Franch.*, 36, 473.  
*sutchuenense*, *Franch.*, 36, 473.  
*taliense*, *Franch.*, 36, 473.  
*tanguticum*, *Maxim.*, 36, 473.  
*tatsienense*, *Franch.*, 36, 473.  
*tongolense*, *Franch.*, 36, 474.  
*trichophorum*, *Franch.*, 36, 474.  
*yunnanense*, *Franch.*, 36, 474.  
*Democritea scrissoides*, *DC.*, 23, 391.  
*Dendrobium aciculare*, *Lindl.*, mentioned, 36, 11.  
*aduncum*, *Wall.*, 36, 9.  
*aggregatum*, *Roxb.*, 36, 9.  
*atropurpureum*, *Miq.*, 36, 9.  
*aurantiacum*, *Reichb. f.*, 36, 9.  
*auriferum*, *Lindl.*, 36, 34; mentioned, 10.  
*bellatulum*, *Rolfe \**, 36, 10.  
*candidum*, *Wall.*, 36, 13.  
*capillipes*, *Reichb. f.*, 36, 474.  
*catenatum*, *Lindl.*, 36, 12.  
*Ceraia*, *Lindl.*, 36, 10.  
*chrysanthum*, *Wall.*, 36, 10.  
*chryseum*, *Rolfe*, 36, 9.  
*chrysostoxum*, *Lindl.*, 36, 474.  
*clavatum*, *Wall.*, 36, 10.  
*Dalhousieanum*, *Wall.*, 36, 12.  
*Fargesii*, *Finet*, 36, 474.  
*fimbriatum*, *Hook.*, 36, 474.  
*formosum*, *Roxb.*, mentioned, 36, 10.  
*Gibsoni*, *Paxt.*, 36, 11.  
*hainanense*, *Rolfe*, 36, 11.  
*Hancockii*, *Rolfe \**, 36, 11.  
*hercoglossum*, *Reichb. f.*, mentioned, 36, 11.  
*japonicum*, *Lindl.*, 36, 12.  
*Linawianum*, *Reichb. f.*, 36, 12.  
*Loddigesii*, *Lindl.*, 36, 12.  
*longicornu*, *Lindl.*, 36, 474.  
*moniliforme*, *Lindl.*, 36, 12.  
*moniliforme*, *Sw.*, 36, 12.  
*nobile*, *Lindl.*, 36, 12.  
— *var. formosanum*, *Reichb. f.*, 36, 13.  
*odiosum*, *Finet*, 36, 474.  
*Pierardi*, *Roxb.*, 36, 474.  
*pulchellum*, *Lodd.*, 36, 12.  
*pulchellum*, *Roxb.*, 36, 12.  
*velutinum*, *Rolfe*, mentioned, 36, 11.  
*yunnanense*, *Finet*, 36, 13, 474.  
*Dendrobium zonatum*, *Rolfe \**, 36, 13.  
*Dendrocalamus affinis*, *Rendle \**, 36, 447.  
*Brandisii*, *Kurz*, mentioned, 36, 447.  
*flagellifer*, *Munro*, mentioned, 36, 448.  
*latiflorus*, *Munro*, 36, 448.  
*Dendrolobium Cephalotes*, *Benth.*, 23, 171.  
*umbellatum*, *Benth.*, 23, 177.  
*Dendropanax japonicum*, *Seem.*, 23, 342.  
*parviflorum*, *Benth.*, 23, 342.  
*proteum*, *Benth.*, 23, 342.  
*Dentaria dasyloba*, *Turcz.*, 23, 43.  
*repens*, *Franch.*, 23, 44.  
*tenella*, *Regel*, 23, 44.  
*tenuifolia*, *Ledeb.*, 23, 44.  
*Dentella repens*, *Forst.*, 23, 372.  
*Dentidia nankinensis*, *Lour.*, 26, 279.  
*Derris chinensis*, *Benth.*, 23, 199.  
*Fordii*, *Oliver*, 36, 474.  
*laxiflora*, *Benth.*, 23, 199.  
*marginata*, *Benth.*, 36, 474.  
*oblonga*, *Hance*, 23, 199.  
*scandens*, *Benth.*, 23, 199.  
*trifoliata*, *Lour.*, 23, 199.  
*uliginosa*, *Benth.*, 23, 199.  
*Deschampsia caespitosa*, *Beauv.*, 36, 399.  
*Desmanthus virgatus*, *Willd.*, 23, 214.  
*Desmochæta prostrata*, *DC.*, 26, 321.  
*Desmodium acrocarpum*, *Hance*, 23, 177.  
*amethystinum*, *Dunn*, 36, 474.  
*Buergeri*, *Miq.*, 23, 175.  
*callianthum*, *Franch.*, 36, 474.  
*Cephalotes*, *Wall.*, 23, 171.  
*cinerascens*, *Franch.*, 36, 474.  
*dubium*, *Lindl.*, 23, 172.  
*elegans*, *Benth.*, 23, 171.  
*floribundum*, *G. Don*, 23, 172.  
*formosum*, *Vogel*, 23, 172.  
*gangeticum*, *DC.*, 23, 172.  
*Gardneri*, *Benth.*, 23, 172.  
*gracillimum*, *Hemsl.*, 36, 474.  
*Griffithianum*, *Benth.*, 36, 474.  
*gyroides*, *DC.*, 23, 173.  
*hamulatum*, *Franch.*, 36, 474.  
*heterocarpum*, *DC.*, 23, 175.  
*heterophyllum*, *DC.*, 23, 173.  
*hispidum*, *Franch.*, 36, 474.  
*japonicum*, *Miq.*, 23, 175.  
*laburnifolium*, *DC.*, 23, 172.  
*latifolium*, *DC.*, 23, 172.  
*laxiflorum*, *DC.*, 23, 172.  
*lutescens*, *DC.*, 23, 173.

- Desmodium microphyllum*, DC., 23, 174.  
*multiflorum*, DC., 23, 172.  
*nervosum*, Vogel, 23, 175.  
*oblatum*, Baker, 23, 174.  
 Oldhami, Oliver, 23, 174.  
*parvifolium*, DC., 23, 174.  
*podocarpum*, DC., 23, 174.  
*podocarpum*, Miq., 23, 172.  
*polycarpum*, DC., 23, 175.  
 — var. *trichocaulon*, Baker, 23, 175.  
*purpureum*, Hook. & Arn., 23, 175.  
*rhabdocladum*, Franch., 36, 474.  
*sinuatum*, Blume, 36, 474.  
 Tashiroi, Matsum., 36, 474.  
*trichocaulon*, DC., 23, 175.  
*yunnanense*, Franch., 36, 474.  
*Desmos chinensis*, Lour., 23, 27.  
*Deutzia albida*, Batalin, 36, 474.  
 Baroniana, Diels, 36, 474.  
*corymbiflora*, Rev. Hort., 36, 474.  
*corymbosa*, R. Br., 23, 276.  
*crenata*, Sieb. & Zucc., 23, 277.  
 — var. *taiwanensis*, Maxim., 23, 277.  
*discolor*, Hemsl. \*, 23, 275.  
*Fargesii*, Franch., 36, 474.  
*Fortunii*, Carr., 23, 277.  
*glaberrima*, Koehne, 36, 474.  
*glabrata*, Komarov, 36, 474.  
*glomerulifolia*, Franch., 36, 474.  
*grandiflora*, Bunge, 23, 276.  
*hamata*, Koehne, 36, 474.  
*longifolia*, Franch., 36, 474.  
*parviflora*, Bunge, 23, 276.  
*scabra*, Thunb., 23, 276.  
*setchuenensis*, Franch., 36, 474.  
*staminea*, R. Br., 23, 277.  
*Deyeuxia collina*, Pilg., 36, 396.  
*effusiflora*, Rendle \*, 36, 392.  
*elatior*, Hook. f., 36, 393.  
*Henryi*, Rendle \*, 36, 393.  
*Hugoniana*, Rendle \*, 36, 393.  
*hupehensis*, Rendle \*, 36, 394; mentioned, 392.  
*japonica*, Spreng., 36, 360.  
*moupinensis*, Pilg., 36, 395.  
*pulchella*, Hook. f., 36, 395.  
*sachalinensis*, Rendle \*, 36, 395.  
*scabrescens*, Munro, 36, 395.  
*sylvatica*, Kunth, 36, 395; mentioned, 393.  
 — var. *borealis*, Rendle \*, 36, 396.  
 — var. *brachytricha*, Hack., 36, 396.  
 — var. *collina*, Rendle \*, 36, 396.  
 — var. *hirsuta*, Hack., 36, 397.  
*Deyeuxia sylvatica*, var. *latifolia*, Rendle \*, 36, 397.  
 — var. *laxiflora*, Rendle \*, 36, 397.  
 — var. *ligulata*, Rendle \*, 36, 398.  
 — var. *sciuroides*, Hack., 36, 398.  
*Diacæcarpum rotundifolium*, Hassk., 23, 344.  
*Dianella ensifolia*, Red., 36, 119.  
*javanica*, Kunth, 36, 119.  
*nemorosa*, Lam., 36, 119.  
*odorata*, Blume, 36, 119.  
*straminea*, Yatabe, 36, 474.  
*Dianthera japonica*, Thunb., 26, 248.  
*Dianthus arbuscula*, Lindl., mentioned, 23, 63.  
*barbatus*, Linn., 23, 63.  
*Caryophyllus*, Linn., 23, 63, 64.  
*chinensis*, Linn., 23, 63.  
 — var. *laciniata*, DC., 23, 63.  
 — var. *sinensis*, Rohrb., 23, 63.  
*Fischeri*, Spreng., 23, 63.  
*japonicus*, Thunb., 23, 63.  
*longicalyx*, Miq., 23, 63.  
*Morrisii*, Hance, 23, 63.  
*Oreadum*, Hance, 23, 64.  
*pulcherrimus*, Loisel., 23, 63.  
*Schraderi*, Reichb., 23, 63.  
*Sequieri*, Chaix, 23, 63.  
*superbus*, Linn., 23, 63.  
*szechuensis*, Williams, 36, 474.  
 DIAPENSIACEÆ, 26, 34.  
*Diaphora cochinchinensis*, Lour., 36, 264.  
*Diarrhena japonica*, Franch., 36, 418.  
*Diarthron linifolium*, Turcz., 26, 401.  
*vesiculosum*, Fisch. & Mey., 26, 402.  
*Dicentra maerantha*, Oliver, 36, 474.  
*spectabilis*, Miq., 23, 35.  
*Dicerma elegans*, DC., 23, 171.  
*pulchellum*, DC., 23, 171.  
*Dichelactina nodicaulis*, Hance, 26, 421.  
*Dichondra repens*, Forst., 26, 167.  
*Dichotomanthes tristaniacarpa*, Kurz, 23, 307.  
*Dichroa cyanitis*, Miq., 23, 275.  
*febrifuga*, Lour., 23, 275.  
*latifolia*, Miq., 23, 275.  
*Dichrocephala Bodinieri*, Vaniot, 36, 474.  
*chrysanthemifolia*, DC., 23, 406.  
 — var. *glabrata*, Hook. f., 23, 406.  
 — var. *Leveillei*, Vaniot, 36, 475.  
 — var. *minutifolia*, Vaniot, 36, 475.  
*Dickinsia*, Franch., 36, 475.  
*hydrocotyloides*, Franch., 36, 475.

- Diclidium ferox*, Schrad., 36, 222.  
*DiCLIPTERA Buergeriana*, Miq., 26, 248.  
*Burmanni*, Nees, 26, 248.  
*cardiocarpa*, Nees, 26, 248.  
*chinensis*, Nees, 26, 247.  
*crinita*, Nees, 26, 248.  
— *var. floribunda*, Hemsl. \*, 26, 248.  
*Roxburghiana*, Nees, 26, 248.  
*Dictytra spectabilis*, DC., 23, 36.  
*Dictamnus Fraxinella* [G. Don], Lindl., 23, 36, 489.  
*Didissandra Delavayi*, Franch., 36, 475.  
*Fargesii*, Franch., 36, 475.  
*glandulosa*, Batalin, 36, 475.  
*lancifolia*, Franch., 36, 475.  
*lanuginosa*, C. B. Clarke, 26, 226.  
*lanuginosa*, Wall., 26, 227.  
*longifolia*, Franch., 36, 475.  
*longipes*, Hemsl., 36, 475.  
*Mihieri*, Franch., 26, 227.  
*Morgani*, Franch., 36, 475.  
*plicata*, Franch., 36, 475.  
*Rosthornii*, Diels, 36, 475.  
*saxatilis*, Hemsl. \*, 26, 227.  
— *var. microcalyx*, Hemsl. \*, 26, 227.  
*sesquifolia*, C. B. Clarke, 26, 228.  
*speciosa*, Hemsl. \*, 26, 228.  
*Didymocarpus aurantiaca*, C. B. Clarke, 26, 231.  
*Auricula*, S. Moore, 26, 226.  
*demissa*, Hance, 26, 229.  
*Fordii*, Hemsl. \*, 26, 229.  
*Hancei*, Hemsl. \*, 26, 229.  
*lanuginosus*, Maxim., 26, 226.  
*Oreocharis*, Hance, 26, 226.  
*rotundifolius*, Hemsl. \*, 26, 230; mentioned, 231.  
*Didymoplexis pallens*, Griff., 36, 47.  
*Dienia congesta*, Lindl., 36, 5.  
*cylindrostachya*, Lindl., 36, 5.  
*Diervilla amabilis*, Carr., 23, 368.  
*floribunda*, Hort., 23, 369.  
*floribunda*, Sieb. & Zucc., 23, 369.  
*florida*, Sieb. & Zucc., 23, 368; mentioned, 369.  
*grandiflora*, Hort., 23, 369.  
*hortensis*, Hort., 23, 369.  
*japonica*, DC., 23, 369.  
*multiflora*, Lem., 23, 369.  
*pauciflora*, Carr., 23, 368.  
*rosea*, Walp., 23, 368.  
*versicolor*, Sieb. et Zucc., 23, 369.  
*Digitalis glutinosa*, Gaertn., 26, 193.  
*sinensis*, Lour., 26, 185.  
*Digitaria barbata*, Willd., 36, 322.  
*Digitaria debilis*, Willd., mentioned, 36, 324.  
*eriantha*, Steud., mentioned, 36, 323.  
*formosana*, Rendle \*, 36, 323.  
*glabra*, Beauv., 36, 324.  
*Henryi*, Rendle \*, 36, 323.  
*humifusa*, Pers., 36, 324.  
*longiflora*, Pers., 36, 324.  
*pertenuis*, Buese, 36, 327.  
*sanguinalis*, Scop., 36, 325.  
— *var. ciliaris*, Doell, 36, 325.  
— *var. cruciata*, Hook. f., 36, 326.  
— *var. extensa*, Hook. f., 36, 326.  
— *var. pruriens*, Hook. f., 36, 326.  
*setifolia*, Stapf, 36, 326.  
*tenuiflora*, Beauv., 36, 324.  
*tenuispica*, Rendle \*, 36, 326.  
*ternata*, Stapf, 36, 327.  
*Dilivaria ilicifolia*, Nees, 26, 242.  
DILLENACEÆ, 23, 22.  
*Dilophia fontana*, Maxim., 23, 49.  
*Dimeria falcata*, Hack., 36, 359.  
*fuscescens*, Munro, 36, 359.  
*pusilla*, Thw., mentioned, 36, 359.  
*sinensis*, Rendle \*, 36, 359.  
*Dimocarpus Longan*, Lour., 23, 139.  
*Dimorphanthus elatus*, Miq., 23, 338.  
*mandshuricus*, Rupr. & Maxim., 23, 338.  
*Dinetus racemosus*, Sweet, 26, 167.  
*Dioscorea acerifolia*, Uline, 36, 475.  
*alata*, Linn., 36, 90.  
*Batatas*, Decne., 36, 91.  
*Buergeri*, Uline, 36, 475.  
*bulbifera*, Linn., 36, 91.  
*dæmona*, *var. reticulata*, Hook. f., 36, 91.  
*Decaisneana*, Carr., 36, 91.  
*doryophora*, Hance, 36, 91; mentioned, 92, 94.  
*Fargesii*, Franch., 36, 475.  
*glabra*, Roxb., 36, 91.  
*Henryi*, Uline, 36, 475.  
*japonica*, Thunb., 36, 92; mentioned, 94.  
*macrocarpa*, Wall., 36, 91.  
*oppositifolia*, Linn., 36, 92.  
*pentaphylla*, Linn., 36, 94, 475.  
*polystachya*, Turcz., 36, 92.  
*quinqueloba*, Thunb., 36, 92.  
*riphogonioides*, Oliver, 36, 93.  
*Rosthornii*, Diels, 36, 475.  
*sagittata*, Royle, 36, 91.  
*sativa*, Linn., 36, 93.  
*sp.*, 36, 94.  
*spinosa*, Roxb., 36, 93.  
*Swinhoei*, Rolfe, 36, 91.  
*tenuipes*, Franch. & Sav., 36, 475.



- Dioscorea tiliæfolia*, Kunth, **36**, 93.  
*tomentosa*, Koen. [mispr. "Linn."],  
 alluded to, **36**, 94.  
*zingiberensis*, C. H. Wright\*, **36**,  
 93.
- DIOSCOREACEÆ, **36**, 90.
- Diospyros armata*, Hemsl.\* , **26**, 69.  
*chinensis*, Blume, **26**, 69.  
*costata*, Carr., **26**, 69.  
*eriantha*, Champ., **26**, 69.  
*japonica*, Sieb. & Zucc., **26**, 70.  
*Kaki*, Linn. f., **26**, 69.  
 — var. *Aurantium*, André, **26**,  
 69.  
 — var. *Bertii*, André, **26**, 69.  
 — var. *costata*, Carr., **26**, 69.  
 — var. *elliptica*, André, **26**, 69.  
 — var. *Sahuti*, André, **26**, 69.  
*Lotus*, Linn., **26**, 70.  
*microcarpa*, Sieb., **26**, 70.  
*Morrisiana*, Hance, **26**, 70.  
*Oldhami*, Maxim., **26**, 70.  
*rhombifolia*, Hemsl.\* , **26**, 70;  
 mentioned, 71.  
*Roxburghii*, Carr., **26**, 69.  
*Schitze*, Bunge, **26**, 69.  
*sinensis*, Hemsl.\* , **26**, 71.  
*utilis*, Hemsl., **36**, 475.  
*vaccinioides*, Lindl., **26**, 71.
- Dipelta elegans*, Batalin, **36**, 475.  
*floribunda*, Maxim., **23**, 357.  
*yunnanensis*, Franch., **36**, 475.
- Diphaca cochinchinensis*, Lour., **23**,  
 170.
- Diphyllia cymosa*, Michx., **36**, 475.
- Diplachne fusca*, Beauv., **36**, 411.  
*serotina*, Link, **36**, 411.  
 — var. *aristata*, Hack., **36**, 411.  
*sinensis*, Hance, **36**, 411.  
*squarrosa*, Richter, **36**, 411.  
 — var. *longe-aristata*, Rendle\*,  
**36**, 411.
- Diplacrum caricinum*, R. Br., **36**,  
 267.
- Diploclinium Evansianum*, Lindl.,  
**23**, 321.
- Diploclisia macrocarpa*, Miers, **23**,  
 28.
- Diplolepis apiculata*, Lindl., **26**, 113.  
*ovata*, Lindl., **36**, 475.
- Diplomeris chinensis*, Rolfe, **36**, 62.
- Diplomorpha canescens*, C. A. Mey.,  
**26**, 397.  
*Chamædaphne*, Meisn., **26**, 397.  
*virgata*, C. A. Mey., **26**, 397.
- Diplopappus asperrimus*, DC., **23**,  
 416.  
*baccharoides*, Benth., **23**, 409  
*chinensis*, Less., **23**, 407.  
*laxus*, Benth., **23**, 416.
- Diploprora Championi*, Hook. f., **36**,  
 34.
- Diplosiphon Oryztorum*, Decne., **36**,  
 2.
- Diplospora fruticosa*, Hemsl.\* , **23**,  
 382.  
*pubescens*, Hook. f., mentioned, **23**,  
 383.  
 sp., **23**, 384.  
*viridiflora*, DC., **23**, 383.
- Dipoma*, Franch., **36**, 475.  
*iberideum*, Franch., **36**, 475.
- DIPSACEÆ, **23**, 399.
- Dipsacus asper*, Wall., **23**, 399.  
*chinensis*, Batalin, **36**, 475.  
*Gmelini*, Maxim., **23**, 400.  
*inermis*, Wall., **36**, 475.  
*japonicus*, Miq., **23**, 400.
- Dipteracanthus calycinus*, Champ.,  
**26**, 239.  
*lanccolatus*, Nees, **26**, 238.  
*subdenticulatus*, Miq., **26**, 249.
- DIPTEROCARPEÆ, **23**, 83.
- Dipteronia*, Oliver, **36**, 475.  
*sinensis*, Oliver, **36**, 475.
- Dischidia chinensis*, Champ., **26**,  
 116.  
*formosana*, Maxim., **26**, 116.
- Disemma Horsfieldii*, Miq., **23**, 312.
- Disporopsis aspersa*, Hua, **36**, 475.  
*fusco-picta*, Hance, **36**, 110.  
*Pernyi*, Hua, **36**, 475.
- Disporum calcaratum*, D. Don, **36**,  
 142.  
*Leschenaultianum*, D. Don, **36**,  
 142.  
*pullum*, Salisb., **36**, 142; men-  
 tioned, 136.  
*sessile*, D. Don, **36**, 143.  
*smilacinum*, A. Gray, **36**, 143.  
 — var. *viridescens*, Maxim., **36**,  
 143.  
*uniflorum*, Baker, **36**, 144.  
*Wallichii*, D. Don, **36**, 142.
- Dissochæta Barthei*, Hance, **23**, 300.
- Dissolana verticillata*, Lour., **26**,  
 266.
- Distylium racemosum*, Sieb. & Zucc.,  
**23**, 289.  
 — var. *chinense*, Franch.\* , **23**,  
 290.
- Dithyrocarpus paniculatus*, Kunth,  
**36**, 158.  
*petiolatus*, Wight, **36**, 158.  
*undulatus*, Wight, **36**, 158.  
*Rothii*, Wight, **36**, 158.
- Diuranthera major*, Hemsl., **36**, 118,  
 475.  
*minor*, C. H. Wright, **36**, 118,  
 475.

- Dobinea Delavayi*, *Baill.*, **36**, 475.  
*Doellingeria scabra*, *Nees*, **23**, 415.  
*trichocarpa*, *DC.*, **23**, 416.  
*Dolichandrone Cauda-felina*, *Benth. & Hook. f.*, **26**, 235.  
*Dolichos albus*, *Lour.*, **23**, 194.  
*biflorus*, *Linn.*, **23**, 194.  
*Catiang*, *Linn.*, **23**, 193.  
*cultratus*, *DC.*, **23**, 195.  
*ensiformis*, *Linn.*, **23**, 192.  
*japonicus*, *Spreng.*, **23**, 162.  
*Lablab*, *Linn.*, **23**, 194.  
*Lagopus*, *Dunn*, **36**, 475.  
*perennans*, [*Hemsl. \**], **23**, 194.  
*phaseoloides*, *Roxb.*, **23**, 191.  
*polystachyos*, *Thunb.*, **23**, 162.  
*purpureus*, *Linn.*, **23**, 194.  
*sinensis*, *Linn.*, **23**, 193.  
*Tetragonolobus*, *Linn.*, **23**, 194.  
*trilobus*, *Linn.*, **23**, 194.  
*unguiculatus*, *Thunb.*, **23**, 195.  
*Dontostemon dentatus*, *Ledeb.*, **23**, 45.  
*eglandulosus*, *C. A. Mey.*, **23**, 45.  
*integrifolius*, *C. A. Mey.*, **36**, 476.  
*micranthus*, *C. A. Mey.*, **23**, 45.  
*oblongifolius*, *Ledeb.*, **23**, 45.  
*Dopatrium junceum*, *Buch.-Ham.*, **26**, 187.  
*Dorcoceras hygrometrica*, *Bunge*, **26**, 234.  
*Doritis Wightii*, *Benth.*, **36**, 34.  
*Doronicum stenoglossum*, *Maxim.*, **23**, 447.  
*Dorstenia chinensis*, *Lour.*, **26**, 456.  
*pubescens*, *G. Forst.*, **26**, 483.  
*Draba alpina*, *Linn.*, **36**, 476.  
*amplexicaulis*, *Franch.*, **36**, 476.  
*bracteata*, *Batalin*, **36**, 476.  
*contorta*, *DC.*, **23**, 41.  
*dentata*, *Ledeb.*, **36**, 476.  
*elata*, *Hook. f. & Thoms.*, mentioned, **23**, 42.  
*incana*, *Linn.*, **23**, 41.  
*moupinensis*, *Franch.*, **36**, 476.  
*nemorosa*, *Linn.*, **23**, 41.  
— *varr. hebecarpa et leiocarpa*, *Ledeb.*, **23**, 42.  
*surculosa*, *Franch.*, **36**, 476.  
*yunnanensis*, *Franch.*, **36**, 476.  
*Dracæna ensata*, *Thunb.*, **36**, 119.  
*ensifolia*, *Linn.*, **36**, 119.  
*ferrea*, *Linn.*, **36**, 118.  
*graminifolia*, *Linn.*, **36**, 79.  
*terminalis*, *Jacq.*, **36**, 118.  
*terminalis*, *Linn.*, **36**, 118.  
*Dracocephalum altaiensis*, *Laxm.*, **26**, 291.  
*Faberii*, *Hemsl. \**, **26**, 291.  
*Dracocephalum fœtidum*, *Bunge*, **26**, 292.  
*grandiflorum*, *Linn.*, **26**, 291.  
*Henryi*, *Hemsl. \**, **26**, 291.  
*imberbe*, *Bunge*, **26**, 292.  
*Moldavica*, *Linn.*, **26**, 292.  
*rupestre*, *Hance*, **26**, 291.  
*Ruyschianum*, *Linn.*, **26**, 292.  
*sinense*, *S. Moore*, **26**, 293.  
*tanguticum*, *Maxim.*, **26**, 293.  
*urticifolium*, *Miq.*, **26**, 293.  
— *var. pedunculatum*, *Hemsl. \**, **26**, 293.  
*Dracontium polyphyllum*, *Thunb.*, **36**, 182.  
*Dracontomelum mangiferum*, *Maxim.*, **23**, 149.  
*sinense*, *Stapf*, **36**, 476.  
*Dregea sinensis*, *Hemsl. \**, **26**, 115.  
*volubilis*, *Benth. & Hook. f.*, **26**, 115.  
*Drosera Burmanni*, *Vahl*, **23**, 288.  
*indica*, *Linn.*, **23**, 289.  
*Loureiri*, *Hook. & Arn.*, **23**, 289.  
*lunata*, *Buch.-Ham.*, **23**, 289.  
*peltata*, *Sm.*, **23**, 289.  
*rotundifolia*, *Lour.*, **23**, 289.  
*umbellata*, *Lour.*, **23**, 289.  
**DROSERACEÆ**, **23**, 288.  
*Dryandra cordata*, *Thunb.*, **26**, 433.  
*oleifera*, *Lam*, **26**, 433.  
*vernicia*, *Correa*, **26**, 433.  
*Drymaria cordata*, *Willd.*, **23**, 71.  
*Dubreuilia peplodes*, *Gaudich.*, **26**, 477.  
*Dubyæa ramosissima*, *Hance*, **23**, 481.  
*Duchesnea fragarioides*, *Sm.*, **23**, 240.  
*sundaica*, *Miq.*, **23**, 240.  
*Duhaldea chinensis*, *DC.*, **23**, 429.  
*Dumartroya fagifolia*, *Gaud.*, **26**, 454.  
*Dumasia cordifolia*, *Hemsl.*, **36**, 476.  
*villosa*, *DC.*, **23**, 489.  
*Dunbaria conspersa*, *Benth.*, **23**, 195.  
*pulchra*, *Benth.*, **36**, 476.  
*subrhombea*, *Hemsl.*, **23**, 195.  
*Dupinia japonica*, *Franch.*, **36**, 476.  
*Dysoda fasciculata*, *Lour.*, **23**, 391.  
*Dysophylla auricularia*, *Blume*, **26**, 276.  
*Benthamiana*, *Hance*, **26**, 277.  
*cruciata*, *Benth.*, **26**, 276.  
*japonica*, *Miq.*, **26**, 277.  
*Sampsoni*, *Hance*, **26**, 276.  
*tetraphylla*, *Wight*, **26**, 276.  
*verticillata*, *Benth.*, **26**, 276.  
*verticillata*, *Benth.*, **26**, 276.

- EBENACEÆ, 26, 69.**  
*Ebermaiera concinnula*, Hance, **26**, 237.  
*Eccoilopus andropogonoides*, Steud., **36**, 351.  
*Ecdysanthera brachiata*, A. DC., **26**, 98.  
     *micrantha*, A. DC., **26**, 98.  
     *pedunculosa*, Miq., **26**, 98.  
     *rosea*, Hook. & Arn., **26**, 98.  
*Echinocarpus sinensis*, Hance, **23**, 94; **36**, 476.  
*Echinocaulon perfoliatum*, Hassk., **26**, 344.  
*Echinochloa Crus-Galli*, Dum., **36**, 329.  
*Echinodorus guianensis*, Griseb., **36**, 190.  
*Echinolytrum dipsaceum*, Desv., **36**, 235.  
*Echinops dahuricus*, Fisch., **23**, 459.  
     *Gmelini*, Ledeb., **26**, 459.  
     *Grijsii*, Hance, **23**, 459.  
     *sphærocephala*, Miq., **23**, 459.  
*Echinopsilon divaricatum*, Kar. & Kir., **26**, 327.  
*Echinospermum anisacanthum*, Turcz., **26**, 150.  
     *anisacanthum*, Turcz., **26**, 150.  
     *javanicum*, Lehm., **26**, 149.  
     *Lappula*, Debeaux, **26**, 150.  
     *Lappula*, Linn., mentioned, **26**, 150.  
*Echites saligna*, Delile, **26**, 100.  
     *sinensis*, Hort. Angl., **26**, 115.  
*Eclipta alba*, Hassk., **23**, 433.  
     *erecta*, Linn., **23**, 433.  
     *marginata*, Boiss., **23**, 433.  
     *prostrata*, Linn., **23**, 433.  
     *thermalis*, Bunge, **23**, 433.  
*Edgeworthia chrysantha*, Lindl., **26**, 401.  
     *chrysantha*, Lindl., **26**, 396; cf. 401.  
     *Gardneri*, Meisn., **26**, 396; mentioned, 401.  
     *papyrifera*, Sieb. & Zucc., **26**, 396.  
*Ehretia acuminata*, R. Br., **26**, 143.  
     *buxifolia*, Roxb., **26**, 144.  
     *corylifolia*, C. H. Wright, **36**, 476.  
     *Dicksoni*, Hance, **26**, 145.  
     *formosana*, Hemsl. \*, **26**, 144.  
     *Hanceana*, Hemsl. \*, **26**, 145.  
     *heterophylla*, Spreng., **26**, 144.  
     *longiflora*, Champ., **26**, 145; mentioned, 149.  
     *macrophylla*, Wall., **26**, 145.  
     *ovalifolia*, Hassk., **26**, 144, 145.  
     *pyrifolia*, D. Don, **26**, 143.  
     *resinosa*, Hance, **26**, 146.  
     *serrata*, Roxb., **26**, 143.  
     — var. *obovata*, DC., **26**, 143.
- ELEAGNACEÆ, 26, 402.**  
*Elæagnus angustifolia*, Linn., **26**, 402.  
     *Bockii*, Diels, **36**, 476.  
     *crispa*, Thunb., **26**, 404.  
     *Davidii*, Franch., **36**, 416.  
     *edulis*, Sieb., **26**, 404.  
     *glabra*, Thunb., **26**, 402.  
     *gonyanthes*, Benth., **26**, 403.  
     *Grijsii*, Hance, **26**, 403.  
     *Henryi*, Warb., **36**, 476.  
     *hortensis*, Bieb., **26**, 402.  
     *lanceolata*, Warb., **36**, 476.  
     *latifolia*, Linn., **26**, 403.  
     *longipes*, A. Gray, **26**, 404.  
     *Loureiri*, Champ., **26**, 403.  
     *macrophylla*, Thunb., **26**, 403.  
     *multiflora*, Thunb., **26**, 404.  
     *Oldhami*, Maxim., **26**, 404.  
     *orientalis*, Linn., **26**, 402.  
     *parvifolia*, Wall., **26**, 404.  
     *pungens*, Thunb., **26**, 404.  
     *reflexa*, Decne., **26**, 404.  
     *Simonii*, Carr., **26**, 403, 404.  
     sp., **26**, 405.  
     *tenuiflora*, Benth., **26**, 402.  
     *umbellata*, Thunb., **26**, 404; mentioned, 405.  
*Elæocarpus chinensis*, Hook. f., **23**, 94.  
     *decipiens*, Hemsl. \*, **23**, 94.  
     *hainanensis*, Oliver, **36**, 476.  
     *Henryi*, Hance, **23**, 95.  
     *japonicus*, Sieb. & Zucc., **23**, 95.  
     *lanceæfolius*, Roxb., **23**, 95.  
     *photiniæfolius*, Sieb. & Zucc., **23**, 95.  
     *serrulatus*, Benth., **23**, 94, 95.  
*Elæocharis*, see *Eleocharis*.  
*Elæococca cordata*, Blume, **26**, 433.  
     *cordata*, A. Juss., **26**, 433.  
     *Vernicia*, A. Juss., **26**, 433.  
     *verrucosa*, A. Juss., **26**, 433.  
*Elæodendron Fortunei*, Turcz., **23**, 124.  
     *glaucum*, Pers., **23**, 125.  
     *japonicum*, Franch. & Sav., **36**, 476.  
**ELATINEÆ, 23, 72.**  
*Elatostema*, Forst., genus mentioned, **26**, 456.  
     *ficoides*, Wedd., **26**, 482.  
     *involveratum*, Franch. & Sav., **26**, 482.  
     *japonicum*, Wedd., **26**, 483.  
     *lævigatum*, Hassk., **26**, 484.  
     *lineolatum*, var. *major*, Thw., **26**, 482.  
     *Mariannæ*, C. B. Clarke, **26**, 482.  
     *oppositifolium*, Dalz., **26**, 480.

- Elatostema ovatum*, Wight, **26**, 480.  
*platyphyllum*, Wedd., **26**, 482.  
*pubescens*, Pers., **26**, 483.  
*sessile*, J. & G. Forst., **26**, 483; mentioned, 482.  
*spp.*, **26**, 483, 484.  
*surculosum*, Wight, mentioned, **26**, 483.  
*umbellatum*, Blume, **26**, 483.  
— *var. majus*, Maxim., **26**, 482.  
*Wightianum*, Wedd., **26**, 484.
- Eleocharis acicularis*, R. Br., **36**, 228.  
*acicularis*, Ledeb., **36**, 225.  
*acutangula*, Schult., **36**, 227.  
*afflata*, Steud., **36**, 226.  
*capitata*, R. Br., **36**, 227.  
*Chaetaria*, Hance, **36**, 225.  
*erythrochlamys*, Miq., **36**, 228.  
*fistulosa*, Link, **36**, 227.  
*interstincta*, R. Br., mentioned, **36**, 228.  
*media*, Schult., **36**, 227.  
*mutata*, R. Br., mentioned, **36**, 227.  
*ochrostachys*, Steud., **36**, 226.  
*palustris*, R. Br., **36**, 227; mentioned, 228.  
*pellucida*, Presl, **36**, 226.  
*pileata*, A. Gray, **36**, 227.  
*plantaginea*, R. Br., **36**, 228; mentioned, 229.  
— *var.*, Nees, **36**, 229.  
*setacea*, R. Br., **36**, 227.  
*spiralis*, Hance, **36**, 230.  
*subprolifera*, Steud., **36**, 226.  
*tetraquetra*, Nees, **36**, 228.  
*tuberosa*, Schult., **36**, 229.  
*uniglumis*, Schult., **36**, 227.  
*variegata*, Kunth, **36**, 229.  
— *var. laxiflora*, Ridl., **36**, 229.
- Eleogenus capitatus*, Nees, **36**, 227.
- Elephantopus scaber*, Linn., **23**, 402; mentioned, 400.
- Eleusine ægyptia*, Desf., **36**, 406.  
*coracana*, Gaertn., **36**, 405.  
*indica*, Gaertn., **36**, 406.
- Eleutherococcus Henryi*, Oliver, **23**, 341.  
*leucorrhizus*, Oliver, **23**, 342.  
*senticosus*, Maxim., **23**, 342.  
*sp.*, **23**, 342.
- Elodea chinensis*, Hance, **23**, 74.
- Elsholtzia blanda*, Benth., mentioned, **26**, 278.  
*calycocarpa*, Diels, **36**, 476.  
*cristata*, Willd., **26**, 277.  
*incisa*, Benth., **26**, 277.
- Elsholtzia incisa*, *var. major*, Hook. f., **26**, 277.  
*integrifolia*, Benth., **26**, 277.  
*Oldhami*, Hemsl. \*, **26**, 277.  
*polystachya*, Benth., **26**, 278.  
*rugulosa*, Hemsl. \*, **26**, 278.  
*Stauntoni*, Benth., **26**, 278.
- Elymus arenarius*, Linn., **36**, 433.  
*dahuricus*, Trin., **36**, 433.  
— *var. cylindricus*, Franch., **36**, 433.  
*dasystachys*, Trin., **36**, 433.  
*excelsus*, Turcz., **36**, 434.  
*Pseudo-agropyrum*, Trin., **36**, 432.  
*sibiricus*, Linn., **36**, 434.
- Elyna stenocarpa*, Kar. & Kir., **36**, 268.
- Embelia Fordii*, Mez, **36**, 476.  
*longifolia*, Hemsl. \*, **26**, 62.  
*myrtifolia*, Mez, **36**, 476.  
*nervosa*, A. DC., **26**, 62.  
*oblongifolia*, Hemsl. \*, **26**, 62.  
*obovata*, Hemsl. \*, **26**, 62.  
*pauciflora*, Diels, **36**, 476.  
*polypodioides*, Hemsl. & Mez, **36**, 476.  
*procumbens*, Hemsl., **36**, 476.  
*prunifolia*, Mez, **36**, 476.  
*Ribes*, Burm. f., **26**, 62.  
*saxatilis*, Hemsl., **36**, 476.
- Embllica officinalis*, Gaertn., **26**, 421.
- Embryopteris Kaki*, G. Don, **26**, 69.
- Emericia sinensis*, Roem. & Schult., **26**, 101.
- Emilia angustifolia*, DC., **23**, 449.  
*flammea*, Cass., **23**, 449.  
*prenanthoidea*, DC., mentioned, **23**, 449.  
*sagittata*, DC., **23**, 449.  
*sinica*, Miq., **23**, 449.  
*sonchifolia*, DC., **23**, 449.
- Emmenopterys Henryi*, Oliver, **36**, 476.
- Empusa paradoxa*, Lindl., **36**, 7.
- Endospermum chinensis*, Benth., **26**, 444.
- Endotropis auriculata*, Decne., **26**, 105.  
*auriculata*, Franch., **26**, 109.  
*auriculata*, Miq., **26**, 105.  
*caudata*, Miq., **26**, 105, 110 [as *Cynanchum caudatum*].
- Enemion Raddeanum*, Regel, **23**, 18.
- Engelhardtia acerifolia*, Blume, **26**, 495.  
*chrysolepis*, Hance, **26**, 495.  
*Colebrookiana*, Lindl., **26**, 495.  
*villosa*, Kurz, **26**, 495.  
*Wallichiana*, Lindl., **26**, 495.

- Engelhardtia Wallichiana*, var. *chrysolepis*, C. DC., 26, 495.  
*sp.*, Benth., 26, 495.  
*Enhalus*, Griff., 36, 3.  
*Enhalus acoroides*, Rich., 36, 3.  
*Koenigii*, Rich., 36, 3.  
*Enkianthus biflorus*, Lour., 26, 18.  
*brachyphyllus*, Franch., 36, 476.  
*chinensis*, Franch., 36, 476.  
*himalaicus*, Hook. f. & Thoms., 36, 476.  
*quinqueflorus*, Lour., 26, 18.  
*reticulatus*, Lindl., 26, 18.  
*Rosthornii*, Diels, 36, 476.  
*uniflorus*, Benth., 26, 18.  
*Enkylia digyna*, Griff., 23, 321.  
*trigyna*, Griff., 23, 321.  
*Euneapogon brachystachyus*, Stapf, 36, 407.  
*Entada Alenantha*, DC., 23, 214.  
*Gigalobium*, DC., 23, 214.  
*monostachya*, DC., 23, 214.  
*Pursætha*, DC., 23, 214.  
*scandens*, Benth., 23, 214.  
*Enteropogon gracilior*, Rendle\*, 36, 403.  
*melicoides*, Nees, mentioned, 36, 403.  
*Eomecon chionantha*, Hance, 23, 34.  
*Eopepon aurantiacus*, Naud., 23, 313.  
*vitifolius*, Naud., 23, 313.  
*Epaltes australis*, Less., 23, 423.  
*Ephedra equisetina*, Bunge, 26, 538.  
*Gerardiana*, Wall., 26, 538.  
— var. *sikkimensis*, Stapf, 26, 539.  
*glauca*, Regel, 26, 539.  
*intermedia*, Schrenk & C. A. Mey., 26, 539.  
— var. *glauca*, Stapf, 26, 539.  
*macrocephala*, Bertol., 26, 539.  
*monosperma*, C. A. Mey., 26, 539.  
*monospermos*, S. G. Gmel., 26, 539.  
*monostachya*, var., Willd., 26, 539.  
*procera*, var. *cernua*, Regel, 26, 538.  
— var. *typica*, Regel, 26, 538.  
*vulgaris*, Brandis, 26, 539.  
*Epicarporus orientalis*, Blume, 26, 454.  
*Epidendrum exsifolium*, Linn., 36, 29.  
*monile*, Thunb., 36, 12.  
*moniliforme*, Linn., 36, 12.  
*nervosum*, Thunb., 36, 7.  
*non-chinense*, Reichb. f., mentioned, 36, 28.  
*pendulum*, Roxb., 36, 31.  
*sinense*, Andr., 36, 31.  
*siucense*, Red., 36, 29.  
*Epidendrum striatum*, Thunb., 36, 19.  
*teres*, Thunb., 36, 33.  
*tuberosum*, Lour., 36, 10.  
*Epilobium affine*, Maxim., 23, 307.  
*angulatum*, Komarov, 36, 477.  
*angustifolium*, Linn., 23, 307.  
*calycinum*, Haussk., 36, 477.  
*cephalostigma*, Haussk., 36, 477.  
*decipiens*, Haussk., 23, 308.  
*himalayense*, Haussk., 36, 477.  
*hirsutum*, Linn., 23, 307.  
*japonicum*, Haussk., 23, 307.  
*latum*, Wall., 36, 477.  
*lividum*, Haussk., 36, 477.  
*minutiflorum*, Haussk., 36, 477.  
*molle*, Lam., 36, 477.  
*nepalense*, Haussk., 36, 477.  
*nervosum*, Boiss., 36, 477.  
*nudicarpum*, Komarov, 36, 477.  
*palustre*, Linn., 23, 308.  
*propinquum*, Haussk., 23, 308.  
*roseum*, Schreb., 23, 308.  
*Rouyanum*, H. Léveillé, 36, 477.  
*sp.*, 23, 308.  
*spicatum*, Lam., 23, 309.  
*tetragonum*, Linn., 23, 308.  
*tetragonum*, Lour., 23, 309.  
*Wallichianum*, Haussk., 36, 477.  
*Epimedium acuminatum*, Franch., 36, 477.  
*brevicornu*, Maxim., 36, 477.  
*Davidi*, Franch., 36, 477.  
*Fargesii*, Franch., 36, 477.  
*macranthum*, Morren & Decne., 23, 32.  
*pubescens*, Maxim., 23, 32.  
*sagittatum*, Baker, 23, 33.  
*sinensis*, Sieb., 23, 33.  
*sutchuenense*, Franch., 36, 477.  
*violaceum*, Morren & Decne., 23, 32.  
*Epipactis americana*, Lindl., 36, 49.  
*cantschatea*, Sw., 36, 49.  
*consimilis*, D. Don, 36, 48.  
*gigantea*, Dougl., 36, 49.  
*Helleborine*, var., Crantz, 36, 477.  
*latifolia*, Sw., 36, 48, 49, 477.  
*macrostachya*, Kränzl., 36, 477.  
*Royleana*, Lindl., 36, 49.  
*Thunbergii*, A. Gray, mentioned, 36, 49.  
*veratrifolia*, Boiss., 36, 48.  
*Epiphanes pallens*, Reichb. f., 36, 47.  
*Epipremnum mirabile*, Schott, 36, 185.  
*Epirhizanthus tenella*, Hook. f., 23, 59.  
*Epistylum leptoclados*, Hance, 26, 422.

- Eragrostis amabilis*, *Wight & Arn.*, 36, 412.  
*atrovirens*, *Hack.*, 36, 413.  
*bahiensis*, *Hance*, 36, 413.  
*Brownei*, *Nees*, 36, 413.  
*cylindrica*, *Nees*, 36, 414.  
*elegantula*, *Steud.*, 36, 412.  
*elongata*, *Jacq.*, 36, 413.  
*ferruginea*, *Beauv.*, 36, 413.  
*geniculata*, *Nees*, 36, 414.  
*harpachnoides*, *Hack.*, 36, 414.  
*Hugoniana*, *Rendle*\*, 36, 414.  
*interrupta*, *Beauv.*, partim, 36, 415.  
*interrupta*, *Beauv.*, var. *tenuissima*, *Stapf*, 36, 415.  
*japonica*, *Trin.*, 36, 415.  
*major*, *Host*, 36, 416; mentioned, 422.  
*Makinoi*, *Hack.*, 36, 416.  
*megastachya*, *Link*, 36, 416.  
*Millettii*, *Nees*, 36, 417.  
*minor*, *Host*, 36, 416.  
*Nevinii*, *Hance*, 36, 414.  
*orientalis*, *Nees*, 36, 413.  
*orientalis*, *Trin.*, 36, 413.  
*pilosa*, *Beauv.*, 36, 417.  
*pilosissima*, *Link*, 36, 417.  
*plumosa*, *Link*, 36, 418.  
*poæoides*, *Beauv.*, 36, 416.  
*poæoides*, *Trin.*, 36, 416.  
*stenophylla*, *Hochst.*, 36, 417.  
*tenella*, *Benth.*, 36, 415.  
*tenella*, *Roem. & Schult.*, 36, 418.  
— var. *plumosa*, *Stapf*, 36, 418.  
*tenuissima*, *Schrad.*, 36, 415.  
*unioloides*, *Nees*, 36, 412.  
*verticillata*, *Beauv.*, 36, 417.  
*Willdenoviana*, *Nees*, 36, 413; mentioned, 415.  
*zeylanica*, *Nees*, 36, 413.
- Eranthemum aspersum*, *Hook. f.*, 26, 243.  
*bicolor*, *Schrank*, 26, 243.  
*crenulatum*, *Wall.*, 26, 243.  
*nervosum*, *R. Br.*, 26, 238.  
*palatiferum*, *Nees*, 26, 243.  
*pulchellum*, *Andrews*, 26, 238.
- Eranthis albiflora*, *Franch.*, 36, 477.  
*stellata*, *Maxim.*, 23, 17.  
*Vaniotiana*, *H. Léveillé*, 36, 477.
- Eremochloa falcata*, *Hack.*, 36, 363.  
*leersioides*, *Hack.*, 36, 363.  
*ophiuroides*, *Hack.*, 36, 363.  
*zeylanica*, *Hack.*, 36, 364; mentioned, 363.
- Eria ambrosia*, *Hance*, 36, 14.  
*armeniaca*, *Lindl.*, mentioned, 36, 16.  
*cæspitosa*, *Rolfe*, 36, 16.
- Eria Corneri*, *Reichb. f.*, 36, 16.  
*formosana*, *Rolfe*, 36, 16.  
*hainanensis*, *Rolfe*\*, 36, 16.  
*luchuensis*, *Yatabe*, 36, 17.  
*mysorensis*, *Lindl.*, mentioned, 36, 171.  
*ornata*, *Lindl.*, mentioned, 36, 16.  
*rosea*, *Lindl.*, 36, 17.  
*pannea*, *Lindl.*, 36, 17.  
*pusilla*, *Benth.*, 36, 17.  
*sinica*, *Lindl.*, 36, 17.  
*tomentosa*, *Hook. f.*, mentioned, 36, 16.
- Eriachne chinensis*, *Hance*, 36, 398.  
*Hookeri*, *Munro*, 36, 398.  
*pallescens*, *R. Br.*, 36, 398.
- Erianthus fastigiatus*, *A. Henry*, 36, 350; mentioned, 351.  
*formosanus*, *Stapf*, 36, 350.  
*fulvus*, *Nees*, 36, 350.  
*Hookeri*, *Hack.*, 36, 350.  
*japonicus*, *Beauv.*, 36, 348.  
*pollinioides*, *Rendle*\*, 36, 350.  
*speciosus*, *Debeaux*, 36, 357.  
*tristachyus*, *Nees*, 36, 356.
- ERICACEÆ, 26, 32.
- Ericala Loureiri*, *G. Don*, 26, 129.
- Erigeron acris*, *Linn.*, 23, 417.  
*alpinus*, *Linn.*, 23, 418.  
*ambiguus*, *Sch.-Bip.*, 23, 418.  
*annuus*, *Pers.*, mentioned, 23, 418.  
*canadensis*, *Linn.*, 23, 418.  
*chinense*, *Jacq.*, 23, 418.  
*ciliatus*, *Ledeb.*, 23, 417.  
*hirsutus*, *Lour.*, 23, 418.  
*kamtschaticus*, *DC.*, 23, 418.  
*latisquamatus*, *Maxim.*, 23, 417.  
*linifolius*, *Willd.*, 23, 418.  
*moupinensis*, *Franch.*, 36, 477.  
*pyrifolius*, *Benth.*, 23, 419.
- Eriobotrya dubia*, *Decne.*, 36, 477.  
*fragrans*, *Champ.*, 23, 261.  
*Griffithii*, *Franch.*, 36, 477.  
*japonica*, *Lindl.*, 23, 261.  
*lasiogyna*, *Franch.*, 36, 477.  
*prionophylla*, *Franch.*, 36, 477.
- ERIOCAULACEÆ, 36, 198.
- Eriocaulon*, *Linn.*, mentioned, 23, 289.  
*alpestre*, *Hook. f. & Thoms.*, 36, 198.  
*australe*, *R. Br.*, 36, 198.  
*Buergerianum*, *Koern.*, 36, 198.  
*cantonense*, *Hook. & Arn.*, 36, 201; mentioned, 202.  
*cinereum*, *R. Br.*, 36, 199.  
*cristatum*, *Benth.*, 36, 199.  
*cristatum*, *Mart.*, var. *brevicalyx*, *C. H. Wright*\*, 36, 190.  
*decemflorum*, *Maxim.*, 36, 477.

- Eriocaulon dieline*, Maxim., 36, 477.  
*echinulatum*, Mart., 36, 199.  
*Faberi*, Ruhl., 36, 199.  
*Henryanum*, Ruhl., 36, 199.  
*heteranthum*, Benth., 36, 199.  
*intermedium*, Koern., 36, 200.  
*kiusianum*, Maxim., 36, 477.  
*longifolium*, Koern., 36, 201.  
*luzulæfolium*, Mart., 36, 199.  
*Miquelianum*, Koern., 36, 200.  
*nipponicum*, Maxim., 36, 200.  
*officinale*, Koern., 36, 200.  
*setaceum*, Linn., 36, 200.  
*sexangulare*, Linn., mentioned, 36, 201.  
*sexangulare*, Miq., 36, 201.  
*sexangulare*, Ruhl., 36, 202.  
 — var.  $\beta$ , Koern., 36, 201.  
*Sieboldianum*, Sieb., 36, 200.  
*sinicum*, Miq., 36, 201.  
*truncatum*, Buch.-Ham., 36, 201.  
*Wallichianum*, Mart., 36, 201.  
*Wightianum*, Mart., 36, 202.
- Eriochloa annulata*, Kunth, 36, 320.  
*polystachya*, H. B. K., 36, 320.  
*villosa*, Kunth, 36, 320.
- Eriochrysis Narenga*, Nees, 36, 349.  
*porphyrocoma*, Hance, 36, 349.
- Eriodendron anfractuosum*, DC., 23, 89.
- Eriophorum alpinum*, Linn., in obs., 36, 249.  
*angustifolium*, Roth, 36, 255.  
*comosum*, Nees, 36, 256.  
*cyperinum*, Linn., 36, 249.  
*latifolium*, Hoppe, 36, 256.  
*polystachion*, Linn., 36, 255, 256.  
 — var., 36, 255.
- Eriosema chinense*, Vogel, 23, 197.
- Erismanthus sinensis*, Oliver, 26, 444.
- Eritrichium japonicum*, Miq., 26, 153.  
*pectinatum*, DC., 26, 150.  
*pedunculare*, A. DC., 26, 153.  
*radicans*, A. DC., 36, 477.
- Erodium Stephanianum*, Willd., 23, 98.
- Eruca sativa*, Lam., 23, 47.
- Ervum hirsutum*, Linn., 23, 184.  
*Lens*, Linn., mentioned, 23, 187.  
*tetraspermum*, Linn., 23, 185.
- Erycibe glaucescens*, Hook. & Arn., 26, 156.  
*obtusifolia*, Benth., 26, 156.
- Eryngium fœtidum*, Linn., 36, 477.
- Erysimum aurantiacum*, Maxim., 36, 477.  
*cheiranthoides*, Linn., 23, 46.  
*hieraciifolium*, Linn., 23, 46.
- Erysimum macilentum*, Bunge, 23, 46.  
*stigmatosum*, Franch., 23, 46.  
*yunnanense*, Franch., 36, 477.
- Erythræa*, genus mentioned, 36, 139.  
*japonica*, Maxim., 26, 122.  
*pulchella*, Hance, 26, 122.  
*ramosissima*, Pers., 26, 122.  
 — var. *altaica*, Pers., 26, 122.
- Erythrina Corallodendron*, Linn., 23, 189.  
*indica*, Lam., 23, 189.
- Erythrochæte palmatifida*, Sieb. & Zucc., 23, 453.
- Erythrophloeum Fordii*, Oliver, 23, 214.
- Erythrospermum hypoleucum*, Oliver, 36, 477.
- Erythrostaphyle vitiginea*, Hance, 23, 115.
- Erythroxylen monogynum*, Roxb., 23, 96.
- Eucapnos spectabilis*, Sieb., 23, 36.
- Euchresta Horsfieldii*, Benn., 23, 200.  
*tenuifolia*, Hemsl. \*, 23, 200.  
*tubulosa*, Dunn, 36, 478.
- Euclea herbacea*, Lour., 26, 71.
- Eucommia ulmoides*, Oliver, 36, 478.
- Eugenia acuminatissima*, Kurz, 23, 296.  
*Bulleckii*, Hance, 23, 296.  
*Championi*, Hemsl. \*, 23, 296.  
*fluvialilis*, Hemsl. \*, 23, 296.  
*gracilentata*, Hance, 23, 296.  
*Grijssii*, Hance, 23, 298.  
*Henryi*, Hance, 23, 297.  
*Jambos*, Linn., 23, 297.  
*malaccensis*, Linn., 23, 297.  
*microphylla*, Abel, 23, 298.  
*Milletiana*, Hemsl. \*, 23, 297.  
*minutiflora*, Hance, 23, 297.  
*myrsinifolia*, Hance, 23, 297; mentioned, 296.  
*operculata*, Roxb., 23, 297.  
*pyxophylla*, Hance, 23, 298.  
*sinensis*, Hemsl. \*, 23, 298.  
*tephrodes*, Hance, 23, 298.
- Eulalia cotulifera*, Munro, 36, 351.  
*densa*, Baker, 36, 348.  
*densa*, Munro, 36, 347.  
*japonica*, Trin., 36, 348.
- Eulophia campestris*, Wall., mentioned, 36, 28.  
*Faberi*, Rolfe, 36, 28.  
*flava*, Hook., 36, 28.  
*formosana*, Rolfe \*, 36, 28.  
*nuda*, Lindl., 36, 29.  
*sinensis*, Miq., 36, 29.  
*yunnanensis*, Rolfe \*, 36, 29.
- Euonymus acanthocarpus*, Franch., 36, 478.

- Euonymus actinocarpus*, *Loes.*, 36, 478.  
*aculeatus*, *Hemsl.*, 36, 478.  
*alatus*, *Regel*, 23, 121.  
*amygdalifolius*, *Franch.*, 36, 478.  
*Bockii*, *Loes.*, 36, 478.  
*Bungeanus*, *Maxim.*, 23, 118.  
*carnosus*, *Hemsl.* \*, 23, 118.  
*chinensis*, *Lindl.*, 23, 119; 36, 478.  
*chinensis*, *Lour.*, 23, 119.  
*cornutus*, *Hemsl.*, 36, 478.  
*Dielsianus*, *Loes.*, 36, 478.  
*echinatus*, *Wall.*, 23, 119.  
*europæus*, *Linn.*, 23, 119.  
*fimbriatus*, *Wall.*, 36, 478.  
*flavescens*, *Loes.*, 36, 478.  
*Forbesianus*, *Loes.*, 36, 478.  
*Forbesii*, *Hance*, 23, 118.  
*gibber*, *Hance*, 23, 119.  
*Giraldii*, *Loes.*, 36, 478.  
*gracillimus*, *Hemsl.* \*, 23, 119.  
*grandiflorus*, *Wall.*, 36, 478.  
*hederaceus*, *Champ.*, 23, 120.  
*Hemsleyana*, *Loes.*, 36, 478.  
*hupehensis*, *Loes.*, 36, 478.  
*ilicifolius*, *Franch.*, 36, 478.  
*japonicus*, *Thunb.*, 23, 120.  
*kiantshovicus*, *Loes.*, 36, 478.  
*lanceifolius*, *Loes.*, 36, 478.  
*latifolius*, *A. Gray*, 23, 121.  
— var. *sachalinensis*, *F. Schmidt*, 23, 121.  
*laxiflorus*, *Blume*, 23, 121.  
*laxiflorus*, *Champ.*, 23, 121.  
*linearifolius*, *Franch.*, 36, 478.  
*longifolius*, *Champ.*, 23, 120.  
*lutchuensis*, *T. Ito & Matsum.*, 36, 478.  
*macropterus*, *Rupr.*, 36, 478.  
*micranthus*, *Bunge*, 23, 118.  
*myrianthus*, *Hemsl.*, 36, 478.  
*nanus*, *Bieb.*, 23, 120.  
*nitidus*, *Benth.*, 23, 119.  
*oxyphyllus*, *Miq.*, 23, 121.  
*patens*, *Rehder*, 36, 478.  
*phellomanus*, *Loes.*, 36, 478.  
*Przewalskii*, *Maxim.*, 23, 121.  
*Rosthornii*, *Loes.*, 36, 478.  
*sachalinensis*, *Maxim.*, 23, 121.  
*salicifolius*, *Loes.*, 36, 478.  
*sanguineus*, *Loes.*, 36, 478.  
*schensianus*, *Maxim.*, 23, 121.  
*subtriflorus*, *Blume*, 23, 121.  
*Tashiroi*, *Maxim.*, 23, 121; 36, 478.  
*theifolia*, *Wall.*, 36, 478.  
*Thunbergianus*, *Blume*, 23, 121.  
*tonkinensis*, *Loes.*, 36, 478.  
*venosus*, *Hemsl.*, 36, 478.  
*verrucosoides*, *Loes.*, 36, 478.
- Euonymus verrucosus*, *Scop.*, var. *tchefouensis*, *Debeaux*, 23, 121.  
*yunnanensis*, *Franch.*, 36, 478.
- Eupatorium cannabinum*, *Linn.*, 23, 404; 36, 478.  
*chinense*, *Linn.*, 23, 404.  
— var. *tripartitum*, *Miq.*, mentioned, 23, 405.  
*Fortunei*, *Turcz.*, 23, 403.  
*japonicum*, *Franch.*, 23, 404.  
*japonicum*, *Thunb.*, 23, 403; mentioned, 404.  
*Kirilowii*, *Turcz.*, 23, 404.  
*Lindleyanum*, *DC.*, 23, 404; mentioned, 405.  
*Lindleyanum*, *Hook.*, fide *Hemsl.*, 23, 503.  
*melanadenium*, *Hance*, 23, 405.  
*punduanum*, *Wall.*, 23, 404.  
*Reevesii*, *Wall.*, 23, 405.  
*scandens*, *Linn.*, 23, 405.  
*sp.*, 23, 405.  
*stœchadosmum*, *Hance*, 23, 405.  
*subtetragonum*, *Miq.*, 23, 404.  
*Wallichii*, *DC.*, 23, 403.
- Euphorbia altaica*, *C. A. Mey.*, 26, 411.  
*Atoto*, *Forst.*, 26, 411.  
*bifida*, *Hook. & Arn.*, 26, 412.  
*Chamæsyce*, *Bunge*, 26, 414.  
*coralloides*, *Thunb.*, 26, 416.  
*Cyparissias*, *Turcz.*, 26, 412.  
*Erythræa*, *Hemsl.* \*, 26, 412; mentioned, 414.  
*Esula*, *Linn.*, 26, 412.  
— var. *cyparissoides*, *Boiss.*, 26, 412.  
*Helioscopia*, *Linn.*, 26, 413.  
*Henryi*, *Hemsl.* \*, 26, 413.  
*hippocrepica*, *Hemsl.* \*, 26, 414.  
*humifusa*, *Willd.*, 26, 414.  
*hypercifolia*, *Linn.*, 26, 414.  
*Jolkini*, *Boiss.*, 26, 415.  
*lasiocaula*, *Boiss.*, 26, 416.  
*Lathyris*, *Linn.*, mentioned, 26, 415.  
*lunulata*, *Bunge*, 26, 415.  
*macrostegia*, *Boiss.*, mentioned, 26, 414.  
*nepalensis*, *Boiss.*, 26, 416.  
*Pallasii*, *Turcz.*, 26, 415.  
*palustris*, *A. Gray*, 26, 415.  
*pekinensis*, *Rupr.*, 26, 415.  
*Peplus*, *Seem.*, 26, 413.  
*pilosa*, *Linn.*, 26, 416.  
*pilulifera*, *Linn.*, 26, 416.  
*prolifera*, *Buch.-Ham.*, 26, 416.  
*Pseudochamæsyce*, *Fisch. & Mey.*, 26, 414.  
*ramosissima*, *Boiss.*, 26, 417.



- Euphorbia Sampsoni*, Hance, **26**, 415.  
*sanguinea*, Hochst. & Steud., **26**, 417.  
*serrulata*, Reinw., **26**, 417.  
*Sparmanni*, Boiss., **26**, 417.  
*Stracheyi*, Boiss., **26**, 417.  
*thymifolia*, Linn., **26**, 417.  
*thymifolia*, Thunb., **26**, 414.  
*Tirucalli*, Linn., **26**, 418.  
*Vachellii*, Hook. & Arn., **26**, 417.  
**EUPHORBACEÆ**, **26**, 411.  
*Euphoria Longana*, Lam., **23**, 140.  
*Euphrasia officinalis*, Linn., **26**, 204.  
*Euptelea Davidiana*, Baill., **36**, 478.  
*Delavayi*, Van Tiegh., **36**, 478.  
*Franchetii*, Van Tiegh., **36**, 478.  
*pleiosperma*, Hook. f. & Thoms., **36**, 478.  
*polyandra*, Sieb. & Zucc., **36**, 478.  
*Eurotia ceratoides*, C. A. Mey., **26**, 326.  
*Eurya chinensis*, R. Br., **23**, 76.  
*distichophylla*, Hemsl. \*, **23**, 77.  
*gynandra*, Vesque, **36**, 478.  
*Henryi*, Hemsl., **36**, 479.  
*japonica*, Thunb., **23**, 77.  
*littoralis*, Sieb., **23**, 76.  
*Macartneyi*, Champ., **23**, 77.  
*obliquifolia*, Hemsl., **36**, 479.  
*Euryale ferox*, Salisb., **23**, 33.  
*Euscaphis staphyleoides*, Sieb. & Zucc., **23**, 143.  
*Eustigma oblongifolium*, Gardn. & Champ., **23**, 291.  
*Euthodon paniculata*, Griff., **26**, 97.  
*Eutrema thibeticum*, Franch., **36**, 479.  
*yunnanense*, Franch., **36**, 479.  
*Euxolus oleraceus*, Moq., **26**, 319.  
*viridis*, Moq., **26**, 321.  
*Evansia chinensis*, Salisb., **36**, 82.  
*dichotoma*, Decne., **36**, 81.  
*fimbriata*, Decne., **36**, 82.  
*vespertina*, Decne., **36**, 81.  
*Evodia Daniellii*, Hemsl. \*, **23**, 104; mentioned, 105.  
*glauca*, Miq., **26**, 104.  
*Lamarekiana*, Benth., **23**, 104.  
*meliaefolia*, Benth., **23**, 104.  
*meiicefolia*, Hook. f., fide Hemsl., **23**, 503.  
*rutæcarpa*, Benth., **23**, 104.  
*triphylla*, DC., **23**, 104.  
*Evolvulus alsinioides*, Linn., **26**, 166.  
*chinensis*, Choisy, **26**, 166.  
*hirsutus*, Lam., **26**, 166.  
*linifolius*, Linn., **26**, 166.  
*pubescens*, Hance, **26**, 166.  
*Evolvulus sinicus*, Miq., **26**, 166.  
*Exacum bellum*, Hance, **26**, 122.  
*tetragonum*, Roxb., **26**, 122.  
*Excoecaria Agallocha*, Linn., **26**, 446.  
*discolor*, Muell. Arg., **26**, 445.  
*japonica*, Muell. Arg., **26**, 446.  
*sebifera*, Muell. Arg., **26**, 445.  
*Exochorda grandiflora*, Lindl., **26**, 228.  
*serratifolia*, S. Moore, **26**, 229.  
*Eyrea vernalis*, Champ., **26**, 143.  
*Fabera*, Sch.-Bip., **23**, 479.  
*Faberia*, Hemsl. \*, **23**, 479.  
*sinensis*, Hemsl. \*, **23**, 479.  
*Fagara* = *Zanthoxylon*, **36**, 479.  
*Avicennæ*, DC., **36**, 479.  
*dimorphylla*, Engl., **36**, 479.  
*dissita*, Engl., **36**, 479.  
*echinocarpus*, Engl., **36**, 479.  
*hasiana*, Engl., **36**, 479.  
*micrantha*, Engl., **36**, 479.  
*piperita*, Lour., **23**, 106.  
*stenophylla*, Engl., **36**, 479.  
*triphylla*, Lam., **23**, 104.  
*volubilis*, E. Pritz., **36**, 479.  
*Fagopyrum cymosum*, Meisn., **26**, 337.  
*emarginatum*, Meisn., **26**, 339.  
— var. *kunawarensense*, Meisn., **26**, 337.  
*esculentum*, Mœnch, **26**, 339.  
*tataricum*, Gaertn., **26**, 351.  
*triangulare*, Meisn., **26**, 337.  
*Fagraea fragrans*, Roxb., **36**, 121.  
*Fagus Castanea*, Linn., **26**, 525.  
*Castanea*, Lour., **26**, 523.  
*Engleriana*, Seemen, **36**, 479.  
*longipetiolata*, Seemen, **26**, 525.  
*Sieboldi*, Endl., **36**, 479.  
*sinensis*, Oliver, **26**, 525.  
*sylvatica*, Linn., **26**, 525.  
— var. *chinensis*, Franch., **26**, 525.  
— var. *longipes*, Oliver, **26**, 525.  
*Falcaria dahurica*, DC., **23**, 328.  
*Fallopia nervosa*, Lour., **23**, 92.  
*Farfugium grande*, Lindl., **23**, 454.  
*Kämpferi*, Benth., **23**, 454.  
*Fargesia spathacea*, Franch., **36**, 438.  
*Fatoua japonica*, Blume, **26**, 454.  
*lanceolata*, Gaudich., **26**, 454.  
*pilosa*, Gaudich., **26**, 454.  
*pilosa*, Gaudich., **26**, 454.  
*subcordata*, Gaudich., **26**, 454.  
*Fatsia papyrifera*, Benth. & Hook. f., **23**, 341.  
*Ferula marathrophylla*, Walp., **23**, 335.  
*Festuca filiformis*, Nees, mentioned, **36**, 405.

- Festuca gigantea*, Vill., 36, 429.  
*modesta*, Steud., 36, 429.  
*ovina*, Linn., 36, 429.  
*parvigluma*, Steud., 36, 429.  
*pauciflora*, Thunb., 36, 430.  
*procumbens*, Kunth, 36, 429.  
*remotiflora*, Steud., 36, 430.  
*Fibraurea tinctoria*, Lour., 23, 30.  
**FICOIDEÆ**, 23, 323.  
*Ficus Abelii*, Miq., 26, 457.  
*altissima*, Blume, 26, 457.  
*angustifolia*, Roxb., 26, 464.  
*angustifolia*, Seem., 26, 464.  
*Beecheyana*, Hook. & Arn., 26, 457;  
 mentioned, 459, 467.  
*Benjamina*, Linn., 26, 457.  
*biglandulosa*, Wall., mentioned, 26,  
 458.  
*Carica*, Linn., 26, 458.  
*Carica*, Sieb. & Zucc., 26, 458.  
*caricoides*, Roxb., 26, 458.  
*caudata*, Wall., 26, 459.  
*Championi*, Benth., 26, 468.  
*chlorocarpa*, Benth., 26, 458.  
*comosa*, Roxb., 26, 458.  
*cuspidifera*, Miq., 26, 460.  
*Decaisneana*, Miq., mentioned, 26,  
 461.  
*diadema*, Hemsl., sphalm., 26, 573,  
 = seq.  
*diadema*, Steud., 26, 458.  
*erecta*, auct. plur., 26, 466.  
*erecta*, Thunb., 26, 459; mentioned,  
 459.  
 — var. *Beecheyana*, King, 26, 457.  
 — var. *Sieboldii*, King, 26, 467.  
*fistulosa*, Reinw., 26, 459.  
*formosana*, Maxim., 26, 459.  
*foveolata*, Wall., 26, 460; men-  
 tioned, 463.  
 — var. *Henryi*, King, 26, 460.  
 — var. *impressa*, King, 26, 463.  
*gibbosa*, Blume, 26, 460; mentioned,  
 461.  
*glabella*, Blume, mentioned, 26,  
 465, 469.  
*glaberrima*, Blume, mentioned, 26,  
 464.  
*grossularioides*, Burm. f., 26, 462.  
*hæmatocarpa*, Decne., 26, 458.  
*Hanceana*, Maxim., 26, 465.  
*Harlandi*, Benth., 26, 461.  
*Henryi*, Warb., 36, 479.  
*heteromorpha*, Hemsl., 26, 461.  
*heterophylla*, Linn. f., 26, 461.  
*hibiscifolia*, Champ., 26, 462.  
*hirta*, Sieb. & Zucc., 26, 458.  
*hirta*, Vahl, 26, 462.  
 — var. *Roxburghii*, King, 26, 462.  
*hispida*, Linn. f., 26, 462.  
*Ficus impressa*, Champ., 26, 463;  
 mentioned, 460, 463.  
*infectoria*, Roxb., 26, 463.  
*insularis*, Miq., mentioned, 26, 460,  
 461.  
*ischnopoda*, Miq., 26, 466.  
*japonica*, Blume, 26, 459.  
*Kingiana*, Hemsl., 26, 463.  
*Kurzii*, King, 26, 463.  
*laccifera*, Roxb., 26, 457.  
*leucantatoma*, Poir., 26, 461.  
*Millesii*, Miq., 26, 466.  
*Millettii*, Miq., 26, 466.  
*neglecta*, Decne., 26, 458.  
*nervosa*, Heyne, 26, 464.  
*nipponica*, Franch. & Sav., 26, 460.  
*nitida*, Thunb., 26, 466.  
*obscura*, Blume, 26, 464.  
*obtusifolia*, Roxb., 26, 465.  
*Oldhami*, Hance, 26, 464.  
*oppositifolia*, Willd., 26, 462.  
*palmata*, Forsk., 26, 458.  
*pandurata*, Hance, 26, 465.  
*parasitica*, Koen., 26, 460.  
*parvifolia*, Miq., 26, 465.  
*pertusa*, Linn., f., mentioned, 26,  
 465.  
*philippensis*, Miq., 26, 461.  
*pomifera*, Wall., 26, 465.  
*Pseudo-sycomorus*, Decne., 26, 458.  
*pumila*, Hook. & Arn., 26, 460.  
*pumila*, Linn., 26, 465.  
*pumila*, Thunb., 26, 459, 465.  
*pyriformis*, Hook. & Arn., 26, 466.  
*pyriformis*, Miq., 26, 457.  
 — var. *Abelii*, King, 26, 457.  
*radiata*, Miq., 26, 464.  
*ramentacea*, Roxb., 26, 466.  
*rapiformis*, Roxb., 26, 464.  
*retusa*, Linn., 26, 466.  
*Roxburghii*, Miq., 26, 462.  
*Roxburghii*, Wall., 26, 467.  
*septica*, Rumph., 26, 464.  
*setosa*, Blume, 26, 462.  
*Sieboldii*, Miq., 26, 467.  
*stenophylla*, Hemsl., 26, 467.  
*stipulata*, Thunb., 26, 465, 466.  
*subpyriformis*, Miq., 26, 466.  
*subulata*, Blume, 26, 467; men-  
 tioned, 461.  
*superba*, Miq., 26, 469.  
 — var. *japonica*, Miq., 26, 469.  
*Swinhoei*, King, 26, 468.  
*Tashiroi*, Maxim., 36, 479.  
*Tikoua*, Bur., 26, 468.  
*trachycarpa*, Miq., 26, 459.  
*vaccinioides*, Hemsl. & King, 26, 468.  
*variegata*, var. *chlorocarpa*, King  
 26, 458.  
*variolosa*, Lindl., 26, 468.

- Ficus vasculosa*, Wall., 26, 468.  
*virgata*, Roxb., 26, 458.  
*Wightiana*, Benth., 26, 469; mentioned, 465, 469.  
*Wrightii*, Benth., 26, 460.  
*Filipendula angustiloba*, Maxim., 23, 223.  
*palmata*, Maxim., 23, 226.  
*Fimbristylis abjiciens*, Steud., 36, 246.  
*acicularis*, R. Br., 36, 230.  
*Actinoschœnus*, C. B. Clarke, 36, 230; mentioned, 247.  
— *var. chinensis*, C. B. Clarke, 36, 230.  
*acuminata*, Vahl, 36, 230.  
*astivalis*, Vahl, 36, 230; mentioned, 231.  
*amblyphylla*, Steud., 36, 231, 238.  
*Arnottii*, Thw., 36, 246.  
*arvensis*, Vahl, 36, 235.  
*autumnalis*, Linn., mentioned, 36, 232.  
— *var. a*, Boeck., 36, 231.  
*barbata*, Benth., 36, 247.  
*bispicata*, Nees, 36, 243, 244, 246.  
*biumbellata*, Boeck., 36, 244.  
*Buergeri*, Miq., 36, 239.  
*cœsia*, Miq., 36, 245.  
*capillaris*, F. Muell., 36, 248.  
*Cinnamometorum*, Hance, 36, 236.  
*Cinnamometorum*, Kunth, 36, 233.  
*communis*, Kunth, 36, 234, 242.  
— (forma), Kunth, 36, 241.  
*complanata*, Benth., 36, 246.  
*complanata*, Link, 36, 231; mentioned, 246.  
— *var. Kraussiana*, C. B. Clarke, 36, 231.  
— *var. microcarpa*, C. B. Clarke, 36, 232.  
*connectens*, Thw., 36, 231.  
*cylindrocarpa*, Kunth, 36, 246.  
*cymosa*, R. Br., 36, 232; mentioned, 242.  
*cyperoides*, R. Br., 36, 232.  
— *var. Cinnamometorum*, Kunth, 36, 233.  
*debilis*, F. Muell., mentioned, 36, 238.  
*decora*, Nees, 36, 243.  
*depauperata*, R. Br., 36, 233.  
*dichotoma*, Vahl, 36, 233; mentioned, 238.  
*Didrichseni*, Boeck., 36, 239.  
— *var. minor*, Boeck., 36, 242.  
*diphylla*, Vahl, 36, 233.  
— *var. leptophylla*, Benth., 36, 234.  
— *var. tomentosa*, Benth., 36, 234.  
*dipsacea*, Benth. & Hook. f., 36, 235.  
*Fimbristylis Eragrostis*, Hance, 36, 235; mentioned, 241.  
*ferruginea*, Vahl, 36, 235.  
*Fordii*, C. B. Clarke\*, 36, 236; mentioned, 237.  
*fulvescens*, Hance, 36, 237.  
*fulvescens*, Thw., mentioned, 36, 237.  
*fusca*, Benth. & Hook. f., 36, 236; mentioned, 237.  
— *var. cantoniensis*, C. B. Clarke\*, 36, 237.  
*globulosa*, Kunth, 36, 237.  
— *var. Torresiana*, C. B. Clarke, 36, 237.  
*glomerata*, Boeck., 36, 232.  
*glomerata*, Nees, 36, 244.  
*gracilentata*, Hance, 36, 237; mentioned, 236.  
*Hanceana*, Boeck., 36, 242.  
*Henryi*, C. B. Clarke\*, 36, 237.  
*insignis*, Thw., 36, 238.  
*japonica*, Sieb., 36, 244.  
*juncea*, Boeck., 36, 241.  
*junciformis*, Munro, 36, 244.  
*Kamphoveneri*, Boeck., 36, 236.  
*Kraussiana*, Krauss, 36, 231.  
*kwantungensis*, C. B. Clarke\*, 36, 238.  
*laxa*, Regel, 36, 231.  
*laxa*, Vahl, 36, 233.  
*leiocarpa*, Maxim., 36, 231.  
*leiocarpa*, Miq., 36, 243.  
*leptoclada*, Benth., 36, 238.  
*leptoclada*, Benth. (sp. div.), 36, 239.  
*limosa*, Kunth, mentioned, 36, 231.  
*longispica*, Steud., 36, 239.  
*Micheliana*, Reichb., 36, 252.  
*micrantha*, Boeck., 36, 239.  
*microcarpa*, F. Muell., 36, 232.  
*miliacea*, Vahl, 36, 239; mentioned, 245.  
*monostachya*, Hassk., 36, 249.  
*multifolia*, Boeck., 36, 232.  
*nigro-brunnea*, Thw., 36, 240.  
*nutans*, Vahl, 36, 241.  
*ovalis*, Nees, 36, 234.  
*pallescens*, Nees, 36, 233.  
*pilosa*, Presl, 36, 233.  
*podocarpa*, Hook. f., 36, 241.  
*podocarpa*, Nees, 36, 234.  
*polymorpha*, Boeck., 36, 234.  
— *forma*, Boeck., 36, 241.  
— *var. depauperata*, Boeck., 36, 243.  
*polytrichoides*, Vahl, 36, 241; mentioned, 236.  
— *var. halophila*, C. B. Clarke, 36, 242.  
*pycnostachya*, Hance, 36, 240.

- Fimbristylis quinquangularis*, Kunth, 36, 242; mentioned, 238.  
*quinquangularis*, Munro, 36, 231, 235.  
*retusa*, Thw., 36, 239.  
*rigida*, Kunth, 36, 244.  
*rigidula*, Kunth, 36, 242.  
*rigidula*, Nees, 36, 242.  
*salbundia*, Boeck., 36, 242.  
*schoenoides*, Vahl, 36, 243; mentioned, 245, 246.  
*sericea*, R. Br., 36, 243.  
*Sieboldii*, Miq., 36, 243.  
*sp.*, F. Muell., 36, 232.  
*spadicea*, Vahl, mentioned, 36, 239.  
— var. *major*, Boeck., 36, 239.  
*spathacea*, Roth, 36, 244; mentioned, 232.  
*squarrosa*, Vahl, 36, 244.  
*Stauntoni*, Debeaux, 36, 244; mentioned, 245.  
*stolonifera*, C. B. Clarke, 36, 244.  
*subbispicata*, Nees & Meyen, 36, 244; mentioned, 239.  
*subtetrastachya*, Boeck., 36, 240.  
*tenera*, Boeck., 36, 232.  
*tetragona*, R. Br., 36, 246.  
*Thomsonii*, Boeck., 36, 246.  
*Thwaitesii*, Boeck., 36, 238.  
*tomentosa*, Vahl, 36, 233.  
*Torresiana*, Gaudich., 36, 237.  
*tristachya*, R. Br., 36, 235.  
*velutina*, Franch., 36, 243.  
*Wightiana*, Nees, 36, 244.  
*yunnanensis*, C. B. Clarke\*, 36, 247.
- Firmiana platanifolia*, Schott & Endl., 23, 90.
- Flacourtia chinensis*, Clos, 23, 57.
- Flagellaria indica*, Linn., 36, 160.  
*repens*, Lour., 36, 186.
- FLAGELLARIACEÆ, 36, 160.
- Flemingia congesta*, Roxb., 23, 197.  
*Grahamiana*, Wight & Arn., 23, 197.  
*Lamontii*, Hance, 23, 197.  
*prostrata*, Roxb., 23, 197.  
*semialata*, Roxb., 23, 197.  
*stricta*, Roxb., 23, 197.  
*vestita*, Benth., 36, 479.  
*yunnanensis*, Franch., 36, 479.
- Fleurya interrupta*, Gaudich., 26, 473.
- Floscopa paniculata*, Benth., 36, 158.  
*scandens*, Lour., 36, 158.
- Flueggea Griffithii*, Baker, 36, 77.  
*Jaburan*, Kunth, 36, 77.  
*japonica*, Rich., 36, 78.
- Flueggea japonica*, var. *umbraticola*, Baker, 36, 78.  
— var. *Wallichiana*, Baker, 36, 78.  
*spicata*, Schult., 36, 79.  
*Wallichiana*, Kunth, 36, 78.
- Fluggea capillipes*, Pax, 36, 479.  
*microcarpa*, Blume, 26, 427.  
*sinensis*, Baill., 26, 427.  
*suffruticosa*, Baill., 26, 426.
- Foeniculum vulgare*, Gaertn., 23, 331.
- Fontanesia chinensis*, Hance, 26, 87.  
*Fortunei*, Carr., 26, 87.  
*phillyreoides*, Labill., 26, 87.  
— var. *sinensis*, Debeaux, 26, 87.
- Fordia*, Hemsl. \*, 23, 160.  
*cauliflora*, Hemsl. \*, 23, 160; mentioned, 489, 491.
- Fordiophyton*, Stapf, 36, 479.  
*cantonense*, Stapf, 36, 479.  
*Faberi*, Stapf, 36, 479.  
*Fordii*, Krasser, 36, 479.
- Forrestia chinensis*, N. E. Br. \*, 36, 158.  
*hispidula*, Less. & Rich., mentioned, 36, 159.
- Forsythia suspensa*, Vahl, 26, 82.  
*viridissima*, Lindl., 26, 82.
- Fortunæa chinensis*, Lindl., 26, 495.
- Fragarea collina*, Maxim., 23, 239.  
*elatior*, Ehrh., 23, 239.  
*filipendula*, Hemsl. \*, 23, 239.  
*indica*, Andr., 23, 240; mentioned, 239.  
*vesca*, Lour., 23, 239.
- Frangula crenata*, Miq., 23, 128.
- Fraxinus bracteata*, Hemsl. \*, 26, 84.  
*Bungeana*, DC., 26, 84; mentioned, 85.  
*chinensis*, Roxb., 26, 85.  
— var. *rhynchophylla*, Hemsl. \*, 26, 86.  
*floribunda*, Bunge, 26, 84, 86.  
*Griffithii*, C. B. Clarke, 26, 84.  
*insularis*, Hemsl. \*, 26, 86.  
*longicuspis*, Sieb. & Zucc., 36, 479.  
*malacophylla*, Hemsl., 36, 479.  
*mandshurica*, Rupr., 26, 86.  
*Mariesii*, Hook. f., 26, 86.  
*obovata*, Blume, 26, 85.  
*Ornus*, var. *Bungeana*, Hance, 26, 85.  
*platypoda*, Oliver, 36, 479.  
*pubinervis*, Blume, 26, 85.  
*retusa*, Champ., 26, 86; mentioned, 84.  
— var. *Henryana*, Oliver, 36, 479.  
*rhynchophylla*, Hance, 26, 85, 86.  
*xanthoxyloides*, Wall., 26, 85, 86.

- Freycinetia formosana*, Hemsl., **36**, 171, 479.  
*Friesia chinensis*, Gardn., **23**, 94.  
*Fritillaria caucasicensis*, Ker-Gawl., **36**, 136.  
     *cantonensis*, Lour., **36**, 136.  
     *ecirrhusa*, D. Don, **36**, 136.  
     — *var. ecirrhusa*, Franch., **36**, 137.  
     *dagana*, Turcz., **36**, 137.  
     *Davidi*, Franch., **36**, 479.  
     *Delavayi*, Franch., **36**, 137.  
     *kamtschatcensis*, Fisch., **36**, 136.  
     *lophophora*, Franch., **36**, 132, 479.  
     *macrophylla*, D. Don, **36**, 133.  
     *Przewalskii*, Maxim., **36**, 137.  
     *Roylei*, Hook., **36**, 479.  
     *Souliei*, Franch., **36**, 137.  
     *Thompsoniana*, D. Don, **36**, 133.  
     *Thunbergii*, Miq., **36**, 137.  
     *verticillata*, Wall., **36**, 137.  
     *verticillata*, Willd., **36**, 137.  
     — *var. ecirrhusa*, Franch., **36**, 138.  
     — *var. Thunbergii*, Baker, **36**, 137.  
*Fuirena glomerata*, Lam., **36**, 256.  
     *pentagona*, Nees, **36**, 257.  
     *pubescens*, Boeck., **36**, 257.  
     *Rottboellii*, Nees, **36**, 256.  
     *umbellata*, Rottb., **36**, 256.  
     *Wallichiana*, Kunth, **36**, 257.  
*Fumaria spectabilis*, Linn., **23**, 36.  
**FUMARIACEÆ**, **23**, 35.  
*Funkia lancifolia*, Spreng., **36**, 116.  
     *ovata*, Spreng., **36**, 116.  
     — *var. α*, Kunth, **36**, 117.  
     — *var. β*, Kunth, **36**, 116.  
     — *var. minor*, Baker, **36**, 117.  
     *Sieboldi*, Lindl., **36**, 117.  
     *Sieboldiana*, Hook., **36**, 117.  
     *subcordata*, Spreng., **36**, 117.  
  
*Gærtnera hongkongensis*, Seem., **26**, 121.  
     *sp.*, Seem., **26**, 94.  
*Gagea fascicularis*, Salisb., **36**, 138.  
     *lutea*, Ker-Gawl., **36**, 138.  
     *pauciflora*, Turcz., **36**, 138.  
     *serotina*, Ker-Gawl., **36**, 139.  
     *triflora*, Schult., **36**, 140.  
*Gahnia javanica*, Moritzi, **36**, 262.  
     *Mariscus*, F. Muell., **36**, 262.  
     *tristis*, Nees, **36**, 263.  
     *Wichurii*, Boeck., **36**, 263.  
*Galactia formosana*, Matsum., **36**, 479.  
     *sp.*, **23**, 190.  
     *Tashiroi*, Maxim., **23**, 190.  
*Galatella Meyendorffii*, Regel, **23**, 412.  
*Galeola Faberi*, Rolfe, **36**, 39.  
  
*Galeola Lindleyana*, Reichb. f., mentioned, **36**, 39.  
*Galium Aparine*, Linn., **23**, 393.  
     *asprellum*, Michx., **23**, 393; mentioned, 394.  
     *boreale*, Linn., **23**, 393.  
     — *var. molle*, Hemsl. \*, **23**, 394.  
     *Bungei*, Steud., **23**, 394.  
     *chinense*, A. Spreng., **23**, 394.  
     *davuricum*, Turcz., **23**, 393.  
     *gracile*, Bunge, **23**, 394.  
     *linearifolium*, Turcz., **23**, 394.  
     *miltorrhizum*, Hance, **23**, 394.  
     *Mollugo*, Linn., **23**, 395.  
     *paradoxum*, Maxim., **36**, 479.  
     *parviflorum*, Maxim., **23**, 393.  
     *pauciflorum*, Bunge, **23**, 393.  
     *pogonanthum*, Franch. & Sav., **23**, 394.  
     *rotundum*, Thunb., **23**, 394.  
     *sororium*, Hance, **23**, 393.  
     *trachyspermum*, A. Gray, **23**, 394.  
     *tricornis*, Stokes, **23**, 395.  
     *trifloriforme*, Komarov, **36**, 479.  
     *triflorum*, Michx., **36**, 479.  
     *tuberosum*, Lour., **23**, 395.  
     *Vaillantii*, DC., **23**, 393.  
     *verum*, Linn., **23**, 395.  
*Garciana cochinchinensis*, Lour., **36**, 150.  
*Garcinia multiflora*, Champ., **23**, 75.  
     *oblongifolia*, Champ., **23**, 75.  
*Gardenia amœna*, Sims, **23**, 383.  
     *daphnoides*, Hance, **23**, 384.  
     *florida*, Linn., **23**, 382.  
     *Fortuniana*, Bot. Reg., **23**, 382.  
     *grandiflora*, Sieb. & Zucc., **23**, 382.  
     *jasminoides*, DC., **23**, 383.  
     *Maruba*, Sieb., **23**, 382.  
     *Pictorum*, Hassk., **23**, 382.  
     *radicans*, Thunb., **23**, 382.  
     *scandens*, Thunb., **23**, 383.  
     *volubilis*, Lour., **23**, 383.  
*Gardneria nutans*, Sieb. & Zucc., **26**, 121.  
*Garnotia drymeia*, Hance, **36**, 386.  
     *patula*, Munro, **36**, 386.  
     — *var. mutica*, Rendle \*, **36**, 387.  
     *stricta*, Benth., **36**, 387.  
*Gastrodia elata*, Blume, **36**, 48.  
*Gaultheria crenulata*, Kurz, **26**, 16.  
     *fragrantissima*, Wall., **36**, 479.  
     *laxiflora*, Diels, **36**, 479.  
     *nummularioides*, D. Don, **36**, 479.  
*Gaura chinensis*, Lour., **23**, 310.  
*Geblera suffruticosa*, Fisch. & Mey., **26**, 426.  
*Geissaspis cristata*, Wight & Arn., **23**, 171.

- Gelonium æquoreum*, *Hance*, 26, 444.  
 — *var. hainanense*, *Hemsl.* \*, 26, 444.  
*Gelsemium elegans*, *Benth.*, 26, 117.  
*Gendarussa vulgaris*, *Nees*, 26, 245.  
*Geniosporum holochilum*, *Hance*, 26, 267.  
*Genipa flava*, *Lour.*, 23, 383.  
*Gentiana algida*, *Pall.*, 36, 480.  
   *alsinoides*, *Franch.*, 26, 123.  
   *Amarella*, *Linn.*, mentioned, 26, 124.  
   *aperta*, *Maxim.*, 26, 123.  
   *aquatica*, *Linn.*, 26, 123; mentioned, 125.  
   *aquatica*, *Lour.*, 26, 129.  
   *argentea*, *Royle. var. albescens*, *Franch.* \*, 26, 124.  
   *aristata*, *Maxim.*, 26, 124.  
   *arrecta*, *Franch.* \*, 26, 124.  
   *barbata*, *Froel.*, 26, 127.  
   *bella*, *Franch.* \*, 26, 124.  
   *bellidifolia*, *Franch.*, 36, 480.  
   *biflora*, *Regel*, 36, 480.  
   *Buergeri*, *Miq.*, 26, 134.  
   *cephalantha*, *Franch.* \*, 26, 125; mentioned, 129.  
   *chinensis*, *Kusnezow*, 36, 480.  
   *contorta*, *Royle*, 26, 125.  
   *crassuloides*, *Bur. & Franch.*, 36, 480.  
   *crassuloides*, *Franch.*, 36, 480.  
   *cyananthiflora*, *Franch.* \*, 26, 126.  
   *dahurica*, *Fisch.*, 36, 480.  
   *Davidi*, *Franch.*, 26, 126; mentioned, 125.  
   *decemfida*, *Buch.-Ham*, 26, 126.  
   *decumbens*, *Linn. f.*, mentioned, 26, 130.  
   *Delavayi*, *Franch.*, 26, 126; mentioned, 129.  
   *delicata*, *Hance*, 26, 127.  
   *detonsa*, *Rottb.*, 26, 127.  
   *Duclouxii*, *Franch.*, 36, 480.  
   *fastigiata*, *Franch.*, 26, 127.  
   *flicaulis*, *Hemsl.* \*, 26, 127.  
   *Fortuni*, *Hook.*, 26, 134.  
   *frigida*, *Haenke*, 36, 480.  
   *gentilis*, *Franch.*, 36, 480.  
   *Grumii*, *Kusnezow*, 36, 480.  
   *Haynaldi*, *Kanitz*, 36, 480.  
   *Henryi*, *Hemsl.* \*, 26, 128.  
   *hexaphylla*, *Maxim.*, 36, 480.  
   *humilis*, *Stev.*, mentioned, 26, 133.  
   *Jamesii*, *Hemsl.* \*, 26, 128.  
   *Jankæ*, *Kanitz*, 36, 480.  
   *Kurroo*, *Royle*, 36, 480.  
   *Kusnezowii*, *Franch.*, 36, 480.  
*Gentiana leucomelæna*, *Maxim.*, 36, 480.  
   *lineolata*, *Franch.*, 26, 128.  
   — *var. verticillaris*, *Forbes & Hemsl.* \*, 26, 128.  
   *linoides*, *Franch.* \*, 26, 129.  
   *Loureiri*, *Griseb.*, 26, 129.  
   *macrophylla*, *Pall.*, 26, 129.  
   *maeulchanensis*, *Franch.*, 36, 480.  
   *Maximowiczii*, *Kusnezow*, 36, 480.  
   *Maximowiczii*, *Kanitz*, 36, 480.  
   *melandrifolia*, *Franch.* \*, 26, 129.  
   *microdonta*, *Franch.* \*, 26, 130.  
   *microphyta*, *Franch.* \*, 26, 130.  
   *myrioclada*, *Franch.*, 36, 480.  
   *napulifera*, *Franch.*, 36, 480.  
   *ornata*, *Wall.*, 36, 480.  
   *otophora*, *Franch.* \*, 26, 130.  
   *papillosa*, *Franch.*, 26, 131; mentioned, 136.  
   *pedicellata*, *Wall.*, mentioned, 26, 126.  
   *Phob*, *Franch.*, 36, 480.  
   *phyllocalyx*, *C. B. Clarke*, 26, 131.  
   *Piasezkii*, *Maxim.*, 26, 131.  
   *picta*, *Franch.* \*, 26, 131; mentioned, 129.  
   *Pneumonanthe*, *Hance*, 26, 134.  
   *Pneumonanthe*, *Linn.*, 26, 134.  
   *praticola*, *Franch.*, 36, 480.  
   *Prattii*, *Kusnezow*, 36, 480.  
   *primulæflora*, *Franch.*, 26, 132.  
   *prostrata*, *Haenke*, mentioned, 26, 125.  
   *Przewalskii*, *Maxim.*, 26, 132.  
   *Pseudo-aquatica*, *Kusnezow*, 36, 480.  
   *pterocalyx*, *Franch.* \*, 26, 132; mentioned, 127.  
   *puberula*, *Franch.* \*, 26, 132.  
   *pulla*, *Franch.* \*, 26, 133.  
   *Pulmonaria*, *Turez.*, mentioned, 26, 126.  
   *purpurata*, *Maxim.*, 36, 480.  
   *quadrifaria*, *Blume*, mentioned, 26, 126.  
   *recurvata*, *C. B. Clarke*, 26, 133.  
   *rhodantha*, *Franch.* \*, 26, 133; mentioned, 127.  
   *rigescens*, *Franch.* \*, 26, 134.  
   *rosularis*, *Franch.*, 36, 480.  
   *rubicunda*, *Franch.*, 26, 134.  
   *samolifolia*, *Franch.*, 36, 480.  
   *scabra*, *Bunge*, 26, 134.  
   *Serra*, *Franch.*, 26, 134; mentioned, 127, 133.  
   *serrata*, *Gunn.*, 26, 127.  
   *sikkimensis*, *C. B. Clarke*, 26, 135.  
   *siphonantha*, *Maxim.*, 36, 480.  
   *Souliei*, *Franch.*, 36, 480.  
   *spathulæfolia*, *Kusnezow*, 36, 480.

- Gentiana squarrosa*, Ledeb., 26, 135; mentioned, 126.  
*stellariæfolia*, Franch. \*, 26, 135.  
*straminea*, Maxim., 26, 136.  
*striata*, Maxim., 26, 136.  
*stylophora*, C. B. Clarke, 26, 136; mentioned, 137.  
*sutchuenensis*, Franch. \*, 26, 136.  
*Szechenyii*, Kanitz, 36, 480.  
*tatsienensis*, Franch., 36, 481.  
*tenella*, Roth., 26, 124, 128, 135.  
*ternifolia*, Franch., 26, 136; mentioned, 129.  
*tetraphylla*, Kusnezow, 36, 481.  
*Thunbergii*, Griseb., 26, 136.  
*Thunbergii*, Sieb. & Zucc., 26, 138.  
*tibetica*, King, 26, 136.  
*tizuensis*, Franch., 36, 481.  
*tongolensis*, Franch., 36, 481.  
*tricholoba*, Franch., 36, 481.  
*trichotoma*, Kusnezow, 36, 481.  
*tubiflora*, Wall., mentioned, 26, 128.  
*vandellioides*, Hemsl. \*, 26, 137.  
*venosa*, Hemsl. \*, 26, 137.  
*volubilis*, D. Don, 26, 122.  
*yunnanensis*, Franch., 26, 138.  
*Zollingerii*, Fawcett, 26, 138.
- GENTIANACEÆ**, 26, 122.
- Geodorum dilatatum*, R. Br., 36, 32.  
*formosanum*, Rolfe, 36, 32.  
*fucatum*, Lindl., 36, 32.  
*purpureum*, R. Br., mentioned, 36, 32.  
*semicristatum*, Lindl., 36, 32; mentioned, 40.
- Geophila reniformis*, D. Don, 23, 288.
- Georchis foliosa*, Lindl., 36, 44.
- GERANIACEÆ**, 23, 97.
- Geranium bifolium*, Maxim., 23, 97.  
*davuricum*, DC., 23, 97.  
*Delavayi*, Franch., 36, 481.  
*eriosomon*, Fisch., 23, 97.  
*koreanum*, Komarov, 36, 481.  
*melanandrum*, Franch., 36, 481.  
*moupinense*, Franch., 36, 481.  
*napuligerum*, Franch., 36, 481.  
*nepalense*, Sweet, 23, 98.  
*ocellatum*, Camb., 36, 481.  
*palustre*, var., Franch., 36, 481.  
*platypetalum*, Franch., 36, 481.  
*pogonanthum*, Franch., 36, 481.  
*pratense*, Linn., 23, 98.  
*pseudosibiricum*, Maxim., 23, 97.  
*Pylzowianum*, Maxim., 23, 98.  
*Robertianum*, Linn., 36, 481.  
*soboliferum*, Komarov, 36, 481.  
*strigosum*, Franch., 36, 481.  
*Thunbergii*, Sieb., 23, 98.
- Geranium umbelliforme*, Franch., 36, 481.  
*Vlassovianum*, DC., 23, 98.  
*Wilfordi*, Maxim., 23, 98.  
*Wlassowianum*, Fisch., 23, 98.  
*yunnanense*, Franch., 36, 481.
- Gerardia glutinosa*, Bunge, 26, 192.
- Gerbera amabilis*, Hance, 23, 473.  
*Anandria*, Sch.-Bip., 23, 472.  
*Delavayi*, Franch., 23, 473.  
*Henryi*, Dunn, 36, 481.  
*ovalifolia*, DC., 23, 473.  
*piloselloides*, Cass., 23, 473.  
*raphanifolia*, Franch., 23, 473.  
*ruficoma*, Franch., 23, 473.  
*Schimperii*, Sch.-Bip., 23, 473.  
*Tananti*, Franch., 36, 481.
- Germainia capitata*, Bal. & Poitr., 36, 379.
- Germaria lutifolia*, Presl, 23, 223.
- GESNERACEÆ**, 26, 224.
- Geum elatum*, Wall., 36, 481.  
*intermedium*, Bess., 23, 239.  
*strictum*, Ait., 23, 239.
- Gilibertia dentigera*, Harms, 36, 481.  
*protea*, Harms, 36, 481.
- Ginkgo biloba*, Linn., 26, 546.
- Giraldia Baroni*, 36, 481.  
*Staplii*, Baroni, 36, 481.
- Giraldinia condensata*, Wedd., 36, 481.  
*vitifolia*, Franch., 36, 481.
- Girardinia cuspidata*, Wedd., 26, 475.  
*heterophylla*, Decne., 26, 475.  
*palmata*, Gaudich., 26, 475.
- Gironniera chinensis*, Benth., 26, 452.  
*nitida*, Benth., 26, 452.  
*subæqualis*, Planch., 26, 452.
- Glabraria tersa*, Linn., 26, 385.
- Glaucium leptopodum*, Maxim., 23, 39.
- Glaux maritima*, Linn., 26, 59.
- Glechoma hederacea*, Linn., 26, 290.
- Gleditschia*, Linn., 23, 207.  
*australis*, Hemsl. \*, 23, 208; mentioned, 209, 491.  
*Delavayi*, Franch., 36, 481.  
*Fontanesii*, Spach, 23, 209.  
*heterophylla*, Bunge, 23, 209.  
*japonica*, Miq., mentioned, 23, 209.  
*macracantha*, Desf., 23, 209.  
*officinalis*, Hemsl., 36, 481.  
*sinensis*, auct., 23, 208.  
*sinensis*, Benth., 23, 208.  
*sinensis*, Lam., 23, 209; mentioned, 208, 210.  
*xylocarpa*, Hance, 23, 209.
- Glehnia littoralis*, F. Schmidt, 23, 331.
- Globba bulbosa*, Gagnep., 36, 67.

- Globba erecta*, Red., 36, 71.  
*japonica*, Thunb., 36, 72.  
*nutans*, Linn., 36, 72.  
*Glochidion Arnottianum*, Muell. Arg., 26, 424.  
*cantoniensis*, Hance, 26, 425.  
*distichum*, Hance, 26, 425.  
*eriocarpum*, Champ., 26, 424.  
*flexuosum*, Muell. Arg., 26, 421.  
*formosanum*, Hayata, 36, 481.  
*Fortuni*, Hance, 26, 424; mentioned, 425.  
*hongkongensis*, Muell. Arg., 26, 424.  
*lanceolatum*, Hayata, 36, 481.  
*littorale*, Benth., 26, 424.  
*macrophyllum*, Benth., 26, 425.  
*molle*, Hook. & Arn., 26, 424.  
*obovatum*, Sieb. & Zucc., 26, 425; mentioned, 425.  
*obscurum*, Blume, 26, 425.  
*philippinense*, Benth., 26, 426.  
*Roxburgianum*, Muell. Arg., 26, 425.  
*sinicum*, Hook. & Arn., 26, 425.  
*villicaule*, Hook. f., 26, 426.  
*Wrightii*, Benth., 26, 426.  
*Glossaspis antennifera*, Reichb., 36, 63.  
*tentaculata*, Spreng., 36, 63.  
*Glossocomia hortensis*, Rupr., 26, 5.  
*lanceolata*, Regel, 26, 5, 6.  
*ussuriensis*, Rupr., 26, 6.  
*Glossogyne tenuifolia*, Cass., 23, 436.  
*Glossostylis arvensis*, Benth., 26, 201.  
*Glossula tentaculata*, Lindl., 36, 63; mentioned, 59.  
*Glyceria aquatica*, Wahlenb., 36, 428.  
*convoluta*, Fries, mentioned, 36, 428.  
*distans*, Wahlenb., 36, 428.  
— var. *convoluta*, Boiss., 36, 428.  
*procumbens*, Dumort., 36, 429.  
*Glycine chinensis*, Sims, 23, 161.  
*floribunda*, Willd., 23, 162.  
*hispida*, Maxim., 23, 188.  
*javanica*, Linn., 23, 189.  
*sinensis*, [Lindl.] Bot. Reg., 23, 161.  
*Soja*, Sieb. & Zucc., 23, 188.  
*tabacina*, Benth., 23, 188.  
*tomentosa*, Benth., 23, 189.  
*ussuriensis*, Regel & Maack, 23, 189.  
*villosa*, Thunb., 23, 195.  
*Glycosmis chinensis*, Oliver, 23, 109.  
*citrifolia*, Lindl., 23, 109.  
*pentaphylla*, Correa, 23, 109.  
*Glycyrrhiza asperima*, Linn., 23, 168.  
*echinata*, Linn., 23, 168.  
*Glycyrrhiza glabra*, Linn., 23, 168; mentioned, 169.  
*glandulifera*, Waldest. & Kit., 23, 168.  
*inflata*, Batalin, 36, 481.  
*pallidiflora*, Maxim., 23, 169.  
*uralensis*, Fisch., 23, 169.  
*Glyptostrobos heterophyllus*, Endl., 26, 544.  
*pensilis*, C. Koch, 26, 544.  
*Gmelina chinensis*, Benth., 26, 257.  
*hainanensis*, Oliver, 26, 257.  
*Gnaphalium amoyense*, Hance, 23, 426.  
*bicolor*, Franch., 36, 481.  
*chrysocephalum*, Franch., 36, 481.  
*confertum*, Benth., 23, 426.  
*confusum*, DC., 23, 427.  
*corymbosum*, Bur. & Franch., 36, 481.  
*Dedekensii*, Bur. & Franch., 36, 481.  
*Delavayi*, Bur. & Franch., 36, 482.  
*hypoleucum*, DC., 23, 426.  
*indicum*, Linn., 23, 427.  
*involutum*, Forst., 23, 427.  
*japonicum*, Thunb., 23, 427.  
*javanum*, DC., 23, 427.  
*leontopodoides*, Willd., 23, 427.  
*likiangense*, Franch., 36, 482.  
*luteo-album*, Linn., var. *multiceps*, Hook. f., 23, 427.  
*margaritaceum*, Linn., 23, 425.  
*multiceps*, Wall., 23, 427; mentioned, 428.  
*nobile*, Bur. & Franch., 36, 482.  
*pellucidum*, Franch., 36, 482.  
*pteroaulon*, Franch. & Sav., 23, 424.  
*purpureum*, Linn., 23, 428.  
*ramigerum*, DC., 23, 427.  
*sericeo-albidum*, Vaniot, 36, 482.  
*Sieboldianum*, Franch. & Sav., 23, 424.  
*sinense*, Franch., 36, 482.  
*Stracheyi*, Franch., 36, 482.  
*subulatum*, Franch., 36, 482.  
*sylvaticum*, Linn., 36, 482.  
*uliginosum*, Linn., 23, 428.  
*yunnanensis*, Franch., 36, 482.  
**GNETACEÆ**, 26, 538.  
*Gnetum elule*, Blume, 26, 536.  
*funiculare*, Wight, 26, 539.  
*gnemonoides*, Brongn., 26, 539.  
*latifolium*, Parl., 26, 539.  
*pyrifolium*, Miq., 26, 539.  
*scandens*, Roxb., 26, 539.  
*Ula*, Brongn., 26, 539.



- Goldbachia lancifolia*, Franch., 36, 482.  
*Goldfussia Cusia*, Nees, 26, 240.  
*pentstemonoides*, Nees, 26, 242.  
*Thomsoni*, Hook., 26, 242.  
*Golowninia japonica*, Maxim., 26, 112.  
*Gomphia neilyherrensis*, Wight, 26, 429.  
*Gomphopetalum* sp., Maxim., 23, 334.  
*Gomphostemma chinensis*, Oliver, 26, 306.  
*Gomphrena globosa*, Linn., 26, 323.  
*sessilis*, Linn., 26, 322.  
*Gongroema Finlaysonii*, Decne., mentioned, 26, 113.  
*Goniocarpus micranthus*, Thunb., 23, 292.  
*scaber*, Kon. & Sims, 23, 292.  
*Gonotheea Blumei*, DC., 23, 377.  
*Gonus amarissimus*, Lour., 23, 112.  
*Gonyanthes nepalensis*, Miers, 36, 4.  
**GOODENOVIÆ**, 26, 1.  
*Goodyera discolor*, A. Rich., mentioned, 36, 44.  
*flabellata*, A. Rich., 36, 43.  
*foliosa*, Benth., 36, 44; mentioned, 45.  
*formosana*, Rolfe, 36, 45.  
*Henryi*, Rolfe, 36, 45.  
*japonica*, Blume, 36, 46.  
*nebularum*, Rolfe \*, 36, 45.  
*procera*, Hook., 36, 45.  
*repens*, R. Br., 36, 45.  
*Schlechtendaliana*, Reichb. f., 36, 46.  
*secundiflora*, Griff., 36, 46.  
*similis*, Blume, 36, 46.  
*spp. un.*, Griff., 36, 45.  
*velutina*, Maxim., 36, 46.  
*Gordonia anomala*, Spreng., 23, 80.  
*axillaris*, var. *acuminata*, E. Pritz., 36, 482.  
*javanica*, Hook., 23, 80.  
*Gorteria Loureiriana*, DC., 23, 459.  
*setosa*, Lour., 23, 459.  
*Gossypium arboreum*, Linn., 23, 88.  
*herbaceum*, Linn., 23, 88.  
*indicum*, Lam., 23, 89.  
*religiosum*, Roxb., 23, 89.  
*Gouffera crassiuscula*, Camb., 23, 70.  
*Gorindooia nervosa*, Wight, 36, 39.  
**GRAMINEÆ**, 36, 319.  
*Grangea maderaspatana*, Poir., 23, 407.  
*Graphistemma pictum*, Benth. & Hook. f., 26, 103.  
*Graptophyllum hortense*, Nees, 26, 247.  
*Gratiola grandiflora*, Roxb., 26, 192.  
*veronicæfolia*, Roxb., 26, 192.  
*Graya elegans*, Nees, 36, 387.  
*Grewia affinis*, Lindl., 23, 92.  
*biloba*, G. Don, 23, 92.  
*glabrescens*, Benth., 23, 92.  
*hirsuta*, Vahl, 23, 92.  
*Microcos*, Hook. & Arn., 23, 92.  
*Microcos*, Linn., 23, 92.  
*nitida*, Juss., 23, 93.  
*parviflora*, Bunge, 23, 93.  
*Piscatorum*, Hance, 23, 93.  
*polygama*, Roxb., 23, 93.  
*Stauntoniana*, G. Don, 23, 92.  
*Grislea tomentosa*, Roxb., 23, 304.  
*Gramilea Reevesii*, Hook. & Arn., 23, 387.  
*Guatteria rufa*, Lindl., 23, 26.  
*suberosa*, Dum., 23, 27.  
*Gueldenstaedtia Delavayi*, Franch., 23, 164.  
*diversifolia*, Maxim., 23, 164.  
*Giraldi*, Harms, 36, 482.  
*Guilloni*, Franch., 23, 164.  
*maritima*, Maxim., 23, 463.  
*multiflora*, Bunge, 23, 164.  
*pauciflora*, Fisch., 36, 482.  
*stenophylla*, Bunge, 23, 164.  
*yunnanensis*, Franch., 36, 482.  
*Guettarda speciosa*, Linn., 23, 384.  
*Guettardella chinensis*, Champ., 23, 384.  
*Guilandina Bonduc*, Linn., 23, 205.  
*Bonducella*, Linn., 23, 205.  
*Gussonea cyperoides*, R. Br., 36, 232.  
*pauciflora*, Brongn., 36, 236.  
*Gutzlaffia africana*, Hance, 26, 23.  
*Gyas humilis*, Salisb., 36, 19.  
*Gymnadenia Chusua*, Lindl., 36, 49.  
*conopsea*, R. Br., 36, 52.  
*crassinervis*, Finet, 36, 482.  
*cucullata*, Rich., 36, 52.  
*Faberi*, Rolfe \*, 36, 52, 53.  
*Galeandra*, Reichb. f., 36, 56.  
*gracilis*, Miq., 36, 52.  
*hemipiliodes*, Finet, 36, 482.  
*lepida*, Reichb. f., 36, 53.  
*obcordata*, Reichb. f., 36, 56.  
*Pinguicula*, S. Moore, 36, 53; mentioned, 52.  
*spathulata*, Lindl., 36, 50.  
*tryphieformis*, Reichb. f., 36, 53.  
*viridis*, A. Rich., 36, 54.  
*Gymnagathis*, Stapf, 36, 482.  
*peperomiifolia*, Stapf, 36, 482.  
*Gymnandra borealis*, Pall., 26, 250.

- Gymnandra Gmelini*, Cham. & Schlecht., 26, 250.  
*Pallasii*, Cham. & Schlecht., 26, 250.  
*Stelleri*, Cham. & Schlecht., 26, 250.
- Gymnema affine*, Decne., 26, 112.  
*Finlaysonii*, Wight, 26, 113.  
*inodorum*, Decne., 26, 107.  
*sylvestre*, var. *chinense*, Benth., 26, 112.  
*tingens*, Wight & Arn., 26, 112.  
— var. *cordifolia*, Wight, 26, 112.
- Gymnocladus chinensis*, Baill., 23, 207.  
*Williamsii*, Hance, 23, 209.
- Gymnopetalum cochinchinensis*.  
*Kurz*, 23, 314.  
sp., 23, 314.
- Gymnopogon digitatus*, Nees, 36, 404.
- Gymnosporia* = *Celastrus*, 36, 482.  
*acuminata*, Hook. f., 36, 482.  
*diversifolia*, Maxim., 23, 123.  
*emarginata*, M. Laws., 36, 482.  
*variabilis*, Loes., 36, 482.
- Gymnotheca chinensis*, Decne., 26, 364.
- Gymnothrix hordeiformis*, Hance, 36, 339.  
*japonica*, Kunth, 36, 338.  
— var. *viridescens*, Miq., 36, 339.
- Gynandropsis pentaphylla*, DC., 23, 50.  
*sinica*, Miq., 23, 5.  
*viscida*, Bunge, 23, 50.
- Gynostemma cardiosperma*, Cogn., 36, 482.  
*cissoides*, Franch. & Sav., 23, 321.  
*pedatum*, Blume, 23, 320.
- Gynura angulosa*, Hance, 23, 447.  
*auriculata*, DC., 23, 448.  
*aurita*, Winkl., 36, 482.  
*bicolor*, DC., 23, 447; mentioned, 448.  
*bulbosa*, Hook. & Arn., 23, 448.  
*divaricata*, DC., 23, 447.  
*ovalis*, DC., 23, 448.  
— var. *pinnatifida*, Hemsl. \*, 23, 448.  
*pinnatifida*, Vaniot, 36, 482.
- Pseudochina*, Benth., mentioned, 23, 448.  
*sarmentosa*, DC., mentioned, 23, 448.  
*sinica*, Diels, 36, 482.
- Gypsophila acutifolia*, Fisch., 23, 64.
- Gypsophila Oldhamiana*, Miq., 23, 64.  
*perfoliata*, Linn., 36, 482.
- Gyrostachys australis*, Blume, 36, 41.
- Habenaria acuifera*, Lindl., 36, 57; mentioned, 58.  
*bracteata*, R. Br., 36, 54.  
*camptoceras*, Rolfe, 36, 58.  
*chlorantha*, Bab., 36, 55.  
*ciliolaris*, Kränzl., 36, 58.  
*commelinifolia*, Wall., mentioned, 36, 58.  
*conopsea*, Benth., 36, 52.  
*cucullata*, Höfft., 36, 52.  
*Davidi*, Franch., 36, 482.  
*Delavayi*, Finet, 36, 482.  
*endothrix*, Miq., 36, 58.  
*enifolia*, Lindl., 36, 58.  
*Faberi*, Rolfe, 36, 52.  
*Fargesii*, Finet, 36, 482.  
*Fordii*, Rolfe, 36, 58.  
*Galeandra*, Reichb. f., 36, 56.  
*geniculata*, D. Don, 36, 58.  
*gigantea*, D. Don, 36, 57.  
*glaucifolia*, Bur., 36, 59.  
*goodyeroides*, D. Don, 36, 54.  
*Hancockii*, Rolfe, 36, 59.  
*Henryi*, Rolfe, 36, 55.  
*herbiola*, R. Br., 36, 55.  
*humidicola*, Rolfe, 36, 59.  
*iantha*, Hook. f., 36, 55.  
*intermedia*, D. Don, 36, 59.  
*japonica*, A. Gray, 36, 56.  
*Keiskei*, Franch., 36, 56.  
*khasiana*, Hook. f., mentioned, 36, 60.  
*lacertifera*, Benth., 36, 53, 63.  
*latilabris*, Hook. f., mentioned, 36, 60.  
*leptoloba*, Benth., 36, 59; mentioned, 63.  
*linearifolia*, Maxim., 36, 61.  
*linguella*, Lindl., 36, 57, mentioned, 58.  
*Miersiana*, Champ., 36, 60.  
— var. *yunnanensis*, Finet, 36, 60.  
*militaris*, Kränzl., 36, 61.  
*militaris*, Reichb. f., mentioned, 36, 61.  
*monophylla*, Rolfe, 36, 50.  
*omeiensis*, Rolfe, 36, 60.  
*pectinata*, Hook. f., 36, 58.  
*peristyloides*, Wight, 36, 57.  
*Pinguicula*, Benth., 36, 53.  
*polytricha*, Rolfe, 36, 60.  
*radiata*, Kränzl., 36, 60.  
*radiata*, Miq., mentioned, 36, 60.  
*reniformis*, Hook. f., 36, 60; mentioned, 59.  
*rhodocheila*, Hance, 36, 61.

- Habenaria sagittifera*, Reichb. f., 36, 61.  
*Sampsoni*, Hance, 36, 54.  
*Sieboldiana*, Miq., 36, 60.  
*stenantha*, Hook. f., 36, 60.  
*stenostachya*, Benth., 36, 56.  
*Susannæ*, R. Br., 36, 57.  
*tentaculata*, Kränzl., 36, 53; mentioned 59.  
*tentaculata*, Reichb. f., 36, 63.  
*ternatea*, Reichb. f., mentioned, 36, 60.  
*virescens*, Spreng., 36, 55.  
*viridis*, R. Br., 36, 54.  
*yunnanensis*, Rolfe \*, 36, 61.
- HEMADORACEÆ, 36, 75.**
- Hæmaria argyroneura*, Miq., 36, 44.  
*discolor*, Lindl., 36, 44.
- Halenia elliptica*, D. Don, 26, 141.  
 — *var. grandiflora*, Hemsl. \*, 26, 141.  
*sibirica*, Borkh., 26, 141.
- Halesia Fortunei*, Hemsl., 26, 75.  
*hispida*, Benth. & Hook. f., 26, 76.
- Halophila ovalis*, Hook. f., 36, 3.  
*ovata*, Gaudich., 36, 3.
- HALORAGÆ, 23, 292.**
- Haloragis micrantha*, R. Br., 23, 292.  
*scabra*, Benth., 23, 292.  
*tetragyna*, Hook. f., 23, 292.
- Haloxyton recurvum*, Bunge, 26, 331.
- HAMAMELIDÆ, 23, 289.**
- Hamamelis arborea*, Mast., 23, 290.  
*chinensis*, R. Br., 23, 290.  
*japonica*, Sieb. & Zucc., 23, 290.  
*mollis*, Oliver, 23, 290.  
*virginiana*, Linn., 23, 290.
- Hamiltonia oblonga*, Franch., 23, 390.  
*suaveolens*, Roxb., 23, 390.
- Hancea*, Hemsl. \*, 26, 309.  
*sinensis*, Hemsl. \*, 26, 310.
- Hancea*, Seem., 26, 310.  
*Hookeriana*, Seem., 26, 440.  
*muricata*, Benth., 26, 440.
- Hancockia*, Rolfe \*, 36, 20.  
*uniflora*, Rolfe \*, 36, 20.
- Hapalosia Laflingiae*, Wall., 23, 71.  
*Haplostylis Meyenii*, Nees, 36, 160.
- Harlandia bryonioides*, Hance, 23, 318.
- Harrisonia Bennettii*, Hook. f., 23, 112.
- Hartia*, Dunn, 36, 482.  
*sinensis*, Dunn, 36, 482.
- Hastingsia coccinea*, Koen., 26, 263.
- Hecatonia pilosa*, Lour., 23, 15.
- Hedeoma nepalensis*, Benth., 26, 280.  
*nepalensis*, Seem., 26, 281.
- Hedera colchica*, C. Koch, 23, 343.  
*Helix*, Linn., 23, 343.
- Hedera hypoglauca*, Hance, 23, 131.  
*japonica*, Jungh., 23, 343.  
*parviflora*, Champ., 23, 343.  
*protea*, Champ., 23, 343.  
*rhombea*, Sieb. & Zucc., 23, 343.  
*senticosa*, Rupr., 23, 342.
- Hedona Davidi*, Williams, 36, 482.  
*sinensis*, Lour., 23, 66.
- Hedycapnos spectabilis*, Planch., 23, 36.
- Hedychium coronarium*, Koen., 36, 69.  
*spicatum*, Buch.-Ham., 36, 482.
- Hedyotis acutangula*, Champ., 23, 372.  
*alata*, Koen., 23, 376.  
*ampliflora*, Hance, 23, 373.  
*angustifolia*, Cham. & Schlecht., 23, 375.  
*Auricularia*, Linn., 23, 373.  
*boerhaavioides*, Hance, 23, 377.  
*borrerioides*, Champ., 23, 376.  
*bracteosa*, Hance, 23, 373.  
*capitellata*, Wall., 23, 373.  
*capituligera*, Hance, 23, 373.  
*cephalophora*, R. Br., 23, 376.  
*consanguinea*, Hance, 23, 373.  
*costata*, R. Br., 23, 373.  
*effusa*, Hance, 23, 374; mentioned, 376.  
*fruticosa*, Rottl., 23, 373.  
*herbacea*, Osbeck, 23, 377.  
*hispida*, Retz., 23, 374.  
*lancea*, Thunb., 23, 374.  
*Lawsoniæ*, Benth., 23, 372.  
*loganioides*, Benth., 23, 374.  
*longidens*, Hance, 23, 372.  
*macrostemon*, Hook. & Arn., 23, 374.  
*ovata*, Thunb., 23, 374.  
*Parryi*, Hance, 23, 374.  
*pinifolia*, Wall., 23, 375.  
*pterita*, Blume, 23, 377.  
*recurva*, Benth., 23, 374, 376.  
*sp.*, 23, 376.  
*tenelliflora*, Blume, 23, 375.  
*tenuipes*, Hemsl. \*, 23, 375.  
*Thwaitesiana*, Hance, 23, 377.  
*uncinella*, Hook. & Arn., 23, 376.  
*Vachellii*, Hook. & Arn., 23, 376.  
*xanthochroa*, Hance, 23, 376.
- Hedysarum brachypterum*, Bunge, 23, 169.  
*diphyllum*, Linn., 23, 171.  
*elegans*, Lour., 23, 171.  
*esculentum*, Ledeb., 23, 169.  
*gangeticum*, Linn., 23, 172.  
*lagopodioides*, Linn., 23, 178.  
*latifolium*, Roxb., 23, 173.

- Hedysarum lutescens*, Poir., 23, 174.  
*multijugum*, Maxim., 23, 169.  
*obscurum*, Linn., 23, 169.  
*pictum*, Jacq., 23, 178.  
*pulchellum*, Linn., 23, 175.  
*reniforme*, Lour., 23, 178.  
 sp., 23, 169.  
*sikkimense*, Benth., 36, 482.  
*tanguticum*, B. Fedtsch., 36, 483.  
*triflorum*, Linn., 23, 176.  
*tuberosum*, B. Fedtsch., 36, 483.  
*umbellatum*, Linn., 23, 177.  
*Vespertilionis*, Linn., 23, 178.  
*Heleocharis acicularis*, Koch, 36, 225.  
 — var. *japonica*, Diels, 36, 226.  
*afflata*, Naves, 36, 226.  
*alta*, Boeck., 36, 229.  
*atropurpurea*, Koch, 36, 226.  
*capitata*, Boeck., 36, 227.  
*chlorocarpa*, Boeck., 36, 226.  
*fistulosa*, Boeck., 36, 227.  
*Græffiana*, Boeck., 36, 230.  
*Kuntzei*, Boeck., 36, 226.  
*ochrostachys*, Boeck., 36, 226.  
*palustris*, Lindl., 36, 228.  
*plantaginea*, Boeck., 36, 228.  
*tetraquetra*, Boeck., 36, 229.  
*Wichurii*, Boeck., 36, 229.  
*Helianthus giganteus*, Lour., 23, 434.  
*Helicandra sinensis*, Hook. & Arn.,  
 26, 96.  
*Helicia cochinchinensis*, Lour., 26,  
 394.  
*cochinchinensis*, Meisn., 26, 394.  
*erratica*, Hook. f., 26, 394.  
*formosana*, Hemsl. \*, 26, 394;  
 36, 483.  
*grandis*, Hemsl., 36, 483.  
*lanceifolia*, Sieb. & Zucc., 26, 394.  
*terminalis*, Kurz, 26, 394.  
*Helicodictyon crinitus*, Schott, men-  
 tioned, 36, 182.  
*Helicteres angustifolia*, Linn., 23,  
 90.  
*spicata*, Colebr., 23, 90.  
 — var. *hainanensis*, Hance, 23,  
 91.  
*undulata*, Lour., 23, 39.  
*Heliotropium brevifolium*, Wall., 26,  
 147.  
*indicum*, DC., 26, 147.  
*indicum*, Linn., 26, 147.  
*strigosum*, Willd., 26, 147.  
*Helleborus chinensis*, Maxim., 36, 483.  
*thibetanus*, Franch., 36, 483.  
*Hellenia abnormis*, Lindl., 36, 71.  
*chinensis*, Willd., 36, 71.  
*Helonias viridis*, Ker-Gawl., 36, 147.  
*Heloniopsis japonica*, Maxim., 36, 140.  
*umbellata*, Baker, 36, 140.  
*Helopus annulatus*, Nees, 36, 320.  
*Helwingia chinensis*, Batalin, 36, 483.  
*himalaica*, Hook. f. & Thoms., 36,  
 483.  
*japonica*, Dietr. & Steud., 23, 341.  
*Hemarthria compressa*, R. Br., 36,  
 361.  
*fasciculata*, Kunth, 36, 361.  
*Hemerocallis alba*, Andr., 36, 117.  
*cærulea*, Andr., 36, 117.  
*citrina*, Baroni, 36, 115.  
*cordata*, Thunb., 36, 130.  
*disticha*, Donn, 36, 115.  
*flava*, Linn., 36, 115.  
*fulva*, Linn., 36, 115.  
 — var. *Kwanso*, Regel, 36, 115.  
 — var. *maculata*, Baroni, 36, 116.  
*graminea*, Andr., 36, 116.  
*graminea*, [P. L. Warren], 36,  
 115.  
*japonica*, Red., 36, 117.  
*japonica*, Thunb., 36, 117.  
*lanceifolia*, Thunb., 36, 116.  
*Middendorffii*, Trautv., 36, 116.  
*minor*, Mill., 36, 116.  
*Sieboldiana*, Lodd., 36, 117.  
*Hemibœa follicularis*, C. B. Clarke,  
 26, 232.  
*gracilis*, Franch., 36, 483.  
*Henryi*, C. B. Clarke, 26, 232.  
*subcapitata*, C. B. Clarke, 26, 232.  
*Hemicicca japonica*, Baill., 26, 421.  
*Hemigraphis chinensis*, T. Anders. \*,  
 26, 238.  
*szechuanica*, Batalin, 36, 483.  
*Hemilophia*, Franch., 36, 483.  
*pulchella*, Franch., 36, 483.  
*Hemiphragma heterophyllum*, Wall.,  
 26, 192.  
*Hemipilia brevicalecarata*, Finet, 36,  
 62.  
*cordifolia*, Lindl., mentioned, 36,  
 62.  
*cruciata*, Finet, 36, 62.  
*flabellata*, Bur., 36, 62.  
*Henryi*, Reichb. f., 36, 62.  
*Hemiptetea Davidii*, Planch., 26, 449.  
*Hemistepta lyrata*, Bunge, 23, 463.  
*Hemsleya chinensis*, Cogn., 23, 490;  
 36, 483.  
*Henrya*, Hemsl. \*, 26, 111.  
*Augustiniana*, Hemsl. \*, 26, 111.  
*Henrya*, Nees, 26, 112.  
*Henslowia frutescens*, Champ., 26,  
 409.  
*sessiliflora*, Hemsl. \*, 26, 409.  
*Hepatica triloba*, Chaix, 23, 11.  
*Heptaca latifolia*, Gardn., 23, 78.  
*Heptapleurum Delavayi*, Franch.,  
 36, 483.

- Heptapleurum Fargesii*, Franch., **36**, 483.  
*Hoi*, Dunn, **36**, 483.  
*macrophyllum*, Dunn, **36**, 483.  
*octophyllum*, Hance, **23**, 342.  
*productum* Dunn, **36**, 483.  
*Heracleum acuminatum*, Franch., **36**, 483.  
*barbatum*, Ledeb., **23**, 336.  
*bivittatum*, Boissieu, **36**, 483.  
*candicans*, Wall., **36**, 483.  
*coreanum*, Boissieu, **36**, 483.  
*dissectum*, Ledeb., **23**, 336.  
*dulce*, Fisch., **23**, 336.  
*Fargesii*, Boissieu, **36**, 483.  
*Hemsleyana*, Diels, **36**, 483.  
*lanatum*, Michx., **23**, 336.  
*Moellendorffii*, Hance, **23**, 336.  
*Rapula*, Franch., **36**, 483.  
*scabridum*, Franch., **36**, 483.  
*Souliei*, Boissieu, **36**, 483.  
*vicinum*, Boissieu, **36**, 483.  
*yunnanense*, Franch., **36**, 483.  
*Heritiera chinensis*, Koen., **36**, 71.  
*littoralis*, Ait., **23**, 90.  
*Herminium alaschanicum*, Maxim., **36**, 52, 483.  
— var. *tangutica*, Maxim., **36**, 52.  
*angustifolium*, Benth., **36**, 50; mentioned, 51.  
*biporosum*, Maxim., **36**, 51.  
*fallax*, Finet, **36**, 483.  
*Monorchis*, R. Br., **36**, 51.  
*reniforme*, Lindl., **36**, 60.  
*Souliei*, Rolfe\*, **36**, 51.  
*tanguticum*, Rolfe\*, **36**, 51.  
*Hernandia peltata*, Meisn., **23**, 393.  
*Herpestis Monniera*, H. B. K., **26**, 186.  
*Hesperis aprica*, Poir., **23**, 44.  
*trichosepala*, Turcz., **23**, 45.  
*Heteropanax fragrans*, Seem., **23**, 343.  
*Heteropappus decipiens*, Maxim., **23**, 412.  
*hispidus*, Less., **23**, 412.  
*incisus*, Sieb. & Zucc., **23**, 412.  
*rigens*, Sieb. & Zucc., **23**, 412.  
*Sampsoni*, Hance, **23**, 415.  
*subserratus*, Sieb. & Zucc., **23**, 412.  
*Heteropogon contortus*, Beauv., **36**, 366.  
*hirtus*, Pers., **36**, 366.  
*Heterosmilax Gaudichaudiana*, Maxim., **36**, 95.  
*japonica*, Kunth, **36**, 96.  
*Hewittia bicolor*, Wight, **26**, 163.  
*Hibiscus Abelmoschus*, Linn., **23**, 87.  
*cancellatus*, Roxb., **36**, 483.  
*Hibiscus chinensis*, DC., **23**, 88.  
*esculentus*, Linn., **23**, 87.  
*ficulneus*, Linn., **23**, 87.  
*mutabilis*, Linn., **23**, 87.  
*Rosa-sinensis*, Linn., **23**, 87.  
*simplex*, Linn., **23**, 90.  
*sinensis*, Mill., **23**, 87.  
*surattensis*, Linn., **23**, 88.  
*syriacus*, Linn., **23**, 88.  
*ternatus*, Cav., **23**, 88.  
*tiliaceus*, Linn., **23**, 88; **36**, 483.  
*Trionum*, Linn., **23**, 88.  
*venustus*, Blume, **36**, 483.  
*Hieracium crocatum*, Fries, mentioned, **23**, 478.  
*sinense*, Vaniot, **36**, 483.  
*umbellatum*, Linn., **23**, 477; mentioned, 478.  
*Hierochloe borealis*, Baker & Moore, **36**, 380.  
*borealis*, Roem. & Schult., **36**, 380.  
*Bungeana*, Trin., **36**, 380.  
*dahurica*, Trin., **36**, 380.  
*glabra*, Trin., **36**, 380.  
*Hookeri*, Maxim., **36**, 380.  
*Himantoglossum cucullatum*, Reichb. f., **36**, 52.  
*viride*, Reichb. f., **36**, 54.  
*Hippocratea obtusifolia*, Roxb., **23**, 125.  
*Hippophaë rhamnoides*, Linn., **26**, 405.  
*Hippuris indica*, Lour., **36**, 229.  
*vulgaris*, Linn., **23**, 292.  
*Hiptage Madablota*, Gaertn., **23**, 96.  
*minor*, Dunn, **36**, 483.  
*obtusifolia*, DC., **23**, 96.  
*sericea*, Hook. f., **26**, 263.  
*Hisingera japonica*, Sieb., **23**, 57.  
*racemosa*, Sieb., **23**, 57.  
*Hisutsua cantoniensis*, DC., **23**, 413.  
*serrata*, Hook. & Arn., **23**, 413.  
*Hoeckia Aschersoniana*, Engl. & Graebn., **36**, 484.  
*Holarrhena affinis*, Hook. & Arn., **26**, 96, 99.  
*Holboellia angustifolia*, Wall., **36**, 484.  
*chinensis*, Franch., **36**, 484.  
*coriacea*, Diels, **36**, 484.  
*cuneata*, Oliver, **36**, 484.  
*latifolia*, Wall., **36**, 484.  
*Holcus fulvus*, R. Br., **36**, 367.  
*latifolius*, Osbeck, **36**, 419.  
*spicatus*, Linn., **36**, 339.  
*Holmskioldia sanguinea*, Willd., **26**, 263.  
*Holostemma pictum*, Champ., **26**, 103.

- Holostemina sinense*, Hemsl. \*, 26, 103.  
*Holosteum sinense*, Henry, 36, 484.  
*Homalanthus Leschenaultianus*, A. Juss., 26, 444.  
*Homalium fagifolium*, Benth., 23, 312.  
*Homœatherum chinense*, Nees, 36, 371.  
*Homoiceltis aspera*, Blume, 26, 452.  
*Homolostachys sinensis*, Boeck., 36, 299.  
*Homonoia riparia*, Lour., 26, 443.  
*Homoplitis crinita*, Trin., 36, 358.  
*Hoppea sibirica*, Reichb., 23, 454.  
*speciosa*, Reichb., 23, 454.  
*Hornemannia bicolor*, Willd., 26, 183.  
*ovata*, Link & Otto, 26, 189.  
*prostrata*, Jacq., 26, 190.  
*Hosta japonica*, Tratt., 36, 117.  
*Hoteia chinensis*, Maxim., 23, 265.  
*Thunbergii*, Sieb. & Zucc., 23, 266.  
*Houttuynia cordata*, Thunb., 26, 364.  
*Hovenia acerba*, Lindl., 23, 131.  
*dulcis*, Thunb., 23, 131.  
*Hoya angustifolia*, Traill, 26, 116.  
*carnosa*, R. Br., 26, 115; mentioned, 116.  
*chinensis*, Traill, 26, 116.  
*crassifolia*, Haw., 26, 116.  
*laxrifolia*, Miq., 26, 116.  
*Motoskei*, Feysm. & Binn., 26, 116.  
*picta*, Hort., 26, 116.  
— var. *argentea*, Hort., 26, 116.  
— var. *aurea*, Hort., 26, 116.  
*Pottsii*, Traill, 26, 116.  
*rotundifolia*, Sieb., 26, 116.  
*trinervis*, Traill, 26, 116.  
*variegata*, Sieb., 26, 116.  
*viridiflora*, R. Br., 26, 115.  
*Humulus aculeatus*, Nutt., 26, 453.  
*cordifolius*, Miq., 26, 453.  
*japonicus*, Sieb. & Zucc., 26, 453.  
*Lupulus*, Linn., 26, 453.  
*Hutchinsia petraea*, Desv., 23, 41.  
*Hybanthera cordifolia*, Link, 26, 115.  
*Hydrangea acuminata*, var. *Buergeri*, Sieb. & Zucc., 23, 273.  
*altissima*, Wall., 36, 484.  
*Arbostiana*, H. Léveillé, 36, 484.  
*aspera*, D. Don, 23, 272.  
— var. *angustifolia*, Hemsl. \*, 26, 273.  
— var. *macrophylla*, Hemsl. \*, 26, 273.  
*Azisai*, Sieb., 23, 273.  
*Belzonii*, Sieb. & Zucc., 23, 273.  
*chinensis*, Maxim., 23, 273.  
*Giraldii*, Diels, 36, 484.  
*Hemsleyana*, Diels, 36, 484.  
*Hydrangea Hortensia*, DC., 23, 273.  
*japonica*, Sieb. & Zucc., 23, 273.  
*Kamienskii*, H. Léveillé, 36, 484.  
*longipes*, Franch., 36, 484.  
*longipes*, Hemsl. \*, 23, 273.  
— var. *lanceolata*, Hemsl. \*, 23, 274.  
*Mandarinorum*, Diels, 36, 484.  
*Maximowiczii*, H. Léveillé, 36, 484.  
*Mœllendorffi*, Hance, 23, 274.  
*oblongifolia*, Blume, 23, 273.  
*Otaksa*, Sieb. & Zucc., 23, 273.  
*petiolaris*, Sieb. & Zucc., 23, 274.  
*Rosthornii*, Diels, 36, 484.  
*scandens*, Maxim., 23, 274.  
*stellata*, Sieb. & Zucc., 23, 273.  
*vestita*, Wall., 23, 274.  
— var. *fimbriata*, Wall., 23, 272.  
*xanthoneura*, Diels, 36, 484.  
*Hydrilla dentata*, Casp., 36, 1.  
*japonica*, Miq., 36, 1.  
*verticillata*, Royle, 36, 1.  
— var. *crispa*, Casp., 36, 1.  
— var. *Roxburghii*, Casp., 36, 1.  
HYDROCHARIDACEÆ, 36, 1.  
*Hydrocharis asiaticus*, Miq., 36, 2.  
*Morsus-ranæ*, Linn., 36, 2.  
*Hydrocotyle abbreviata*, A. Rich., 23, 325.  
*asiatica*, Linn., 23, 324.  
— var. *crispata*, Maxim., 23, 325.  
*Batrachium*, Hance, 23, 325.  
*burmanica*, Kurz, 23, 325.  
*chinensis*, Linn., 23, 326.  
*javanica*, Thunb., 23, 325.  
*lurida*, Hance, 23, 325.  
*nepalensis*, Hook., 23, 325.  
*nitidula*, A. Rich., 23, 325.  
*perexigua*, Hance, 23, 325.  
*polycephala*, Wight & Arn., 23, 325.  
*rotundifolia*, Roxb., 23, 325.  
*rubescens*, Franch., 36, 484.  
*sibthorpioides*, Lam., 23, 325.  
*sinensis*, Thunb., 36, 326.  
*Wilfordi*, Maxim., 23, 326.  
*Hydrolea inermis*, Lour., 26, 142.  
*zeylanica*, Vahl, 26, 142.  
HYDROPHYLLACEÆ, 26, 142.  
*Hydropyrum latifolium*, Griseb., 36, 345.  
*Hydroschænus kyllingioides*, Moritzi, 36, 209.  
*Hydrosme Rivieri*, Engl., 36, 182.  
*Hygrophila assurgens*, Nees, 26, 237.  
*salicifolia*, Nees, 26, 237.  
*Hylomecon japonicum*, Prantl, 36, 484.  
*sutchuense*, Franch., 36, 484.  
*vernalis*, Maxim., 36, 484.

- Hymenachne amplexicaulis*, Nees, **36**, 328.  
*Myuros*, Beauv., **36**, 328.  
*Hymenochæte grossa*, Nees, **36**, 250.  
*Hyoseyamus niger*, Linn., **26**, 177.  
*physaloides*, Linn., **26**, 176.  
*Hypælyptum argenteum*, Vahl, **36**, 257.  
*microcephalum*, R. Br., **36**, 257.  
*Hypocoum chinense*, Franch., **23**, 35.  
*erectum*, Linn., **23**, 35.  
*leptocarpum*, Hook. f. & Thoms., **23**, 35.  
**HYPERICINÆ**, **23**, 72.  
*Hypericum Ascyron*, Linn., **23**, 72.  
*attenuatum*, Choisy, **23**, 72; mentioned, 73.  
*aureum*, Lour., **23**, 72.  
*biflorum*, Lam., **23**, 74.  
*chinense*, Hook., **23**, 72.  
*chinense*, Linn., **23**, 72.  
*chinense*, Retz., **23**, 74.  
*cochinchinense*, Lour., **23**, 74.  
*electrocarpum*, Maxim., **23**, 74.  
*elodeoides*, Choisy, **36**, 484.  
*erectum*, Thunb., **23**, 73.  
*formosanum*, Maxim., **23**, 73.  
*geminiflorum*, Hemsl., **36**, 484.  
*japonicum*, Thunb., **23**, 73.  
*Lalandei*, Choisy, **36**, 484.  
*longistylum*, Oliver, **36**, 484.  
*monanthemum*, Hook. f. & Thoms., **36**, 484.  
*monogynum*, Linn., **23**, 72.  
— var. *salicifolium*, Choisy, **23**, 72.  
*mutilum*, Maxim., **23**, 73.  
*nepaulense*, Choisy, **36**, 484.  
*nervatum*, Hance, **23**, 73.  
*oblongifolium*, Wall., **23**, 73.  
*olympicum*, Lour., **23**, 73.  
*patulum*, Thunb., **23**, 73.  
*perforatum*, Linn., **23**, 74.  
*petiolatum*, Lour., **23**, 73.  
*petiolulata*, Hook. f. & Thoms., **36**, 484.  
*Prattii*, Hemsl., **36**, 484.  
*Przewalskii*, Maxim., **23**, 74.  
*salicifolium*, Sieb. & Zucc., **23**, 72.  
*Sampsoni*, Hance, **23**, 74.  
*Seniawini*, Maxim., **23**, 74.  
*trinervium*, Hemsl., **36**, 484.  
*uralum*, Buch.-Ham., **23**, 73.  
*yunnanense*, Franch., **36**, 484.  
*Hypochoeris*, Linn., genus mentioned, **23**, 479.  
*grandiflora*, Ledeb., **23**, 478.  
*Hypoestes Cumingiana*, Benth. & Hook. f., **26**, 249.  
*Hypoestes purpurea*, R. Br., **26**, 249.  
*sinica*, Miq., **26**, 249.  
*sp. nova?*, **26**, 249.  
*Hypolytrum latifolium*, Rich., **36**, 258.  
*nemorum*, Beauv., mentioned, **36**, 258.  
*pergracile*, Nees, **36**, 266.  
*Hypopithys lanuginosa*, Nutt., mentioned, **26**, 34.  
*Hypoporum annulare*, Nees, **36**, 263.  
*Hypoxis aurea*, Lour., **36**, 86.  
*curculigoides*, Wall., **36**, 86.  
*farinosa*, Thunb., **36**, 76.  
*Franquevillei*, Miq., **36**, 86.  
*minor*, D. Don, **36**, 86.  
*minor*, Seem., **36**, 87.  
*orchioides*, Kurz, **36**, 87.  
*spicata*, Franch., **36**, 76.  
*Hypserpa nitida*, Miers, **23**, 28.  
*Hyptis capitata*, Jacq., **26**, 275.  
*suaveolens*, Poit., **26**, 275.  
*Hyrtanandra hirta*, Miq., **26**, 489.  
*Hyssopus Lophanthus*, Linn., **26**, 288.  
*Ichnanthus pallens*, Munro, **36**, 334.  
*Idesia polycarpa*, var., Diels, **36**, 485.  
*Ilex Aquifolium*, Linn., **23**, 118; **36**, 485.  
*Aquifolium*, Thunb., **26**, 87.  
*ardisioides*, Loes., **36**, 485.  
*asprella*, Champ., **23**, 115.  
*buxifolia*, Hance, **23**, 116.  
*Championii*, Loes., **36**, 485.  
*chinensis*, Sims, **23**, 115.  
*cinerea*, Champ., **23**, 115.  
*corallina*, Franch., **36**, 485.  
*cornuta*, Lindl., **23**, 115.  
*Delavayi*, Franch., **36**, 485.  
*dipyrena*, Wall., **36**, 485.  
*dubia*, var. *hupehensis*, Loes., **36**, 485.  
*Fargesii*, Franch., **36**, 485.  
*ficoidea*, Hemsl. \*, **23**, 116.  
*formosana*, Maxim., **23**, 116.  
*Fortunei*, Lindl., **23**, 116.  
*graciliflora*, Champ., **23**, 116.  
*Hanceana*, Maxim., **23**, 116.  
*Henryi*, Loes., **36**, 485.  
*integra*, Thunb., **23**, 116.  
*intermedia*, Loes., **36**, 485.  
*japonica*, Thunb., **23**, 31.  
*kelungensis*, Loes., **36**, 485.  
*laevigata*, Blume, **23**, 118.  
*leptacantha*, Lindl., **23**, 117.  
*liukiensis*, Loes., **36**, 485.  
*macrocarpa*, Oliver, **36**, 485.  
*Maximowicziana*, Loes., **36**, 485.

- Ilex inemecylifolia*, *Champ.*, 23, 117.  
*Mertensii*, *Maxim.*, 23, 117.  
*metabaptista*, *Loes.*, 36, 485.  
*microcarpa*, *Lindl. & Paxt.*, 23, 118.  
*myriadenia*, *Hance*, 23, 117.  
*nummularia*, *Champ.*, 23, 117.  
*Oldhami*, *Miq.*, 23, 117.  
*oxyphylla*, *Miq.*, 23, 117.  
*pedunculosa*, *Miq.*, 36, 485.  
*Pernyi*, *Franch.*, 23, 117.  
*Pseudo-Godajam*, *Franch.*, 36, 485.  
*pubescens*, *Hook. & Arn.*, 23, 117.  
*purpurea*, var., *Miq.*, 36, 485.  
*ramosa*, *Oliver*, 36, 485.  
*Reevesiana*, *Fortune*, 23, 116.  
*rotunda*, *Thunb.*, 23, 118.  
*szechwanensis*, *Loes.*, 36, 485.  
*viridis*, *Champ.*, 23, 118.  
*Warburgii*, *Loes.*, 36, 485.  
*yunnanense*, *Franch.*, 36, 485.
- ILICINÆ, 23, 115.
- Illecebrum sessile*, *Linn.*, 26, 322.
- Illicium anisatum*, *Linn.*, 23, 23.  
*anisatum*, *Lour.*, 23, 23.  
*Griffithii*, *Hook. f. & Thoms.*, 36, 485.  
*Henryi*, *Diels*, 36, 485.  
*micranthum*, *Dunn*, 36, 485.  
*religiosum*, *Sieb. & Zucc.*, 23, 23.  
*Tashiroi*, *Maxim.*, 36, 485.  
*verum*, *Hook.*, 36, 485.
- Illigera rhodantha*, *Hance*, 23, 295.
- Ilysanthes hyssopoides*, *Benth.*, 26, 191.
- Impatiens angustiflora*, *Hook. f.*, 36, 485.  
*arguta*, *Hook. f. & Thoms.*, 23, 100.  
*Balsamina*, *Linn.*, 23, 100.  
*blepharosepala*, *E. Pritz.*, 36, 485.  
*chætodonta*, *Franch.*, 36, 485.  
*chinensis*, *Linn.*, 23, 100.  
*cochleata*, *Lour.*, 23, 101.  
*corchorifolia*, *Franch.*, 36, 485.  
*Cornucopia*, *Franch.*, 36, 485.  
*Davidi*, *Franch.*, 23, 101.  
*Delavayi*, *Franch.*, 36, 485.  
*dimorphophylla*, *Franch.*, 36, 485.  
*divaricata*, *Franch.*, 36, 485.  
*dolichoceras*, *E. Pritz.*, 36, 485.  
*fasciculata*, *Lam.*, 23, 101.  
*fissicornis*, *Maxim.*, 36, 485.  
*furcellata*, *Hemsl. \**, 23, 101.  
*Hancockii*, *C. H. Wright*, 36, 485.  
*Henryi*, *E. Pritz.*, 36, 485.  
*longialata*, *E. Pritz.*, 36, 485.  
*mutila*, *Lour.*, 23, 101.  
*Nolitangere*, *Linn.*, 23, 101.  
*notolopha*, *Maxim.*, 36, 486.
- Impatiens odontopetala*, *Maxim.*, 36, 486.  
*parviflora*, *Linn.*, mentioned, 23, 101.  
*platyceras*, *Maxim.*, 36, 486.  
*plebeja*, *Hemsl. \**, 23, 101.  
*Potanini*, *Maxim.*, 36, 486.  
*procumbens*, *Franch.*, 36, 486.  
*radiata*, *Hook. f.*, 36, 486.  
*recurvicornis*, *Maxim.*, 36, 486.  
*rostellata*, *Franch.*, 36, 486.  
*stenantha*, *Hook. f.*, 36, 486.  
*stenosepala*, *E. Pritz.*, 36, 486.  
*Textori*, *Miq.*, 23, 102.  
*tubulosa*, *Hemsl. \**, 23, 102.  
*uliginosa*, *Franch.*, 36, 486.  
*vittata*, *Franch.*, 36, 486.  
*yunnanensis*, *Franch.*, 36, 486.
- Imperata arundinacea*, *Cyrill.*, 36, 346.  
— var. *Koenigii*, *Hack.*, 36, 346.  
— var. *pedicellata*, *Debeaux*, 36, 346.  
*eulalioides*, *Miq.*, 36, 347.  
*Koenigii*, *Beauv.*, 36, 346.  
*pedicellata*, *Steud.*, 36, 346.  
*sacchariflora*, *Maxim.*, 36, 347.
- Imperialis dagana*, *Turcz.*, 36, 137.
- Incarvillea Beresovskii*, *Batalin*, 36, 486.  
*compacta*, *Maxim.*, 26, 236.  
*Delavayi*, *Bur. & Franch.*, 36, 486.  
*grandiflora*, *Bur. & Franch.*, 36, 486.  
*grandiflora*, *Poir.*, 26, 235.  
*lutea*, *Bur. & Franch.*, 36, 486.  
*sinensis*, *Lam.*, 26, 236.  
*variabilis*, *Batalin*, 36, 486.
- Indigofera Anil*, *Linn.*, 23, 155.  
*atropurpurea*, *Buch.-Ham.*, 23, 155.  
*Benthamiana*, *Hance*, 23, 156.  
*Bungeana*, *Walp.*, 23, 156.  
*caudata*, *Dunn*, 36, 486.  
*chætodonta*, *Franch.*, 36, 486.  
*chinensis*, *Vogel*, 23, 156.  
*cinerascens*, *Franch.*, 36, 486.  
*coccinea*, *Lour.*, 23, 158.  
*decora*, *Lindl.*, 23, 156.  
*Delavayi*, *Franch.*, 36, 486.  
*endecaphylla*, *Jacq.*, 23, 156.  
*hedysaroides*, *Linn.*, 23, 158.  
*heterantha*, *Wall.*, 36, 486.  
*hirsuta*, *Linn.*, 23, 157.  
*Kirilowi*, *Maxim.*, 36, 486.  
*linifolia*, *Retz.*, 36, 486.  
*liukuensis*, *Makino*, 36, 486.  
*macrostachya*, *Vent.*, 23, 157.  
*melilotoides*, *Hance*, 23, 167.  
*micrantha*, *Bunge*, 23, 156.  
*mollis*, *Franch.*, 36, 486.  
*pendula*, *Franch.*, 36, 486.  
*reticulata*, *Franch.*, 36, 486.



- Indigofera rotundifolia*, Lour., **23**, 157.  
*scabrida*, Dunn, **36**, 486.  
*sensitiva*, Franch., **36**, 486.  
*sericophylla*, Franch., **36**, 486.  
*subulata*, Vahl, **36**, 486.  
*tinctoria*, Linn., **23**, 157.  
*trifoliata*, Linn., **23**, 157.  
*trita*, Linn., **23**, 158.  
*venulosa*, Champ., **23**, 158.  
*virgata*, DC., **23**, 158.  
*Inga bigemina*, Hook. & Arn., **23**, 217.  
*dimidiata*, Hook. & Arn., **23**, 217.  
*duleis*, Willd., mentioned, **23**, 217.  
*Inula ammophila*, Bunge, **23**, 428.  
*Britannica*, Linn., **23**, 428.  
— var. *japonica*, Franch. & Sav., **23**, 429.  
— var. *linariæfolia*, Regel, **23**, 429.  
*Cappa*, DC., **23**, 429.  
*chinensis*, Rupr., **23**, 428.  
*chrysantha*, Diels, **36**, 486.  
*cordata*, Boiss., **23**, 430.  
*cuspidata*, C. B. Clarke, **36**, 486.  
*Giraldii*, Diels, **36**, 486.  
*involutrata*, Miq., **23**, 430.  
*linariæfolia*, Turcz., **23**, 429.  
*Pseudo-cappa*, DC., **23**, 429.  
*pterocaula*, Franch., **36**, 486.  
*racemosa*, Hook., **23**, 430.  
*repanda*, Turcz., **23**, 429.  
*Rouleana*, C. B. Clarke, **23**, 430.  
*salicina*, Bunge, **23**, 428.  
*salicina*, Linn., **23**, 430.  
*sericophylla*, Franch., **36**, 486.  
*serrata*, Bur. & Franch., **36**, 486.  
*Iodes vitiginea*, Hance, **23**, 115.  
*Ionidium suffruticosum*, Ging., **23**, 57.  
*Ioniris ruthenica*, Klatt, **36**, 84.  
*tenuifolia*, Klatt, **36**, 85.  
*Iozoste chinensis*, Blume, **26**, 379.  
*lancifolia*, Blume, **26**, 383.  
*pedunculata*, Blume, **26**, 383.  
*rotundifolia*, var. *oblongifolia*, Nees, **26**, 379.  
*Ipomœa anceps*, Roem. & Schult., **26**, 163.  
*angustifolia*, Jacq., **26**, 156.  
*aquatica*, Forsk., **26**, 157.  
*barbata*, Roth., **26**, 160.  
*Batatas*, Linn., **26**, 157, 160.  
*biflora*, Pers., **26**, 157.  
*biloba*, Forsk., **26**, 157.  
*Bona-nox*, Linn., **26**, 158.  
— var. *purpurascens*, [Ker-Gawl.] Bot. Reg., **26**, 161.  
*cœrulea*, Koen., **26**, 160.  
*calycina*, Benth., **26**, 160.  
*capitellata*, Choisy, **26**, 161.  
*cardiosepala*, Hochst., **26**, 160.  
*Ipomœa carnosâ*, R. Br., **26**, 158.  
*chyseides*, Ker-Gawl., **26**, 158.  
*coccinea*, Linn., mentioned, **26**, 163.  
*congesta*, R. Br., **26**, 158; mentioned, 163.  
*cynosa*, Roem. & Schult., **26**, 159.  
*dasysperma*, Jacq., **26**, 159.  
*digitata*, Linn., **26**, 159.  
*dissecta*, Willd., **26**, 159.  
*edulis*, Choisy, **26**, 157.  
*fastigiata*, Sweet, **26**, 159; mentioned, 157.  
*fastigiata*, Roxb., **26**, 159.  
*flicaulis*, Blume, **26**, 156.  
*Hardwickii*, Hemsl. \*, **26**, 160; mentioned, 157, 163.  
*hederacea*, Jacq., **26**, 160; mentioned, 158, 163.  
*hepaticifolia*, Linn., **26**, 161.  
*insignis*, Andrews, **26**, 159.  
*insularis*, Steud., **26**, 158.  
*involutrata*, Hance, **26**, 162.  
*linifolia*, Blume, mentioned, **26**, 161.  
*maritima*, R. Br., **26**, 157.  
*Nil*, Roth, **26**, 160.  
*obscura*, Ker-Gawl., **26**, 161.  
*palmata*, Forsk., mentioned, **26**, 159, 161.  
*paniculata*, R. Br., **26**, 159.  
*Pes-capræ*, Sweet, **26**, 157.  
*Pes-tigridis*, Linn., **26**, 161.  
*philippinensis*, Choisy, **26**, 161.  
*pileata*, Roxb., **26**, 162.  
*pilosa*, Sweet, **26**, 162.  
*platensis*, [Ker-Gawl.] Bot. Reg., **26**, 159.  
*purpurea*, Lam., **26**, 162.  
*Quamocht*, Linn., **26**, 162.  
*quinata*, R. Br., **26**, 162.  
*repens*, Roth, **26**, 157.  
*reptans*, Hook. & Arn., **26**, 158.  
*reptans*, Poir., **26**, 157.  
*sagittata*, Roxb., **26**, 160.  
*sibirica*, Pers., **26**, 162.  
*sinensis*, Choisy, **26**, 162.  
*sinensis*, Fisch., **26**, 163.  
*sinuata*, Ortega, **26**, 163.  
*sp. n. ?*, **26**, 163.  
*Tashirol*, Matsum., **36**, 487.  
*triloba*, Thunb., **26**, 160.  
*tuberculata*, [Ker-Gawl.] Bot. Reg., **26**, 159.  
*tuberculata*, Roem. & Schult., **26**, 161.  
*Turpethum*, R. Br., **26**, 163.  
*Iranthera discolor*, Maxim., **26**, 225.  
**IRIDEE**, **36**, 80.

- Iris biglumis*, Vahl, **36**, 81.  
*Bungei*, Maxim., **36**, 81.  
*cæspitosa*, Pall., **36**, 84.  
*caricifolia*, Pall., **36**, 81.  
*chinensis*, Bunge, **36**, 85.  
*chinensis*, Curt., **36**, 82.  
*cristata*, Miq., **36**, 85.  
*Delavayi*, Micheli, **36**, 80.  
*dichotoma*, Pall., **36**, 81.  
*Domiana*, Spach, **36**, 81.  
*ensata*, Thunb., **36**, 81.  
— var. *chinensis*, Maxim., **36**, 81.  
*fimbriata*, Klatt, **36**, 85.  
*fimbriata*, Vent., **36**, 82.  
*flavissima*, var. *rupestris*, Ledeb., **36**, 83.  
*fragrans*, Lindl., **36**, 81.  
*gracilis*, Maxim., **36**, 82.  
*graminea*, Thunb., **36**, 81.  
*Grijsi*, Maxim., **36**, 82.  
*hæmatophylla*, Fisch., **36**, 84.  
*Henryi*, Baker, **36**, 82.  
*Itsihatsi*, Hassk., **36**, 83.  
*japonica*, Thunb., **36**, 82.  
*Kæmpferi*, Sieb., **36**, 83.  
*levigata*, Fisch., **36**, 83.  
— var. *Kæmpferi*, Maxim., **36**, 83.  
*Loczyi*, Kanitz, **36**, 83.  
*longifolia*, Royle, **36**, 81.  
*longispatha*, Fisch., **36**, 81.  
*Moorcroftiana*, Wall., **36**, 81.  
*Nertchinskia*, Lodd., **36**, 84.  
*orientalis*, Thunb., **36**, 84.  
*oxypetala*, Bunge, **36**, 81.  
*oxypetala*, Hance, **36**, 82.  
*Pallasii*, Fisch., **36**, 81.  
*pandurata*, Maxim., **36**, 83.  
*pomeridiana*, Fisch., **36**, 81.  
*Potanini*, Maxim., **36**, 83.  
*Rossii*, Baker, **36**, 83.  
*Rosthornii*, Diels, **36**, 487.  
*ruthenica*, Dryand., **36**, 84.  
— var. *nana*, Maxim., **36**, 84.  
*sibirica*, var. *orientalis*, Baker, **36**, 84.  
— var. *sanguinea*, Ker-Gawl., **36**, 84.  
*songarica*, var. *gracilis*, Maxim., **36**, 84.  
*speculatrix*, Hance, **36**, 84.  
*squalens*, Thunb., **36**, 82.  
*tectorum*, Maxim., **36**, 85.  
*tenuifolia*, Pall., **36**, 85; mentioned, 83.  
*Tigridia*, Bunge, **36**, 85.  
*tomiolopha*, Hance, **36**, 85.  
*triflora*, Balb., **36**, 81.  
— var. *chinensis*, Fisch., **36**, 81.  
*ventricosa*, Maxim., **36**, 80.  
*violacea*, Klatt, **36**, 83.
- Isachne albens*, Trin., **36**, 321.
- Isachne altissima*, Debeaux, **36**, 321.  
*australis*, R. Br., **36**, 321.  
*debilis*, Rendle\*, **36**, 322.  
*firmula*, Buese, **36**, 322.  
— var. *nudiglumis*, Hack., **36**, 322.  
*miliacea*, Roth, mentioned, **36**, 322.  
*muricata*, Nees, **36**, 321.  
*Myosotis*, Benth., **36**, 322.  
*pulchella*, Roth, **36**, 307.
- Isatis indigotica*, Fortune, **23**, 49.  
*tinctoria*, Linn., mentioned, **23**, 49.
- Ischæmum angustifolium*, Hack., **36**, 364.  
*antephoroides*, Miq., **36**, 364.  
*aristatum*, Linn., **36**, 364.  
— subsp. *barbatum*, Hack., **36**, 364.  
— subsp. *imberbe*, Hack., **36**, 364.  
*aureum*, Hack., **36**, 365.  
*barbatum*, Debeaux, **36**, 365.  
*barbatum*, Miq., **36**, 364.  
*barbatum*, Retz., **36**, 364.  
*ciliare*, Retz., **36**, 365.  
— var. *genuinum*, Hack., **36**, 365.  
*eristachyum*, Hack., **36**, 365.  
*latifolium*, Miq., **36**, 362.  
*laxum*, R. Br., **36**, 365.  
*leersioides*, Munro, **36**, 363.  
*ophiouroides*, Munro, **36**, 363.  
*pectinatum*, Nees, **36**, 363.  
*rugosum*, Salisb., **36**, 366.  
*Segetum*, Trin., **36**, 366.  
*Sieboldi*, Miq., **36**, 366.
- Isis fimbriata*, Tratt., **36**, 82.
- Isolepis atropurpurea*, Roem. & Schult., **36**, 226.  
*barbata*, R. Br., **36**, 247.  
*capillaris*, Debeaux, **36**, 247.  
*dipsucea*, Roem. & Schult., **36**, 235.  
*Haenkei*, Presl, **36**, 244.  
*longispica*, Steud., **36**, 236.  
*margaritifera*, Nees, **36**, 261.  
*maxima*, A. Dietr., **36**, 250.  
*Micheliana*, Roem. & Schult., **36**, 252.  
*Micheliana*, Stev., **36**, 207.  
*rigescens*, Presl, **36**, 244.  
*setacea*, R. Br., **36**, 253.  
*squarrosa*, Miq., **36**, 258.  
*supina*, R. Br., **36**, 254.  
*trifida*, Nees, **36**, 248.  
*verrucifera*, Maxim., **36**, 235.
- Isolobus campanuloides*, A. DC., **26**, 3.  
*Kerii*, A. DC., **26**, 3.  
*radicans*, A. DC., **26**, 3.  
*Roxburghianus*, A. DC., **26**, 3.  
*Isonema ovata*, Cass., **23**, 401.

- Isopyrum adoxoides*, DC., 23, 18.  
*anemonoides*, Kar. & Kir., 23, 18.  
*auriculatum*, Franch., 36, 487.  
*Delavayi*, Franch., 36, 487.  
*Fargesii*, Franch., 36, 487.  
*grandiflorum*, Fisch., 23, 18.  
*Henryi*, Oliver, 36, 487.  
*japonicum*, Sieb., 23, 18.  
*peltatum*, Franch., 36, 487.  
*sutchuenense*, Franch., 36, 487.  
*vaginatam*, Maxim., 36, 487.  
*Itea chinensis*, Hook. & Arn., 23, 278.  
*ilicifolia*, Oliver, 23, 278.  
*parviflora*, Hemsl., 36, 487.  
*Rosmarinus*, Roem. & Schult., 23, 278.  
*yunnanensis*, Franch., 36, 487.  
*Itoa*, Hemsl., 36, 487.  
*orientalis*, Hemsl., 36, 487.  
*Ixeris debilis*, A. Gray, 23, 480.  
*fontinalis*, DC., 23, 483.  
*Gregorii*, Hance, 23, 485.  
*polycephala*, Cass., 23, 483.  
*ramosissima*, A. Gray, 23, 481.  
*repens*, A. Gray, 23, 483.  
*Thunbergiana*, A. Gray, 23, 483.  
*versicolor*, DC., 23, 485.  
*Ixia chinensis*, Linn., 36, 86.  
*Ixonanthes chinensis*, Champ., 23, 96.  
*Ixora blanda*, Ker-Gawl., 23, 385.  
*Brunonis*, Wall., mentioned, 23, 385.  
*chinensis*, Lam., 23, 385.  
*coccinea*, Curt., mentioned, 23, 385.  
*crocata*, Lindl., 23, 385.  
*grandiflora*, Ker-Gawl., 23, 385.  
*Pavetta*, Roxb., 23, 386.  
*rosea*, Sims, 23, 385.  
*stricta*, Roxb., mentioned, 23, 385.  
  
*Jacquemontia violacea*, Benth., 26, 166.  
*Jambolifera pedunculata*, Linn., 23, 109.  
*resinosa*, Lour., 23, 109.  
*Jambosa acuminatissima*, Hassk., 23, 296.  
*malaccensis*, DC., 23, 297.  
*vulgaris*, DC., 23, 297.  
*Jasminum angulare*, Bunge, 26, 79.  
*angustifolium*, Ker-Gawl., 26, 79.  
*arborescens*, Roxb., mentioned, 26, 82.  
*caudatum*, Wall., mentioned, 26, 81.  
*discolor*, Franch., 36, 487.  
  
*Jasminum dispernum*, Wall., mentioned, 26, 81.  
*floribundum*, 36, 487, sphalm. = seq.  
*floridum*, Bunge, 26, 78; 36, 487.  
*Giraldii*, Diels, 36, 487.  
*grandiflorum*, Linn., mentioned, 26, 79.  
*hirsutum*, Willd., 26, 80.  
*humile*, Linn., mentioned, 26, 78.  
*inornatum*, Hemsl. \*, 26, 78.  
*lanceolaria*, Roxb., 26, 78.  
 — *var. puberulum*, Hemsl. \*, 26, 78.  
*laurifolium*, Roxb., 26, 79.  
*Mesnyi*, Hance, 26, 79.  
*microcalyx*, Hance, 26, 79; mentioned, 78.  
*multiflorum*, Andrew, 26, 80.  
*nudiflorum*, Lindl., 26, 79.  
*officinale*, Linn., mentioned, 26, 79.  
*pachyphyllum*, Hemsl. \*, 26, 79.  
*paniculatum*, Roxb., 26, 80.  
*polyanthum*, Franch., 36, 487.  
*pubescens*, Willd., 26, 80.  
*primulinum*, Hemsl., 36, 487.  
*revolutum*, Sims, 26, 78.  
*Sambac*, Ait., 26, 80.  
*Sieboldianum*, Blume, 26, 79.  
*sinense*, Hemsl. \*, 26, 80.  
*subtriplinerve*, Blume, 26, 81.  
*subulatum*, Lindl., 26, 78.  
*trinerve*, Hance, 26, 81.  
*undulatum*, Ker-Gawl., 26, 81; mentioned, 80.  
 — *var. elegans*, Hemsl. \*, 26, 81.  
*Jatropha Curcas*, Linn., mentioned, 26, 433.  
*Jeffersonia dubia*, Benth. & Hook. f., 23, 33.  
*manchuriensis*, Hance, 23, 33.  
*Johnia Wightii*, Wight & Arn., mentioned, 23, 189.  
 JUGLANDACEÆ, 26, 493.  
*Juglans Camirium*, Lour., 26, 434.  
*japonica*, Sieb., 26, 493.  
*mandshurica*, Maxim., 26, 493.  
*pterococca*, Roxb., 26, 495.  
*regia*, Linn., 26, 493.  
 — *var. sinensis*, C. DC., 26, 493.  
*regia*, Lour., 26, 493.  
*Sieboldiana*, Maxim., 36, 487.  
*villosa*, Wall., 26, 495.  
 JUNCACEÆ, 36, 160.  
*Juncellus inundatus*, C. B. Clarke, 36, 207.  
*pygmaeus*, C. B. Clarke, 36, 207; mentioned, 208.  
*serotinus*, C. B. Clarke, 36, 208.

- Juncus alatus*, Franch., 36, 162.  
*allioides*, Franch., 36, 162.  
*bottnicus*, Wahlenb., 36, 163.  
*bufonius*, Linn., 36, 162.  
*bulbosus*, Linn., 36, 163.  
*compressus*, Jacq., 36, 163.  
*diastrophanthus*, Buchen., 36, 163.  
*effusus*, Linn., 36, 163.  
*glaucus*, Sibth., 36, 164.  
*Hancockii*, Hance, 36, 166.  
*himalensis*, Klotzsch, 36, 164.  
*inflexus*, Huds., 36, 164.  
*khasiensis*, Buchen., 36, 164.  
*lamprocarpus*, Buchen., 36, 166.  
*Leersii*, Buchen., 36, 163.  
*Leschenaultii*, J. Gay, 36, 165.  
*leucomelas*, Royle, mentioned, 36, 166.  
*luzuliformis*, Franch., 36, 164.  
*macranthus*, Buchen., 36, 162.  
*modestus*, Buchen., 36, 164.  
*modicus*, N. E. Br. \*, 36, 165.  
*pauciflorus*, R. Br., 36, 165.  
*Potanini*, Buchen., 36, 165.  
*prismatocarpus*, R. Br., 36, 165.  
— var. *Leschenaultii*, Buchen., 36, 166.  
*Przewalskii*, Buchen., 36, 166.  
*setchuenensis*, Buchen., 36, 166.  
*sinensis*, J. Gay, 36, 165.  
*sphenostemon*, Buchen., mentioned, 36, 165.  
*Thompsoni*, Buchen., 36, 166.  
*triglumis*, Linn., mentioned, 36, 166.  
*Juniperus Cabianca*, Vis., 26, 541.  
*cernuus*, Roxb., 26, 541.  
*chinensis*, Linn., 26, 541.  
— var. *pendula*, Franch., 26, 542.  
— var. *procumbens*, Endl., 26, 542.  
— var. *Smithii*, Loud., 26, 543.  
*communis*, Linn., 26, 542; mentioned, 543.  
— var. *nana*, Baumg., 26, 542.  
*conferta*, Parl., 26, 542.  
*densa*, Gord., 26, 543.  
*flagelliformis*, Reeve, 26, 541.  
*Fortunei*, Hort., 26, 543.  
*japonica*, Hort., 26, 541.  
*litoralis*, Maxim., 26, 542.  
*mongolica*, Hort., 26, 541.  
*nana*, Willd., 26, 542.  
*nepalensis*, Hort., 26, 541.  
*oblonga pendula*, Hort., 26, 543.  
*recurva*, Buch.-Ham., 26, 542.  
— var. *squamata*, Parl., 26, 543.  
*Reevesiana*, Hort., 26, 541.  
*rigida*, Sieb. & Zucc., 26, 543.  
*Juniperus sphaerica*, Lindl., 26, 543.  
*squamata*, Buch.-Ham., 26, 543.  
*struthiacea*, Knight, 26, 542.  
*taxifolia*, Hook. & Arn., 26, 543.  
*Thunbergii*, Hook. & Arn., 26, 542.  
*triangularis*, Hort., 26, 543.  
*Jurinea edulis*, Franch., 36, 487.  
*Souliei*, Franch., 36, 487.  
*Jussiaea Fauriei*, H. Léveillé, 36, 487.  
*fruticosa*, DC., 23, 309.  
*japonica*, H. Léveillé, 36, 487.  
*octofila*, DC., 23, 309.  
*Parmentieri*, H. Léveillé, 36, 487.  
*Philippiana*, H. Léveillé, 36, 487.  
*repens*, Linn., 23, 309.  
*suffruticosa*, Linn., 23, 309.  
*tetragona*, Spreng., 23, 309.  
*villosa*, Lam., 23, 309.  
*Justicia Adhatoda*, Linn., 26, 244.  
*Atkinsoni*, T. Anders., mentioned, 26, 245.  
*bicolor*, Sims, 26, 243.  
*Championi*, T. Anders., 26, 244.  
*chinensis*, Linn., 26, 248.  
*crinita*, Thunb., 26, 248.  
*diffusa*, Willd., 26, 245.  
*Gendarussa*, Linn., 26, 245.  
*latiflora*, Hemsl. \*, 26, 245.  
*leptostachya*, Hemsl. \*, 26, 245.  
*mollissima*, Wall., 26, 247.  
*nervosa*, Vahl, 26, 238.  
*palatifera*, Wall., 26, 244.  
*picta*, Vahl, 26, 247.  
*procumbens*, Linn., 26, 246; mentioned, 247.  
*purpurea*, Lour., 26, 248.  
*quadrifaria*, Wall., 26, 246.  
*simplex*, D. Don, 26, 247.  
*tinctoria*, Lour., 26, 248.  
*ventricosa*, Wall., 26, 247.  
*Kadsura Championi*, C. B. Clarke, 36, 487.  
*chinensis*, Hance, 23, 25.  
*chinensis*, Turcz., 23, 25.  
*japonica*, Benth., 23, 25.  
*Kaempferia cœnobialis*, C. H. Wright, 36, 68.  
*elegans*, Wall., 36, 68.  
*fongyuensis*, Gagnep., 36, 69.  
*Galanga*, Linn., 36, 69.  
*pandurata*, Roxb., 36, 69.  
*rotunda*, Linn., 36, 69.  
*yunnanensis*, Gagnep., 36, 69.  
*Kalanchoë gracilis*, Hance, 23, 280.  
*laciniata*, DC., 23, 280.  
*macrosepala*, Hance, 23, 281.

- Kalanchoë pinnata*, Pers., **23**, 280.  
*spathulata*, DC., **23**, 281.  
*Kalidium gracile*, Fenzl, **26**, 328.  
*Kalopanax divaricatum*, Miq., **23**, 339.  
*ricinifolium*, Miq., **23**, 340.  
*Kandelia Rheedii*, Wight & Arn., **23**, 293.  
*Karivia longicirra*, Miq., **23**, 320.  
*umbellata*, Arn., **23**, 318.  
*Keiskea*, Miq., genus mentioned, **26**, 313.  
*japonica*, Miq., **26**, 279.  
*Kelloggia chinensis*, Franch., **36**, 487.  
*Kerria japonica*, DC., **23**, 229; its fruits not edible, **23**, 489.  
*Keteleeria Davidiana*, Beissner, **26**, 554.  
*Evelyniana*, Mast., **36**, 487.  
*Fabri*, Mast. \*, **26**, 555.  
*Fortunei*, Carr., **26**, 555.  
*sacra*, Beissner, **26**, 555.  
*sp. n.*, **26**, 556.  
*Kirganelia sinensis*, Baill., **26**, 422.  
*Kleinhovia Hospita*, Linn., **23**, 90.  
*Knesebeckia discolor*, Klotzsch, **23**, 321.  
*Knoxia corymbosa*, Willd., **23**, 384.  
*Kobresia cercostachys*, C. B. Clarke \*, **36**, 267.  
*fragilis*, C. B. Clarke \*, **36**, 267.  
*graminifolia*, C. B. Clarke \*, **36**, 268.  
*Prattii*, C. B. Clarke \*, **36**, 268.  
*Royleana*, Boeck., **36**, 268.  
*trinervis*, Boeck., **36**, 269.  
*uncinoides*, C. B. Clarke, **36**, 269.  
*Kochia Scoparia*, Schrad., **26**, 328.  
*Koeleria cristata*, Pers., **36**, 410.  
*Koelreuteria bipinnata*, Franch., **36**, 487.  
*minor*, Hemsl., **36**, 487.  
*paniculata*, Laxm., **23**, 138.  
*paullinioides*, L'Hérit., **23**, 138.  
*Koenigia fertilis*, Maxim., **26**, 345.  
*islandica*, Linn., **26**, 345.  
*pilosa*, Maxim., **26**, 345.  
*Kolkwitzia*, Graeb., **36**, 487.  
*amabilis*, Graeb., **36**, 487.  
*Krascheninikowia Davidi*, Franch., **23**, 67.  
*heterophylla*, Miq., **23**, 68.  
*raphanorrhiza*, Palib., **36**, 487.  
*rupestris*, Turcz., **23**, 69.  
*sylvatica*, Maxim., **36**, 487.  
*Kyllinga albescens*, Steud., **36**, 257.  
*brevifolia*, Boeck., **36**, 223.  
*brevifolia*, Moritzi, **36**, 224.  
*brevifolia*, Rottb., **36**, 223.  
*cylindrica*, Nees, **36**, 223.  
*Kyllinga cyperina*, Retz., **36**, 220.  
*fuscescens*, Boeck., **36**, 224.  
*gracilis*, Kunth, **36**, 223.  
*gracilis*, Steud., **36**, 224.  
*intermedia*, R. Br., **36**, 223.  
— var. *oligostachya*, C. B. Clarke \*, **36**, 224.  
*intermedia*, Seem., **36**, 224.  
*melanosperma*, Nees, **36**, 224.  
*mendorensis*, Steud., **36**, 224.  
*monocephala*, Nees, **36**, 225.  
*monocephala*, Rottb., **36**, 224.  
*monocephala*, Seem., **36**, 224.  
— var., **36**, 224.  
*nana*, Nees, **36**, 225.  
*odorata*, Vahl, mentioned, **36**, 223.  
*oligostachya*, Boeck., **36**, 224.  
*sororia*, Kunth, **36**, 223.  
*sororia*, Miq., **36**, 224.  
*triceps*, Linn. f., **36**, 224.  
*triceps*, Rottb., **36**, 225.  
*triceps*, Thunb., **36**, 223.  
*umbellata*, Roxb., **36**, 221.  
*vaginata*, Steud., **36**, 224.  
LABIATE, **26**, 266.  
*Lablab nankinensis*, DC., **23**, 194.  
*perennans*, DC., **23**, 194 (sphalm. *Dolichos*).  
*vulgaris*, Savi, **23**, 194.  
*Lactuca amurensis*, Herder, **23**, 483.  
*amurensis*, Regel, **23**, 480.  
*atropurpurea*, Franch., **36**, 488.  
*brevirostris*, Champ., **23**, 479.  
*crepidioides*, Vaniot, **36**, 488.  
*debilis*, Maxim., **23**, 480.  
*denticulata*, Maxim., **23**, 480.  
— var. *flexuosa*, Hemsl. \*, **23**, 481.  
*diversifolia*, Vaniot, **36**, 488.  
*elata*, Hemsl. \*, **23**, 481.  
*elegans*, Franch., **36**, 488.  
*erythrocarpa*, Vaniot, **36**, 488.  
*Faberia*, Franch., **36**, 488.  
*formosana*, Maxim., **23**, 482.  
*graciliflora*, DC., **23**, 482.  
*gracilis*, DC., **23**, 482; mentioned, 483.  
*grandiflora*, Franch., **36**, 488.  
*hastata*, Franch., **36**, 488.  
*Hemslayi*, Franch., **36**, 488.  
*Henryi*, Dunn, **36**, 488.  
*hirsuta*, Franch., **36**, 488.  
*humifusa*, Dunn, **36**, 488.  
*lignea*, Vaniot, **36**, 488.  
*likiangensis*, Franch., **36**, 488.  
*melanantha*, Franch., **36**, 488.  
*napifera*, Franch., **36**, 488.

- Lactuca ochroleuca*, Franch., 36, 488.  
*Oldhami*, Maxim., 23, 485.  
*polyccephala*, C. B. Clarke, 23, 483.  
*polypodiifolia*, Franch., 36, 488.  
*Prattii*, Dunn, 36, 488.  
*Pseudosenecio*, Vaniot, 36, 488.  
*Raddeana*, Maxim., 23, 483; 36, 488.  
*ramosissima*, Baker & S. Moore, 23, 481.  
*repens*, Maxim., 23, 483.  
*Roborowskii*, Maxim., 23, 483.  
*rubrolutea*, Vaniot, 36, 488.  
*sativa*, Linn., 23, 483.  
*Scariola*, Linn., 23, 483.  
*sibirica*, Maxim., 23, 484.  
*sonchifolia*, Debeaux, 23, 481.  
*sororia*, Miq., 23, 484; mentioned, 485.  
*Souliei*, Franch., 36, 488.  
*spp. nn.*, 23, 486.  
*squarrosa*, Miq., 23, 480.  
*stolonifera*, Maxim., 23, 484; mentioned, 486.  
*taliensis*, Franch., 36, 488.  
*tatarica*, C. A. Mey., 23, 484.  
*Tatarinowii*, Franch., 36, 488.  
*thibetica*, Franch., 36, 488.  
*Thunbergiana*, Maxim., 23, 484.  
*triangulata*, Maxim., 23, 482; mentioned, 482, 486.  
*triflora*, Hemsl. \*, 23, 485.  
*umbrosa*, Dunn, 36, 488.  
*versicolor*, Sch.-Bip., 23, 485.  
*yunnanensis*, Franch., 36, 488.  
*Lagarosiphon Roxburghii*, Benth., 36, 2.  
*Lagenaria dasystemon*, Miq., 23, 315.  
*vulgaris*, Ser., 23, 314.  
*Lagenophora Billardieri*, Cass., 23, 407.  
*Lagerstroemia floribunda*, Jack, 23, 305.  
*Flos-Reginæ*, Retz., 23, 305.  
*Fordii*, Koehne, 36, 488.  
*indica*, Linn., 23, 306.  
*intermedia*, Koehne, 36, 488.  
*microcarpa*, Hance, 23, 306.  
*Munchausia*, Lam., 26, 306.  
*Reginæ*, Roxb., 23, 306.  
*speciosa*, Pers., 23, 306.  
*subcostata*, Koehne, 23, 306.  
*Laggera alata*, Sch.-Bip., 23, 422.  
*Lagotis brachystachya*, Maxim., 26, 250.  
*brevituba*, Maxim., 26, 250.  
*glauca*, Gært., 26, 250.  
*Lagunea cochinchinensis*, Lour., 26, 343.  
*Lamium album*, Linn., 26, 302.  
— var. *barbatum*, Hook. f., 26, 303.  
*amplexicaule*, Linn., 26, 303.  
*barbatum*, Sieb. & Zucc., 26, 303.  
*chinese*, Benth., 26, 303.  
— var. *parviflora*, Hemsl. \*, 26, 303.  
*garganicum*, Thunb., 26, 303.  
*petiolatum*, Royle, 26, 303.  
*Lampsana apogonoides*, Maxim., 23, 474.  
*Lancea tibetica*, Hook. f. & Thoms., 26, 184.  
*Languas chinensis*, Koen., 36, 71.  
*Lantana Camara*, Linn., 26, 251.  
*Laportea bulbifera*, Wedd., 36, 488.  
*evittata*, Franch., 36, 488.  
*Giraidiana*, E. Pritz., 36, 488.  
*grossedentata*, C. H. Wright \*, 26, 474.  
*oleracea*, Wedd., 26, 474; mentioned, 475.  
*pterostigma*, Wedd., 26, 474.  
*sinensis*, C. H. Wright \*, 26, 474.  
*terminalis*, Wight, 36, 488.  
*Lappa edulis*, Sieb., 23, 460.  
*major*, Gaertn., 23, 460.  
*Lappago racemosa*, Honck., 36, 343.  
*Larbrea aquatica*, Ser., 23, 67.  
*Larix chinensis*, Beissner, 26, 558.  
*davurica*, Turcz., 26, 558.  
*Griffithii*, Hook. f. & Thoms., 26, 558.  
*Kæmpferi*, Carr., 26, 558.  
*leptolepis*, Hort., 26, 558.  
*Potanini*, Batalin, 26, 558.  
*sibirica*, Ledeb., 26, 558.  
*thibetica*, Franch., 26, 559.  
*Lasiagrostis mongholica*, Trin., 36, 383.  
*Lasianthus chinensis*, Benth., 23, 388.  
*cyanocarpus*, Jack, 23, 388.  
*Fordii*, Hance, 23, 388; mentioned, 389.  
*formosensis*, Matsum., 36, 488.  
*Hartii*, Finet & Franch., 36, 488.  
*japonicus*, Miq., mentioned, 23, 389.  
*plagiophyllus*, Hance, 23, 389.  
*sp.*, 23, 389.  
*trichophlebus*, Hemsl. \*, 23, 388.  
*Wallichii*, Wight, 23, 389.  
*Lasiolapis Bennettii*, Planch., 23, 112.  
*multijuga*, Benn., 23, 112.  
*paucijuga*, Benn., 23, 112.

- Lasiolytrum hirtum*, Steud., **3**, 360.  
*hispidum*, Steud., **36**, 360.  
*japonicum*, Steud., **36**, 360.  
*Latania chinensis*, Jacq., **36**, 168.  
*Lathyrus Davidii*, Hance, **23**, 186.  
*Dielsianus*, Harms, **36**, 488.  
*humilis*, Fisch., **36**, 488.  
*linearifolius*, Ser., **23**, 187.  
*maritimus*, Bigel., **23**, 186.  
*palustris*, Linn., **23**, 186.  
*pratensis*, Linn., **36**, 488.  
*vernus*, Koch, **23**, 185.  
*Latouchea*, Finet & Franch., **36**, 488.  
*fokienensis*, Finet & Franch., **36**, 488.  
*Launæa glabra*, Franch., **36**, 488.  
*lampsanoides*, Franch., **36**, 488.  
*pinnatifida*, Cass., **23**, 488.  
**LAURINEÆ**, **26**, 370.  
*Laurus Camphora*, Linn., **26**, 371.  
*chinensis*, Blume, **26**, 379.  
— Hort. Berol., **26**, 372.  
*Cinnamomum*, André, **26**, 371.  
*Cubeba*, Lour., **26**, 380.  
*glauca*, Thunb., **26**, 381.  
*indica*, Thunb., **26**, 378.  
*pedunculata*, Thunb., **26**, 372.  
*Sassafras*, Lour., **26**, 372.  
*umbellata*, Thunb., **23**, 265.  
*Lawsonia alba*, Lam., **23**, 305.  
*inermis*, Linn., **23**, 305.  
*Layia emarginata*, Hook. & Arn., **23**, 204.  
*Lecanthus major*, Wedd., **26**, 480.  
*peduncularis*, Wedd., **26**, 480.  
*Wallichii*, Wedd., **26**, 480.  
*Wightii*, Hook. f., **26**, 480.  
*Wightii*, Wedd., **26**, 480.  
*Lechea chinensis*, Lour., **36**, 156.  
*Leea spinosa*, Spreng., **23**, 338.  
*Leersia hexandra*, Sw., **36**, 345.  
**LEGUMINOSÆ**, **23**, 150.  
*Leibnitzia cryptogama*, Cass., **23**, 473.  
*phænogama*, Cass., **23**, 473.  
*Leimanthium sibiricum*, Schult. f., **36**, 148.  
*Lemna bannatica*, Waldst., **36**, 189.  
*major*, Griff., **36**, 189.  
*minor*, Linn., **36**, 188.  
*orbiculata*, Roxb., **36**, 189.  
*paucicostata*, Hegelm., **36**, 188.  
*polyrhiza*, Linn., **36**, 188.  
**LEMNACEÆ**, **36**, 188.  
**LENTIBULARIÆ**, **26**, 222.  
*Leontice microrhyncha*, S. Moore, **23**, 32.  
*venosa*, S. Moore, **23**, 32.  
*Leontodon sinense*, Lour., **23**, 478.  
*Taraxacum*, Linn., **23**, 478.  
*Leontoglossum scabrum*, Hance, **23**, 22.  
*Leontopodium alpinum*, Cass., **23**, 423.  
— var. *Stracheyi*, Hook. f., **23**, 244.  
*japonicum*, Miq., **23**, 424.  
*sibiricum*, Cass., **23**, 424.  
*sinense*, Hemsl. \*, **23**, 424.  
*Leonurus japonicus*, Miq., **26**, 302.  
*macranthus*, Maxim., **26**, 302.  
*sibiricus*, Linn., **26**, 302.  
*supinus*, Steph., **26**, 299.  
*Lepidagathis hyalina*, Nees, **26**, 244.  
*Lepidium affine*, Ledeb., **23**, 48.  
*chinense*, Franch., **23**, 48.  
*chinense*, Stokes, **23**, 41.  
*incisum*, Roth, **23**, 48.  
*latifolium*, Linn., **23**, 48.  
*micranthum*, Ledeb., **23**, 49.  
*petræum*, Lour., **23**, 41.  
*ruderales*, Linn., **23**, 48.  
*Lepidopelma podocarpifolia*, Klotzsch, **26**, 418.  
*Lepidosperma chinense*, Nees, **36**, 261.  
*Lepironia mucronata*, Rich., **36**, 258.  
*Lepta triphylla*, Lour., **23**, 104.  
*Leptatherum japonicum*, Franch., **36**, 356.  
*Leptochloa chinensis*, Nees, **36**, 407.  
*filiformis*, Roem. & Schult., **36**, 407.  
— var. *humilior*, Presl, **36**, 407.  
*tenerrima*, Miq., **36**, 407.  
*tenerrima*, Roem. & Schult., **36**, 407.  
*Leptodermis diffusa*, Batalin, **36**, 489.  
*oblonga*, Bunge, **23**, 390.  
*Potanini*, Batalin, **36**, 489.  
*umbellata*, Batalin, **36**, 489.  
*vestita*, Hemsl. \*, **23**, 390.  
*Leptopteris sumatrana*, Blume, **26**, 117.  
*Leptopus cordifolius*, Decne., **26**, 420.  
*Leptopyrum fumaroides*, Reichb., **36**, 489.  
*Lepturus repens*, R. Br., **36**, 433.  
*Lepyrodiclis holosteoides*, Fenzl, **23**, 70.  
*quadridentata*, Maxim., **36**, 489.  
*Lespedeza bicolor*, Turcz., **23**, 179; mentioned, 183.  
*Buergeri*, Miq., **23**, 179.

- Lespedeza capillipes*, Franch., 36, 489.  
*Caraganæ*, Bunge, 23, 179.  
*chinensis*, G. Don, 23, 180.  
*ciliata*, Benth., 23, 180.  
*cuneata*, G. Don, 23, 181.  
*cyclobotrya*, Miq., 23, 180.  
*Davidii*, Franch., 23, 180.  
*Delavayi*, Franch., 36, 489.  
*diversifolia*, Hemsl., 36, 489.  
*elliptica*, Benth., 23, 180.  
*eriocarpa*, Benth., 36, 489.  
*fasciculiflora*, Franch., 36, 489.  
*floribunda*, Bunge, 23, 180.  
*Gerardiana*, R. Grah., 36, 489.  
*hirtella*, Franch., 36, 489.  
*juncea*, Pers., 23, 181; mentioned, 180.  
— var. *inschanica*, Maxim., 23, 181.  
— var. *juncea*, Pers., 23, 181.  
— var. *kanaoriensis*, Maxim., 23, 181.  
— var. *latifolia*, Maxim., 23, 182.  
— var. *sericea*, Maxim., 23, 181; mentioned, 180.  
*kandoriensis*, Miq., 23, 507, sphalm. = *juncea*, var. *kauaoriensis*.  
*lanceolata*, Dunn, 36, 489.  
*latifolia*, Dunn, 36, 489.  
*macrocarpa*, Bunge, 23, 182; mentioned, 180.  
*macrophylla*, Bunge, 23, 183.  
*medicaginoides*, Bunge, 23, 182.  
*Oldhami*, Miq., 23, 179.  
*pilosa*, Sieb. & Zucc., 23, 182.  
*sericea*, Miq., 23, 180, 181.  
*Sieboldi*, Miq., 23, 179.  
*sp. n.*, 23, 184.  
*stipulacea*, Maxim., 23, 182.  
*striata*, Hook. & Arn., 23, 182.  
*Swinhoei*, Hance, 23, 183.  
*tomentosa*, Sieb., 23, 183.  
*trichocarpa*, Pers., 23, 183.  
*trigonoclada*, Franch., 36, 489.  
*variegata*, Camb., 36, 489.  
*velutina*, Dunn, 36, 489.  
*Viatorum*, Champ., 23, 183.  
*villosa*, Pers., 23, 183.  
*virgata*, DC., 23, 183.  
*yunnanensis*, Franch., 36, 489.  
*Lethia* (err. cf. *Sethia*).  
*Lettsomia Chalmersii*, Hance, 26, 156.  
*Championi*, Benth. & Hook. f., 26, 26, 156.  
*Leucæna glauca*, Benth., 23, 215.  
*Leucanthemum palmatum*, Lam., 23, 438.  
*sibiricum*, DC., 23, 438.  
*Leucas Benthamiana*, Hook. & Arn., 26, 304.  
*Leucas chinensis*, R. Br., 26, 303; mentioned, 304.  
*javanica*, Benth., 26, 304.  
*javanica*, Hook. & Arn., 26, 304.  
*lanata*, Benth., 26, 304.  
*mollissima*, Wall., 26, 304.  
*zeylanica*, R. Br., 26, 304.  
*Leucococcus vimineus*, Wedd., 26, 491.  
*Leucojum capitulatum*, Lour., 36, 87.  
*Leuconotis sp.*, Benth., 26, 94, 121.  
*Leucorchis sylvatica* Blume, 36, 47.  
*Leucosceptrum canum*, Sm., 26, 310.  
*sinense*, Hemsl. \*, 26, 310.  
*Leucothoë sp. ?*, 26, 16.  
*Leuzea dahurica*, Bunge, 23, 470.  
*Leveillea Vaniot*, 36, 489.  
*chinensis*, Vaniot, 36, 489.  
*Martini*, Vaniot, 36, 489.  
*Leycesteria formosa*, Wall., 23, 368.  
*sinensis*, Hemsl., 36, 489.  
*Libanotis sibirica*, Koch, 23, 331.  
*vulgaris*, DC., 23, 330.  
*Libocedrus macrolepis*, Benth., 26, 540.  
*Licuala Fordiana*, Becc., 36, 168.  
*Ligularia* = *Senecio*, 36, 489.  
*altaica*, DC., 23, 450.  
*Clivorum*, Maxim., 23, 451.  
*euodon*, Miq., 23, 454.  
*Fargesii*, Franch., 36, 489.  
*gigantea*, Sieb. & Zucc., 23, 454.  
*jaluense*, Komarov, 36, 489.  
*japonica*, Less., 23, 453.  
*Kæmpferi*, Sieb. & Zucc., 23, 454.  
*mongolica*, DC., 23, 455.  
*Przewalskii*, Maxim., 36, 489.  
*racemosa*, DC., 23, 454.  
*sibirica*, Cass., 23, 454.  
— var. *speciosa*, DC., 23, 454.  
*tenuipes*, Franch., 36, 489.  
*villifera*, Franch., 36, 489.  
*yesoënsis*, var., Franch., 36, 489.  
*Ligusticum acuminatum*, Franch., 36, 489.  
*acutilobum*, Sieb. & Zucc., 23, 332.  
*angelicæfolium*, Franch., 36, 489.  
*austriacum*, Linn., 23, 333.  
*brachilobum*, Franch., 36, 489.  
*daucoides*, Franch., 36, 489.  
*Delavayi*, Franch., 36, 489.  
*ferulaceum*, Franch., 36, 489.  
*glaucescens*, Franch., 36, 489.  
*involucratum*, Franch., 36, 489.  
*multivittatum*, Franch., 36, 489.  
*Pseudo-angelica*, Boissieu, 36, 489.  
*pteridophyllum*, Franch., 36, 489.  
*scoticum*, Linn., 23, 332.  
*sinense*, Oliver, 36, 490.  
*tenuifolium*, Franch., 36, 490.



- Ligusticum tenuisectum*, Boissieu, **36**, 490.  
     Thomsoni, C. B. Clarke, **36**, 490.  
     Wallichii, Franch., **36**, 490.  
*Ligustridium japonicum*, Spach, **26**, 91.  
*Ligustrina amurensis*, Rupr., **26**, 82.  
     — var. *pekinensis*, Maxim., **26**, 82.  
*Ligustrum amurense*, Carr., **26**, 91.  
     *brachystachyum*, Decne., **26**, 89.  
     *californicum*, Hort., **26**, 91.  
     *Calleryanum*, Decne., **26**, 89.  
     *chinense*, Carr., **26**, 93.  
     *ci'iatum*, Sieb., **26**, 91.  
     *confusum*, Decne., **26**, 89.  
     *coriaceum*, Carr., **26**, 90.  
     *deciluum*, Hemsl. \*, **26**, 90.  
     *Delavayanum*, Hariot, **36**, 490.  
     *glabrum*, Hort., **26**, 91.  
     *Henryi*, Hemsl. \*, **26**, 90.  
     *Ibota*, Sieb., **26**, 91.  
     — var. *angustifolium*, Blume, **26**, 91.  
     — var. *obovatum*, Blume, **26**, 91.  
     — var. *velutinum*, Blume, **26**, 91.  
     *japonicum*, Thunb., **26**, 91.  
     *Kellerianum*, Vis., **26**, 91.  
     *lucidum*, Ait., **26**, 91; mentioned, 92.  
     — var., **26**, 90.  
     *mellosum*, Decne., **36**, 490.  
     *molliculum*, Hance, **26**, 92.  
     *myrianthum*, Diels, **36**, 490.  
     *obtusifolium*, Sieb. & Zucc., **26**, 91.  
     *ovalifolium*, Hassk., **26**, 91.  
     *parviflorum*, Vis., mentioned, **26**, 93.  
     *patulum*, Palib., **36**, 490.  
     *Quihoui*, Carr., **26**, 92.  
     *reticulatum*, Blume, **26**, 91.  
     *robustum*, Blume, **26**, 92.  
     *Roxburghii*, Blume, **26**, 92.  
     *Sieboldii*, Hort., **26**, 91.  
     *sinense*, Lour., **26**, 92; mentioned, 89, 93.  
     — var., **26**, 89.  
     *spicatum*, Hort., **26**, 91.  
     *Stauntoni*, DC., mentioned, **26**, 93.  
     *strongylophyllum*, Hemsl. \*, **26**, 93.  
     *syringæflorum*, Hort., **26**, 91.  
     *tibeticum*, Decne., **36**, 490.  
     *Tschonoskii*, Decne., mentioned, **26**, 90.  
     *villosum*, May, **26**, 93.  
     *vulgare*, Thunb., **26**, 91.  
     *vulgare*, Linn., mentioned, **26**, 90.  
     *yunnanense*, L. Henry, **36**, 490.  
**LILIACEÆ**, **36**, 95.
- Lilium amabile*, Palibin, **36**, 127.  
     *apertum*, Franch., **36**, 127.  
     — var. *thibetica*, Franch., **36**, 128.  
     *Bakerianum*, Collett & Hemsl., **36**, 128.  
     *Biondii*, Baroni, **36**, 133.  
     *Brownii*, F. E. Brown, **36**, 128.  
     — var. *chloraster*, Baker, **36**, 128.  
     — var. *leucanthum*, Baker, **36**, 1-8.  
     *Buschianum*, Lodd., **36**, 129.  
     *callosum*, Sieb., **36**, 129.  
     *camschatcense*, Linn., **36**, 136.  
     *caudidum*, Thunb., **36**, 131.  
     *cernuum*, Komarov, **36**, 490.  
     *chinense*, Baroni, **36**, 134.  
     — var. *atropurpureum*, Baroni, **36**, 134.  
     *Christi*, Baroni, **36**, 490.  
     *concolor*, [Ker-Gawl.] Bot. Mag., **36**, 1-9.  
     *concolor*, Elwes, **36**, 129.  
     *concolor*, Salisb., **36**, 129.  
     — var. *Buschianum*, Baker, **36**, 129.  
     — var. *pulchellum*, Baker, **36**, 129.  
     — var. *sinicum*, Hook. f., **36**, 124.  
     *cordifolium*, D. Don, **36**, 131.  
     *cordifolium*, Thunb., **36**, 130.  
     *Davidi*, Duch., **36**, 490.  
     *Delavayi*, Franch., **36**, 130.  
     *Duchartrei*, Franch., **36**, 130.  
     *Fargesii*, Franch., **36**, 130.  
     *formosum*, Franch., **36**, 130.  
     *giganteum*, Wall, **36**, 130.  
     *Henrici*, Franch., **36**, 131.  
     *Henryi*, Baker, **36**, 131.  
     *Jama-Juri*, Sieb., **36**, 131.  
     *japonicum*, Sims, **36**, 128.  
     *lanceifolium*, var. *gloriosides*, Bretsch., **36**, 134.  
     *lankongense*, Franch., **36**, 131.  
     *longiflorum*, Franch., **36**, 128.  
     *longiflorum*, Thunb., **36**, 131.  
     — var. *chloraster*, Baker, **36**, 128.  
     — var. *suaveolens*, [Ker-Gawl.] Bot. Reg., **36**, 131.  
     *lophophorum*, Franch., **36**, 132.  
     *Lowii*, Baker, **36**, 132.  
     *medeoloides*, A. Gray, **36**, 132.  
     *mirabile*, Franch., **36**, 132.  
     *myriophyllum*, Franch., **36**, 132.  
     *nepalense*, Collett & Hemsl., **36**, 132.  
     *nepalense*, D. Don, **36**, 133.  
     *ochraceum*, Franch., **36**, 133.  
     *oxypetalum*, Franch., **36**, 127.  
     *Palibinianum*, Yabe, **36**, 490.  
     *papilliferum*, Franch., **36**, 133.  
     *partheneion*, Sieb., **36**, 129.  
     *pomponicum*, Thunb., **36**, 129.  
     *Pseudotogrinum*, Carrière, **36**, 133.

- Lilium pulchellum*, Fisch., 36, 129.  
*roseum*, Wall., 36, 133.  
*Rosthornii*, Diels, 36, 490.  
*shensiense*, Baroni, 36, 490.  
*sinicum*, Fl. des Serres, 36, 129.  
*speciosum*, Andr., 36, 135.  
*speciosum*, Thunb., 36, 134.  
— var. *gloriosoides*, Baker, 36, 134.  
*superbum*, Thunb., 36, 134.  
*sutchuenense*, Franch., 36, 134.  
— var. *atropurpureum*, Baroni, 36, 134.  
*taliense*, Franch., 36, 134.  
*tenuifolium*, Fisch., 36, 135.  
— var. *punctatum*, Bur., 36, 134.  
*Thompsonianum*, Lindl., 36, 133.  
*tigrinum*, Ker-Gawl., 36, 132.  
*tsingtanense*, Gilg, 36, 490.  
*yunnanense*, Franch., 36, 136.
- Limacia cuspidata*, Hook. f. & Thoms., 23, 28.  
*sagittata*, Oliver, 36, 490.
- Limnanthemum cristatum*, Griseb., 26, 142.  
*nymphoides*, Hoffm. & Link, 26, 142.
- Limnochloa acutangula*, Nees, 36, 227.  
*media*, Nees, 36, 227.  
*plantaginea*, Nees, 36, 228.  
*tumida*, Nees, 36, 228.
- Limnophila erecta*, Benth., 26, 185.  
*gratioloides*, R. Br., 26, 186.  
*gratissima*, Blume, 26, 185.  
*heterophylla*, Benth., 26, 186.  
*hirsuta*, Benth., 26, 186.  
*hypericifolia*, Benth., 26, 186.  
*Menthastrum*, Benth., 26, 186.  
*punctatum*, Blume, 26, 185.  
*Roxburghii*, G. Don, 26, 186.  
*sessiliflora*, Blume, 26, 186.
- Limodorum falcatum*, Thunb., 36, 38.  
*Incarvillei*, Sw., 36, 18.  
*striatum*, Thunb., 36, 19.  
*Tancarvilleæ*, Banks, 36, 18.  
*Tankervilleæ*, Dryand., 36, 18.
- Limonia bilocularis*, Roxb., 23, 110.  
*monophylla*, Lour., 23, 110.  
*parvifolia*, Sims, 23, 109.  
*trichocarpa*, Hance, 23, 109.  
*trifoliata*, Linn., 23, 109.
- Linaria buriatica*, Turcz., 26, 177.  
*Elatine*, Mill., mentioned, 26, 177.  
*linifolia*, Bunge, 26, 178.  
*quadrifolia*, Hance, 26, 178.  
*spuria*, Mill., mentioned, 26, 177.  
*thibetica*, Franch., 36, 490.  
*vulgaris*, Mill., 26, 178.
- Lindenbergia macrostachya*, Hance, 26, 184.  
*philippensis*, Benth., 26, 184.  
*urticæfolia*, Lehm., 26, 184.
- Lindera*, Thunb., mentioned, 26, 370.  
*bifaria*, Benth., 26, 386.  
*caudata*, Diels, 36, 490.  
*cercidifolia*, Hemsl. \*, 26, 387.  
*citridiora*, Hemsl. \*, 26, 387.  
*communis*, Hemsl. \*, 26, 387.  
*fragrans*, Oliver, 26, 388.  
*fruticosa*, Hemsl. \*, 26, 388.  
*glauca*, Blume, 26, 388, 391; 36, 490.  
*megaphylla*, Hemsl. \*, 26, 389; mentioned, 390.  
*membranacea*, Maxim., 26, 389.  
*mollis*, Oliver, 26, 390.  
*obovata*, Franch., 36, 490.  
*obtusiloba*, Blume, 26, 390.  
*Oldhami*, Hemsl. \*, 26, 390.  
*pedunculata*, Diels, 36, 490.  
*populifolia*, Hemsl. \*, 26, 390.  
*præcox*, Blume, 26, 391.  
*puberula*, Franch., 36, 495.  
*reflexa*, Hemsl. \*, 26, 391.  
*Rosthornii*, Diels, 36, 490.  
*sericea*, Blume, 26, 389; mentioned, 391.  
*sinensis*, Hemsl. \*, 26, 392.  
*strychnifolia*, Villar, 26, 392; 36, 490.  
— var., 26, 392.  
*triloba*, Blume, 26, 392; mentioned, 23, 142.  
*Tzumu*, Hemsl. \*, 26, 392.  
*umbellata*, Thunb., 26, 393.
- Lindernia japonica*, Thunb., 26, 183.
- LINEÆ, 23, 95.
- Linnæa*, partim = *Abelia*.  
*adenotricha*, Graebn., 36, 490.  
*angustifolia*, Bur. & Franch., 36, 490.  
*Aschersoniana*, Graebn., 36, 490.  
*biflora*, Koehne, 36, 490.  
*borealis*, Linn., 23, 359.  
*chinensis*, A. Br. & Vatke, 23, 358.  
*Dielsii*, Graebn., 36, 490.  
*Engleriana*, Graebn., 36, 490.  
*Koehneana*, Graebn., 36, 490.  
*macrotera*, Graebn. & Buchw., 36, 490.  
*onkocarpa*, Graebn., 36, 490.  
*parviflora*, Graebn., 36, 490.  
*rupestris*, A. Br. & Vatke, 23, 358; 36, 490.  
*Schumannii*, Graebn., 36, 490.  
*serrata*, Graebn., 36, 490.  
*teraticalyx*, Graebn. & Buchw., 36, 491.

- Linnaea umbellata*, Graebn., **36**, 491.  
*uniflora*, A. Br. & Vatke, **23**, 358.  
*Zanderi*, Graebn., **36**, 491.
- Linociera cambodiana*, Hance, **26**, 89.  
*chinensis*, Fisch., **26**, 89.
- Linum davuricum*, Miq., **23**, 95.  
*nutans*, Maxim., **23**, 95.  
*perenne*, Linn., **23**, 95.  
*salsuginosum*, Franch., **23**, 95.  
*stelleroides*, Planch., **23**, 95.  
*trigynum*, Roxb., **23**, 96.
- Liparis acuminata*, Hook. f., mentioned, **36**, 6.  
*bituberculata*, var. *formosana*, Ridl., **36**, 7.  
*bootanensis*, Griff., **36**, 491.  
*chloroxantha*, Hance, **36**, 6.  
*cornicaulis*, Makino, **36**, 491.  
*formosana*, Reichb. f., **36**, 7.  
*Henryi*, Rolfe, **36**, 6.  
*latilabris*, Rolfe\*, **36**, 6; mentioned, 8.  
*longipes*, Lindl., **36**, 7.  
*macrantha*, Rolfe, **36**, 7; mentioned, 6.  
*nervosa*, Lindl., **36**, 7.  
*odorata*, Lindl., **36**, 7; mentioned, 8.  
*paradoxa*, Reichb. f., **36**, 7.  
*pauciflora*, Rolfe, **36**, 8.  
*plantaginea*, Lindl., mentioned, **36**, 6, 8.  
*plicata*, Franch., **36**, 8.  
*rostrata*, Reichb. f., mentioned, **36**, 8.  
*yunnanensis*, Rolfe\*, **36**, 8.
- Lipocarpa foliosa*, Miq., **36**, 259.  
*microcephala*, Hance, **36**, 253.
- Lippia nodiflora*, Michx., **26**, 251.
- Liquidambar acerifolia*, Maxim., **23**, 291.  
*chinensis*, Champ., **23**, 291.  
*formosana*, Hance, **23**, 291.  
*Maximowiczii*, Miq., **23**, 291.  
*Rosthornii*, Diels, **36**, 491.  
*sp.*, **23**, 292.
- Liriodendron chinense*, Sargent, **36**, 491.  
*Coco*, Lour., **23**, 24.  
*Figo*, Lour., **23**, 24.  
*liliifera*, Linn., **23**, 24.  
*Tulipifera*, Linn., **23**, 25.
- Liriope graminifolia*, Baker, **36**, 79.  
 — var. *densiflora*, **36**, 80.  
*kansuensis*, C. H. Wright\*, **36**, 79.  
*spicata*, Lour., **36**, 79.  
 — var. *densiflora*, C. H. Wright\*, **36**, 79.
- Liriope spicata*, var. *intermedia*, Maxim., **36**, 80.  
 — var. *koreana*, C. H. Wright\*, **36**, 80.  
 — var. *latifolia*, Franch., **36**, 80.  
 — var. *minor*, C. H. Wright\*, **36**, 80.
- Liriopsis fuscata*, Spach, **23**, 24.
- Listera grandiflora*, Rolfe, **36**, 40.  
*puberula*, Maxim., **36**, 41.  
*reniformis*, D. Don, **36**, 60.
- Litchi chinensis*, Sonnerat, **23**, 139.
- Lithospermum arvense*, Linn., **26**, 154.  
*chinense*, Hook. & Arn., **26**, 154.  
*confertiflorum*, Miq., **26**, 155.  
*erythrorhizon*, Sieb. & Zucc., **26**, 154.  
*Hancockianum*, Oliver, **36**, 491.  
*japonicum*, A. Gray, **26**, 155.  
*officinale*, Linn., **26**, 154; mentioned, 155.  
*Zollingeri*, A. DC., **26**, 155.
- Litsæa*, see *Litsea*.
- Litsea*, Lam., **26**, 370.  
*ceylanica*, Benth., **26**, 384.  
 — var. *chinensis*, Benth., **26**, 384.  
*chinensis*, Blume, **26**, 379.  
*chinensis*, Heyne, **26**, 379.  
*chinensis*, Lam., **26**, 379, 385.  
*citrata*, Blume, **26**, 379.  
*citriifolia*, Juss., **26**, 379.  
*confertifolia*, Hemsl. \*, **26**, 379; mentioned, 381.  
*Cubeba*, Pers., **26**, 380.  
*cupularis*, Hemsl. \*, **26**, 380.  
*elongata*, [Benth. &] Hook. f., **26**, 381.  
*Faberi*, Hemsl. \*, **26**, 381.  
*glauca*, Sieb., **26**, 381.  
*gracilipes*, Hemsl. \*, **26**, 381.  
*hupehana*, Hemsl. \*, **26**, 382.  
*japonica*, Juss., **26**, 382.  
*lancifolia*, Villar, **26**, 382.  
*laxiflora*, Hemsl. \*, **26**, 383.  
*ligustrina*, Hook. f., **26**, 379.  
*Meissneri*, Hook. f., mentioned, **26**, 332.  
*mollis*, Hemsl. \*, **26**, 383.  
*monopetala*, Pers., **26**, 384.  
*piperita*, Juss., **26**, 380, 385.  
*Playfairi*, Hemsl. \*, **26**, 384.  
*polyantha*, Juss., **26**, 384.  
*pulchella*, Meissn., **26**, 384.  
*pungens*, Hemsl. \*, **26**, 384; mentioned, 380, 385.  
*rotundifolia*, Hemsl. \*, **26**, 385.  
*sebifera*, Pers., **26**, 385; mentioned, 379.  
*sericea*, Hook. f., **26**, 386.

- Litsea variabilis*, Hemsl. \*, 26, 386.  
*verticillata*, Hance, 26, 386.
- Livistona chinensis*, Mart., 36, 168.  
*sinensis*, Griff., 36, 168.
- Lloydia alpina*, Salisb., 36, 139.  
*Delavayi*, Franch., 36, 139.  
*filiformis*, Franch., 36, 139.  
*ixiolirioides*, Baker, 36, 139.  
*oxycarpa*, Franch., 36, 139.  
*serotina*, Sweet, 36, 139.  
*tibetica*, Baker, 36, 140.  
— var. *lutescens*, Franch., 36, 140.  
— var. *purpurascens*, Franch., 36, 140.  
*triflora*, Baker, 36, 140.  
*yunnanensis*, Franch., 36, 491.
- Lobelia affinis*, Wall., 26, 2; mentioned, 3.  
*begoniæfolia*, Wall., 26, 2.  
*campanuloides*, Thunb., 26, 3.  
*chinensis*, Lour., 26, 2.  
*Davidii*, Franch., mentioned, 26, 3.  
*dolichothyrsa*, Diels, 36, 491.  
*pyramidalis*, Wall., 26, 3.  
*radicans*, Thunb., 26, 3.  
*sessilifolia*, Lamb., 26, 4.  
*trialata*, Buch.-Ham., 26, 3.  
*trigona*, Benth., mentioned, 26, 2, 3.
- Lodhra caudata*, Miers, 26, 74.  
*crassifolia*, Miers, 26, 72.  
*cratægoides*, Decne., 26, 72.  
*ferruginea*, Miers, 26, 72.  
*microcarpa*, Miers, 26, 73.  
*spicata*, Miers, 26, 75.
- LOGANIACEÆ, 26, 117.
- Lonicera acuminata*, Wall., 36, 491.  
*adenophora*, Franch., 36, 491.  
*æmulans*, Rehder, 36, 491.  
*affinis*, Hook. & Arn., 23, 359.  
— var. *pubescens*, Maxim., 23, 359.  
*alseuosmoides*, Graebn., 36, 491.  
*barbinervis*, Komarov, 36, 491.  
*Bournei*, Hemsl. \*, 23, 360.  
*brachypoda*, DC., 23, 364.  
— var. *foliis aureo-reticulatis*, Ill. Hort., 23, 364.  
— var. *reticulata*, Belg. Hort., 23, 364.  
*cærulea*, Linn., 23, 360; mentioned, 368.  
*calcarata*, Hemsl., 36, 491.  
*Caprifolium*, Linn., mentioned, 23, 367.  
*chinensis*, Wats., 23, 364.  
*chrysantha*, Turcz., 26, 360.
- Lonicera chrysantha*, var. *longipes*, Maxim., 23, 360.  
*confusa*, DC., 23, 361.  
*confusa*, Miq., 23, 364.  
*crassifolia*, Batalin, 36, 491.  
*cyanocarpa*, Franch., 36, 491.  
*decipiens*, Hook. f. & Thoms., 23, 361.  
*deflexicalyx*, Batalin, 36, 491.  
*Delavayi*, Franch., 36, 491.  
*diversifolia*, Wall., 23, 365.  
*Elisæ*, Franch., 23, 361.  
*Fargesii*, Franch., 36, 491.  
*Ferdinandi*, Franch., 23, 361.  
*ferruginea*, Rehder, 36, 491.  
*flexuosa*, Thunb., 23, 364.  
*fragrans*, Carr., mentioned, 23, 362.  
*fragrantissima*, Lindl. & Paxt., 23, 361; mentioned, 362.  
*fuchsioides*, Hemsl. \*, 23, 362; mentioned, 492.  
*Giraldii*, Rehder, 36, 491.  
*glabrata*, Wall., mentioned, 23, 363 [sphalm. 359].  
*gracilipes*, Miq., 36, 491.  
*gynochlamydea*, Hemsl. \*, 23, 362.  
*Hemsleyana*, Rehder, 36, 491.  
*Henryi*, Hemsl. \*, 23, 363 [sphalm. 359].  
*heteroloba*, Batalin, 36, 491.  
*Hildebrandiana*, Rehder, 36, 491.  
*hirtiflora*, Champ., 23, 365.  
*hispida*, Pall., 23, 363 [sphalm. 359].  
*hypoglauca*, Miq., 23, 360.  
*inconspicua*, Batalin, 36, 491.  
*Infundibulum*, Franch., 36, 491.  
*japonica*, Andr., 23, 361.  
*japonica*, Hook. f. & Thoms., 23, 365.  
*japonica*, Thunb., 23, 364.  
— var. *chinensis*, Baker, 23, 364.  
*Koebneana*, Rehder, 36, 491.  
*leiantha*, Kurz, 23, 364.  
*Leschenaultii*, Miq., 23, 360.  
*ligustrina*, Wall., mentioned, 23, 363 [sphalm. 359].  
— var., 36, 491.  
*longa*, Rehder, 36, 491.  
*longiflora*, DC., 23, 364.  
*Maackii*, Maxim., 23, 364.  
*macrantha*, DC., 23, 365; mentioned, 368.  
*Maximowiczii*, Maxim., mentioned, 36, 368.  
*microphylla*, Willd., 23, 365.  
*minuta*, Batalin, 36, 491.  
*mollissima*, Blume, 23, 359.

- Lonicera Morrowii*, A. Gray, **23**, 360.  
*mucronata*, *Rehder*, **36**, 491.  
*multiflora*, *Champ.*, **23**, 361.  
*nervosa*, *Maxim.*, **23**, 365.  
*odoratissima*, *Lindl.*, **23**, 361.  
*orientalis*, *Lam.*, **36**, 491.  
*ovalis*, *Batalin*, **36**, 491.  
*pekinensis*, *Rehder*, **36**, 491.  
*Periclymenum*, *Lour.*, **23**, 361.  
*phyllocarpa*, *Maxim.*, **23**, 365.  
*pileata*, *Oliver*, **23**, 365; mentioned, 363 [sphalm. 359].  
*præflorens*, *Batalin*, **36**, 491.  
*quinquelocularis*, *Hardw.*, **23**, 365.  
*reticulata*, *Champ.*, **23**, 366.  
*retusa*, *Franch.*, **36**, 491.  
*rhododendroides*, *Graebn.*, **36**, 491.  
*Ruprechtiana*, *Regel*, **23**, 366.  
*saccata*, *Rehder*, **36**, 491.  
*scabrida*, *Franch.*, **36**, 491.  
*serpyllifolia*, *Rehder*, **36**, 491.  
*setchuenensis*, *Rehder*, **36**, 492.  
*setifera*, *Franch.*, **36**, 492.  
*similis*, *Hemsl.* \*, **23**, 366.  
*spp. nn.*, **23**, 368.  
*Standishii*, *Hook. f.*, **23**, 367.  
*stenosiphon*, *Franch.*, **36**, 492.  
*stephanocarpa*, *Franch.*, **36**, 492.  
*subæqualis*, *Rehder*, **36**, 492.  
*syringantha*, *Maxim.*, **23**, 367.  
*szechuanica*, *Batalin*, **36**, 492.  
*tangutica*, *Maxim.*, **23**, 367.  
*Tatarinowii*, *Maxim.*, **23**, 367.  
*tatsienensis*, *Franch.*, **36**, 492.  
*Telfairii*, *Hook. & Arn.*, **23**, 361.  
*thibetica*, *Bur. & Franch.*, **36**, 492.  
*tragophylla*, *Hemsl.* \*, **23**, 367.  
*trichopoda*, *Franch.*, **36**, 492.  
*trichosantha*, *Bur. & Franch.*, **36**, 492.  
*vegeta*, *Rehder*, **36**, 492.  
*vesicaria*, *Komarov*, **36**, 492.  
*Webbiana*, *Sargent*, **36**, 492.  
*Xylosteum*, *Regel*, **23**, 360.  
*yunnanensis*, *Franch.*, **36**, 492.  
*Lophanthus chinensis*, *Benth.*, **26**, 288.  
*rugosus*, *Fisch. & Mey.*, **26**, 288.  
*Lophatherum elatum*, *Zoll.*, **36**, 420.  
*gracile*, *Brongn.*, **36**, 420; mentioned, 221.  
*japonicum*, *Steud.*, **36**, 420.  
*Lehmanni*, *Nees*, **36**, 420.  
*sinense*, *Rendle* \*, **36**, 421.  
*Lophiocarpus guyanensis*, *Micheli*, **36**, 190.  
*Lophocarpus guyanensis*, *Durand*, **36**, 189.  
 LORANTHACEÆ, **26**, 405.  
*Loranthus ampullaceus*, *Roxb.*, **26**, 405.  
*bibracteolatus*, *Hance*, **26**, 405.  
*chinensis*, *DC.*, **26**, 405.  
*cuneatus*, *Heyne*, mentioned, **26**, 407.  
*Delavayi*, *Engl.*, **36**, 492.  
*Delavayi*, *Van Tiegh.*, **36**, 492.  
*estipitatus*, *Stapf*, **26**, 405.  
*europæus*, *Linn.*, **26**, 406.  
*Fordii*, *Hance*, **26**, 406.  
*Hemsleyanus*, *King*, **26**, 406.  
*Jodoniki*, *Sieb. & Zucc.*, **26**, 407.  
*nigraus*, *Hance*, **26**, 406.  
*notothixiodes*, *Hance*, **26**, 406.  
*odoratus*, *Wall.*, **26**, 406.  
*pentapetalus*, *Roxb.*, **26**, 406.  
*Sampsoni*, *Hance*, **26**, 406.  
*Scurula*, *Linn.*, **26**, 407; mentioned, 405.  
 — var. *obtecta*, *Kurz*, **26**, 405.  
*Yadoriki*, *Sieb. & Zucc.*, **26**, 407; mentioned, 406.  
*Loropetalum chinense*, *R. Br.*, **23**, 290.  
*subcordatum*, *Oliver*, **26**, 291.  
*Lotus corniculatus*, *Linn.*, **23**, 155.  
*Lourea obcordata*, *Desv.*, **23**, 178.  
*reniformis*, *DC.*, **23**, 178.  
*Vespertilionis*, *Desv.*, **23**, 178.  
*Loxocalyx*, *Hemsl.* \*, **26**, 308.  
*urticifolius*, *Hemsl.*, **26**, 309.  
*Loxostemon Delavayi*, *Franch.*, **36**, 492.  
*Lubinia spathulata*, *Vent.*, **26**, 53.  
*Luculia Pinceana*, *Hook.*, **23**, 371.  
*Ludisia discolor*, *A. Rich.*, **36**, 44.  
*Furetii*, *Blume*, **36**, 44.  
*odorata*, *Blume*, **36**, 44.  
*Ludwigia diffusa*, *Buch.-Ham.*, **23**, 309.  
*parviflora*, *Roxb.*, **23**, 309.  
*prostrata*, *Roxb.*, **23**, 309.  
*Luffa acutangula*, *Roxb.*, **23**, 314.  
*egyptiaca*, *Mill.*, **23**, 315.  
*cylindrica*, *Roem.*, **23**, 315.  
*Luisia bicordata*, *Thw.*, **36**, 34.  
*Hancockii*, *Rolfe*, **36**, 33.  
*Morsei*, *Rolfe* \*, **36**, 33.  
*teres*, *Blume*, mentioned, **36**, 37, 38.  
*teres*, *Maxim.*, **36**, 33.  
*teretifolia*, *Gaudich.*, **36**, 33.  
*spp.*, **36**, 34; mentioned, 33, 34, 38.  
*Lupinus cochinchinensis*, *Lour.*, **23**, 152.

- Luzula campestris*, DC., 36, 160.  
 — var. *capitata*, Buchen., 36, 161.  
 — var. *congesta*, Buchen., 36, 160.  
*chinensis*, N. E. Br. \*, 36, 161.  
*effusa*, Buchen., 36, 161.  
*gigantea*, Desv., mentioned, 36, 161.  
*multiflora*, Lej., 36, 160.  
*multiflora*, Syme, 36, 161.  
*pilosa*, Willd., 36, 162.  
*pilosa*, Willd., var. *plumosa*, Franch., 36, 161.  
*plumosa*, E. Mey., 36, 161.  
*vernalis*, DC., 36, 162.
- Lychnis apetala*, Linn., 23, 65.  
*Bungeana*, Fisch., 23, 65.  
*cognata*, Maxim., 23, 66.  
*coronata*, Thunb., 23, 66.  
*Davidi*, Franch., 36, 492.  
*fulgens*, Fisch., 23, 66.  
*glandulosa*, Maxim., 36, 492.  
*grandiflora*, Jacq., 23, 66.  
*Haageana* ×, Lemaire, mentioned, 23, 66.  
*laciniata*, Maxim., 36, 492.  
*Senno*, Sieb. & Zucc., 23, 66.  
*Sieboldi*, Van Houtte, 23, 66.  
*yunnanensis*, Baker, 36, 492.
- Lycimnia suaveolens*, Hance, 26, 94.
- Lycium barbarum*, Linn., 26, 175.  
 — var. *chinense*, Ait., 26, 175.  
*chinense*, Mill., 26, 175.  
*megistocarpum*, Dun., 26, 175.  
*ovatum*, Loisel., 26, 176.  
*Trewianum*, G. Don, 26, 175.  
*turbinatum*, Loisel., 26, 175.
- Lycopersicum esculentum*, Mill., mentioned, 26, 169.
- Lycopsis picta*, Bunge, 26, 155.
- Lycopus europæus*, Linn., 26, 282.  
*lucidus*, Turcz., 26, 282.
- Lycoris aurea*, Herb., 36, 89.  
*radiata*, Herb., 36, 89.  
*sanguinea*, Maxim., 36, 89.  
*squamigera*, Maxim., 36, 89.  
*straminea*, Lindl., 36, 90.
- Lysidice rhodostegia*, Hance, 23, 213.
- Lysimachia acroadenia*, Maxim., 26, 51.  
*albescens*, Franch., 36, 492.  
*Alfredi*, Hance, 26, 47.  
*alpestris*, Champ., 26, 47; mentioned, 51.  
*alternifolia*, Wall., mentioned, 26, 49.  
*anagalloides*, Sibth. & Sm., mentioned, 26, 49.
- Lysimachia auriculata*, Hemsl. \*, 26, 47; mentioned, 55.  
*barystachys*, Bunge, 26, 47.  
*barystachys*, Klatt, 26, 52.  
*brachystachys*, Carr., 26, 48.  
*candida*, Lindl., 26, 48.  
*capillipes*, Hemsl. \*, 26, 48.  
*Christinæ*, Hance, 26, 49; mentioned, 50.  
*circæoides*, Hemsl. \*, 26, 49.  
*clethroides*, Duby, 26, 49.  
*congestiflora*, Hemsl. \*, 26, 50.  
*consobrina*, Hance, 26, 51.  
*crispidens*, Hemsl. \*, 26, 50.  
*cuspidata*, Klatt, 26, 53.  
*davurica*, Ledeb., 26, 51.  
*decurrens*, G. Forst., 26, 51.  
*Delavayi*, Franch., 36, 492.  
*deltoides*, Franch., 36, 492.  
*drynarifolia*, Franch., 36, 492.  
*evalvis*, Wall., mentioned, 26, 52.  
*Fargesii*, Franch., 36, 492.  
*ferruginea*, Edgew., 26, 53.  
*Fœnumgræcum*, Hance, 26, 51; mentioned, 57.  
*Fordiana*, Oliver, 36, 492.  
*Fortunei*, Maxim., 26, 52.  
*glaucescens*, Wall., 26, 48.  
*glaucina*, Franch., 36, 492.  
*grammica*, Hance, 26, 52.  
*Hemsleyana*, Oliver, 36, 492.  
*Hemsleyi*, Franch., 36, 492.  
*Henryi*, Hemsl. \*, 26, 52.  
*heterogenea*, Klatt, 26, 52; mentioned, 47.  
*hypericoides*, Hemsl., 36, 492.  
*inconspicua*, Miq., 26, 53.  
*insignis*, Hemsl., 36, 492.  
*involuta*, Hemsl., 36, 492.  
*japonica*, Thunb., 26, 53; mentioned, 50.  
*javanica*, Blume, 26, 51.  
*Klattiana*, Hance, 26, 53; mentioned, 52.  
*Leschenaultii*, Duby, 26, 57.  
*lineariloba*, Hook. & Arn., 26, 53.  
*lobelioides*, Wall., mentioned, 26, 49.  
*longipes*, Hemsl., 36, 492.  
*lubinioides*, Sieb. & Zucc., 26, 53.  
*mauritanica*, Lam., 26, 53.  
*multandra*, Franch., 36, 492.  
*multiflora*, Wall., 26, 51.  
*Nebeliana*, Gilg, 36, 492.  
*nemorum*, Linn., 36, 492.  
*nigrolineata*, Hemsl., 36, 493.  
*obovata*, Buch.-Ham., 26, 48.  
*omeiensis*, Hemsl., 36, 493.  
*ophelioides*, Hemsl. \*, 26, 54.

- ysimachia paludicola*, Hemsl. \*, 26, 54.  
*paridiformis*, Franch., 26, 55.  
 — var. *elliptica*, Franch., 26, 55.  
 — var. *stenophylla*, Franch., 26, 55.  
*parvifolia*, Franch. \*, 26, 55.  
*pentapetala*, Bunge, 26, 55.  
*platypetala*, Franch., 36, 493.  
*prolifera*, Klatt, mentioned, 26, 55.  
*pterantha*, Hemsl. \*, 26, 56.  
*pumila*, Franch., 36, 493.  
*ramosa*, Wall., 36, 493.  
*rubiginosa*, Hemsl. \*, 26, 56.  
*samolina*, Hance, 26, 48.  
*sikokiana*, Miq., 26, 57.  
*simulans*, Hemsl. \*, 26, 57.  
*sinica*, Miq., 26, 57.  
*spathulata*, Klatt, 26, 53.  
*stenosepala*, Hemsl. \*, 26, 57.  
*thyrsiflora*, Linn., 26, 58.  
*trichopoda*, Franch., 36, 493.  
*trientalioides*, Hemsl., 36, 493.  
*unguiculata*, Diels, 36, 493.  
*violascens*, Franch., 36, 493.  
*vulgaris*, Linn., 26, 58; mentioned, 51.  
*yunnanensis*, Franch., 36, 493.
- Lysionotus carnosus*, Hemsl., 36, 493.  
*heterophylla*, Franch., 36, 493.  
*involverata*, Franch., 36, 493.  
*ophiorrhizoides*, Hemsl. \*, 26, 224.  
*pauciflorus*, Maxim., 26, 224.  
*serratus*, D. Don, mentioned, 26, 225.
- LYTHRACEÆ**, 23, 302.
- Lythrum fruticosum*, Linn., 23, 304.  
*Salicaria*, Linn., 23, 304.  
*virgatum*, Linn., 23, 305.
- Maba Ebenus*, Spreng., 26, 71.  
*elliptica*, Forst., 26, 71.
- Macaranga Henricorum*, Hemsl. \*, 26, 442.  
*indica*, Wight, 26, 443.  
*Sampsoni*, Hance, 26, 443.  
*sinensis*, Muell. Arg., 26, 443.  
*Tanarius*, Muell. Arg., 26, 443.
- Machilus*, genus mentioned, 26, 370.  
*Bournei*, Hemsl. \*, 26, 373.  
*breviflora*, Hemsl. \*, 26, 374.  
*chinensis*, Hemsl. \*, 26, 374.  
*Faberi*, Hemsl. \*, 26, 374.  
*Grijsii*, Hance, 26, 375.  
*Henryi*, Hemsl. \*, 26, 375.
- Machilus japonica*, Sieb. & Zucc., mentioned, 26, 378.  
*longifolia*, Blume, 26, 375.  
*macrophylla*, Hemsl. \*, 26, 375.  
*microcarpa*, Hemsl. \*, 26, 376.  
*Nanmu*, Hemsl., 26, 376; mentioned, 374, 376.  
*neurantha*, Hemsl. \*, 26, 376; mentioned, 377.  
*oreophila*, Hance, 26, 377; mentioned, 376.  
*platyphylla*, Diels, 36, 493.  
*rimosa*, Blume, 26, 378.  
*Sheareri*, Hemsl. \*, 26, 377.  
*Thunbergii*, Sieb. & Zucc., 26, 377; mentioned, 376, 378; 36, 493.  
 — var. *glaucescens*, Blume, 26, 378.  
*velutina*, Champ., 26, 378.  
*villosus*, Hook. f., mentioned, 26, 375.
- Machlis hemisphærica*, DC., 23, 439.  
*Maclura tricuspidata*, Carr., 26, 470.  
*Macodes argyroneura*, Rolfe, 36, 44.  
*Macrotropis emarginata*, Walp., 23, 204.  
*fatida*, DC., 23, 204.
- Maddenia himalaica*, Hook., 36, 493.
- Maesa castaneifolia*, Mez, 36, 493.  
*coriacea*, Champ., 26, 60.  
*depauperata*, Diels, 36, 493.  
*Doræna*, Blume, 26, 59.  
*formosana*, Mez, 36, 493.  
*indica*, Benth., 26, 60.  
*indica*, Wall., mentioned, 26, 60.  
*montana*, Benth., 26, 60.  
*sinensis*, A. DC., 26, 60.  
*sp. n.*, 26, 60.  
*tenera*, Mez, 36, 493.
- Magnolia Championi*, Benth., 23, 24.  
*Coco*, DC., 23, 24.  
*conspicua*, Salisb., 23, 23.  
*Delavayi*, Franch., 36, 493.  
*denudata*, Lam., 23, 23.  
*discolor*, Vent., 23, 23.  
*fuscata*, Andr., 23, 24.  
*glauca*, Thunb., 23, 23.  
*Henryi*, Dunn, 36, 493.  
*hypoleuca*, Sieb. & Zucc., 36, 493.  
*inodora*, DC., 23, 24.  
*liliflora*, [Desr.], Lam., 23, 23.  
*obovata*, Thunb., 23, 23.  
*parviflora*, Blume, 23, 24.  
*parviflora*, Sieb. & Zucc., 36, 493.  
*Precia*, Correa, 23, 23.  
*pumila*, Andr., 23, 24.  
*purpurea*, Curt., 23, 23.  
*Yulan*, Desf., 23, 23.  
 — var. *Soulangiana*, [Lindl.] Bot. Reg., 23, 23.

- MAGNOLIACEÆ, 23, 23.  
*Mahonia*, Nutt. = *Berberis*, 36, 493.  
*eurybracteata*, Fedde, 36, 493.  
*Fortunei*, Fedde, 36, 493.  
*gracilipes*, Fedde, 36, 493.  
*japonica*, DC., 23, 31.  
— var., 36, 493.  
*napaulensis*, DC., 23, 31.  
*polyodonta*, Fedde, 36, 493.  
*subtriplinervis*, Franch., 36, 493.  
*trifurca*, Loudon, 36, 493.  
*Maianthemum bifolium*, DC. 36, 112.  
*Convallaria*, Wigg., 36, 112.  
*Malachium aquaticum*, Fries 23, 67.  
*Malachra Urena*, DC., 23, 86.  
*Malaisia acuminata*, Planch., 26, 454.  
*scandens*, Planch., 26, 454.  
*tortuosa*, Blanco, 26, 454.  
*viridescens*, Planch., 26, 454.  
*Malaxis iridifolia*, Reichb. f., 36, 5.  
*lancifolia*, Sm., 36, 7.  
*monophyllos*, Sw., 36, 5.  
*nervosa*, Sw., 36, 7.  
*odorata*, Willd., 36, 7.  
*Malcolmia africana*, R. Br., 23, 45.  
*Mallotus*, Lour., 26, 310.  
*albus*, Muell. Arg., 36, 493.  
*Apelta*, Muell. Arg., 26, 439.  
*barbatus*, Muell. Arg., 26, 439.  
*chinensis*, Muell. Arg., 26, 439.  
*cochinchinensis*, Lour., 26, 439.  
*contubernalis*, Hance, 26, 441.  
*Furetianus*, Muell. Arg., 26, 439.  
*Hookerianus*, Muell. Arg., 26, 440.  
*japonicus*, Muell. Arg., 26, 440.  
*moluccanus*, Muell. Arg., 26, 440.  
*paniculatus*, Muell. Arg., 26, 439.  
*philippinensis*, Muell. Arg., 26, 440.  
*Playfairii*, Hemsl. \*, 26, 441.  
*populifolius*, Hemsl. \*, 26, 441.  
*repandus*, Muell. Arg., 26, 441.  
*ricinoides*, Muell. Arg., 26, 442.  
*tenuifolius*, Pax, 36, 493.  
*tiliæfolius*, Muell. Arg., 26, 442.  
*Malouetia asiatica*, Sieb. & Zucc., 26, 99.  
MALPIGHIACEÆ, 23, 96.  
*Malus baccata*, Desf., 23, 255.  
*cerasifera*, Spach, 23, 255.  
*floribunda*, Sieb., 23, 258.  
*microcarpa*, var. *Torrino*, Carr., 23, 258, 490.  
*spectabilis*, Desf., 23, 258.  
*Malva chinensis*, Mill., 23, 84.  
*mauritiana*, Linn., 23, 84.  
*Malva parviflora*, Linn., 23, 83.  
*pulchella*, Bernh., 23, 84.  
*sinensis*, DC., 23, 84.  
*sylvestris*, Linn., 23, 84.  
*verticillata*, Linn., 23, 84.  
MALVACEÆ, 23, 83.  
*Malvastrum carpinifolium*, A. Gray, 23, 84, 85.  
*ruderale*, Hance, 23, 84.  
*tricuspidatum*, A. Gray, 23, 84.  
*Mandragora caulescens*, C. B. Clarke, 26, 175.  
*Mangifera indica*, Linn., 23, 148.  
*Manglietia Fordiana*, Oliver, 36, 493.  
*Manisuris granularis*, Linn., 36, 363.  
*Mappa sinensis*, Baill., 26, 443.  
*Tanaria*, Spreng., 26, 443.  
*tomentosa*, Blume, 26, 443.  
*Mappia pittosporoides*, 36, 493.  
*Maranta Galanga*, Sw., 36, 72.  
*parviflora*, A. Dietr., 36, 73.  
*Mariscus albescens*, Gaudich., 36, 220.  
*biglumis*, Gaertn., mentioned, 36, 220.  
*capitatus*, Steud., 36, 258.  
*cyperinus*, Presl, 36, 221.  
*cyperinus*, Vahl, 36, 220.  
*dilutus*, Nees, 36, 221.  
*ferax*, C. B. Clarke, 36, 222.  
*maritimus*, Miq., 36, 259.  
*microcephalus*, Presl, 36, 221.  
*philippensis*, Steud., 36, 221; mentioned, 220.  
*Pohlianus*, Nees, 36, 222.  
*pungens*, Miq., 36, 259.  
*Sieberianus*, Nees, 36, 221; mentioned, 220.  
*umbellatus*, Presl, 36, 221.  
*umbellatus*, Vahl, 36, 221.  
*Marquartia tomentosa*, Vogel, 23, 159.  
*Marrubium incisum*, Benth., 26, 299.  
*vulgare*, Linn., mentioned, 26, 284.  
*Marsdenia eriostoma*, Benth. MS., 26, 113.  
*lachnostoma*, Benth., 26, 113.  
*sinensis*, Hemsl. \*, 26, 113.  
*tinctoria*, R. Br., 26, 114.  
*tomentosa*, Morr. & Decne., mentioned, 26, 113.  
*urceolata*, Decne., 26, 114.  
*Martinia*, Vaniot, 36, 493.  
*polymorpha*, Vaniot, 36, 493.  
*Mastacanthus sinensis*, Endl., 26, 263.  
*Matricaria cantoniensis*, Lour., 23, 413.



- Matricaria inodora*, Linn., 23, 439.  
*monspeliensis*, Desv., 23, 438.  
*Maximowiczia chinensis*, Rupr., 23, 25.  
*Mazus caducifer*, Hance, 26, 181.  
*dentatus*, Wall., mentioned, 26, 182.  
*gracilis*, Hemsl. \*, 26, 181.  
*lanceifolius*, Hemsl. \*, 26, 181.  
*macranthus*, Diels, 36, 493.  
*pulchellus*, Hemsl. \*, 26, 182.  
*procumbens*, Hemsl. \*, 26, 182.  
*rugosus*, Lour., 26, 183.  
*stachydifolius*, Maxim., 26, 183.  
*vandellioides*, Hance, 26, 183.  
*villosus*, Hemsl., 26, 184.  
*Meconopsis betonicifolia*, Franch., 36, 494.  
*chelidonifolia*, Bur. & Franch., 36, 494.  
*Henrici*, Bur. & Franch., 36, 494.  
*horridula*, Hook. f. & Thoms., 36, 494.  
*integrifolia*, Franch., 36, 494.  
*punicea*, Maxim., 36, 494.  
*quintuplinervia*, Regel, 36, 494.  
*racemosa*, Maxim., 23, 34; 36, 494.  
*Medicago denticulata*, Willd., 23, 153.  
*falcata*, Linn., 23, 153.  
*lappacea*, Lam., mentioned, 23, 153.  
*lupulina*, Linn., 23, 154.  
*minima*, Batalin, 36, 494.  
*ruthenica*, Ledeb., 23, 153, 489.  
*sativa*, Linn., mentioned, 23, 154.  
*Medicia elegans*, Gardn., 26, 117.  
*Megabotrya meliæfolia*, Hance, 23, 104.  
*Megacarpæa Delavayi*, Franch., 36, 494.  
*Megacodon* (§ *Gentianæ*), Hemsl. \*, 26, 137.  
*Megadenia*, Maxim., 36, 494.  
*pygmæa*, Maxim., 36, 494.  
*Melaleuca Leucadendron*, Linn., 23, 295.  
*Melampyrum ciliare*, Miq., 26, 220.  
*jedoense*, Miq., 26, 220.  
*pratense*, Hemsl., 26, 220.  
*roseum*, Maxim., 26, 220.  
*Melandryum apicum*, Rohrb., 23, 65.  
*cæspitosum*, Williams, 36, 494.  
*firmum*, Rohrb., 23, 65.  
*glandulosum*, Williams, 36, 494.  
*kialense*, Williams, 36, 494.  
*Oldhamianum*, Rohrb., 23, 65.  
*platypetalum*, Williams, 36, 494.  
*Melandryum Soulici*, Williams, 36, 494.  
*Melanocarpa alata*, Turcz., 23, 121.  
*Melanolepis multiglandulosus*, Reichb. & Zoll., 26, 440.  
*Melanosciadum*, Boissieu, 36, 494.  
*pimpinelloideum*, Boissieu, 36, 494.  
*Melanthesa cernua*, Benth., 26, 427.  
*chinensis*, Blume, 26, 427.  
*glaucescens*, Miq., 26, 427.  
*rectinata*, Muell. Arg., 26, 427.  
*rhamnoides*, Wight, 26, 428.  
*Melanthesopsis fruticosa*, Muell. Arg., 26, 427.  
*patens*, Muell. Arg., 26, 428.  
*Melanthium cochinchinense*, Lour., 36, 102, 146.  
*nigrum*, Thunb., 36, 148.  
*sibiricum*, Linn., 36, 148.  
*viride*, Thunb., 36, 147.  
*Melastoma calycinum*, Benth., 23, 299.  
*candidum*, D. Don, 23, 299.  
*decemfidum*, Roxb., 23, 300.  
*dodecandrum*, Lour., 23, 300.  
*macrocarpum*, D. Don, 23, 299.  
*malabathricum*, Linn., 23, 300.  
*malabathricum*, [Sims], 23, 299.  
*Nobotan*, Blume, 23, 299.  
*repens*, Desr., 23, 300.  
*sanguineum*, Sims, 23, 300.  
**MELASTOMACEÆ**, 23, 298.  
*Melia Azedarach*, Linn., 23, 113.  
*composita*, Willd., 23, 113.  
*dubia*, Cav., 23, 113.  
*japonica*, G. Don, 23, 113.  
*Toosendan*, Sieb. & Zucc., 36, 494.  
**MELIACEÆ**, 23, 113.  
*Melica Gmelini*, Turcz., 36, 418.  
*nutans*, Linn., 36, 418.  
*Oncei*, Franch., 36, 419.  
*Radula*, Franch., 36, 419.  
*scabra*, Linn., mentioned, 36, 419.  
*scabrosa*, Trin., 36, 419.  
*Melidora pellucida*, Noronha, 26, 18.  
*Melilotus alba*, Desr., 23, 154.  
*cærulea*, Desr., 23, 154.  
*dentata*, Willd., 23, 154.  
*graveolens*, Bunge, 23, 155.  
*leucantha*, Koch, 23, 154.  
*macrorhiza*, Pers., 23, 154.  
*officinalis*, Desr., 23, 154.  
*palustris*, Kit., 23, 155.  
*parviflora*, Desf., 23, 155.  
*suaveolens*, Ledeb., 23, 155.  
*Meliosma cuneifolia*, Franch., 36, 494.  
*dillenifolia*, Hook. f., 23, 145.  
*Fordii*, Hemsl. \*, 23, 144.

- Meliosma Henryi*, Diels, **36**, 494.  
*myriantha*, Sieb. & Zucc., **23**, 145.  
*Oldhami*, Maxim., **23**, 145.  
*patens*, Hemsl. \*, **23**, 145.  
*pungens*, Hook. f., **23**, 145.  
*rigida*, Sieb. & Zucc., **23**, 145.  
*rhoifolia*, Maxim., **23**, 146.  
*simplicifolia*, Roxb., **23**, 145.  
*squamulata*, Hance, **23**, 146.  
*Wallichii*, Planch., **23**, 146.  
*yunnanensis*, Franch., **36**, 494.  
*Melissa cretica*, Lour., **26**, 279.  
*umbrosa*, Bieb., **26**, 281.  
*Melochia concatenata*, Linn., **23**, 91.  
*corchorifolia*, Linn., **23**, 91.  
*Melodinus Cumingianus*, A. DC.,  
 mentioned, **26**, 94.  
*fusiformis*, Champ., **26**, 93; men-  
 tioned, 94.  
*Hemsleyanus*, Diels, **36**, 494.  
*lætus*, Champ., **26**, 94.  
*monogynus*, Roxb., **26**, 94.  
*sp. n.*, **26**, 94.  
*suaveolens*, Champ., **26**, 94.  
*Melodorum glaucescens*, Hance, **23**,  
 27.  
*Oldhami*, Hemsl. \*, **23**, 27.  
*verrucosum*, Hook. f. & Thoms., **23**,  
 28.  
*Melothria Baueriana*, F. Muell., **23**,  
 318.  
*cucumerina*, Naud., **23**, 319.  
*indica*, Lour., **23**, 319.  
*leucocarpa*, Cogn., **23**, 319.  
*maderaspata*, Cogn., **23**, 319.  
*mucronata*, Cogn., **23**, 318.  
*odorata*, Hook. f. & Thoms., **23**,  
 319.  
*pendula*, Linn., mentioned, **23**,  
 319.  
*Memecylon ligustrifolium*, Champ.,  
**23**, 302.  
*nigrescens*, Hook. & Arn., **23**, 302.  
*scutellatum*, Hook. & Arn., **23**,  
 302.  
*Memorialis hirta*, Wedd., **26**, 489.  
*hispida*, Buch.-Ham., **26**, 489.  
*pentandra*, var. *hypericifolia*, Wedd.,  
**26**, 490.  
*quinquenervis*, Buch.-Ham., **26**,  
 489.  
**MENISPERMACEÆ**, **23**, 28.  
*Menispermum dauricum*, DC., **23**,  
 29.  
*Mentha aquatica*, Linn., mentioned,  
**26**, 281.  
*arvensis*, Linn., **26**, 281.  
*canadensis*, Linn., mentioned, **26**,  
 281.  
*crispa*, Linn., **26**, 281.  
*Mentha Pulegium*, Linn., mentioned,  
**26**, 281.  
*reticulosa*, Hance, **26**, 279.  
*stellata*, Lour., **26**, 277.  
*sylvestris*, Linn., mentioned, **26**,  
 281.  
*Meoschium lodiculare*, Nees, **36**, 364  
*Meyerianum*, Nees, **36**, 364.  
*rugosum*, Nees, **36**, 366.  
*Mephitidia chinensis*, Champ., **23**,  
 388.  
*cyanocarpa*, DC., **23**, 388.  
*Wallichii*, Wight & Arn., **23**, 389.  
*Mercurialis leiocarpa*, Sieb. & Zucc.,  
**26**, 436.  
*Mesona chinensis*, Benth., **26**, 267.  
*procumbens*, Hemsl., **36**, 494.  
*prunelloides*, Hemsl. \*, **26**, 267.  
*Mespilus chinensis*, Blume, mentioned,  
**23**, 259.  
*Cotoneaster*, Linn., **23**, 260.  
*glabra*, Colla, **23**, 262.  
*japonica*, Thunb., **23**, 261.  
*pinnatifida*, C. Koch, **23**, 259.  
*Pyracantha*, Linn., **23**, 260.  
*Sieboldii*, Blume, **23**, 265.  
*sinensis*, Poir., **23**, 264.  
*spiralis*, Blume, mentioned, **23**,  
 259, 264.  
*Metaplexis chinensis*, Decne., **26**,  
 110.  
*Hemsleyana*, Oliver, **36**, 494.  
*rostrata*, Turcz., **26**, 111.  
*Stauntoni*, Roem. & Schult., **26**,  
 110.  
*Mezoneurum brachycarpum*, Benth.,  
 mentioned, **23**, 205.  
*cucullatum*, Wight & Arn., men-  
 tioned, **23**, 205.  
*sinense*, Hemsl. \*, **23**, 204.  
 — var. *parvifolium*, Hemsl. \*, **23**,  
 205.  
*Michelia Champaca*, Linn., **23**, 24.  
*fuscata*, Blume, **23**, 24.  
*Microcala filiformis*, Hoffmgg. & Link,  
 mentioned, **26**, 130.  
*Microcarpæa muscosa*, R. Br., **26**,  
 192.  
*Microchloa setacea*, R. Br., **36**, 402.  
*Microdesmis caseariæfolia*, Planch.,  
**26**, 433.  
*Microglossa albescens*, C. B. Clarke,  
**23**, 419.  
*salicifolia*, Diels, **36**, 494.  
*volubilis*, DC., **23**, 419; mentioned,  
 420.  
*Micromeles alnifolia*, Koehne, **36**,  
 494.  
*Microptelea parvifolia*, Spach, **26**,  
 448.

- Microrhamnus franguloides*, Maxim., 23, 127.  
*Microrhynchus sarmentosus*, DC., 23, 488.  
*Microstachys Chamælea*, A. Juss., 26, 446.  
*Microstylis congesta*, Reichb. f., 36, 5.  
     *cylindrostachya*, Reichb. f., 36, 5.  
     *monophyllos*, Lindl., 36, 5.  
*Microtinus odoratissimus*, Oerst., 23, 354.  
*Microtis parviflora*, R. Br., 36, 46.  
*Microtæna cymosa*, Prain, 26, 306; mentioned, 308.  
     *Delavayi*, Prain, 36, 494.  
     *moupinensis*, Franch., 36, 494.  
     *Prainiana*, Diels, 36, 494.  
     *robusta*, Hemsl. \*, 26, 307.  
     *urticifolia*, Hemsl. \*, 26, 308.  
*Microula tangutica*, Maxim., 26, 151.  
*Miegia maritima*, Willd., 36, 258.  
*Mikania scandens*, Willd., 23, 405.  
     *volubilis*, Willd., 23, 405.  
*Milium cimicinoides*, Roxb., 36, 342.  
     *effusum*, Linn., 36, 383.  
*Millettia Bockii*, Harms, 36, 494.  
     *Championi*, Benth., 23, 159.  
     *cinerea*, Thunb., mentioned, 23, 159.  
     *cognata*, Hance, 23, 159.  
     *Dielsiana*, Harms, 36, 494.  
     *nitida*, Benth., 23, 159.  
     *pachycarpa*, Benth., 36, 494.  
     *purpurea*, Yatabe, 36, 494.  
     *reticulata*, Benth., 23, 159.  
     *scabricaulis*, Franch., 36, 494.  
     *sericosema*, Hance, 23, 159.  
     *speciosa*, Champ., 23, 159.  
     *spp. ?*, 23, 159.  
*Millingtonia pungens*, Wall., 23, 159.  
*Mimosa arborea*, Thunb., 23, 216.  
     *corniculata*, Lour., 23, 202, 216.  
     *sepiaria*, Benth., 23, 214.  
     *speciosa*, Jacq., 23, 216.  
**MIMOSÆ**, 23, 214.  
*Mimulus nepalensis*, Benth., 26, 181.  
     — *var. japonica*, Miq., 26, 181.  
     *tenellus*, Bunge, 26, 181.  
*Miquelia barbulata*, Nees, 36, 387.  
*Mirabilis Jalapa*, Linn., mentioned, 26, 317.  
*Miscanthus japonicus*, Anderss., 36, 347.  
     *nepalensis*, Hack., 36, 347.  
     *purpurascens*, Anderss., 36, 348.  
     *sacchariflorus*, Benth. & Hook. f., 36, 347.  
     *sinensis*, Anderss., 36, 348; mentioned, 347.  
*Miscanthus sinensis*, *var. purpurascens*, Rendle \*, 36, 348.  
*Mischospora efoliata*, Boeck., 36, 246.  
*Missiessya parvifolia*, Wedd., 26, 492.  
*Mitella nuda*, Linn., 36, 494.  
*Mitostigma gracile*, Blume, 36, 53.  
*Mitrasacme alsinoides*, R. Br., 26, 117.  
     *capillaris*, Wall., 26, 118.  
     *chinensis*, Griseb., 26, 118.  
     *indica*, Wight, 26, 117.  
     *malaccensis*, Wight, 26, 118.  
     *nudicaulis*, Reinw., 26, 117.  
     *polymorpha*, R. Br., 26, 118.  
     — *var. grandiflora*, Hemsl. \*, 26, 118.  
*Mitreola pedicellata*, Benth., 26, 117.  
*Mitrosicyos lobatus*, Maxim., 23, 320.  
     *paniculatus*, Maxim., 23, 320.  
     *racemosus*, Maxim., 23, 320.  
*Mochringia lateriflora*, Fenzl, 23, 70.  
     *linearifolia*, Williams, 36, 494.  
*Molineria capitulata*, Herb., 36, 87.  
     *plicata*, Colla, 36, 87.  
     *recurvata*, Herb., 36, 87.  
*Molinia Fauriei*, Hack., 36, 412; 495.  
     *japonica*, Hack., 36, 412.  
     *serotina*, Mert. & Koch, 36, 411.  
     *squarrosa*, Trin., 36, 411.  
*Mollugo Spergula*, Linn., 23, 324.  
     *stricta*, Linn., 23, 324.  
     *triphylla*, Lour., 23, 324.  
*Momordica Charantia*, Linn., 23, 315.  
     *cochinchinensis*, Spreng., 23, 315.  
     *cylindrica*, Linn., 23, 315.  
     *dioica*, Roxb., 23, 315.  
     *heterophylla*, Cogn., 23, 318.  
     *mixta*, Roxb., 23, 315.  
     *sicyoides*, Ser., 23, 215.  
*Moneses grandiflora*, Salish., 26, 33.  
*Monochasma monantha*, Hemsl. \*, 26, 203.  
     *Savatieri*, Franch., 26, 203.  
     *Sheareri*, Maxim., 26, 203.  
*Monochoria dilatata*, Kunth, 36, 148.  
     *hastæfolia*, Presl, 36, 148.  
     *hastata*, Solms, 36, 148.  
     *Junghuhniana*, Hassk., 36, 150.  
     *Korsakowii*, Regel, 36, 149.  
     *linearis*, Miq., 36, 150.  
     *ovata*, Kunth, 36, 149.  
     *plantaginea*, Kunth, 36, 150.  
     *sagittata*, Kunth, 36, 148.  
     *vaginalis*, Presl, 36, 149.  
     — *var. Korsakowii*, Solms, 36, 140.  
     — *var. plantaginea*, Solms, 36, 150.  
*Monocystis abnormis*, Lindl., 36, 71.

- Monolophus cœnobialis*, Hance, **36**, 68.  
*elegans*, Wall., **36**, 68.  
*Monotropa uniflora*, Linn., **26**, 31.  
**MONOTROPEÆ**, **26**, 34.  
*Moquinia eriosematoides*, Walp., **23**, 429.  
*Moraea chinensis*, Thunb., **36**, 86.  
*Morella rubra*, Lour., **26**, 496.  
*Moricandia sonchifolia*, Hook. f., **23**, 47.  
— var. *homœophylla*, Hance, **23**, 47.  
*Morina betonicoides*, Benth., **23**, 399.  
*Delavayi*, Franch., **23**, 399.  
*Morinda citrifolia*, Linn., **23**, 386.  
*umbellata*, Linn., **23**, 286.  
*villosa*, Hook., **23**, 386.  
*Morisia Wallichii*, Nees, **36**, 260.  
*Morocarpus edulis*, Sieb. & Zucc., **26**, 492.  
*Morus alba*, Linn., **26**, 455; mentioned, 456.  
*cathayana*, Hemsl. \*, **26**, 456.  
*constantinopolitana*, Poir., **26**, 455.  
*indica*, Linn., **26**, 455, 456.  
*laevigata*, Wall., mentioned, **26**, 456.  
*nigra*, Linn., mentioned, **26**, 456.  
*Moschosma polystachyum*, Benth., **26**, 269.  
*Moseleya*, Hemsl., **36**, 495.  
*pinnata*, Hemsl., **36**, 495.  
*Mosla chinensis*, Maxim., **26**, 280.  
*dianthera*, Maxim., **26**, 280.  
*Fordii*, Maxim., **26**, 280; **36**, 495.  
*formosana*, Maxim., **26**, 280.  
*grosseserrata*, Maxim., **26**, 280; **36**, 495.  
*lanceolata*, Maxim., **26**, 281.  
*punctata*, Maxim., **26**, 281.  
*Mucuna Championi*, Benth., **23**, 189.  
*macrobotrys*, Hance, **23**, 190.  
*macrocarpa*, Wall., mentioned, **23**, 190.  
*sempervirens*, Hemsl. \*, **23**, 190.  
*Muehlenbeckia platyclada*, Meisn., mentioned, **26**, 358.  
*Muehlenbergia Huegelii*, Trin., **36**, 335.  
*japonica*, Steud., **36**, 385.  
*viridissima*, Nees, **36**, 385.  
*Mukia scabrella*, Arn., **23**, 319.  
*Mulgedium sibiricum*, Less., **23**, 484.  
*tataricum*, DC., **23**, 484.  
*Munchausia speciosa*, Linn., **23**, 306.  
*Munronia Delavayi*, Franch., **36**, 495.  
*sinica*, Diels, **36**, 495.  
*unifoliata*, Oliver, **36**, 495.  
*yunnanensis*, Franch., **36**, 495.  
*Muricia cochinchinensis*, Lour., **23**, 315.  
*Murraya exotica*, Linn., **23**, 109.  
*Musa Basjoo*, Sieb., **36**, 74.  
*Cavendishii*, Lamb., **36**, 74.  
*chinensis*, Sweet, **36**, 74.  
*coccinea*, Andr., **36**, 74.  
*corniculata*, Lour., **36**, 75.  
*lasiocarpa*, Franch., **36**, 74.  
*nana*, Lour., **36**, 75.  
*nepalensis*, Wall., alluded to, **36**, 75.  
*odorata*, Lour., **36**, 75.  
*sapientum*, Linn., **36**, 74.  
*seminifera*, Lour., **36**, 75.  
*sinensis*, Sagot, **36**, 74.  
*sp.*, **36**, 75.  
*superba*, Roxb., mentioned, **36**, 75.  
*Uranoscopos*, Lour., **36**, 74.  
*Wilsoni*, Tutch., **36**, 75.  
*Mussaenda chinensis*, Lour., **23**, 379.  
*erosa*, Champ., **23**, 378; mentioned, 379.  
*frondosa*, Linn., **23**, 379.  
*glabra*, Hook. & Arn., **23**, 379.  
*hirsutula*, Miq., **23**, 379.  
*pubescens*, Ait. f., **23**, 379.  
**MYOPORINÆ**, **26**, 250.  
*Myoporum bontioides*, A. Gray, **26**, 250.  
*chinense*, A. Gray, **26**, 250.  
*Myosotis chinensis*, A. DC., **26**, 153.  
*intermedia*, Link, **36**, 495.  
*peduncularis*, Roem. & Schult., **26**, 154.  
*peduncularis*, Trev., **26**, 153.  
*sylvatica*, Hoffm., **36**, 495.  
*Myosoton aquaticum*, Moench, **23**, 67.  
*Myriactis nepalensis*, Less., **36**, 495.  
*Myrica adenophora*, Hance, **26**, 496.  
*Nagi*, C. DC., **26**, 496.  
*nana*, Cheval., **26**, 496.  
*rubra*, Lour., **26**, 496.  
*rubra*, Sieb. & Zucc., **26**, 496.  
*sapida*, Wall., **26**, 496.  
**MYRICACEÆ**, **26**, 496.  
*Myricaria alopecuroides*, Schrenk, **23**, 347.  
*bracteata*, Royle, **23**, 347.  
*germanica*, Desv., **23**, 347.  
*pulcherrima*, Batalin, **36**, 495.  
*Myriogyne minuta*, Less., **23**, 440.  
*Myrioneuron Faberii*, Hemsl. \*, **23**, 380.  
*nutans*, Wall., mentioned, **23**, 380.  
*Myriophyllum spiratum*, Linn., **23**, 293.  
*verticillatum*, Linn., **23**, 293.

- Myriopsis dioica*, Bunge, 23, 472.  
*Maximowiczii*, Winkl., 36, 495.  
*uniflora*, Maxim., 23, 472.  
*Myrmecis chinensis*, Rolfe\*, 36, 44.  
*gracilis*, Franch. & Sav., 36, 44.  
*japonica*, Rolfe\*, 36, 44.  
*Myroxylon controversum*, Kuntze, 36, 495.  
*Myrrhis chærophyllodes*, Hance, 23, 330.  
*Myrsine africana*, Linn., 26, 60.  
*bifaria*, Wall., 26, 60.  
*buxifolia*, Hance, 26, 61.  
*capitellata*, Wall., 26, 61.  
*glabra*, Gaertn., 26, 60.  
*marginata*, Mez, 36, 495.  
*neriifolia*, Sieb. & Zucc., 26, 61.  
*Playfairii*, Hemsl. \*, 26, 61.  
*retusa*, Ait., 26, 60.  
*scabra*, Gaertn., 26, 60.  
*semiserrata*, Wall., 26, 61.  
*sp.*, 26, 62.  
 MYRSINEÆ, 26, 59.  
 MYRTACEÆ, 23, 295.  
*Myrtus acuminatissima*, Blume, 23, 296.  
*canescens*, Lour., 23, 295.  
*chinensis*, Lour., 26, 75.  
*tomentosa*, Ait., 23, 295.  
  
*Nabalus ochroleucus*, Maxim., 23, 486.  
*repens*, Ledeb., 23, 483.  
 NAIADACEÆ, 36, 192.  
*Najas graminea*, Delile, 36, 197.  
*major*, Allioni, 36, 198.  
*marina*, Linn., 36, 198.  
 — form.  $\alpha$ , Rendle, 36, 198.  
 — —  $\beta$ , Rendle, 36, 198.  
*minor*, Benth., 36, 197.  
*Nandina domestica*, Thunb., 23, 32.  
*Nannoglottis carpesioides*, Maxim., 23, 430.  
*Nanocnide japonica*, Blume, 26, 473.  
*lobata*, Wedd., 26, 473.  
*Naravelia pitulifera* Hance, 23, 8.  
*Narcissus Tazetta*, var. *chinensis*, M. Roem., 36, 87.  
*Nardostachys chinensis*, Batalin, 36, 495.  
*Jatamansi*, DC., 23, 398.  
*Nardurus filiformis*, var. *chinensis*, Franch., 36, 404.  
*Nardus ciliaris*, Linn., 36, 363.  
*indica*, Linn., 36, 402.  
*Nasturtium amphibium*, R. Br., 23, 39.  
*atrovirens*, DC., 23, 40.  
*austriacum*, Crantz, 23, 39.  
*Nasturtium barbaræfolium*, Franch., 36, 495.  
*benghalense*, Hance, 23, 40.  
*Camelina*, Fisch., 23, 39.  
*cantonense*, Hance, 23, 39.  
*globosum*, Turcz., 23, 39.  
*Henryi*, Oliver, 36, 495.  
*indicum*, DC., 23, 40.  
*microspermum*, DC., 23, 40.  
*montanum*, Wall., 23, 40.  
*officinale*, R. Br., 23, 40.  
*palustre*, DC., 23, 41.  
*sinense*, DC., 23, 41.  
*sylvestre*, R. Br., 23, 41.  
*Natsiatum sinense*, Oliver, 36, 495.  
*Nauclea Adina*, Sm., 23, 370.  
*adinoides*, Lindl., 23, 370.  
*pitulifera*, Baill., 23, 370.  
*purpurea*, Roxb., mentioned, 23, 370.  
*racemosa*, Sieb. & Zucc., 23, 371.  
*sinensis*, Oliver, 36, 495.  
*Naumburgia thyrsiflora*, Reichb., 26, 58.  
*Nechamandra alternifolia*, Thw., 36, 2.  
*Roxburghii*, Planch., 36, 2.  
*Neillia affinis*, Hemsl., 36, 495.  
*gracilis*, Franch., 36, 495.  
*longiracemosa*, Hemsl., 36, 495.  
*rubiflora*, D. Don, 36, 495.  
*sinensis*, Oliver, 23, 228.  
*thibetica*, Bur. & Franch., 36, 495.  
*thyrsiflora*, D. Don, 36, 495.  
*Neottia australis*, R. Br., 36, 41.  
 —  $\beta$  *chinensis*, Edwards, 36, 41.  
*camtschatea*, Reichb., 36, 40.  
*camtschatica*, Spreng., 36, 40.  
*kamtschatica*, Lindl., 36, 40.  
*mierantha*, Lindl., 36, 40.  
*procera*, Ker-Gawl., 36, 45.  
*reniformis*, Spreng., 36, 60.  
*sinensis*, Pers., 36, 41.  
 NEPENTHACEÆ, 26, 358.  
*Nepenthes Phyllamphora*, Willd., 26, 358.  
*Nepeta botryoides*, Miq., 26, 290.  
*carulescens*, Maxim., 26, 289.  
*calaminthoides*, Benth., mentioned, 26, 290.  
*Cataria*, Linn., 26, 288; mentioned, 290.  
*Everardi*, S. Moore, 26, 289.  
*Fordii*, Hemsl. \*, 26, 289.  
*Glechoma*, Benth., 26, 290; mentioned, 289.  
 — var. *sinensis*, Miq., 26, 290.  
*incana*, Thunb., 26, 263.  
*japonica*, Willd., 26, 263.

- Nepeta lavandulacea*, Linn., 26, 290.  
*leucophylla*, Benth., 26, 290.  
*tenuifolia*, Benth., 26, 290.
- Nephelaphyllum chinense*, Rolfe, 36, 19.  
*cordifolium*, Lindb., mentioned, 36, 19.  
*cristatum*, Rolfe, 36, 19.
- Nephelium Litchi*, Camb., 23, 139.  
*Longana*, Camb., 23, 139, 492.
- Nephrocodum malaccense*, C. Muell., 36, 4.
- Nephrocœlium malaccense*, Turcz., 36, 4.
- Nephroica cuneifolia*, Miers, 23, 28.  
*cynanchoides*, Miers, 23, 28.  
*dilatata*, Miers, 23, 28.  
*hastata*, Miers, 23, 28.  
*hexagyna*, Miers, 23, 28.  
*ovalifolia*, Miers, 23, 28.  
*pycnantha*, Miers, 23, 28.  
*sarmentosa*, Lour., 23, 28.  
*Thunbergii*, Miers, 23, 28.  
*triloba*, Miers, 23, 28.
- Nerine aurea*, Bury, 36, 89.  
*japonica*, Miq., 36, 89.
- Nerium odorum*, Soland., 26, 97.  
*Oleander*, Linn., mentioned, 26, 97.
- Nertera sinensis*, Hemsl. \*, 23, 391.
- Neubeckia tenuifolia*, Alefeld, 36, 85.
- Neustanthus chinensis*, Benth., 23, 191.  
*phaseoloides*, Benth., 23, 191.
- Neyraudia madagascariensis*, Hook. f., 36, 409.
- Nicandra physaloides*, Gaertn., mentioned, 26, 175.
- Nicotiana chinensis*, Fisch., 26, 169.  
*rustica*, Linn., 26, 169.  
*Tabacum*, Linn., 26, 169.
- Nigrinia serrata*, Thunb., 26, 369.  
*spicifera*, Lam., 26, 368.
- Nitraria Schoberi*, Linn., 23, 97.
- Nomocharis meleagrina*, Franch., 36, 136.  
*pardanthina*, Franch., 36, 136.
- Norysca aurea*, Blume, 23, 72.  
*chinensis*, Spach, 23, 72.  
*punctata*, Blume, 23, 72.
- Nothopanax Bockii*, Harms, 36, 495.  
*Davidi*, Franch., 36, 495.  
*Rosthornii*, Harms, 36, 495.
- Nothoscordum nerinifolium*, Benth., 36, 120.
- Nothosmyrnum japonicum*, Miq., 23, 120.
- Notopterygium*, Boissieu, 36, 495.  
*Forbesii*, Boissieu, 36, 495.  
*Francheti*, Boissieu, 36, 495.
- Nouelia insignis*, Franch., 23, 472.
- NYCTAGINEÆ, 26, 317.
- Nyctanthes grandiflora*, Lour., 26, 82.  
*Sambac*, Linn., 26, 80.
- Nymphæa acutifolia*, DC., distrib., 23, 483.  
*acutiloba*, DC., 23, 33.  
*Nelumbo*, Lour., 23, 34.  
*pygmæa*, Ait., 23, 33.  
*tetragona*, Georgi, 23, 33; distrib., 489.
- NYMPHÆACEÆ, 23, 33.
- Nymphanthus chinensis*, Lour., 26, 423.
- Nyssa sessiliflora*, Hook. f. & Thoms., mentioned, 23, 346.  
*sinensis*, Oliver, 36, 495.
- Oberonia iridifolia*, Lindl., 36, 5.  
*pyrulifera*, Lindl., mentioned, 36, 6.  
*yunnanensis*, Rolfe \*, 36, 6.
- Obione arenaria*, Moq., 26, 325.  
*muricata*, Gaertn., 26, 326.
- Ochranthe arguta*, Lindl., 23, 143.
- Ocimum Basilicum*, Linn., 26, 266.  
*canum*, Sims, 26, 266.  
*capitellatum*, Linn., 26, 269.  
*grandiflorum*, Blume, 26, 269.  
*sanctum*, Linn., 26, 266.  
*stamineum*, Sims, 26, 266.
- Ocimum punctatum*, Thunb., 26, 281.  
*scabrum*, Thunb., 26, 281.
- Odontochilus Clarkei*, Hook. f., 36, 43.  
*Elwesii*, C. B. Clarke, 36, 43.  
*yunnanensis*, Rolfe \*, 36, 43.
- Oeceoclades falcata*, Lindl., 36, 38.
- Oenanthe benghalensis*, Benth. & Hook. f., 23, 331.  
*Hookeri*, C. B. Clarke, 36, 495.  
*linearis*, Wall., 36, 496.  
*rivularis*, Dunn, 36, 496.  
*Rosthornii*, Diels, 36, 496.  
*sinensis*, Dunn, 36, 496.  
*sp.*, 23, 332.  
*species diversa*, 23, 490.  
*stolonifera*, DC., 23, 331; mentioned, 329.  
*Thomsoni*, C. B. Clarke, 23, 332.
- Oenothera fruticosa*, Hance, 23, 310.  
*longiflora*, Jacq., mentioned, 23, 310.
- OLACINEÆ, 23, 114.
- Oldenlandia alata*, Hook. f., 23, 376.  
*angustifolia*, Benth., 23, 375.  
 — var. *pedicellata*, Miq., 23, 377.

- Oldenlandia corymbosa*, Linn., 23, 377.  
*diffusa*, Roxb., 23, 377.  
*herbacea*, DC., 23, 377.  
— var. *uniflora*, Benth., 23, 377.  
*hispida*, Benth., 23, 374.  
*paniculata*, Linn., 23, 377.  
*pterita*, Miq., 23, 376.  
*repens*, Linn., 23, 372.  
*rubroides*, Miq., 23, 373.  
*umbellata*, Osb., 23, 377.  
*Olea Aquifolium*, Sieb. & Zucc., 26, 87.  
*clavata*, D. Don, 26, 92.  
*consanguinea*, Hance, 26, 93.  
*fragrans*, Thunb., 26, 88.  
*marginata*, Champ., 26, 88.  
*ovalis*, Miq., 26, 88.  
*Walpersiana*, Hance, 26, 93.  
OLEACEÆ, 26, 78.  
*Oligobotrya Henryi*, Baker, 36, 109.  
*Olyra orientalis*, Lour., 36, 265.  
*Omphalodes blepharolepis*, Maxim., 26, 148.  
*cordata*, Hemsl. \*, 26, 148.  
*diffusa*, Maxim., 26, 148.  
*moupinensis*, Franch., 36, 496.  
*sericea*, Maxim., 26, 148.  
*trichocarpa*, Maxim., 26, 148.  
*Omphalogramma*, Franch., 36, 496.  
*Delavayi*, Franch., 36, 496.  
*Soulieri*, Franch., 36, 496.  
*vincæflora*, Franch., 36, 496.  
ONAGRARIÆ, 23, 307.  
*Onœa japonica*, Franch., 36, 418.  
*Onosma exsertum*, Hemsl., 36, 496.  
*paniculata*, Bur. & Franch., 36, 496.  
*sinica*, Diels, 36, 496.  
*Onychium japonicum*, Blume, 36, 12.  
*Opa Metrosideros*, Lour., 23, 264.  
*odorata*, Lour., 23, 297.  
*Ophelia angustifolia*, D. Don, 26, 138.  
*bimaculata*, Sieb. & Zucc., 26, 139, 140.  
*chinensis*, Bunge, 26, 139.  
*diluta*, Ledeb., 26, 139.  
*pulchella*, D. Don, 26, 140.  
*tetragona*, Edgew., 26, 141.  
*vacillans*, Hance, 26, 141.  
*Ophiopogon Bockiana*, Diels, 36, 496.  
*clavatus*, C. H. Wright, 36, 77.  
*dracænoides*, Hook. f., mentioned, 36, 77.  
*gracilis*, Kunth, 36, 79.  
*Griffithii*, Hook. f., 36, 77.  
*Jaburan*, Lodd., 36, 77.  
*Ophiopogon japonicus*, Ker-Gawl., 36, 78, 153, 496.  
— var. *genuinus*, Maxim., 36, 78.  
— var. *intermedius*, Maxim., mentioned, 36, 78, 80.  
— var. *umbraticola*, C. H. Wright \*, 36, 78.  
— var. *umbrosus*, Maxim., 36, 78.  
— var. *Wallichianus*, Maxim., 36, 78.  
*japonicus*, Wall., 36, 79.  
*kansuensis*, Batalin, 36, 79.  
*longifolius*, Decne., 36, 79.  
*Muscari*, Decne., 36, 79.  
*spicatus*, Hook., 36, 79.  
*spicatus*, Ker-Gawl., 36, 79.  
— var., 36, 496.  
— var. *communis*, Maxim., 36, 79.  
— var. *gracilis*, Miq., 36, 79.  
— var. *koreanus*, Palibin, 33, 80.  
— var. *Kunthianus*, Maxim., 36, 79.  
— var. *minor*, Maxim., 36, 80.  
*spicatus*, Lodd., 36, 80.  
*umbraticola*, Hance, 36, 78.  
*Wallichianus*, Hook. f., 36, 78.  
*Ophiorrhiza cantonensis*, Hance, 23, 378.  
*Eyrei*, Champ., 23, 378.  
*inflata*, Maxim., 36, 496.  
*japonica*, Blume, 23, 371.  
*Kuroiwai*, Makino, 36, 496.  
*pumila*, Champ., 23, 378.  
*Tashiroi*, Maxim., 36, 496.  
*Ophiospermum sinense*, Lour., 26, 402.  
*Ophioxylon chinense*, Hance, 26, 95.  
*Ophiurus monostachyus*, J. S. Presl, 36, 362.  
*undatus*, Nees, 36, 363.  
*Ophrys camtschatea*, Linn., 36, 40.  
*Corallorhiza*, Linn., 36, 9.  
*Monorchis*, Linn., 36, 51.  
*nervosa*, Thunb., 36, 7.  
*Oplismenus compositus*, Beauv., 36, 337.  
*Crus-Galli*, Dum., 36, 329.  
*frumentaceus*, Kunth, 36, 329.  
*lohiaceus*, Beauv., 36, 337.  
*undulatifolius*, Beauv., 36, 338.  
*Opuntia Dillenii*, Haw., mentioned, 23, 323.  
ORCHIDACEÆ, 36, 5.  
*Orchis aristata*, Fisch., 36, 49.  
*bracteata*, Willd., 36, 54.  
*camtschatica*, Cham., mentioned, 36, 49.  
*Chusua*, D. Don, 36, 49.  
*conopsea*, Linn., 36, 52.  
*cucullata*, Linn., 36, 52.

- Orchis dentata*, Sw., **36**, 59.  
*falcata*, Thunb., **36**, 38.  
*fuscescens*, Linn., **36**, 55.  
*geniculata*, Finet, **36**, 496.  
*gigantea*, Sm., **36**, 37.  
*incarnata*, var. *lanceolata*, Reichb. f.,  
**36**, 50.  
*japonica*, Thunb., **36**, 56.  
*latifolia*, Linn., mentioned, **36**, 49.  
— var. *Beeringiana*, Cham., **36**,  
49.  
*monophylla*, Rolfe, **36**, 50.  
*obcordata*, Buch.-Ham., **36**, 56.  
*Roborovskii*, Maxim., **36**, 50.  
*salina*, Turcz., **36**, 50.  
*spathulata*, Reichb. f., **36**, 50.  
*Susannæ*, Linn., **36**, 57.  
*Szechenyiana*, Reichb. f., **36**, 50.  
*viridis*, Crantz, **36**, 54.  
*Oreocharis aurantiaca*, Franch., **26**,  
226.  
*Auricula*, C. B. Clarke, **26**, 226.  
*Benthami*, C. B. Clarke, **26**, 226.  
*Delavayi*, Franch., **26**, 226.  
*filipes*, Hance, **26**, 226.  
*fokienensis*, Franch., **36**, 496.  
*Henryana*, Oliver, **36**, 496.  
*Maximowiczii*, C. B. Clarke, **26**,  
226.  
*tubicella*, Franch., **36**, 496.  
*Oreocnide Fargesii*, Finet, **36**, 496.  
*frutescens*, Miq., **26**, 491.  
*Oreorchis foliosa*, Lindl., **36**, 9.  
*Oresitrophe rupifraga*, Bunge, **23**, 271.  
*Origanum creticum*, Lour., **26**, 282.  
*Dictamnus*, Linn., mentioned, **26**,  
282.  
*heracleoticum*, Lour., **26**, 282.  
*Marjorana*, Linn., mentioned, **26**,  
282.  
*normale*, D. Don, **26**, 282.  
*vulgare*, Linn., **26**, 282.  
*Orithyia edulis*, Miq., **36**, 138.  
*Orixa japonica*, Thunb., **36**, 496.  
*Ormocarpum sennoides*, DC., **23**, 170.  
*Ormosia emarginata*, Benth., **23**, 204.  
*Fordiana*, Oliver, **36**, 496.  
*Henryi*, Prain, **36**, 496.  
*pachycarpa*, Champ., **23**, 204.  
*semicastrata*, Hance, **23**, 204.  
*striata*, Dunn, **36**, 496.  
*yunnanensis*, Prain, **36**, 496.  
*Ornithocephalochloa arenicola*, Kurz,  
**36**, 340.  
*Ornithochilus Delavayi*, Finet, **36**,  
496.  
*eublepharon*, Hance, **36**, 38.  
*fuscus*, Wall., **36**, 38.  
*Ornithogalum japonicum*, Buerger, **36**,  
192.  
*Ornithogalum japonicum*, Thunb., **36**,  
127.  
*luteum*, Linn., **36**, 138.  
*pauciflorum*, Turcz., **36**, 139.  
*triflorum*, Ledeb., **36**, 140.  
**OROBANCHACEÆ**, **26**, 220.  
*Orobanche albolanata*, Steud., **26**,  
221; mentioned, 222.  
*ammophila*, C. A. Mey, **26**, 221;  
mentioned, 222.  
*canescens*, Bunge, **26**, 221.  
*caryophyllacea*, Bunge, **26**, 222.  
*ombrochares*, Hance, **26**, 222.  
*pycnostachya*, Hance, **26**, 222.  
*Orobis lathyroides*, Linn., **23**, 186.  
*venosus*, Willd., mentioned, **23**,  
185.  
*vernus*, Linn., mentioned, **23**,  
185.  
*Orontium cochinchinense*, Lour., **36**,  
187.  
*japonicum*, Thunb., **36**, 115.  
*Oroxylum indicum*, Vent., **26**, 234.  
*Orthopogon compositus*, R. Br., **36**,  
337.  
*Orthosiphon debilis*, Hemsl. \*, **26**, 267.  
*sinensis*, Hemsl. \*, **26**, 268.  
*stamineus*, Benth., **26**, 268.  
*Orycophragmus sonchifolius*, Bunge,  
**23**, 47.  
*Oryza communissima*, Lour., **36**, 344.  
*glutinosa*, Lour., **36**, 344.  
*montana*, Lour., **36**, 344.  
*præcox*, Lour., **36**, 344.  
*sativa*, Linn., **36**, 344.  
*Oryzopsis obtusa*, Stapf, **36**, 383.  
*Osbeckia chinensis*, Linn., **23**, 298.  
*crinita*, Benth., **23**, 299.  
*heteranthera*, Spreng., **23**, 300.  
*repens*, DC., **23**, 300.  
*stellata* β, DC., **23**, 299.  
*stellata*, Naud., **23**, 299.  
*stellata*, Wall., mentioned, **23**, 299.  
*Osmanthus Aquifolium*, Benth. &  
Hook. f., **26**, 87.  
*armatus*, Diels, **36**, 496.  
*bracteatus*, Matsum., **36**, 496.  
*Cooperi*, Hemsl., **36**, 496.  
*Delavayi*, Franch., **26**, 87.  
*Fordii*, Hemsl. \*, **26**, 88.  
*fragrans*, Lour., **26**, 88.  
*ilicifolius*, Hort., **26**, 87.  
*marginatus*, Benth. & Hook. f., **26**,  
88.  
*Osmorrhiza longistylis*, DC., men-  
tioned, **23**, 300.  
*Osmothamnus fragrans*, DC., **26**, 23.  
*pallidus*, DC., **26**, 24.  
*Osteomeles anthyllidifolia*, Lindl., **23**,  
265.



- Osteomeles Pyracantha*, Decne., 23, 262.  
*subrotunda*, C. Koch, 23, 265.  
*Ostrya carpinifolia*, Franch., 26, 503.  
*japonica*, Sargent, 26, 503.  
*virginica*, Willd., 26, 503.  
— var. *japonica*, Maxim., 26, 503.  
*Ostryopsis Davidiana*, Decne., 26, 502.  
— var. *cinerascens*, Franch., 26, 503.  
*Davidiana*, Franch., 26, 503.  
*Osyris japonica*, Thunb., 23, 341.  
*Otanthra Fordii*, Hance, 23, 299.  
*Ottelia alismoides*, Pers., 36, 3.  
*indica*, Planch., 36, 3.  
*japonica*, Miq., 36, 3.  
*Ourouparia* = *Nauclea*.  
*sinensis*, Diels, 36, 496.  
*Oxalis Acetosella*, Linn., 23, 99.  
*corniculata*, Linn., 23, 99.  
*fontana*, Bunge, 23, 99.  
*Griffithii*, Edgew. & Hook. f., 23, 99.  
*japonica*, Franch. & Sav., 36, 496.  
*Martiana*, Zucc., 23, 99.  
*obtriangulata*, Maxim., 23, 99.  
*Reinwardtii*, Zucc., 23, 100.  
*sensitiva*, Linn., 23, 100.  
*stricta*, Linn., 23, 99.  
*Oxyceros sinensis*, Lour., 23, 382.  
*Oxygraphis Cymbalaria*, Pursh, 36, 496.  
*Delavayi*, Franch., 36, 496.  
*glacialis*, Bunge, 36, 497.  
*Oxyria sinensis*, Hemsl., 36, 497.  
*Oxyspora pauciflora*, Benth., 23, 301.  
*Oxytropis Beketowii*, Krassn., 36, 497.  
*bicolor*, Bunge, 23, 167.  
*Delavayi*, Franch., 36, 497.  
*falcata*, Bunge, 36, 497.  
*hirta*, Bunge, 23, 167.  
*kansuensis*, Bunge, 36, 497.  
*melanocalyx*, Bunge, 36, 497.  
*Mœllendorffi*, Bunge, 23, 167.  
*ochrocephala*, Bunge, 36, 497.  
*oxyphylla*, DC., 23, 168.  
*psammocharis*, Hance, 23, 168.  
*strobilacea*, Bunge, 36, 497.  
*subfalcata*, Hance, 23, 168.  
*sylinchanensis*, Franch., 36, 497.  
*trichophora*, Franch., 36, 497.  
*uratensis*, Franch., 23, 168.  
*yunnanensis*, Franch., 36, 497.  
  
*Pachyne spectabilis*, Salisb., 36, 18.  
*Pachyrhizus angulatus*, Rich., 23, 194.  
*Pachyrhizus Thunbergianus*, Sieb. & Zucc., 23, 191.  
*trilobus*, DC., 23, 191, 194.  
*Pachysandra axillaris*, Franch., 26, 419; 36, 497.  
*coriacea*, Hook., 26, 418.  
*terminalis*, Sieb. & Zucc., 26, 419.  
*Pachystoma chinense*, Reichb. f., 36, 17.  
*Fortunei*, Reichb. f., 36, 18.  
*Paderia chinensis*, Hance, 23, 389.  
*fatida*, Thunb., 23, 389.  
*tomentosa*, Blume, 23, 389.  
*Paderota axillaris*, Sieb. & Zucc., 26, 195.  
*bracteata*, Sieb. & Zucc., 26, 195.  
*villosula*, Miq., 26, 195.  
*Pæonia anomala*, Linn., 36, 49.  
*Delavayi*, Franch., 36, 497.  
*lutea*, Franch., 36, 497.  
*Paliurus aculeatus*, Lam., 23, 126.  
*Aubletia*, Schultz, 23, 126.  
*australis*, Gaertn., 23, 126.  
*hirsutus*, Hemsl., 36, 497.  
*orientalis*, Hemsl., 36, 497.  
*ramosissimus*, Poir., 23, 126.  
**PALMÆ**, 36, 169.  
*Palura sinica*, Miers, 26, 75.  
*Panax aculeatum*, Ait., 23, 339.  
*Davidi*, Franch., 36, 497.  
*Delavayi*, Franch., 36, 497.  
*divaricatum*, Sieb. & Zucc., 23, 339.  
*fallax*, Miq., 23, 339.  
*fragrans*, Roxb., 23, 343.  
*Ginseng*, C. A. Mey., 23, 338.  
*Loureirianum*, DC., 23, 339.  
*quinquefolium*, Linn., 23, 338.  
*ricinifolium*, Sieb. & Zucc., 23, 340.  
*sessiliflorum*, Rupr. & Maxim., 23, 340.  
*spinosum*, Linn., 23, 341.  
*Pancovia Delavayi*, Franch., 36, 497.  
*Pancreatium biflorum*, Roxb., 36, 90.  
*zeylanicum*, Linn., mentioned, 36, 90.  
**PANDANACEÆ**, 36, 171.  
*Pandanus fascicularis*, Lam., 36, 171.  
*furcatus*, Roxb., 36, 171.  
*odoratissimus*, Linn., 36, 171.  
*sp.*, Hance, 36, 171.  
*sp.*, Matsum., 36, 497.  
*urophyllus*, Hance, 36, 171.  
*Panicum acerescens*, Trin., 36, 332.  
*acroanthum*, Steud., 36, 327.  
*albens*, Steud., 36, 321.  
*ambiguum*, Trin., 36, 327.  
*amplexicaule*, Rudge, 36, 327.  
*angustum*, Trin., 36, 330.  
*arborescens*, Linn., 36, 328.  
*asperum*, Roem., 36, 339.

- Panicum barbatum*, Willd., 36, 322.  
*brevifolium*, Linn., 36, 328.  
*cæsiuum*, Nees, 36, 329.  
*ciliare*, Retz., 36, 325.  
*cimicinum*, Retz., 36, 334.  
*cocospermum*, Steud., 36, 333.  
*colonum*, Linn., 36, 328.  
*commutatum*, Nees, 36, 325.  
*compositum*, Linn., 36, 337.  
*concinnum*, Nees, 36, 333.  
*Crus-Corvi*, Linn., 36, 329.  
*Crus-Galli*, Linn., 36, 328.  
— *var. frumentaceum*, Trimen, 36, 329.  
*decompositum*, R. Br., 36, 330.  
*distachyum*, Linn., 36, 330.  
*excurrens*, Trin., 36, 336.  
*fibrosum*, Hack., 36, 326.  
*Forbesianum*, Nees, 36, 334.  
*frumentaceum*, Roxb., 36, 329.  
*glabrum*, Gaudin, 36, 324.  
*glaucum*, Linn., 36, 335.  
*heteranthum*, Nees, 36, 323.  
*hispidulum*, Hook., 36, 329.  
*humifusum*, Kunth, 36, 324.  
*humile*, Nees, 36, 330.  
*incomtum*, Trin., 36, 333.  
*indicum*, Linn., 36, 330.  
— *var. angustum*, Hook. f., 36, 330.  
— *var. contractum*, Miq., 36, 330.  
*ischæmoides*, Retz., 36, 333.  
*italicum*, Linn., 36, 335.  
*hasianum*, Munro, 36, 331.  
*mandshuricum*, Maxim., 36, 341.  
*maximum*, Jacq., 36, 331.  
*micrognostum*, Steud., 36, 333.  
*miliaceum*, Linn., 36, 331.  
*montanum*, Roxb., 36, 331.  
*multinode*, Presl, 36, 331.  
*Myurus*, H. B. K., 36, 327.  
*natans*, Roem., 36, 332.  
*nervosum*, Roxb., 36, 336.  
*nilagiricum*, Steud., 36, 333.  
*nodosum*, Kunth, 36, 331.  
*ovalifolium*, Poir., 36, 328.  
*pallens*, Sw., 36, 334.  
*palmafolium*, Koen., 36, 336.  
*paludosum*, Roxb., 36, 332.  
*parvulum*, Trin., 36, 324.  
*patens*, Linn., 36, 332.  
*patens*, Osbeck, 36, 321.  
*penicillatum*, Nees, 36, 335.  
*plicatum*, Lam., 36, 336.  
*proliferum*, Lam., 36, 332.  
*prostratum*, Lam., 36, 332.  
*psilopodium*, Trin., 36, 332.  
*pulchellum*, Spreng., 36, 387.  
*radicans*, Retz., 36, 332.  
*Panicum repens*, Linn., 36, 332.  
*Rhignon*, Steud., 36, 332.  
*sanguinale*, Linn., 36, 325.  
*sarmentosum*, Roxb., 36, 333.  
*semialatum*, R. Br., 36, 334.  
*simpliciusculum*, Wight & Arn., 36, 399.  
*spicatum*, Roxb., 36, 339.  
*spinescens*, R. Br., 36, 339.  
*stagninum*, Hance, 36, 329.  
*tenuiflorum*, R. Br., 36, 324.  
*ternatum*, Hochst., 36, 327.  
*trypheron*, Schult., 36, 333.  
*undulatifolium*, Arduino, 36, 338.  
*vacillans*, Steud., 36, 333.  
*villosum*, Lam., 36, 333.  
*viride*, Linn., 36, 337.  
*Williamsii*, Hance, 36, 341.  
*Papaver alpinum*, Linn., 23, 34.  
*nudicaule*, Linn., 23, 34.  
*Rhœas*, Linn., mentioned, 23, 34.  
*somniferum*, Linn., 23, 34.  
PAPAVERACEÆ, 23, 34.  
*Paphiopedilum Parishii*, Pfitz., 36, 67.  
*purpuratum*, Pfitz., 36, 67.  
PAPILIONACEÆ, 23, 150.  
*Pappophorum brachystachyus*, Jaub. & Spach, 36, 407.  
*boreale*, Griseb., 36, 407.  
*Papyrus odoratus*, Willd., 36, 222.  
*Paradisia minor*, C. H. Wright, 36, 497.  
*Paradombeya*, Stapf, 36, 497.  
*sinensis*, Stapf, 36, 497.  
*Parapottsia*, Miq., 26, 97.  
*Paratropia cantoniensis*, Hook. & Arn., 23, 342.  
*Pardanthus chinensis*, Ker-Gawl., 36, 86.  
*dichotoma*, Ledeb., 36, 81.  
*Parechites adnascens*, Hance, 26, 100.  
*Bowringii*, Hance, 26, 99.  
*Thunbergii*, A. Gray, 26, 99.  
*Parietaria debilis*, G. Forst., 26, 492.  
— *var. micrantha*, Wedd., 26, 492.  
*debilis*, G. Forst. partim, 26, 492.  
*micrantha*, Ledeb., 26, 492.  
*microphylla*, Liebm., 26, 476.  
*Paris Bockiana*, Diels, 36, 497.  
*chinensis*, Franch., 36, 144.  
*Christii*, H. Léveillé, 36, 497.  
*Delavayi*, Franch., 36, 144.  
*Fargesii*, Franch., 36, 145.  
*Franchetiana*, H. Léveillé, 36, 497.  
*Henryi*, Diels, 36, 497.  
*hexaphylla*, Cham., 36, 146.  
*incompleta*, Bieb., 36, 145.  
*Mercieri*, H. Léveillé, 36, 497.  
*obovata*, Ledeb., 36, 146.

- Paris petiolata*, Baker, **36**, 145.  
 — *var. membranacea*, C. H. Wright \*, **36**, 145.  
*polyphylla*, Sm., **36**, 145.  
 — *var. brachystemon*, Franch., **36**, 146.  
 — *var. platypetala*, Franch., **36**, 146.  
*quadrifolia*, Linn., **36**, 146.  
 — *var. obovata*, Regel, **36**, 146.  
 — *var. setchuenensis*, Franch., **36**, 146.  
*thibetica*, Franch., **36**, 497.  
*yunnanensis*, Franch., **36**, 146.  
*Parnassia amoena*, Diels, **36**, 497.  
*appendiculata*, Batalin, **36**, 497.  
*chinensis*, Franch., **36**, 497.  
*crassifolia*, Franch., **36**, 497.  
*Davidii*, Franch., **36**, 497.  
*Delavayi*, Franch., **36**, 497.  
*Faberi*, Oliver, **36**, 497.  
*foliosa*, Hook. f. & Thoms., **23**, 271.  
*Laxmanni*, Pall., **36**, 497.  
*lutea*, Batalin, **36**, 498.  
*monochorifolia*, Franch., **36**, 498.  
*mysorensis*, Heyne, **23**, 271.  
*Noemiæ*, Franch., **36**, 498.  
*nummularia*, Maxim., **23**, 271.  
*oreophila*, Hance, **23**, 272.  
*palustris*, Linn., **23**, 272.  
*perciliata*, Diels, **36**, 498.  
*pusilla*, Hook. f. & Thoms., **36**, 498.  
*setchuenensis*, Franch., **36**, 498.  
*viridiflora*, Batalin, **36**, 498.  
*Wightiana*, Wall., **23**, 272; **36**, 498.  
 — *var. microblephara*, Franch., **23**, 272.  
*yunnanensis*, Franch., **36**, 498.  
*Parochætus communis*, Buch.-Ham., **36**, 498.  
*Parophiorrhiza khasiana*, C. B. Clarke, **26**, 117.  
*Parrya Beketowi*, Krassn., **36**, 498.  
*prolifera*, Maxim., **36**, 498.  
*siliquosa*, Krassn., **36**, 498.  
*villosa*, Maxim., **36**, 498.  
*Parsonia Helicandra*, Hook. & Arn., mentioned, **26**, 96.  
*ovata*, Wall., **26**, 96.  
*spiralis*, Wall., **26**, 96.  
*Parthenocissus Henryana*, Hemsl., **36**, 498.  
*himalayana*, Gilg, **36**, 498.  
*sinensis*, Diels & Gilg, **36**, 498.  
*tricuspidata*, Planch., **36**, 498.  
*Parthenoxylon porrectum*, Blume, **26**, 372.  
*Pseudosassafras*, Blume, **26**, 372.  
*Parvattia chinensis*, Franch., **36**, 498.  
*decora*, Dunn, **36**, 498.  
*Paspalum ambiguum*, DC., **36**, 324.  
*annulatum*, Fluegge, **36**, 320.  
*brevifolium*, Fluegge, **36**, 324.  
*chinense*, Nees, **36**, 324.  
*conjugatum*, Berg, **36**, 319.  
*dissectum*, Linn., **36**, 320.  
*distichum*, Linn., **36**, 319.  
*heteranthum*, Hook. f., **36**, 323.  
*longiflorum*, Retz., **36**, 324.  
*sanguinale*, Lam., **36**, 325.  
*serobiculatum*, Linn., **36**, 320.  
*spp.*, A. Henry, **36**, 319, 320.  
*ternatum*, Hook., **36**, 327.  
*Thunbergii*, Kunth, **36**, 320.  
*vaginatum*, Sw., **36**, 419.  
*villosum*, Thunb., **36**, 320.  
*Passerina Chamædaphne*, Bunge, **26**, 397.  
*vesiculosa*, Fisch. & Mey., **26**, 402.  
*Passiflora cærulea*, Linn., mentioned, **23**, 312.  
*cupiformis*, Mast., **36**, 498.  
*foetida*, Linn., mentioned, **23**, 312.  
*Franchetiana*, Hemsl., **36**, 498.  
*hainanensis*, Hance, **23**, 312.  
*Henryi*, Hemsl., **36**, 498.  
*Horsfieldii*, Blume, **23**, 312.  
*ligulifolia*, Mast., **23**, 312.  
*Seguini*, H. Lévêillé & Vaniot, **36**, 498.  
 PASSIFLORACEÆ, **23**, 312.  
*Patrinia angustifolia*, Hemsl. \*, **23**, 396.  
*Dielsii*, Graebn., **36**, 498.  
*graveolens*, Hance, **23**, 396.  
*heterophylla*, Bunge, **23**, 396.  
*hispida*, Bunge, **23**, 397.  
*intermedia*, Roem. & Schult., mentioned, **23**, 396.  
*monandra*, C. B. Clarke, var., **36**, 498.  
*ovata*, Bunge, **23**, 396, 398.  
*palmata*, Maxim., mentioned, **23**, 397.  
*parviflora*, Sieb. & Zucc., **23**, 397.  
*rupestris*, Juss., **23**, 396.  
*saniculæfolia*, Hemsl. \*, **23**, 397.  
*scabiosæfolia*, Fisch., **23**, 397.  
*scabra*, Bunge, **23**, 398.  
*serratulæfolia*, Fisch., **23**, 397.  
*villosa*, Juss., **23**, 398; mentioned, 396.  
*Paullinia asiatica*, Linn., **23**, 108.  
*japonica*, Thunb., **23**, 136.  
*Paulownia Fargesii*, Franch., **36**, 498.  
*Fortunei*, Hemsl. \*, **26**, 180.  
*imperialis*, Hance, **26**, 180.

- Paulownia imperialis*, Sieb. & Zucc., 26, 180.  
*Pavetta arenosa*, Lour., 23, 386.  
     *Brunonis*, Wall., mentioned, 23, 385.  
     *indica*, Linn., 23, 386.  
     *Kroneana*, Miq., mentioned, 23, 385.  
     *sinica*, Miq., mentioned, 23, 385.  
     *stricta*, Roxb., mentioned, 23, 385.  
**PEDALINEÆ**, 26, 236.  
*Pedicularis alaschanica*, Maxim., 26, 204.  
     *Alopecurus*, Franch., 26, 205; mentioned, 207.  
     *Anas*, Maxim., 26, 205.  
     *armata*, Maxim., 26, 205.  
     *Artselaeri*, Maxim., 26, 205.  
     *axillaris*, Franch., 26, 205; mentioned, 208.  
     *bantangensis*, Bur. & Franch., 36, 498.  
     *bicolor*, Diels, 36, 498.  
     *bidentata*, Maxim., 26, 205.  
     *Bietii*, Franch., 36, 498.  
     *binaria*, Maxim., 26, 205.  
     *Biondiana*, Diels, 36, 498.  
     *birostris*, Bur. & Franch., 36, 498.  
     *brevifolia*, D. Don, 26, 205.  
     *brevilabris*, Franch., 36, 498.  
     *carnosa*, Wall., 26, 205.  
     *cephalantha*, Franch., 26, 206.  
     *cheilanthifolia*, Schrenk, 26, 206.  
     *chinensis*, Maxim., 26, 206.  
     *cinerascens*, Franch., 36, 498.  
     *comptoniæfolia*, Franch., 26, 206.  
     *conifera*, Maxim. \*, 26, 206.  
     *cranolopha*, Maxim., 26, 207.  
     *craspedotricha*, Maxim., 26, 207.  
     *crenata*, Maxim., 26, 207.  
     *cristata*, Maxim., 26, 207.  
     *curvituba*, Maxim., 26, 207.  
     *cyathophylla*, Franch., 36, 498.  
     *Davidi*, Franch., 26, 207; 36, 498.  
     *debilis*, Franch., 26, 207.  
     *decora*, Franch., 36, 498.  
     *Delavayi*, Franch., 26, 208.  
     *deltoides*, Franch., 26, 208.  
     *densispica*, Franch., 26, 208.  
     *elegans*, Franch., 36, 499.  
     *Elwesii*, Hook., 26, 208.  
     *Fargesii*, Franch., 36, 499.  
     *fastigiata*, Franch., 36, 499.  
     *filicifolia*, Hemsl. \*, 26, 208.  
     *Filicula*, Franch., 26, 209.  
     *flaccida*, Prain, 36, 499.  
     *floribunda*, Franch., 36, 499.  
     *Franchetiana*, Maxim., 26, 209.  
     *Pedicularis goniantha*, Bur. & Franch., 36, 499.  
     *gracilis*, Wall., 26, 209; mentioned, 36, 217.  
     *gruina*, Franch., 26, 209.  
     *gyrorhyncha*, Franch., 26, 209.  
     *Hemsleyana*, Prain, 36, 499.  
     *Henryi*, Maxim., 26, 209.  
     *hirtella*, Franch. \*, 26, 209.  
     *imperialis*, Franch., 36, 499.  
     *ingens*, Maxim., 26, 210.  
     *integrifolia*, Hook. f., 26, 210.  
     *kansuensis*, Maxim., 26, 210.  
     *kialensis*, Franch., 36, 499.  
     *labellata*, Jacquem., 26, 210.  
     *lachnoglossa*, Hook. f., 26, 210.  
     *lasiophrys*, Maxim., 26, 211.  
     *laxiflora*, Franch., 36, 499.  
     *leiandra*, Franch. \*, 26, 211; mentioned, 220.  
     *lineata*, Franch., 26, 211.  
     *likiangensis*, Franch., 26, 211.  
     *longepetiolata*, Franch., 26, 211.  
     *longicaulis*, Franch., 26, 212.  
     *longiflora*, Rudolphi, 26, 206; 36, 499.  
     *longipes*, Maxim., 26, 212.  
     *lutescens*, Franch., 26, 212.  
     *lyrata*, Prain, 36, 499.  
     *macilentata*, Franch. \*, 26, 212.  
     *macrosiphon*, Franch., 26, 208, 212; 36, 499.  
     *Maximowiczii*, Krassn., 36, 499.  
     *melampyriflora*, Franch., 26, 212.  
     *microchila*, Franch., 26, 212.  
     *microphyton*, Bur. & Franch., 36, 499.  
     *moupinensis*, Franch., 26, 213; 36, 499.  
     *musciicola*, Maxim., 26, 213.  
     *Mussoti*, Franch., 36, 499.  
     *myriophylla*, Pall., 26, 213; mentioned, 217.  
     *nasturtiifolia*, Franch., 36, 499.  
     *oligantha*, Franch., 26, 213.  
     *oxycarpa*, Franch., 26, 213; mentioned, 212.  
     *phaceliæfolia*, Franch., 36, 499.  
     *pilostachya*, Maxim., 26, 213.  
     *plicata*, Maxim., 26, 213.  
     *polyphylla*, Franch., 26, 213; mentioned, 207.  
     *porrecta*, Wall., mentioned, 26, 211.  
     *Potanini*, Maxim., 26, 214.  
     *Princeps*, Franch., 36, 499.  
     *Provoti*, Franch., 36, 499.  
     *Przewalskii*, Maxim., 26, 214.  
     *ramalana*, Britt., 36, 499.  
     *recurva*, Maxim., 26, 214.

- Pedicularis refracta*, Maxim., 26, 214.  
*resupinata*, Linn., 26, 214.  
*Rex*, C. B. Clarke, 26, 214.  
*rhodotricha*, Maxim., 26, 215.  
*rhynchodonta*, Franch., 36, 499.  
*rigida*, Franch., 26, 214.  
*Roborowskii*, Maxim., 26, 215.  
*rudis*, Maxim., 26, 215.  
*rupicola*, Franch., 26, 215.  
*salviaeflora*, Franch. \*, 26, 215.  
*Scolopax*, Maxim., 26, 216.  
*semitorta*, Maxim., 26, 216.  
*sigmoidea*, Franch., 26, 216.  
*Sima*, Maxim., 26, 216.  
*siphonantha*, D. Don, 26, 216.  
*Souliei*, Franch., 36, 499.  
*spicata*, Pall., 26, 216.  
*stenantha*, Franch., 36, 499.  
*stenocorys*, Franch., 36, 499.  
*striata*, Pall., 26, 216.  
*strobilacea*, Franch. \*, 26, 216.  
*superba*, Franch., 26, 217.  
*szetschuanica*, Maxim., 26, 217.  
*Tartarinowi*, Maxim., 26, 217.  
*tatsienensis*, Franch., 36, 499.  
*tenuisecta*, Franch., 26, 217.  
*ternata*, Maxim., 26, 218.  
*tibetica*, Franch., 36, 499.  
*tongolensis*, Franch., 36, 499.  
*torta*, Maxim., 26, 218.  
*trichoglossa*, Franch., 26, 218.  
*tristis*, Linn., 26, 218.  
*tsangchanensis*, Franch., 26, 218.  
*vagans*, Hemsl. \*, 26, 218.  
*verbenæfolia*, Franch., 26, 219; mentioned, 211.  
*veronicifolia*, Franch., 36, 499.  
*versicolor*, Wahlenb., 26, 219.  
*verticillata*, Linn., 26, 219.  
*Viali*, Franch. \*, 26, 219.  
*villosula*, Franch. \*, 26, 220.  
*yunnanensis*, Franch., 26, 220.
- Peganum Nigellastrum*, Bunge, 23, 103.
- Peliosanthes Delavayi*, Franch., 36, 77.  
*macrostegia*, Hance, 36, 77.
- Pellionia*, genus referred to, 26, 482.  
*ambigua*, Wedd., 26, 482.  
*brevifolia*, Benth., 26, 481.  
*Griffithiana*, Wedd., 26, 481.  
*Grijssii*, Hance, 26, 481.  
*radicans*, Wedd., 26, 481; mentioned, 484.  
*scabra*, Benth., 26, 481.  
*umbellata*, Wedd., 26, 483.  
*viridis*, C. H. Wright \*, 26, 481.
- Peltophorum ferrugineum*, Benth., 23, 204.
- Pemphis acidula*, Forst., 23, 305.
- Penicillaria spicata*, Willd., 36, 339.
- Pennisetum chinense*, Nees, 36, 338.  
*compressum*, R. Br., 36, 338.  
 — var. *viridescens*, Rendle \*, 36, 339.  
*flaccidum*, Griseb., 36, 339.  
*japonicum*, Trin., 36, 338.  
 — var. *viridescens*, Palib., 36, 339.  
*spicatum*, Roem. & Schult., 36, 339.  
*typhoideum*, Rich., 36, 339.
- Pentacelium bontioides*, Sieb. & Zucc., 26, 250.
- Pentaneura khasiana*, Kurz, 26, 100.
- Pentapanax Henryi*, Harms, 36, 499.  
*verticillatum*, Dunn, 36, 499.  
*yunnanensis*, Franch., 36, 499.
- Pentapetes phœnicea*, Linn., 23, 91.
- Pentaphylax euryoides*, Gardn. & Champ., 23, 77.
- Pentasachme caudatum*, Wall., 26, 112.  
*Championi*, Benth., 26, 112.  
*glaucescens*, Decne., 26, 112.  
*Stauntoni*, Decne., 26, 112.
- Pentatropis officinalis*, Hemsl. \*, 26, 110.  
*spiralis*, Decne., mentioned, 26, 110.
- Penthorum chinense*, Pursh, 23, 288.  
*intermedium*, Turcz., 23, 288.  
*sedoides*, Linn., 23, 288.  
 — forma *angustifolia*, Miq., 23, 288.
- Peperomia dindygulensis*, Miq., 26, 366.  
*polystachya*, A. Dietr., mentioned, 26, 366.  
*reflexa*, A. Dietr., 26, 366.
- Peplis indica*, Willd., 23, 303.
- Perdicium Anandria*, R. Br., 23, 473.  
*tomentosum*, Turcz., 23, 473.
- Pergularia divaricata*, Lour., 26, 97.  
*japonica*, Thunb., 26, 111.  
*minor*, Andr., mentioned, 26, 114.  
*odoratissima*, Sm., 26, 114.  
*odoratissima*, Sm., 26, 114.  
*pallida*, Wight & Arn., 26, 114.  
*sinensis*, Lour., 26, 101.
- Pericampylus aduncus*, Miers, 23, 29.  
*assamicus*, Miers, 23, 29.  
*incanus*, Miers, 23, 29.  
*membranaceus*, Miers, 23, 29.
- Perilla arguta*, Benth., 26, 279.  
*lanceolata*, Benth., 26, 281.  
*nankinensis*, Decne., 26, 279.  
*oeymoides*, Linn., 26, 279.  
 — var. *crispa*, Benth., 26, 279.

- Periploca calophylla*, *Falconer*, **26**, 101.  
*chinensis*, *Lour.*, **26**, 101.  
*chinensis*, *Spreng.*, **26**, 101.  
*Sepium*, *Bunge*, **26**, 101.  
*Peristrophe chinensis*, *Nees*, **26**, 248.  
*Cumingiana*, *Nees*, **26**, 249.  
*fera*, *C. B. Clarke*, **26**, 249.  
*montana*, *Nees*, **26**, 249.  
*tinctoria*, *Nees*, **26**, 248.  
*Peristylis grandis*, *Blume*, **36**, 54.  
*Peristylus bracteatus*, *Lindl.*, **36**, 54.  
*chloranthus*, *Lindl.*, **36**, 53.  
*cœloceras*, *Finet*, **36**, 499.  
*Forceps*, *Finet*, **36**, 499.  
*goodyeroides*, *Lindl.*, **36**, 54.  
*monanthus*, *Finet*, **36**, 499.  
*Sampsoni*, *Hance*, **36**, 54.  
*tetralobus*, *Finet*, **36**, 499.  
*viridis*, *Lindl.*, **36**, 54.  
*Perotis latifolia*, *Ait.*, **36**, 343.  
*longiflora*, *Nees*, **36**, 343.  
*patula*, *Nees*, **36**, 343.  
*Perrottetia racemosa*, *Oliver*, **36**, 499.  
*Persea*, *Plum.*, **26**, 370.  
*Camphora*, *Spreng.*, **26**, 371.  
*Cubeba*, *Spreng.*, **26**, 380.  
*Nanmu*, *Oliver*, **26**, 376.  
*Persica Davidiana*, *Carr.*, **23**, 220.  
*vulgaris*, *Mill.*, **23**, 220.  
*Pertya Bodinieri*, *Vaniot*, **36**, 499.  
*sinensis*, *Oliver*, **36**, 499.  
*Perularia fuscescens*, *Lindl.*, **36**, 55.  
*Petasites japonicus*, *Miq.*, **23**, 446; mentioned, 447.  
*spurius*, *Miq.*, **23**, 446.  
*tricholobus*, *Franch.*, **23**, 447.  
*Petrocodon dealbatus*, *Hance*, **26**, 229.  
*Petrocosmea grandiflora*, *Hemsl.*, **36**, 499.  
*iodioides*, *Hemsl.*, **36**, 500.  
*minor*, *Hemsl.*, **36**, 500.  
*sinensis*, *Oliver*, **26**, 229.  
*Petroselinum sativum*, *Hoffm.*, mentioned, **23**, 328.  
*Peucedanum cartilagineo-marginatum*, *Makino*, **36**, 500.  
*decursivum*, *Maxim.*, **23**, 335.  
*Delavayi*, *Franch.*, **36**, 500.  
*elegans*, *Komarov*, **36**, 500.  
*Giraldii*, *Diels*, **36**, 500.  
*graveolens*, *Benth. & Hook. f.*, mentioned, **23**, 335; **36**, 500.  
*heterophyllum*, *Franch.*, **36**, 500.  
*macilentum*, *Franch.*, **36**, 500.  
*medicum*, *Dunn*, **36**, 500.  
*Podagraria*, *Boissieu*, **36**, 500.  
*Peucedanum Præruptorum*, *Dunn*, **36**, 500.  
*reptans*, *Diels*, **36**, 500.  
*rigidum*, *Bunge*, **23**, 335.  
*spp.*, **23**, 335, 336.  
*terebinthaceum*, *Fisch.*, **23**, 335.  
*torilifolium*, *Boissieu*, **36**, 500.  
*Phaca Salsula*, *Pall.*, **23**, 162.  
*trifoliata*, *Linn.*, **23**, 167.  
*villosa*, *Nutt.*, mentioned, **23**, 167.  
*Phacellanthus* sp., **26**, 221.  
*tubulosus*, *Sieb. & Zucc.*, **26**, 221.  
— var., **26**, 221.  
*Phænosperma globosa*, *Munro*, **36**, 340.  
*Phaius grandifolius*, *Lour.*, **36**, 18; mentioned, 28.  
*Tankervillii*, *Blume*, **36**, 18.  
*Phakellanthus multiflorus*, *Steud.*, **36**, 263.  
*Phalænopsis amabilis*, *Lindl.*, **36**, 34.  
*Aphrodite*, *Reichb. f.*, **36**, 34.  
*Wightii*, *Reichb. f.*, **36**, 34.  
*Phalangium parviflorum*, *Wight*, **36**, 119.  
*Phalaris arundinacea*, *Linn.*, **36**, 379.  
*Phanera Championi*, *Benth.*, **23**, 212.  
*corymbosa*, *Benth.*, **23**, 212.  
*glauca*, *Benth.*, **23**, 212.  
*variegata*, *Benth.*, **23**, 213.  
*Pharbitis diversifolia*, *Lindl.*, **26**, 160.  
*hederacea*, *Choisy*, **26**, 160.  
*hispida*, *Choisy*, **26**, 162.  
*insularis*, *Choisy*, **26**, 158.  
*Nil*, *Choisy*, **26**, 160.  
*triloba*, *Miq.*, **26**, 160.  
*Phaseolus aconitifolius*, *Jacq.*, **36**, 500.  
*anguinus*, *Bun* **3**, 1  
*chrysanthus*, *Savi*, **23**, 193.  
*grandis*, *Dalz. & Gibs.*, **36**, 500.  
*humilis*, *Hassk.*, **23**, 193.  
*minimum*, *Roxb.*, **23**, 193.  
*Mungo*, *Linn.*, **23**, 193; **36**, 500.  
*radiatus*, *Linn.*, **23**, 193; **36**, 500.  
*Ricciardianus*, *Ten.*, **36**, 500.  
— var. *hysterinus*, *Dur.*, **23**, 193.  
*trilobus*, *Ait.*, **23**, 193.  
*vexillatus*, *Linn.*, **23**, 192.  
*vulgaris*, *Linn.*, mentioned, **23**, 193.  
*Phelipæa salsa*, *C. A. Mey.*, **26**, 222.  
*Phellodendron amurense*, *Rupr.*, **23**, 108.  
*japonicum*, *Maxim.*, **36**, 500.  
*Phellopterus littoralis*, *Benth.*, **23**, 331.

- Philadelphus coronarius*, Linn., 23, 277.  
 — *var.*, *pekinensis*, Maxim., 23, 277.  
 — *va* *tenuifolius*, Maxim., 23, 278.  
 — *var.* *tomentosus*, Hook. f. & Thoms., 23, 277.  
*neanus*, Koehne, 36, 500.  
*parviflorus*, Carr., 23, 278.  
*pekinensis*, Rupr., 23, 277.  
*rubricaulis*, Carr., 23, 278.  
*Schrenkii*, Rupr., 36, 500.  
*sericanthus*, Koehne, 36, 500.  
*tenuifolius*, Rupr., 23, 278.  
*tomentosus*, Wall., 23, 277.  
*Phillyrea paniculata*, Roxb., 26, 92.  
*robusta*, Roxb., 26, 92.  
*Philoxerus Wrightii*, Hook. f., 26, 323.  
 PHILYDRACEÆ, 36, 150.  
*Philydrum lanuginosum*, Banks, 36, 150.  
*Phlebophyllum apricum*, Benth., 26, 239.  
*Phleum alpinum*, Linn., 36, 384.  
*asperum*, Jacq., 36, 384.  
*Phlomis albiflora*, Hemsl. \*, 26, 304.  
*chinensis*, Retz., 26, 303.  
*gracilis*, Hemsl. \*, 26, 305.  
*medicinalis*, Diels, 36, 500.  
*mongolica*, Turcz., 26, 306.  
*oreophila*, Kar. & Kir., mentioned, 26, 306.  
*rugosa*, Benth., 26, 305, 306.  
*setifera*, Bur. & Franch., 36, 500.  
*stenocalyx*, Diels, 36, 500.  
*tatsienensis*, Bur. & Franch., 36, 500.  
*tuberosa*, Linn., 26, 306.  
*umbrosa*, Turcz., 26, 306.  
 — *var.* *australis*, Hemsl. \*, 26, 306.  
*Phylarodoxa leucantha*, S. Moore, 26, 92, 93.  
*Phoberos Arnottianus*, Hance, 23, 57.  
*chinensis*, Lour., 23, 57.  
*sævus*, Hance, 23, 57.  
*Phabe latifolia*, Champ., 26, 372.  
*Phœnix acaulis*, Benth., 36, 168.  
*farinifera*, Hance, 36, 168.  
*Hanceana*, Naud., 36, 168.  
*humilis*, Royle, *var.* *Hanceana*, Becc., 36, 168.  
*Pholidota articulata*, Lindl., 36, 24.  
*cantonensis*, Rolfe, 36, 23.  
*chinensis*, Lindl., 36, 24.  
*Lugardi*, Rolfe, 36, 24.  
*Pholas*, Reichb. f., 36, 24.  
*Pholidota yunnanensis*, Rolfe \*, 36, 24.  
*Photinia Benthamiana*, Hance, 23, 262.  
*crenatoserata*, Hance, 23, 262.  
*deflexa*, Hemsl., 36, 500.  
*Fortuneana*, Maxim., 23, 262.  
*glabra*, Hort., 23, 263.  
*glabra*, Maxim., 23, 262.  
 — *var.* *chinensis*, Maxim., 23, 263.  
*lævis*, DC., 23, 263.  
*melanostigma*, Hance, 23, 262.  
*prunifolia*, Lindl., 23, 262.  
*pustulata*, Lindl., 23, 263.  
*pustulata*, S. Moore, 23, 263.  
*serrulata*, Lindl., 23, 263; mentioned, 262.  
 — *var.* *prunifolia*, Hook. & Arn., 23, 262.  
*serrulata*, Sieb. & Zucc., 23, 263.  
*variabilis*, Hemsl. \*, 23, 263; mentioned, 264.  
*villosa*, DC., 23, 263.  
 — *var.* *formosana*, Hance, 26, 263.  
*Wrightiana*, Maxim., 36, 500.  
*Phragmites communis*, Trin., 36, 409.  
*Karka*, Trin., 36, 410.  
*longivalvis*, Steud., 36, 409.  
*Roxburghii*, Steud., 36, 410.  
*Phreatia elegans*, Lindl., 36, 17.  
*formosana*, Rolfe, 36, 17.  
*Phryma Leptostachya*, Linn., 26, 251.  
*Phrynium capitatum*, Willd., 36, 73.  
*parviflorum*, Roxb., 36, 73.  
*sindicum*, Miq., 36, 73.  
*Phtheirospermum chinense*, Bunge, 26, 204.  
*tenuisectum*, Bur. & Franch., 36, 500.  
*Phyla chinensis*, Lour., 26, 251.  
*Phyllamphora mirabilis*, Lour., 26, 358.  
*Phyllanthus anceps*, Benth., 26, 420.  
*anceps*, Vahl, 26, 423.  
*Arnottianus*, Muell. Arg., 26, 424.  
*bacciformis*, Linn., 26, 420.  
*Benthamianus*, Muell. Arg., 26, 425.  
*cantonensis*, Hornem., 26, 423.  
*cantonensis*, Schweigg., sp. dub., 26, 421.  
*cantonensis*, Zipp., 26, 421.  
*cinerascens*, Hook. & Arn., 26, 421.  
*cochinchinensis*, Spreng., 26, 421.  
*Emblica*, Linn., 26, 421.  
*eriocarpus*, Muell. Arg., 26, 424.  
*fasciculatus*, Muell. Arg., 26, 421.

- Phyllanthus flexuosus*, *Muell. Arg.*, 26, 421.  
*fluggeiformis*, *Muell. Arg.*, 26, 422.  
*glaucus*, *Wall.*, 26, 421.  
*hongkongensis*, *Muell. Arg.*, 26, 424.  
*japonicus*, *Muell. Arg.*, 26, 421.  
*lepidocarpus*, *Sieb. & Zucc.*, 26, 423.  
*leprocarpus*, *Wight*, 26, 423.  
*leptoclados*, *Benth.*, 26, 422.  
*lucens*, *Poir.*, 26, 427.  
*maderaspatensis*, *Linn.*, 26, 422.  
*microcarpus*, *Muell. Arg.*, 26, 422.  
*myrtifolius*, *Wall.*, 26, 422.  
*Niruri*, *Linn.*, 26, 422.  
*obovatus*, *Muell. Arg.*, 26, 425.  
*obscurus*, *Willd.*, 26, 425.  
*philippinensis*, *Muell. Arg.*, 26, 426.  
*puberulus*, *Muell. Arg.*, 26, 425.  
— var. *Fortunei*, *Muell. Arg.*, 26, 424.  
*ramiflorus*, *Pers.*, 26, 426.  
*reticulatus*, *Poir.*, 26, 422.  
*rhamnoides*, *Retz.*, 26, 428.  
*simplex*, *Retz.*, 26, 423; mentioned, 421.  
— var. *chinensis*, *Muell. Arg.*, 26, 420, 423.  
— var. *ussuriensis*, *Muell. Arg.*, 26, 423.  
*sinensis*, *Muell. Arg.*, 26, 422.  
*turbinatus*, *Sims*, 26, 427.  
*urinaria*, *Linn.*, 26, 423; mentioned, 424.  
*ussuriensis*, *Rupr.*, 26, 422, 423.  
*villosus*, *Poir.*, mentioned, 26, 423.  
*Wrightii*, *Muell. Arg.*, 26, 426.  
*Zippelianus*, *Muell. Arg.*, 26, 421.  
*Phyllobœa sinensis*, *Oliver*, 26, 225.  
*Phyllodes placentaria*, *Lour.*, 36, 73.  
*Phyllodesmis*, *Van Tiegh.*, 36, 500.  
*coriacea*, *Van Tiegh.*, 36, 500.  
*Delavayi*, *Van Tiegh.*, 36, 500.  
*paucifolia*, *Van Tiegh.*, 36, 500.  
*Phyllodium elegans*, *Desv.*, 23, 171.  
*pulchellum*, *Desv.*, 23, 175.  
*Phyllolobium chinense*, *Fisch.*, 23, 165.  
*Phyllostachys bambusoides*, *Sieb. & Zucc.*, 36, 438.  
*congesta*, *Rendle \**, 36, 438.  
*Faberi*, *Rendle \**, 36, 439.  
*flexuosa*, *A. & C. Rivière*, 36, 440.  
*Henonis*, *Mitford*, 36, 443.  
*Henryi*, *Rendle \**, 36, 440.  
*heteroclada*, *Oliver*, 36, 441.  
*montana*, *Rendle \**, 36, 441.  
*nana*, *Rendle \**, 36, 441.  
*Phyllostachys Nevinii*, *Hance*, 36, 442; mentioned, 440.  
— var. *hupehensis*, *Rendle \**, 36, 442.  
*nidularia*, *Munro*, 36, 442.  
*nigra*, *Munro*, 36, 443; mentioned, 442.  
— var. *Henonis*, *Stapf \**, 36, 443.  
*nigra*, *Pilger*, 36, 442.  
*quadrangularis*, *Rendle \**, 36, 443.  
*Stauntoni*, *Munro*, 36, 443; mentioned, 440.  
*Veitchiana*, *Rendle \**, 36, 443; mentioned, 441.  
*viridi-glaucescens*, *A. & C. Rivière*, 36, 444.  
*Physalis Alkekengi*, *Linn.*, 26, 173.  
*angulata*, *Linn.*, 26, 173.  
*ciliata*, *Sieb. & Zucc.*, 26, 173.  
*Francheti*, *Mast.*, 36, 500.  
*minima*, *Linn.*, 26, 174; mentioned, 173.  
*peruviana*, *Linn.*, mentioned, 26, 169.  
*pubescens*, *Linn.*, mentioned, 26, 174.  
*Physocarpus anomala*, *Batalin*, 36, 500.  
*Physochlaina physaloides*, *G. Don*, 26, 176.  
*Physolychnis apetala*, *Rohrb.*, 23, 65.  
*Physurus chinensis*, *Rolfe*, 36, 41.  
*Phyteuma bipinnata*, *Lour.*, 23, 348.  
*campanuloides*, *Bieb.*, mentioned, 26, 10.  
*japonicum*, *Miq.*, 26, 10.  
*Phytolacca acinosa*, *Roxb.*, 26, 331.  
*decandra*, *Maxim.*, 26, 331.  
— var. *acinosa*, *Moq.*, 26, 331.  
*Kæmpferi*, *A. Gray*, 26, 331.  
*octandra*, *Bunge*, 26, 331.  
*pekinensis*, *Hance*, 26, 332.  
*polyandra*, *Batalin*, 36, 500.  
*Picea ajanensis*, *Fisch.*, 26, 553.  
*Alcockiana*, *Carrière*, 26, 553.  
*brachytila*, *Mast.*, 26, 553.  
*likiangensis*, *Mast.*, 26, 554.  
*Maximowiczii*, *Regel*, 26, 554.  
*Neoveitchii*, *Mast.*, 36, 500.  
*obovata*, *Ledeb.*, 26, 554.  
— subvar. *Loczyi*, *Kanitz*, 26, 554.  
— var. *Schrenkiana*, *Mast. \**, 26, 554.  
*Schrenkiana*, *Fisch. & Mey.*, 26, 554.  
*Veitchii*, *A. Murr.*, 26, 557.  
*Wilsoni*, *Mast.*, 36, 500.



- Pierasma ailanthoides*, Planch., 23, 112.  
*japonica*, A. Gray, 23, 112.  
*quassioides*, Benn., 23, 112.  
*Picria Fel-terrae*, Lour., 26, 187.  
*Pieris divaricata*, Vaniot, 36, 500.  
*hieracioides*, Linn., 23, 474; 36, 500.  
*japonica*, Thunb., 23, 474.  
*repens*, Lour., 23, 474.  
*Piddingtonia Nummularia*, DC., 26, 6.  
*Pieris Bodinieri*, H. Léveillé, 36, 501.  
*divaricata*, H. Léveillé, 36, 501.  
*Duclouxii*, H. Léveillé, 36, 501.  
*formosa*, D. Don, 26, 16.  
*Gagnepainiana*, H. Léveillé, 36, 501.  
*japonica*, D. Don, 26, 17.  
*konyangensis*, H. Léveillé, 36, 301.  
*Martini*, H. Léveillé, 36, 501.  
*ovalifolia*, D. Don, 26, 17.  
— *var. hebecarpa*, Franch. \*, 26, 17.  
— *var. lanceolata*, C. B. Clarke, 26, 17.  
*repens*, H. Léveillé, 36, 501.  
*Swinhoei*, Hemsl. \*, 26, 17.  
*Pilea angulata*, Blume, mentioned, 26, 478, 479.  
*anisophylla*, Wedd., 26, 475.  
*bracteosa*, Wedd., 26, 475.  
*crassifolia*, Hance, 26, 476.  
*fasciata*, Franch., 36, 501.  
*Goglado*, Wedd., 26, 478.  
*Hookeriana*, Wedd., 26, 476.  
*microphylla*, Liebm., 26, 476.  
*mongolica*, Wedd., 26, 476.  
*mongolica*, Wedd., 26, 477.  
*muscosa*, Lindl., 26, 476.  
*notata*, C. H. Wright \*, 26, 476.  
*peltata*, Hance, 26, 477; mentioned, 480.  
*peploides*, Hook. & Arn., 26, 477.  
*petiolaris*, Franch. & Sav., 26, 478.  
*petiolaris*, Wedd., 26, 478.  
*platyflora*, C. H. Wright \*, 26, 478.  
*pumila*, A. Gray, 26, 477.  
*rubriflora*, C. H. Wright \*, 26, 478.  
*scripta*, Wedd., 26, 478.  
*spp.*, 26, 479, 480.  
*stipulosa*, Miq., 26, 479.  
*Synmeria*, Wedd., 26, 479.  
*trinervia*, Wight, 36, 501.  
*umbrosa*, Wedd., 26, 479.  
*Wattersii*, Hance, 26, 479.  
*Pilea Wightii*, Wedd., mentioned, 26, 479.  
*Pileostegia viburnoides*, Hook. f. & Thoms., 23, 275.  
*Pimela alba*, Lour., 23, 113.  
*nigra*, Lour., 23, 113.  
*Pimpinella*, Diels, = *Carum*, 36, 501.  
*arguta*, Diels, 36, 501.  
*diversifolia*, DC., 23, 329.  
*Dunnii*, Boissieu, 36, 501.  
*Fargesii*, Boissieu, 36, 501.  
*Fauriei*, Boissieu, 36, 501.  
*flicina*, Franch., 36, 501.  
*helosciadoidea*, Boissieu, 36, 501.  
*Henryi*, Diels, 36, 501.  
*nikoensis*, Yabe, 36, 501.  
*rhomboidea*, Diels, 36, 501.  
*Rosthornii*, Diels, 36, 501.  
*silatifolia*, Boissieu, 36, 501.  
*sinica*, Hance, 23, 329.  
*Souliei*, Boissieu, 36, 501.  
*sutchuensis*, Boissieu, 36, 501.  
*Tanakæ*, Franch. & Sav., 36, 501.  
*trichomanifolia*, Franch., 36, 501.  
*triternata*, Diels, 36, 501.  
*Pinellia cordata*, N. E. Br. \*, 36, 173.  
*integrifolia*, N. E. Br., 36, 174.  
*pedatisecta*, Schott, 36, 174.  
*tripartita*, Schott, 36, 174.  
*tuberifera*, Tenore, 36, 174.  
— *var. pedatisecta*, Engl., 36, 174.  
— *var. subpandurata*, Engl., 36, 175.  
*Wawrae*, Engl., 36, 174.  
*Pinguicula vulgaris*, Linn., 36, 501.  
*Pinus*, genus referred to, 26, 548.  
*Armandi*, Franch., 26, 549.  
*Bungeana*, Zucc., 26, 549.  
*Cembra*, var. *pumila*, Fall., 26, 551.  
*chinensis*, Beissner, 26, 559.  
*densiflora*, Hort., 26, 553.  
*densiflora*, Sieb. & Zucc., 26, 549; mentioned, 550.  
— *var. tabuliformis*, Fortune \*, 26, 549.  
*excorticata*, Hort. 26, 549.  
*Fortunei*, Parl., 26, 555.  
*funebri*, Komarov, 36, 501.  
*Gerardiana*, Wall., 26, 550.  
*Griffithiana*, Parl., 26, 558.  
*Henryi*, Hemsl. \*, 26, 550.  
*holophylla*, Parl., 26, 557.  
*koraiensis*, Sieb. & Zucc., 26, 550.  
*lanceolata*, Lamb., 26, 548.  
*luchuensis*, Mayr, 26, 551.

- Pinus mandshurica*, Rupr., 26, 551.  
*Massoniana*, Lamb., 26, 551; mentioned, 552.  
 — *var. planiceps*, A. Murr. \*, 26, 551, 552.  
*Massoniana*, Sieb. & Zucc., 26, 553.  
*parviflora*, Sieb. & Zucc., mentioned, 26, 550.  
*scipioniformis*, Mast., 26, 552.  
*selenolepis*, Parl., 26, 557.  
*silvestris*, Thunb., 26, 551, 553.  
*sinensis*, Benth., 26, 551.  
*sinensis*, Lamb., 26, 552, 553.  
*Thunbergiana*, Parl., 26, 552.  
*Thunbergii*, Parl., 26, 552.
- Piper arcuatum*, Seem., 26, 366.  
 — *var.*, Miq., 26, 365.  
*aurantiacum*, Wall., 26, 364.  
*Betle*, Linn., mentioned, 26, 364.  
*Chaba*, Hunter, 26, 364.  
*chandocanum*, C. DC., 36, 501.  
*chinense*, Miq., 26, 365.  
*Ferriei*, C. DC., 36, 501.  
*fragile*, Benth., 26, 366.  
*Futokadsura*, Sieb., 26, 365.  
*hainanense*, Hemsl. \*, 26, 365.  
*Hancei*, Maxim., 26, 365.  
*Harmandi*, C. DC., 36, 501.  
*Henryi*, C. DC., 36, 501.  
*hongkongense*, C. DC., 26, 366.  
*laosanum*, C. DC., 36, 501.  
*longum*, Linn., mentioned, 26, 364.  
*nigrum*, Linn., mentioned, 26, 364.  
*officinarum*, C. DC., 26, 364.  
*philippinum*, Miq., mentioned, 26, 365.  
*pinnatum*, Lour., 23, 108.  
*puberulum*, Maxim., 26, 365.  
*sarmentosum*, Roxb., 26, 366.  
*sinense*, C. DC., 26, 366.  
*sp. n. ?*, 26, 366.
- PIPERACEÆ, 26, 363.
- Piptanthus nepalensis*, D. Don, 36, 501.  
*tomentosus*, Franch., 36, 501.
- Pircunia Latbenia*, Moq., 26, 332.
- Pisonia aculeata*, Linn., 26, 317.
- Pistacia weinmannifolia*, J. Poisson, 36, 501.
- Pistia Stratiotes*, Linn., 36, 175.
- Pisum maritimum*, Linn., 23, 186.  
*sativum*, Linn., mentioned, 23, 187.
- Pithecolobium Clypearia*, Benth., 23, 216.  
*dulce*, Benth., 23, 217.
- Pithecolobium lucidum*, Benth., 23, 217.
- PITTOSPOREÆ, 23, 58.
- Pittosporum Fortunei*, Turcz., 23, 59.  
*glabratum*, Lindl., 23, 58; 36, 501.  
*heterophyllum*, Franch., 36, 501.  
*nilghirensense*, Wight & Arn., 36, 502.  
*pauciflorum*, Hook. & Arn., 23, 58.  
*Tobira*, Ait., 23, 58.  
*truncatum*, E. Pritz., 36, 502.  
*yunnanensis*, Franch., 36, 502.
- Plagiorhegma dubium*, Maxim., 23, 33.
- Plagiospermum sinense*, Oliver, 36, 502.
- Plagiostigma pumila*, Zucc., 26, 466.  
*stipulata*, Zucc., 26, 466.
- Planera acuminata*, Lindl., 26, 449.  
*Davidii*, Hance, 26, 449.  
*japonica*, Miq., 26, 449.  
*parvifolia*, Sweet, 26, 448.  
*sp. nov. ?*, Maxim., 26, 449.
- PLANTAGINEÆ, 26, 316.
- Plantago asiatica*, Linn., 26, 316.  
*depressa*, Willd., 26, 316.  
*exaltata*, Hornem., 26, 316.  
*lanceolata*, Linn., 26, 316.  
*Loureiri*, Roem. & Schult., 26, 316.  
*major*, Linn., 26, 316.  
*media*, Linn., 26, 316.
- Platanthera Championi*, Lindl., 36, 56.  
*chlorantha*, Cust., 36, 55.  
*dentata*, Lindl., 36, 59.  
*flava*, Lindl., 36, 55.  
*Galeandra*, Reichb. f., 36, 56.  
*Henryi*, Rolfe \*, 36, 55.  
*hologlottis*, Maxim., 36, 55.  
*iantha*, Wight, 36, 55.  
*interrupta*, Maxim., 36, 56.  
*japonica*, Lindl., 36, 56.  
*Keiskei*, Franch., 36, 56.  
*Mandarinorum*, Reichb. f., 36, 56; mentioned, 55.  
*manubriata*, Kränzl., 36, 502.  
*obcordata*, Lindl., 36, 56.  
*oreades*, Franch., 36, 56.  
*robusta*, Lindl., 36, 57.  
*setchuenica*, Kränzl., 36, 502.  
*stenostachya*, Lindl., 36, 56.  
*Susannæ*, Lindl., 36, 57.  
*tipuloides*, var. *ussuriensis*, Regel, 26, 57.  
*ussuriensis*, Maxim., 26, 57.  
*viridis*, Lindl., 26, 54.

- Platanthera viridis*, var. *bracteata*, Reichb. f., **36**, 54.  
*Platycarya strobilacea*, Sieb. & Zucc., **26**, 495.  
*Platycodon autumnalis*, Decne., **26**, 5.  
     *chinensis*, Lindl., **26**, 5.  
     *grandiflorus*, A. DC., **26**, 5.  
     — var. *Candolleanus*, Franch., **26**, 5.  
     — var. *chinensis*, Franch., **26**, 5.  
     *homallanthinum*, A. DC., **26**, 5.  
     *sinensis*, Lem., **26**, 5.  
*Platyrrhapha japonica*, Miq., **23**, 329.  
*Plecosperrum cuneifolium*, Thw., **26**, 469.  
*Plecostigma pauciflorum*, Turcz., **36**, 138.  
*Plectranthus amethystoides*, Benth., **26**, 269; mentioned, 274.  
     *Buergeri*, Miq., **26**, 271.  
     *calcaratus*, Hemsl., **36**, 502.  
     *cardiophyllus*, Hemsl. \*, **26**, 269.  
     *carnosifolius*, Hemsl. \*, **26**, 270.  
     *cordifolius*, D. Don, **26**, 270.  
     *dichromophyllus*, Diels, **36**, 502.  
     *excisus*, Maxim., **26**, 270.  
     *Gerardianus*, Benth., **26**, 270; mentioned, 274.  
     *glaucocalyx*, Maxim., **26**, 271; mentioned, 272.  
     *graciliflorus*, Benth., **26**, 271.  
     *Henryi*, Hemsl. \*, **26**, 271.  
     *incanus*, Link, mentioned, **26**, 270.  
     *inconspicuus*, Miq., **26**, 272.  
     *inflexus*, Vahl, **26**, 272.  
     *leptobotrys*, Diels, **36**, 502.  
     *marmoritis*, Hance, **26**, 272.  
     *Maximowiczii*, Miq., **26**, 271.  
     *nankinensis*, Spreng., **26**, 279.  
     *nervosus*, Hemsl. \*, **26**, 272.  
     *nudipes*, Hemsl. \*, **26**, 272.  
     *Patchouli*, C. B. Clarke, **26**, 306.  
     *racemosus*, Hemsl. \*, **26**, 273.  
     *Rosthornii*, Diels, **36**, 502.  
     *rubescens*, Hemsl. \*, **26**, 273.  
     *sinensis*, Miq., **26**, 274.  
     *striatus*, Benth., mentioned, **26**, 271.  
     *Tatei*, Hemsl. \*, **26**, 274.  
     *ternifolius*, D. Don, **26**, 275.  
     *Websteri*, Hemsl. \*, **26**, 275.  
*Plectronia chinensis*, Lour., **23**, 339.  
     *didyma*, Kurz, **23**, 385.  
     *parvifolia*, Kurz, **23**, 385.  
*Pleiogyne anthemoides*, C. Koch, **23**, 439.  
     *cardiosperma*, Edgew., **23**, 439.  
*Pleione bulbocodioides*, Rolfe, **36**, 502.  
     *Delavayi*, Rolfe, **36**, 502.  
     *grandiflora*, Rolfe, **36**, 502.  
     *pogonioides*, Rolfe, **36**, 502.  
     *yunnanensis*, Rolfe, **36**, 502.  
*Pleurobotrys* (Teucris n. §), Hemsl. \*, **26**, 311; mentioned, 314.  
*Pleurogyna carinthiaca*, G. Don, var., **36**, 502.  
     *rotata*, Griseb., **26**, 138.  
     *rotata*, Sieb. & Zucc., **26**, 139.  
*Pleuroplitis Langsdorffii*, Trin., **36**, 360.  
     — var. *brevisetata*, Regel, **36**, 360.  
     — var. *japonica*, Regel, **36**, 360.  
     — var. *typica*, Regel, **36**, 360.  
*Pleuropterus cordatus*, Turcz., **26**, 343.  
*Pleurospermum austriacum*, Hoffm., **23**, 333.  
     *Davidi*, Franch., **36**, 502.  
     *decurrens*, Franch., **36**, 502.  
     *dentatum*, Benth., mentioned, **36**, 333.  
     *foetens*, Franch., **36**, 502.  
     *Franchetianum*, Hemsl., **36**, 502.  
     *Giraldii*, Diels, **36**, 502.  
     *Govanianum*, Benth., **36**, 502.  
     *kamtschaticum*, Hoffm., **23**, 333.  
     *meoides*, Diels, **36**, 502.  
     *nanum*, Franch., **36**, 502.  
     *Pulzkyi*, Kanitz, **36**, 502.  
     sp. ?, **23**, 333.  
     *Szechenyi*, Kanitz, **36**, 502.  
     *uralense*, Hoffm., **23**, 333.  
     *Wrightianum*, Boissieu, **36**, 502.  
     *yunnanense*, Franch., **36**, 502.  
*Plocoglottis* ? sp., **36**, 32.  
*Pluchea Dioscoridis*, DC., **23**, 422.  
     *hirsuta*, Less., mentioned, **23**, 422.  
     *indica*, Less., **23**, 422.  
     *pteropoda*, Hemsl. \*, **23**, 422.  
*Plumbagidium roseum*, Spach, **26**, 36.  
**PLUMBAGINEÆ**, **26**, 34.  
*Plumbago ceylanica*, Bretschn., **26**, 36.  
     *coccinea*, Boiss., **26**, 36.  
     *Larpenæ*, Lindl., **26**, 36.  
     *rosea*, Linn., **26**, 36.  
     — var. *coccinea*, Hook. f., **26**, 36.  
     *zeylanica*, Linn., **26**, 36.  
*Plumeria acutifolia*, Poir., mentioned, **26**, 95.  
     *obtusata*, Lour., **26**, 95.  
*Poa acroleuca*, Steud., **36**, 422.  
     *annua*, Linn., **36**, 422.  
     — var. *nepalensis*, Griseb., **36**, 422.

- Poa attenuata*, Trin., 36, 423.  
 — *var. vivipara*, Rendle\*, 36, 423.  
*chinensis*, Linn., 36, 407.  
*Faberi*, Rendle\*, 36, 424.  
 — *var. ligulata*, Rendle\*, 36, 424.  
*ferruginea*, Thunb., 36, 413.  
*gracillima*, Rendle\*, 36, 424.  
*japonica*, Thunb., 36, 415.  
*linearis*, Trin., 36, 427.  
*Mariesii*, Rendle\*, 36, 425.  
*megastachya*, Koel., 36, 416.  
*nemoralis*, Linn., 36, 425.  
 — *var.*, Hance, 36, 423.  
 — *var. ligulata*, Stapf, 36, 426.  
 — *var. mongolica*, Rendle\*, 36, 426.  
*nepalensis*, Wall., 36, 423.  
*orientalis*, Franch., 36, 413.  
*palustris*, Linn., *var. strictula*, Hack., 36, 427.  
*pilosa*, Linn., 36, 417.  
*plumosa*, Retz., 36, 418.  
*pratensis*, Linn., 36, 426; mentioned, 425.  
 — *var. angustifolia*, Gaud., 36, 426.  
*procumbens*, Curt., 36, 429.  
*prolixior*, Rendle\*, 36, 427.  
*sphondylodes*, Trin., 36, 427.  
*strictula*, Steud., 36, 427.  
*unioloides*, Retz., 36, 412.  
*viridula*, Palibin, 36, 428.  
*Podanthum*, Boiss., 26, 10.  
*Podocarpus argotaenia*, Hance, 26, 547.  
*bracteata*, Blume, 26, 548.  
*chinensis*, Wall., 26, 547.  
*insignis*, Hemsl., 26, 547.  
*koraiana*, Sieb., 26, 546.  
*koreana*, Hort., 26, 546.  
*latifolia*, Wall., 26, 547.  
*macrophylla*, Wall., 26, 548; mentioned, 547.  
*Maki*, Sieb. & Zucc., 26, 547.  
*Makoya*, Hort., 26, 547.  
*Makoyæ*, Hort., 26, 547.  
*Makoyi*, Hort., 26, 547.  
*Miquelia*, Hort., 26, 547.  
*Nageia*, Henry, 26, 548.  
*neriifolia*, D. Don, 26, 548.  
*sinensis*, Teysm. & Binn., 26, 547.  
*sutchuenensis*, Franch., 26, 548.  
*Podochilus cornutus*, Schlechter, 36, 502.  
*Podoon Delavayi*, Baill., 36, 502.  
 PODOONACEÆ, H. Baill., 36, 502.  
*Podophyllum Delavayi*, Franch., 36, 502.  
*pleianthum*, Hance, 23, 33.  
*Pogonatherum crinitum*, Kunth, 36, 357.  
*polystachyum*, Nees, 36, 358.  
*polystachyum*, Roem. & Schult., 36, 357.  
*refractum*, Nees, 36, 358.  
*rufobarbatum*, Griff., 36, 358 in obs.  
*saccharoideum*, Beauv., 36, 357.  
 — *var. monandrum*, Hack., 36, 357.  
*saccharoideum*, Benth., 36, 358.  
*Pogonia Fordii*, Hance, 36, 47.  
*japonica*, Reichb. f., 36, 47.  
*lanceolata*, Kränzl., 36, 502.  
*ophioglossoides*, Ker, 36, 47.  
*pleionoides*, Kränzl., 36, 502.  
*pulchella*, Hook. f., 36, 47.  
*similis*, Blume, 36, 47.  
*sp.*, Hance, 36, 23.  
*yunnanensis*, Finet, 36, 47.  
*Pogonopsis tenera*, Presl, 36, 358.  
*Pogonostylis squarrosa*, Bertol., 36, 214.  
*Pogostemon formosanus*, Oliver, 36, 502.  
*ianthinus*, Kanitz, 36, 503.  
*parviflorus*, Benth., 26, 276.  
*Polanisia viscosa*, DC., 23, 50.  
 POLEMONIACEÆ, 26, 142.  
*Polemonium cæruleum*, Linn., 26, 142.  
*Poliothyrsis*, Oliver, 36, 503.  
*sinensis*, Oliver, 36, 503.  
*Pollia japonica*, Hance, 36, 159.  
*sorzogonensis*, Endl., 36, 159.  
 — *var. gigantea*, C. B. Clarke, 36, 159.  
*Zollingeri*, C. B. Clarke, 36, 159.  
*Pollinia articulata*, Trin., 36, 353.  
*cantonensis*, Rendle\*, 36, 354.  
*ciliata*, Trin., 36, 354.  
 — *var. breviaristata*, Rendle\*, 36, 355.  
*Cumingii*, Nees, 36, 355.  
*eriopoda*, Hance, 36, 364.  
*glabrata*, Trin., 36, 355.  
*imberbis*, Nees, 36, 355; mentioned, 354.  
 — *var. Willdenowiana*, Hack., 36, 355.  
*imberbis*, Nees, partim, 36, 355.  
*japonica*, Franch., 36, 355.  
*japonica*, Miq., 36, 356.  
*monantha*, Nees, 36, 355.  
*nuda*, Trin., 36, 356.  
*pallens*, Hack., 36, 356.  
*phæothrix*, Hack., 36, 356.  
*quadrinervis*, Hack., 36, 356.  
 — *var. latifolia*, Rendle\*, 36, 357.

- Pollinia speciosa*, Hack., **36**, 357.  
*tenuis*, Benth., **36**, 354.  
*vagans*, Nees, mentioned, **36**, 354.  
*velutina*, Hack., **36**, 357.  
*villosa*, Munro, **36**, 356.  
— var. *chefuensis*, Franch., **36**, 356.
- Polyalthia suberosa*, Benth. & Hook. f., **23**, 27.
- Polycarpaea corymbosa*, Lam., **23**, 71.
- Polycarpon Loetlingiae*, Edgew. & Hook. f., **23**, 71.
- Polychroa repens*, Lour., **26**, 482.
- Polycælium chinense*, A. DC., **26**, 250.
- Polygala arillata*, Buch.-Ham., **23**, 59.  
*chinensis*, Linn., mentioned, **23**, 59.  
*erotalarioides*, Buch.-Ham., **36**, 503.  
*cyanolopha*, Hance, **23**, 62.  
*elegans*, Benth., **23**, 62.  
*fallax*, Hemsl. \*, **23**, 59.  
*floribunda*, Dunn, **36**, 503.  
*Forbesii*, Chod., **36**, 503.  
*globulifera*, Dunn, **36**, 503.  
*glomerata*, Lour., **23**, 60.  
*hongkongensis*, Hemsl. \*, **23**, 60, 490; mentioned, 61.  
*japonica*, Houtt., **23**, 62.  
*Karensium*, Kurz, **23**, 61.  
*Latouchei*, Franch. & Finet, **36**, 503.  
*Loureirii*, Gardn., **23**, 62.  
*Mariesii*, Hemsl. \*, **23**, 61, 490.  
*persicarioides*, DC., **36**, 503.  
*saxicola*, Dunn, **36**, 503.  
*sibirica*, Linn., **23**, 61; **36**, 503.  
— var. *angustifolia*, Ledeb., **23**, 62.  
— var. *stricta*, Debeaux, **23**, 62.  
— var. *tenuifolia*, Baker & S. Moore, **23**, 62.  
*sibirica*, Lour., var., **23**, 61.  
*Tatarinowii*, Regel, **23**, 62.  
*telephioides*, Willd., **23**, 62.  
*tenuifolia*, Willd., **23**, 62.  
*triphylla*, Buch.-Ham., **23**, 62.  
*venenosa*, Juss., **23**, 60.  
*Wattersii*, Hance, **23**, 63; mentioned, 61.
- POLYGALEÆ**, **23**, 59.
- POLYGONACEÆ**, **26**, 332.
- Polygonatum*, Tourn., **23**, 395.  
*agglutinatum*, Hua, **36**, 104.  
*anomalum*, Hua, **36**, 104.  
*Bodinieri*, H. Léveillé, **36**, 503.  
*canaliculatum*, Miq., **36**, 106.  
*chinense*, Kunth, **36**, 104, 109.
- Polygonatum cirrhifolium*, Royle, **36**, 104; mentioned, 109.  
*commutatum*, Dietr., **36**, 106.  
*curvistylum*, Hua, **36**, 104.  
*cyrtoneura*, Hua, **36**, 104.  
*Delavayi*, Hua, **36**, 104.  
*ensifolium*, H. Léveillé, **36**, 503.  
*erythrocarpum*, Hua, **36**, 105.  
*falcatum*, A. Gray, **36**, 503.  
*Fargesii*, Hua, **36**, 105.  
*Franchetii*, Hua, **36**, 105.  
*fuscum*, Hua, **36**, 105.  
*giganteum*, Dietr., **36**, 106.  
*Govanianum*, Royle, **36**, 106.  
*Henryi*, Diels, **36**, 503.  
*Hookeri*, Baker, **36**, 105.  
*humile*, Fisch., **36**, 503.  
*inflatum*, Komarov, **36**, 503.  
*japonicum*, Morren, **36**, 107.  
*kansuense*, Maxim., **36**, 105.  
*Kingianum*, Collett, **36**, 105.  
*latifolium*, Desf., 106, 107.  
— var. *commutatum*, Baker, **36**, 106.  
*macropodium*, Turcz., **36**, 106.  
*maiale*, Allioni, **36**, 112.  
*Martini*, H. Léveillé, **36**, 503.  
*Maximowiczii*, F. Schmidt, **36**, 108.  
*multiflorum*, Allioni, **36**, 106, 107.  
*nodosum*, Hua, **36**, 107.  
*officinale*, Allioni, **36**, 107.  
— var. *humile*, Baker, mentioned, **36**, 104.  
— var. *japonicum*, Maxim., **36**, 107.  
— var. *macranthum*, Hua, **36**, 108.  
— var. *Maximowiczii*, Maxim., **36**, 108.  
— var. *papillosum*, Franch., **36**, 108.  
*platyphyllum*, Franch., **36**, 108.  
*Prattii*, Baker, **36**, 108.  
*pumilum*, Hua, **36**, 108.  
*roseum*, Kunth, **36**, 108.  
*roseum*, Maxim., **36**, 108.  
*sibiricum*, Baker, **36**, 104; mentioned, 105.  
*sibiricum*, Red., **36**, 109.  
*Souliei*, Hua, **36**, 109.  
*sp.*, Maxim., **36**, 108.  
*Thunbergii*, Morren, **36**, 106.  
*trinerve*, Hua, **36**, 109.  
*umbellatum*, Baker, **36**, 106.  
*verticillatum*, Allioni, **36**, 109; mentioned, 108.  
*vulgare*, Desf., **36**, 107.  
— var. *japonicum*, Franch. & Sav., **36**, 108.

- Polygonum acaule*, Franch. & Sav., 36, 503.  
*acutatum*, Lehm., 26, 332.  
*alatum*, Buch.-Ham., 26, 332.  
*alpinum*, Allioni, 26, 332; mentioned, 342.  
— var., 36, 503.  
*ambiguum*, Meisn., 26, 333.  
*amphibium*, Linn., 26, 333; 36, 503.  
— var. *vestitum*, Hemsl. \*, 26, 333.  
*anplexicaule*, D. Don, 26, 333.  
— var. *sinense*, Hook. f., 26, 333.  
*arifolium*, Linn., mentioned, 26, 338.  
*aviculare*, Linn., 26, 334.  
— var. *minutiflorum*, Franch., 26, 346.  
*aviculare*, Lour., 26, 346.  
*Babingtonii*, Hance, 26, 349.  
*barbatum*, Linn., 26, 334; mentioned, 349, 353.  
*Bistorta*, Linn., 26, 334; mentioned, 336.  
— var. *angustifolium*, Meisn., 26, 336.  
— var. *minor*, Franch. & Sav., 26, 350.  
*bistortoides*, Boiss., 26, 336.  
*Blumei*, Meisn., 26, 334; mentioned, 349.  
*Bodinieri*, H. Lévillé & Vaniot, 36, 503.  
*Bungeanum*, Turcz., 26, 335.  
*cæspitosum*, Blume, 26, 346.  
*campanulatum*, Hook. f., mentioned, 26, 346.  
*capitatum*, Buch.-Ham., 26, 335.  
*carneum*, Koch, 26, 336.  
*chinense*, Houtt., 26, 343.  
*chinense*, Linn., 26, 335.  
— var. *Thunbergianum*, Meisn., 26, 335.  
*ciliatum*, Lour., 26, 336.  
*ciliosum*, Meisn., 26, 346.  
*cliffortioides*, Meisn., 26, 346.  
*cochinchinensis*, Meisn., 26, 343.  
*compactum*, Hook. f., mentioned, 26, 337.  
*confusum*, Meisn., 26, 336; mentioned, 345.  
*conspersum*, Meisn., 26, 350.  
*criopolitanum*, Hance, 26, 336; mentioned, 347.  
*cuspidatum*, Sieb. & Zucc., 26, 336; mentioned, 337, 349.  
*cymosum*, Trev., 26, 337.  
*cynanchoides*, Hemsl. \*, 26, 338.  
*dentato-alatum*, F. Schmidt, 26, 348.
- Polygonum dibotrys*, D. Don, 26, 337.  
*dissitiflorum*, Hemsl. \*, 26, 338.  
*divaricatum*, Linn., 26, 339.  
*Donii*, Meisn., 26, 349.  
*Dumetorum*, Linn., 26, 339; mentioned, 348.  
*effusum*, Meisn., 26, 346.  
*emarginatum*, Roth, 26, 339.  
*Fagopyrum*, Linn., 26, 339; mentioned, 337, 340, 351.  
*filiforme*, Thunb., 26, 352.  
*flaccidum*, Meisn., 26, 339; mentioned, 332, 347, 349.  
— var. *ripularum*, Hance, 26, 339.  
*Forbesii*, Hance, 26, 339.  
*Gilesii*, Hemsl., mentioned, 26, 340.  
*glabrum*, Willd., 26, 340.  
— var. *fucatum*, Hance, 26, 340.  
*glutinosum*, Wall., 26, 341.  
*gracilipes*, Hemsl. \*, 26, 340.  
*hastato-trilobum*, Meisn., 26, 351.  
*hastatum*, Murr., 26, 349.  
*herniarioides*, Meisn., 26, 346.  
*Hydropiper*, Linn., 26, 340.  
*illecebroides*, Meisn., 26, 346.  
*interruptum*, Bunge, 26, 341.  
*japonicum*, Meisn., 26, 341; mentioned, 349.  
*juvendum*, Meisn., 26, 341.  
*Labordei*, H. Lévillé & Vaniot, 36, 503.  
*lanigerum*, R. Br., 26, 341.  
— var. *cristatum*, Hemsl. \*, 26, 342.  
— var. *glutinosum*, Hook. f., 26, 341.  
*lapathifolium*, Linn., 26, 342; mentioned, 345, 353.  
— var. *nodosum*, Hook. f., 26, 343.  
*Lourieri*, Poir., 26, 336.  
*Maackianum*, Regel, 26, 351.  
*Martini*, H. Lévillé & Vaniot, 36, 503.  
*Meyeni*, C. Koch, 26, 346.  
*minus*, Huds., 26, 342; mentioned, 332, 341.  
*Miquelianum*, Meisn., 26, 346.  
*multiflorum*, Thunb., 26, 342.  
*muricatum*, Meisn., 26, 343.  
*Myosurus*, Franch., 36, 503.  
*nepalense*, Meisn., 26, 332; 36, 503.  
*nodosum*, Pers., 26, 343; mentioned, 345.  
— var. *incanum*, Ledeb., 26, 343.

- Polygonum orientale*, Linn., 26, 343; mentioned, 350.  
 — var. *discolor*, Benth., 26, 342.  
*paleaceum*, Wall., mentioned, 26, 336.  
*panduriforme*, H. Léveillé & Vaniot, 36, 503.  
*pauciflorum*, Maxim., 36, 503.  
*pedunculare*, Wall., 26, 344.  
*pensylvanicum*, Bunge, 26, 335.  
*pensylvanicum*, Hance, 26, 342.  
*perfoliatum*, Linn., 26, 344.  
*perfoliatum*, Thunb., 26, 349.  
*pergracilis*, Hemsl. \*, 26, 344.  
*Perrotteti*, Meisn., 36, 346.  
*Persicaria*, Linn., 26, 345.  
*pilosum*, Maxim., 26, 345.  
*pilosum*, Roxb., 26, 343.  
*Pinetorum*, Hemsl. \*, 26, 345.  
*plebeium*, R. Br., 26, 346.  
*polymorphum*, Ledeb., 26, 333.  
 — var. *arenarium*, Debeaux, 26, 333.  
 — var. *salignum*, Franch., 26, 333.  
*polystachyum*, Wall., 26, 342.  
*Posumbu*, Buch.-Ham., 26, 346; mentioned, 332, 339, 349.  
*prætermisum*, Hook. f., 26, 347.  
*pteropus*, Hance, 26, 351.  
*punctatum*, Buch.-Ham., 26, 332.  
*radicans*, Hemsl. \*, 26, 347.  
*Roxburghii*, Meisn., 26, 343.  
 — var. *longifolium*, Meisn., 26, 346.  
*runcinatum*, Buch.-Ham., 26, 347.  
 — var. *sinense*, Hemsl. \*, 26, 347.  
*sagittatum*, Linn., 26, 348.  
*sagittifolium*, H. Léveillé & Vaniot, 36, 503.  
*scandens*, Linn., 26, 348.  
*senticosum*, Franch. & Sav., 26, 349.  
*serrulatum*, Lag., 26, 349; mentioned, 332, 335, 347.  
*sibiricum*, Laxm., 26, 349.  
*sibiricum*, Linn., 26, 333.  
*Sieboldi*, Hort., 26, 337.  
*Sieboldi*, Meisn., 26, 348.  
*Sinarum*, Desv., 26, 351.  
*sinense*, Hook. & Arn., 26, 335.  
*sinuatum*, R. Br., var., 36, 503.  
*sp.*, 26, 352.  
*Spaethii*, Dammer, 36, 503.  
*speciosum*, Meisn., 26, 333.  
*splendens*, Klotzsch, 26, 336.  
*stagninum*, Buch.-Ham., 26, 350.  
*stoloniferum*, F. Schmidt, 26, 351.  
*strigosum*, Benth., 26, 347.  
*Polygonum strigosum*, R. Br., 26, 350.  
*subcordatum*, Miq., 26, 350.  
*suffultum*, Maxim., 26, 350; 36, 503.  
*tataricum*, Linn., 26, 350.  
*Thunbergii*, Sieb. & Zucc., 26, 351.  
*tinctorium*, Lour., 26, 351.  
*triangulare*, Wall., 26, 337.  
*typhonifolium*, Hance, 26, 349.  
*urophyllum*, Bur. & Franch., 36, 503.  
*virginianum*, Linn., 26, 352.  
*viscosum*, Buch.-Ham., 26, 352.  
*viviparum*, Linn., 26, 352.  
*volubile*, Turcz., 26, 337.  
*Polyozus lanceolata*, Lour., 23, 395.  
*Polypara cochinchinense*, Lour., 26, 364.  
*Polyphema Jaca*, Lour., 26, 470.  
*Polypogon littoralis*, Sm., 36, 386.  
*monspeliensis*, Desf., 36, 386.  
*Polyspora axillaris*, Sweet, 23, 80.  
*Pomasterium japonicum*, Miq., 23, 320.  
*Pongamia chinensis*, DC., 23, 200.  
*glabra*, Vent., 23, 200.  
*Pongatium indicum*, Lam., 26, 8.  
*Pontederia dilatata*, Buch.-Ham., 36, 149.  
*hastata*, Linn., 36, 149.  
*linearis*, Hassk., 36, 150.  
*ovata*, Hook. & Arn., 36, 149.  
*ovata*, Linn., 36, 73.  
*plantaginea*, Roxb., 36, 150.  
*sagittata*, Roxb., 36, 149.  
*vaginalis*, Burm. f., 36, 149.  
 PONTEDERIACEÆ, 36, 148.  
*Populus acuta*, David, 26, 534.  
*adenopoda*, Maxim., 26, 537.  
*alba*, Linn., 26, 535.  
 — var. *tomentosa*, Wesmael, 26, 535.  
*balsamifera*, Linn., 26, 535.  
 — var. *laurifolia*, Wesmael, 26, 535.  
 — var. *Simonii*, Wesmael, 26, 536.  
 — var. *suaveolens*, Loud., 26, 535.  
*diversifolia*, Schrenck, 26, 536.  
*euphratica*, Oliv., 26, 536; mentioned, 537.  
*Fargesii*, Franch., 26, 536.  
*lasiocarpa*, Oliver, 26, 536.  
*laurifolia*, Ledeb., 26, 535.  
*nigra*, Linn., 26, 536.  
*pekinensis*, L. Henry, 36, 503.  
*Przewalskii*, Maxim., 26, 537.  
*Simonii*, Carr., 26, 536.

- Populus spinosa*, F. P. Smith, sp. dub., 26, 537.  
*suaveolens*, Fisch., 26, 535.  
*tremula*, Linn., 26, 537; mentioned, 538.  
— var. *adenopoda*, Burkill\*, 26, 537.  
— var. *villosa*, Wesm., 26, 537.  
*Porana racemosa*, Roxb., 26, 166.  
*spectabilis*, Kurz, mentioned, 26, 167.  
*Porophyllum hieracioides*, DC., 23, 448.  
*Porphyra dichotoma*, Lour., 26, 254.  
*Porphyroscias decursiva*, Miq., 23, 335.  
*Portulaca oleracea*, Linn., 23, 77.  
*villosa*, Cham., var., 36, 503.  
PORTULACÆ, 23, 71.  
*Potamogeton angustifolius*, Bercht., 36, 193.  
*crispus*, Linn., 36, 193.  
— var. *serratus*, S. F. Gray, 36, 193.  
*cristatus*, Regel, 36, 193.  
*Delavayi*, A. Benn., 36, 194.  
*Gaudichaudii*, Cham., 36, 194.  
*heterophyllum*, Buch.-Ham., 36, 194.  
— var. *latifolius*, Mert. & Koch, 36, 193.  
*Hillii*, Morong, 36, 194.  
*hybridus*, Hook., 36, 194.  
*hybridus*, Makino, 36, 193.  
*indicus*, Roth, 36, 193.  
*japonicus*, Franch., 36, 194.  
*javanicus*, Hassk., 36, 194.  
*limosellifolius*, Maxim., 36, 195.  
*lucens*, Linn., 36, 195.  
*lucens*, Vidal, 36, 194.  
*malaiiana*, Miq., 36, 194.  
*Miduhikimo*, Makino, mentioned, 36, 195.  
*mucronatus*, Presl, 36, 195.  
*natans*, Linn., 36, 195.  
*oxyphyllum*, Miq., 36, 195.  
*parvifolius*, Buchen., 36, 194.  
*pectinatus*, Linn., 36, 195.  
*perfoliatus*, Linn., 36, 196.  
*polygonifolius*, Pourr., 36, 196.  
*pusillus*, Linn., 36, 196.  
*Robbinsii*, Oakes, var. *japonicus*, A. Benn., 36, 196.  
*serrulatus*, Regel, 36, 196.  
*sp.*, 36, 196.  
*tenuicaulis*, F. Muell., 36, 194.  
*Tepperi*, A. Benn., 36, 196; mentioned, 195.  
*tretocarpus*, Maxim., 36, 194.  
*Potamogeton tricarinatus*, F. Muell. & A. Benn., mentioned, 36, 195.  
*Wrightii*, Morong, 36, 194.  
*Zizii*, Roth, 36, 193.  
*Potamosace Filicula*, Maxim., 26, 36.  
*Potentilla albiflora*, Wall., 38, 503.  
*ambigua*, Cambess., 36, 504.  
*amurensis*, Maxim., 23, 240; mentioned, 245.  
*ancistrifolia*, Bunge, 23, 240.  
*Anserina*, Linn., 23, 240.  
*articulata*, Franch., 36, 504.  
*betonicaefolia*, Poir., 23, 240.  
*bifurca*, Linn., 23, 241.  
*centigrana*, Maxim., 23, 241.  
*chinensis*, Ser., 23, 241; mentioned, 244.  
*coriandrifolia*, D. Don, 36, 504.  
*Cryptotæniæ*, Maxim., 23, 241.  
*Davidi*, Franch., 36, 504.  
*davurica*, Nestl., 23, 243.  
*Delavayi*, Franch., 36, 504.  
*discolor*, Bunge, 23, 241.  
*eriocarpa*, Wall., 36, 504.  
*exaltata*, Bunge, 23, 241.  
*flagellaris*, Willd., 23, 242.  
*formosana*, Hance, 23, 242.  
*fragarioides*, Linn., 23, 242.  
*fragiformis*, Willd., 23, 242; mentioned, 243.  
*fruticosa*, Linn., 23, 243.  
*fulgens*, Wall., 36, 504.  
*gelida*, C. A. Mey., mentioned, 23, 243.  
*glabra*, Lodd., 23, 243.  
*grandiflora*, Linn., mentioned, 23, 243.  
*Griffithii*, Hook. f., var., 36, 504.  
*hololeuca*, D. Don, 36, 504.  
*Kleiniana*, Wight & Arn., 23, 243.  
*Leschenaultiana*, Ser., 36, 504.  
*leuconota*, D. Don, 36, 504.  
*moupinensis*, Franch., 36, 504.  
*multicaulis*, Bunge, 23, 244.  
*multifida*, Baker & S. Moore, 23, 241.  
*multifida*, Linn., 36, 504.  
*nivea*, Linn., 23, 243.  
*paradoxa*, Nutt., 23, 245.  
*peduncularis*, D. Don, var., 36, 504.  
*pensylvanica*, Linn., 23, 243; mentioned, 241.  
*poterioides*, Franch., 36, 504.  
*purpurea*, Royle, 36, 504.  
*reptans*, Linn., 23, 244.  
— var. *trifoliolata*, A. Gray, 23, 241.  
*Sanguisorba*, Willd., 23, 244.



- Potentilla sericea*, Linn., 23, 244.  
*Sibbaldi*, Hall. f., 36, 504.  
*sischanensis*, Bunge, 23, 244.  
*songarica*, var. *chinensis*, Bunge, 23, 244.  
 sp. n. ?, 23, 246.  
*subacaulis*, Linn., 23, 245.  
*supina*, Linn., 23, 245; mentioned, 245.  
*tanacetifolia*, Willd., 23, 245.  
*verticillaris*, Steph., 23, 245.  
*viscosa*, Donn, 23, 245.  
*Wallichiana*, DC., 23, 245.
- Poterium canadense*, A. Gray, 23, 246.  
*filiforme*, Hook. f., 23, 247.  
*officinale*, Benth. & Hook. f., 23, 247.  
*tenuifolium*, Franch. & Sav., 23, 247.
- Pothos Loureiri*, Hook. & Arn., 36, 186.  
*microphyllus*, Schott, 36, 186.  
*scandens*, Benth., 36, 187.  
*Seemanni*, Schott, 36, 186.  
*terminalis*, Hance, 36, 186.  
*Warburgii*, Engl., 36, 504.
- Pottsia cantonensis*, Hook. & Arn., 26, 96.  
*Hookeriana*, Wight, 26, 96.  
*ovata*, A. DC., 26, 96.
- Poupartia Fordii*, Hemsl., 36, 504.
- Pourthiæa arguta*, Decne., 36, 504.  
*Calleryana*, Decne., 23, 263.  
*coreana*, Decne., 23, 263.  
*lucida*, Decne., 23, 263.  
*Oldhami*, Decne., 23, 263.  
*parvifolia*, E. Pritz., 36, 504.  
*Thunbergii*, Decne., 23, 263.  
*variabilis*, Hemsl., 36, 504.  
*villosa*, Decne., 23, 263.
- Pouzolzia alienata*, Gaudich., 26, 490.  
*borbonica*, Wight, 26, 491.  
*cochinchinensis*, Lour., 26, 490.  
*elegans*, Wedd., 26, 489.  
*glomerata*, Hassk., 26, 490.  
*hirta*, Hassk., 26, 489.  
*hispida*, Benn., 26, 489.  
*hypericifolia*, Blume, 26, 490.  
*indica*, Gaudich., 26, 490.  
 — *subvar. microphylla*, Wedd., 26, 490.  
 — *subvar. procumbens*, Wedd., 26, 490.  
 — *var. alienata*, Wedd., 26, 490.  
 — *var. angustifolia*, Wedd., 26, 490.  
*ovalis*, Miq., 26, 491.  
*procumbens*, Wight, 26, 490.
- Pouzolzia* sp., 26, 491.  
*viminea*, Wedd., 26, 491.  
*zeylanica*, Benn., 26, 490.
- Pratia begonifolia*, Lindl., 26, 2.  
*radicans*, G. Don, 26, 3.  
*Thunbergii*, G. Don, 26, 3.
- Premna formosana*, Maxim., 26, 255.  
*glabra*, A. Gray, 26, 255.  
*integrifolia*, Linn., 26, 255.  
*japonica*, Miq., 26, 256.  
*ligustroides*, Hemsl. \*, 26, 256.  
*microphylla*, Turcz., 26, 256.  
*serratifolia*, Linn., 23, 351; 26, 256.  
 sp. n. ?, Hemsl., 26, 256.  
*staminea*, Maxim., 26, 257.
- Prenanthes Brunoniana*, Wall., 36, 504.  
*chinensis*, Thunb., 23, 485.  
*debilis*, Thunb., 23, 480.  
*dentata*, Thunb., 23, 484.  
*denticulata*, Houtt., 23, 480.  
*Faberii*, Hemsl. \*, 23, 486.  
*glandulosa*, Dunn, 36, 504.  
*hastata*, Thunb., 23, 481.  
*Henryi*, Dunn, 36, 504.  
*humilis*, Thunb., 23, 476.  
*integra*, Thunb., 23, 475.  
*japonica*, Linn., 23, 475.  
*laciniata*, Houtt., 23, 480.  
*lanceolata*, Houtt., 23, 475.  
*lyrata*, Thunb., 23, 475.  
*macrophylla*, Franch., 36, 504.  
*ochroleuca*, Hemsl. \*, 23, 486.  
*repens*, Linn., 23, 483.  
*sonchifolia*, Bunge, 23, 481.  
*Tatarinowii*, Maxim., 23, 486.  
*versicolor*, Fisch., 23, 485.  
*vitifolia*, Diels, 36, 504.
- Primula amethystina*, Franch., 26, 36.  
*argutidens*, Franch., 36, 504.  
*auriculata*, Lam., 26, 36; mentioned, 37.  
 — *var. polyphylla*, Franch., 26, 37.  
*barbicalyx*, C. H. Wright, 36, 504.  
*bella*, Franch., 26, 37.  
*blattariformis*, Franch., 26, 37.  
*bracteata*, Franch., 26, 37.  
*breviscapa*, Franch., 36, 504.  
*bullata*, Franch., 26, 37.  
*calliantha*, Franch., 26, 37.  
*cernua*, Franch., 26, 37.  
*chartacea*, Franch., 36, 504.  
*cinerascens*, Franch., 36, 504.  
*Cockburniana*, Hemsl., 36, 504.

- Primula cortusoides*, Linn., 26, 37.  
*Davidi*, Franch., 36, 504.  
*Delavayi*, Franch., 26, 38.  
*denticulata*, Sm., 26, 38.  
*diantha*, Franch., 36, 504.  
*dryadifolia*, Franch., 26, 38.  
*Faberi*, Oliver, 26, 38.  
*Fargesii*, Franch., 36, 504.  
*farinosa*, Linn., mentioned, 26, 41.  
— var. *denudata*, Ledeb., 26, 41.  
*filipes*, Watt, mentioned, 26, 41.  
*flava*, Maxim., 26, 38.  
*Forbesii*, Franch., 26, 38.  
*gemmifera*, Batalin, 36, 504.  
*Gillii*, Britt., 36, 504.  
*glacialis*, Franch., 26, 39.  
*heucherifolia*, Franch., 36, 504.  
*incisa*, Franch., 26, 39.  
*japonica*, A. Gray, 26, 39.  
— var. *angustidens*, Franch., 26, 39.  
*kialensis*, Franch., 36, 504.  
*lineariloba*, Winkl., 36, 505.  
*Listeri*, King, 26, 39.  
*Loczii*, Kanitz, 36, 505.  
*malacoides*, Franch., 26, 39.  
*malvacea*, Franch., 26, 39.  
*mandorina*, Hoffm., 26, 42.  
*Maximowiczii*, Regel, 26, 40.  
*membranifolia*, Franch., 26, 40.  
*moupinensis*, Franch., 36, 505.  
*mutabilis*, Lour., 23, 273.  
*neurocalyx*, Franch., 36, 505.  
*nivalis*, Pall., 26, 40; 36, 505.  
— var. *pumila*, Pax, 26, 40.  
— var. *purpurea*, Pax, 26, 40.  
*nutans*, Delavay, 26, 40.  
*nutantifolia*, Hemsl., 36, 505.  
*obconica*, Hance, 26, 40.  
— var. *glabrescens*, Franch., 26, 40.  
— var. *rotundifolia*, Franch., 26, 40.  
*oreocharis*, Hance, 26, 40.  
*oreodoxa*, Franch., 36, 505.  
*ovalifolia*, Franch., 26, 41.  
*Paxiana*, Gilg, 36, 505.  
*pellucida*, Franch., 26, 41.  
*petiolaris*, Wall., var., 36, 505.  
*pinnatifida*, Franch., 26, 41.  
*poculiformis*, Hook. f., 26, 40.  
*Poissoni*, Franch., 26, 41.  
*polyneura*, Franch., 36, 505.  
*prænitens*, Ker-Gawl., 26, 42, 43.  
*Prattii*, Hemsl., 36, 505.  
*pulchella*, Franch. \*, 26, 41.  
*Pumilio*, Maxim., 26, 41.  
*purpurea*, Royle, 26, 40.
- Primula pycnoloba*, Bur. & Franch., 36, 505.  
*ramalina*, Bretsch., 36, 505 (a blunder for *Pedicularis ramalana*, Britt.).  
*Rosthornii*, Diels, 36, 505.  
*saxatilis*, Komarov, 36, 505.  
*secundiflora*, Franch., 26, 42.  
*semperflorens*, Loisel., 26, 42.  
*septemloba*, Franch., 26, 42.  
*serratifolia*, Franch., 26, 42.  
*sertulosa*, Kickx, 26, 42.  
*sertulum*, Franch., 36, 505.  
*sibirica*, Jacq., mentioned, 26, 41.  
*sikkimensis*, Hook., 26, 42.  
*sinensis*, Sabine, 26, 42; mentioned, 43.  
*sinuata*, Franch., 36, 505.  
*sonchifolia*, Franch., 26, 43.  
*Souliei*, Franch., 36, 505.  
*spicata*, Franch., 26, 43.  
*stenocalyx*, Maxim., 26, 43.  
*Stuartii*, Franch., 26, 41.  
— var. *typica*, Franch., 26, 40.  
*Stuartii*, Hook. f., 26, 40.  
*tibetica*, Watt, 26, 41.  
*tongolensis*, Franch., 36, 505.  
*urticifolia*, Maxim., 26, 43.  
*Viali*, Delav., 36, 505.  
*vincæflora*, Franch., 26, 43.  
*vittata*, Bur. & Franch., 36, 505.  
*Wilsoni*, Dunn, 36, 505.  
*yunnanensis*, Franch., 26, 43.
- PRIMULACEÆ**, 26, 36.
- Primulina Tabacum*, Hance, 26, 228.
- Prinos asprellus*, Hook. & Arn., 26, 115.
- Prinsepia utilis*, Royle, 36, 505.
- Procris australis**, Spreng., 36, 483.
- cantoniensis*, Spreng., 26, 470.  
*chinensis*, Spreng., 26, 456.  
*ficoidea*, Wall., 26, 482.  
*lævigata*, Blume, 26, 484.  
*obtusata*, Royle, 26, 480.  
*peduncularis*, Wall., 26, 480.  
*sessilis*, Hook. & Arn., 26, 483.  
*umbellata*, Sieb. & Zucc., 26, 483.  
*Wightiana*, Wall., 26, 484.
- Prosartes viridescens*, Regel, 36, 144.
- PROTEACEÆ**, 26, 394.
- Proteinophallus Rivieri*, Hook. f., 36, 182.
- Prunella vulgaris*, Linn., 26, 299.
- Prunopsis Lindleyi*, André, 23, 222.
- Prunus Amygdalus*, Baill., mentioned, 23, 217.  
*Armeniaca*, Linn., 23, 217.  
*aspera*, Thunb., 26, 452.

- Prunus Avium*, Linn., **36**, 505.  
*brachypoda*, Batalin, **36**, 505.  
*Bungei*, Walp., **23**, 219.  
*campanulata*, Maxim., **23**, 218.  
*caudatus*, Franch., **36**, 505.  
*Cerasus*, Linn., mentioned, **23**, 218.  
*chinensis*, Blume, **23**, 219.  
*cinerascens*, Franch., **36**, 505.  
*communis*, Huds., **23**, 218.  
*communis*, Linn., **23**, 222.  
*Davidiana*, Franch., **23**, 218, 220.  
*domestica*, Linn., **23**, 218.  
*glandulosa*, Thunb., **23**, 219.  
*hirtipes*, Hemsl. \*, **23**, 218.  
*humilis*, Bunge, **23**, 218.  
*insititia*, Linn., **23**, 218.  
*japonica*, Carr., **23**, 219.  
*japonica*, Thunb., **23**, 219.  
*macrophylla*, Sieb. & Zucc., **23**, 219, 489.  
*Maximowiczii*, Rupr., **23**, 219; **36**, 505.  
*Mume*, Sieb. & Zucc., **36**, 505.  
*nepalensis*, Hort., var., **36**, 505.  
*oxycarpa*, Maxim., **23**, 219, 489.  
*Padus*, Linn., **23**, 220; **36**, 505.  
*paniculata*, Ker-Gawl., **23**, 221.  
*pauciflora*, Bunge, **23**, 220.  
*Persica*, Sieb. & Zucc., **23**, 220; mentioned, 218.  
*phæosticta*, Maxim., **23**, 221.  
*pogonostyla*, Maxim., **23**, 221; mentioned, **26**, 450.  
*Pseudo-Cerasus*, Lindl., **23**, 221.  
*Puddum*, Roxb., **36**, 505.  
*punctata*, Hook. f., **23**, 221.  
*salicina*, Lindl., **23**, 221.  
*serrula*, Franch., **36**, 505.  
*serrulata*, Lindl., **23**, 221.  
*setulosa*, Batalin, **36**, 505.  
*sibirica*, Linn., mentioned, **23**, 217.  
*Sieboldi*, Maxim., **23**, 221.  
*sinensis*, Pers., **23**, 219.  
*spinulosa*, Sieb. & Zucc., **36**, 505.  
*Ssiori*, F. Schmidt, **23**, 221.  
*stipulacea*, Maxim., **23**, 222.  
*szechuanica*, Batalin, **36**, 505.  
*tatsienensis*, Batalin, **36**, 505.  
*thibeticus*, Franch., **36**, 505.  
*tomentosa*, Thunb., **26**, 222.  
*trichocarpa*, Bunge, **23**, 222.  
*triflora*, Roxb., **23**, 222.  
*trifolia*, Kurz, **23**, 222.  
*triloba*, Lindl., **23**, 222.  
*undulata*, Wall., **36**, 505.  
*velutina*, Batalin, **36**, 505.  
*virgata*, Hort., **23**, 222.  
*xerocarpa*, Hemsl., **36**, 505.  
*Prunus yunnanensis*, Franch., **36**, 505.  
*Przewalskia tangutica*, Maxim., **26**, 177.  
*Psammanthe marina*, Hance, **23**, 323.  
*Pseudægle sepiaria*, Miq., **23**, 111.  
*Pseudanthistiria heteroclita*, Hook. f., **36**, 370.  
*Pseudo-Carex plantaginea*, Miq., **36**, 310.  
*Pseudolarix Fortunei*, Mayr, **26**, 557.  
*Kæmpferi*, Gord., **26**, 558.  
*Pseudostachyum polymorphum*, Munro, **36**, 448.  
*Pseudotsuga Davidiana*, Bertrand, **26**, 554.  
*jezoënsis*, Bertrand, **26**, 555.  
*Kæmpferi*, Gord., **26**, 558.  
*Psidium caninum*, Lour., **23**, 295.  
*Cattleianum*, Sabine, mentioned, **23**, 295.  
*Guyava*, Linn., mentioned, **23**, 295.  
*pomiferum*, Linn., **23**, 295.  
*pumilum*, Vahl, mentioned, **23**, 295.  
*pyriferum*, Linn., **23**, 295.  
*Psilopeganum*, Hemsl. \*, **23**, 103, 491; colour of flowers, 489.  
*sinense*, Hemsl. \*, **23**, 103.  
*Psophocarpus tetragonolobus*, DC., **23**, 194.  
*Psoralea corylifolia*, Franch., **36**, 506.  
*Psychotria elliptica*, Ker-Gawl., **23**, 387.  
*Reevesii*, Wall., **23**, 387.  
*scandens*, Hook. & Arn., **23**, 387.  
*serpens*, Linn., **23**, 387.  
*Ptarmica alpina*, Ledeb., **23**, 437.  
*mongolica*, DC., **23**, 437.  
*sibirica*, Ledeb., **23**, 437.  
*Pterilema acerifolium*, Reinw., **26**, 495.  
*Pternopetalum*, Franch., **36**, 506.  
*Davidi*, Franch., **36**, 506.  
*Pterocarpus flavus*, Lour., **23**, 199.  
*indicus*, Willd., **23**, 199.  
*Pterocarya chinensis*, Hort., **26**, 494.  
*Delavayi*, Franch., **26**, 493; mentioned, 494.  
*hupehensis*, Skan \*, **26**, 493.  
*laevigata*, Hort., **26**, 494.  
*macroptera*, Batalin, **26**, 494.  
*rhoifolia*, Sieb. & Zucc., **36**, 506.  
*stenoptera*, C. DC., **26**, 494.  
— *forma tonkinensis*, Forbes & Hemsl. \*, **26**, 494.

- Pteroceltis Tatarinowii*, Maxim., **26**, 451.  
*Pterocephalus Bretschneideri*, Batalin, **36**, 506.  
*Pterolobium indicum*, Hance, **23**, 207.  
   *punctatum*, Hemsl. \*, **23**, 207.  
   *Rosthornii*, Harms, **36**, 506.  
   *subvestitum*, Hance, **23**, 207.  
*Pteroloma triquetrum*, Benth., **23**, 177.  
*Pteronia tomentosa*, Lour., **23**, 406.  
*Pterospermum acerifolium*, Benth., **23**, 91.  
   *acerifolium*, Willd., **23**, 91.  
   *formosanum*, Matsum., **36**, 506.  
   *heterophyllum*, Hance, **23**, 91.  
   *Proteus*, Burkill, **36**, 506.  
*Pterostigma capitatum*, Benth., **26**, 185.  
   *grandiflorum*, Benth., **26**, 185.  
   *rubiginosum*, Walp., **26**, 185.  
*Pterostyrax volubilis*, Maxim., **26**, 123.  
*Pterygiella Duclouxii*, Franch., **36**, 506.  
   *nigrescens*, Oliver, **36**, 506.  
*Ptilagrostis mongolica*, Griseb., **36**, 383.  
*Ptychochilus septemnervis*, Schau., **36**, 40.  
*Pueraria calycina*, Franch., **36**, 506.  
   *peduncularis*, R. Grah., **36**, 506.  
   *phaseoloides*, Benth., **23**, 190.  
   *Thunbergiana*, Benth., **23**, 191.  
   *tuberosa*, DC., **36**, 506.  
   *yunnanensis*, Franch., **36**, 506.  
*Pulicaria konyangensis*, Vaniot, **36**, 506.  
*Pulsatilla cernua*, Spreng., **36**, 506.  
   *chinensis*, Bunge, **36**, 506.  
   *daurica*, Spreng., **36**, 506.  
*Punica Granatum*, Linn., **23**, 306.  
*Pycnospora hedysaroides*, R. Br., **23**, 177.  
   *nervosa*, Wight & Arn., **23**, 177.  
*Pycnostelma chinensis*, Bunge, **26**, 102; mentioned, 103.  
   *lateriflorum*, Hemsl. \*, **26**, 102.  
*Pycreus angulatus*, Nees, **36**, 202.  
   — *var. β. Wightii*, C. B. Clarke, **36**, 202.  
   *Baccha*, Nees, **36**, 203.  
   *capillaris*, Nees, **36**, 203.  
   *Cooperi*, C. B. Clarke, mentioned, **36**, 203.  
   *Delavayi*, C. B. Clarke \*, **36**, 203.  
   *globosus*, Reichb., **36**, 203.  
   — *var. β. nilagirica*, C. B. Clarke, **36**, 204.  
   — *var. γ. stricta*, C. B. Clarke, **36**, 205.  
*Pycreus globosus*, *var. δ. erecta*, C. B. Clarke \*, **36**, 205.  
   *latespicatus*, C. B. Clarke, **36**, 205.  
   *Linneanus*, Nees, **36**, 205.  
   *nitens*, Nees, **36**, 206.  
   *polystachyus*, Beauv., **36**, 205.  
   *pulvinatus*, Nees, **36**, 206.  
   *puncticulatus*, Nees, **36**, 203.  
   *sanguinolentus*, Nees, **36**, 206.  
*Pygeum Henryi*, Dunn, **36**, 506.  
   *latifolium*, Miq., **23**, 223.  
   *oxycarpum*, Hance, **23**, 219.  
   *phaeostictum*, Hance, **23**, 221.  
*Pyracantha coccinea*, M. Roem., **23**, 260.  
   *crenulata*, M. Roem., **23**, 260.  
*Pyrenaria*, Blume, **23**, 83.  
*Pyrethrum indicum*, Cass., **23**, 437.  
   *sinense*, DC., **23**, 439.  
*Pyrola asarifolia*, Miq., **26**, 33.  
   *atropurpurea*, Franch., **36**, 506.  
   *Corbieri*, H. Lévêillé, **36**, 506.  
   *japonica*, Sieb., **26**, 33.  
   *renifolia*, Maxim., mentioned, **26**, 33.  
   *rotundifolia*, Linn., **26**, 32.  
   *secunda*, Linn., mentioned, **26**, 33.  
   *uniflora*, Linn., **26**, 33.  
*Pyrus alnifolia*, Decne., **36**, 506.  
   *Aria*, Ehrh., **23**, 254.  
   *Aucuparia*, Gaertn., **23**, 255.  
   *baccata*, Linn., **23**, 255.  
   *betulaefolia*, Bunge, **23**, 256.  
   *Calleryana*, Decne., **23**, 256.  
   *cathayensis*, Hemsl. \*, **23**, 256.  
   *chinensis*, Lindl., **23**, 257.  
   *chinensis*, Roxb., **23**, 258.  
   *coccinea*, Forbes & Hemsl., lapsu, **23**, 513 = *Pyracantha coccinea*.  
   *communis*, Linn., **23**, 257.  
   *communis*, Thunb., **23**, 258.  
   — *var. sinensis*, C. Koch, **23**, 258.  
*Cydonia*, Linn., mentioned, **23**, 256.  
   *Delavayi*, Franch., **36**, 506.  
   *discolor*, Maxim., **23**, 255.  
   *floribunda*, Lindl., **23**, 258.  
   *foliolosa*, Wall., **36**, 506.  
   *granulosa*, Bertol., mentioned, **23**, 258.  
   *indica*, Wall., **23**, 257.  
   *Jacquemontiana*, Maxim., **23**, 256.  
   *japonica*, Thunb., **23**, 257.  
   *kansuensis*, Batalin, **36**, 506.  
   *Malus*, Linn., **23**, 257.  
   *Maulei*, Mast., **23**, 257.  
   *microphylla*, Wall., **36**, 506.  
   *Pashia*, Buch.-Ham., **23**, 256; **36**, 506.  
   *pohuashanensis*, Hance, **23**, 255.

- Pyrus Prattii*, Hemsl., 36, 506.  
*prunifolia*, Willd., mentioned, 23, 255.  
*scabrifolia*, Franch., 36, 506.  
*Simonii*, Carr., 23, 256.  
*sinensis*, Dum., 23, 258.  
*sinensis*, Lindl., 23, 257; mentioned, 256.  
*sinensis*, Poir., 23, 256.  
*spp.*, 23, 258, 259.  
*spectabilis*, Ait., 23, 258.  
*transitoria*, Batalin, 36, 506.  
*ussuriensis*, Maxim., 23, 258.  
*variolosa*, Wall., mentioned, 23, 256.  
*vestita*, Wall., 23, 259; 36, 506.  
*Wallichii*, Hook. f., 36, 505.  
*yunnanensis*, Franch., 36, 506.  
*Pyxipoma polyandrum*, Fenzl, 23, 323.
- Quadriala lanceolata*, Sieb. & Zucc., 26, 409.
- Quamoclit vulgaris*, Choisy, 26, 162.
- Quercus*, host of a *Loranthus*, 26, 406.
- acrodonta*, O. Seemen, 26, 516.  
*acutissima*, Carruth., 26, 508.  
*aliena*, Blume, 26, 505.  
 — *var. urticifolia*, Skan\*, 26, 506.  
*amygdalifolia*, Skan\*, 26, 506.  
*annulata*, Sm., 26, 515.  
*attenuata*, Skan\*, 26, 506; mentioned, 512.  
*Augustinii*, Skan\*, 26, 507.  
*bambusæfolia*, Hance, 26, 519.  
*bambusæfolia*, Fortune, 26, 522.  
*bancana*, Kurz, 26, 518.  
*Baronii*, Skan\*, 26, 507.  
*Blakei*, Skan, 36, 506.  
*brevicaudata*, Skan\*, 26, 508.  
*bullata*, O. Seemen, 26, 516.  
*Bungeana*, F. B. Forbes, 26, 508.  
*calathiformis*, Skan\*, 26, 508.  
*callicarpifolia*, Griff., 26, 511.  
*calliprinos*, Webb, mentioned, 26, 507.  
*canescens*, *var. urticæfolia*, Miq., 26, 522.  
*Carlesii*, Hemsl., 26, 509.  
*Carolinæ*, Skan, 36, 506.  
*Championi*, Benth., 26, 509.  
*chinensis*, Abel, 26, 509.  
*chinensis*, Bunge, 26, 508.  
*cleistocarpa*, O. Seemen, 26, 510.  
*cornea*, Lour., 26, 510; mentioned, 518.  
*cuspidata*, Thunb., 26, 510.  
 — *var. sinensis*, DC., 26, 520.
- Quercus dealbata*, Hook. f. & Thoms., 26, 510.  
*Delavayi*, Franch., 26, 511.  
*deusifolia*, Abel, 26, 511.  
*dentata*, Thunb., 26, 511.  
 — *var. McCormickii*, Skan\*, 26, 511.  
 — *var. oxyloba*, Franch., 26, 512.  
*dentosa*, Lindl., 26, 515.  
*Dielsiana*, Seemen, 36, 506.  
*Edithæ*, Skan, 36, 506.  
*elæagnifolia*, O. Seemen, 26, 512.  
*Engleriana*, O. Seemen, 26, 512.  
*Eyreii*, Champ., 26, 512.  
*Eyreii*, Hance, 26, 506, 512.  
*Fabri*, Hance, 26, 512.  
*Fargesii*, Franch., 26, 517.  
*fenestrata*, *var. dealbata*, Wenzig, 26, 511.  
*fissa*, Champ., 26, 512; mentioned, 509.  
*Fordiana*, Hemsl., 36, 506.  
*formosana*, Skan\*, 26, 513.  
*fragifera*, Franch., 26, 510.  
*Franchetii*, Skan\*, 26, 513.  
*gilva*, Blume, 26, 514.  
*glabra*, Thunb., mentioned, 26, 513.  
*glandulifera*, Blume, 26, 514.  
*glauca*, Thunb., 26, 515, 507; 36, 506.  
 — *var. lineata*, Franch., 26, 517.  
 — *var. villosa*, Skan\*, 26, 515.  
*Griffithii*, Hook. f. & Thoms., 26, 505.  
 — *var. urticæfolia*, Franch., 26, 506.  
*grosseserrata*, Blume, 36, 507.  
*Hancei*, Benth., 26, 515; mentioned, 522.  
*Harlandi*, Hance, 26, 515.  
*Henryi*, O. Seemen, 26, 521.  
*Ilex*, Linn., 26, 516.  
 — *var. acrodonta*, Skan\*, 26, 516.  
 — *var. bullata*, Franch., 26, 516.  
 — *var. phillyræoides*, Franch., 26, 516.  
 — *var. rufescens*, Franch., 26, 516.  
 — *var. spinosa*, Franch., 26, 516.  
*ilicioides*, A. Gray, 26, 516.  
*inversa*, Lindl., 26, 521.  
*Irwinii*, Hance, 26, 517.  
*iteaphylla*, Hance, 26, 517.  
*lamellosa*, Sm., 36, 507.  
*lanuginosa*, Franch., mentioned, 26, 514.  
*laxiflora*, Lindl., 26, 513.  
*lineata*, Blume, 26, 517; 36, 507.  
 — *var. Fargesii*, Skan\*, 26, 517.

- Quercus lineata*, var. *grandifolia*,  
*Skan*\*, 26, 517.  
 — var. *oxyodon*, *Wenzig*, 23, 517.  
*litseifolia*, *Hance*, 26, 518.  
*Lycoperdon*, *Skan*\*, 26, 518.  
*McCormickii*, *Carruth.*, 26, 511.  
*mongolica*, *Fisch.*, 26, 518.  
*Moulei*, *Hance*, 26, 519.  
*Naiadarum*, *Hance*, 26, 519.  
*neriifolia*, *O. Seemen*, 26, 518.  
*obovata*, *Bunge*, 26, 511.  
*obscura*, *O. Seemen*, 26, 519.  
*oxyodon*, *Miq.*, 26, 517.  
*pachyloma*, *O. Seemen*, 26, 519.  
*pachyphylla*, *Kurz*, mentioned, 26,  
 518.  
*phillyræoides*, *A. Gray*, 26, 516.  
*Phyllata*, *Buch.-Ham.*, 26, 515.  
*polystachya*, *Wall.*, 26, 519.  
*reversa*, *Lindl.*, 26, 521.  
*Rex*, *Hemsl.*, 36, 507.  
*Robur*, *Pall.*, 26, 518.  
*salicina*, *Blume*, 26, 519.  
*sclerophylla*, *Lindl.*, 26, 520.  
*semecarpifolia*, *Sm.*, 26, 520.  
 — var. *glabra*, *Franch.*, 26, 520.  
*serrata*, *Thunb.*, 26, 520.  
 — var. *chinensis*, *Wenzig*, 26, 508.  
*sessiliflora*, var. *mongolica*, *Franch.*,  
 26, 518.  
*sessilifolia*, *Blume*, 26, 521.  
*Sieboldiana*, *Blume*, 26, 521.  
*Silvicularum*, *Hance*, 26, 521.  
*spicata*, *Sm.*, 26, 521; mentioned,  
 508, 522.  
*spinosa*, *A. David*, 26, 516.  
*sutchuenensis*, *Franch.*, 26, 512.  
*synbalanos*, *Hance*, 26, 521.  
*thalassica*, *Hance*, 26, 521.  
 — var. *Irwinii*, *Wenzig*, 26, 517.  
 — var. *vestita*, *Franch.*, 26, 522.  
*truncata*, *King*, 26, 506.  
*urticæfolia*, *Blume*, 26, 522.  
*uvariifolia*, *Hance*, 26, 522.  
*variolosa*, *Franch.*, 26, 522.  
*Vibrayeana*, *Franch. & Sav.*, 26,  
 522.  
*yunnanensis*, *Franch.*, 26, 511.  
*Quinaria Lansium*, *Lour.*, 23, 110.  
*tricuspidata*, *Koehne*, 36, 507.  
*Quisqualis grandiflora*, *Miq.*, 23, 294.  
*indica*, *Linn.*, 23, 294.  
*sinensis*, *Lindl.*, 23, 294.
- Radermachera pentandra*, *Hemsl.*, 36,  
 507.  
*Randia accedens*, *Hance*, 23, 381.  
*canthioides*, *Champ.*, 23, 381.  
*densiflora*, *Benth.*, 23, 381.
- Randia Dumetorum*, *Lam.*, 23, 381.  
*Henryi*, *E. Pritz.*, 36, 507.  
*leucocarpa*, *Champ.*, 23, 382.  
*racemosa*, *Maxim.*, 23, 382.  
*sinensis*, *Roem. & Schult.*, 23, 382.  
*Wallichii*, *Hook. f.*, 23, 382.
- RANUNCULACEÆ, 23, 1.
- Ranunculus acris*, *Linn.*, 23, 13.  
*affinis*, var., *Maxim.*, 36, 507.  
*aquatilis*, *Linn.*, 23, 13.  
 — var. *brevifolius*, *Rossm.*, 23, 13.  
*auricomus*, *Linn.*, 23, 13; men-  
 tioned, 17.  
*Bungei*, *Steud.*, 23, 13.  
*cantoniensis*, *DC.*, 23, 15.  
*chinensis*, *Bunge*, 23, 14.  
*Cymbalaria*, *Pursh*, 23, 14.  
*diffusus*, *DC.*, 23, 14.  
 — forma *subpinnatus*, *Wight &*  
*Arn.*, 23, 14.  
*extorris*, *Hance*, 23, 16.  
*fibrosus*, *Wall.*, 23, 14.  
*flaccidus*, *Hook. f. & Thoms.*, 36,  
 507.  
*hirtellus*, *Royle*, 23, 13.  
*holophyllus*, *Hance*, 23, 16.  
*hydrocharis*, *Spenn.*, 23, 13.  
*hydrophyllus*, *Bunge*, 23, 13.  
*hyperboreus*, *Rotth.*, 36, 507.  
*japonicus*, *Langsd.*, 23, 14; men-  
 tioned, 15, 16.  
*japonicus*, *Thunb.*, 23, 13.  
*Labordei*, *H. Lévillé & Vaniot*, 36,  
 507.  
*Mællendorffii*, *Hance*, 23, 14.  
*Oryztorum*, *Bunge*, 23, 16.  
*pensylvanicus*, *Linn.*, 23, 14.  
*plantaginifolius*, *Murr.*, 23, 14.  
*Polii*, *Franch.*, 23, 15.  
*propinquus*, *C. A. Mey.*, 23, 13.  
*pulchellus*, *C. A. Mey.*, 23, 15.  
*repens*, *Linn.*, 23, 15.  
*reptans*, *Linn.*, 36, 507.  
*salsuginosus*, *Pall.*, 23, 14.  
*scleratus*, *Linn.*, 23, 16.  
*Sieboldii*, *Miq.*, 23, 16.  
*sp.*, 23, 17.  
*stenorhynchus*, *Franch.*, 36, 507.  
*subpinnatus*, *Wight & Arn.*, 23, 14.  
*ternatus*, *DC.*, 23, 14.  
*ternatus*, *Thunb.*, 23, 16; men-  
 tioned, 14.  
*trichophyllus*, *Chaix*, mentioned,  
 23, 13.  
*tricuspis*, *Maxim.*, 36, 507.  
*yunnanensis*, *Franch.*, 23, 17.  
*Zuccarini*, *Miq.*, 23, 16.
- Rapanea buxifolia*, *Mez*, 36, 507.  
*Faberi*, *Mez*, 36, 507.  
*Playfairii*, *Mez*, 36, 507.

- Rapanea yunnanensis*, Mez, **36**, 507.  
*Raphanus chinensis*, Mill., **23**, 50.  
     *Raphanistrum*, Linn., **23**, 49.  
     *sativus*, Linn., **23**, 50.  
*Raphiolepis indica*, Lindl., **23**, 264.  
     *integerrima*, Hook. & Arn., **23**, 264.  
     *japonica*, Sieb. & Zucc., **23**, 264.  
     *phæostemon*, Lindl., **23**, 264.  
     *rubra*, Lindl., **23**, 264.  
     *salicifolia*, Lindl., **23**, 264.  
*Rapinia herbacea*, Lour., **26**, 8.  
*Rapuntium cæspitosum*, Presl, **26**, 3.  
     *campanuloides*, Presl, **26**, 3.  
     *pyramidale*, Presl, **26**, 3.  
     *radicans*, Presl, **26**, 3.  
     *Wallichianum*, Presl, **26**, 3.  
*Rauwolfia chinensis*, Hemsl. \*, **26**,  
     95.  
*Reevesia thyrsoidea*, Lindl., **23**, 90.  
*Rehmannia angulata*, Hemsl. \*, **26**,  
     193; mentioned, 194.  
     *chinensis*, Fisch. & Mey., **26**, 193.  
     *glutinosa*, Libosch., **26**, 193; men-  
     tioned, 194.  
     — var. *angulata*, Oliver, **26**, 193.  
     *glutinosa*, Hance, **26**, 193.  
     *Oldhami*, Hemsl. \*, **26**, 194;  
     **36**, 507.  
     *Piasezkii*, Maxim., **26**, 194.  
     *rapestris*, Hemsl. \*, **26**, 195.  
*Reineckia carnea*, Kunth, **36**, 113.  
*Reinwardtia sinensis*, Hemsl., **36**,  
     507.  
     *trigyna*, Planch., **23**, 96.  
*Remirea maritima*, Aubl., **36**, 258.  
*Renanthera coccinea*, Lour., **36**, 35.  
*Renealmia calcarata*, Haw., **36**, 71.  
     *nutans*, Andr., **36**, 72.  
*Reseda chinensis*, Lour., **23**, 51.  
     *odorata*, Linn., mentioned, **23**, 51.  
 RFSEDACEÆ, **23**, 51.  
*Restio articulatus*, Retz., **36**, 258.  
*Rhabdothamnopsis*, Hemsl., **36**, 507.  
     *sinensis*, Hemsl., **36**, 507.  
 RHAMNACEÆ, **23**, 125.  
*Rhamnella franguloides*, Weberb., **36**,  
     507.  
     *japonica*, Miq., **23**, 127.  
*Rhamnus argutus*, Maxim., **23**, 128.  
     *argutus*, Maxim., **23**, 129.  
     *catharticus*, Linn., mentioned, **23**,  
     128.  
     *chlorophorus*, Decne., **23**, 130.  
     *crenatus*, Baker & S. Moore, **23**,  
     129.  
     *crenatus*, Sieb. & Zucc., **23**, 127,  
     130.  
     *crenatus*, Sieb. & Zucc., **23**, 128.  
     *dahuricus*, M. A. Laws., **23**, 129,  
     489.  
*Rhamnus davuricus* Pull., **23**, 128;  
     mentioned, 130.  
     *formosanus*, Matsum., **36**, 507.  
     *globosus*, Bunge, **23**, 129, 130.  
     *heterophyllus*, Oliver, **36**, 507.  
     *infectorius*, var. *pubescens*, Griseb.,  
     **23**, 129.  
     *Jujuba*, Linn., **23**, 126.  
     *lineatus*, Linn., **23**, 127.  
     *lineatus*, Lour., **23**, 127.  
     *oreigenes*, Hance, **23**, 128.  
     *parvifolius*, Bunge, **23**, 129; men-  
     tioned, 130.  
     *persicus*, Boiss., **36**, 507.  
     *Rosthornii*, E. Pritz., **36**, 507.  
     *rugulosus*, Hemsl. \*, **23**, 129.  
     *sinensis*, Ser., **23**, 128.  
     *soporifer*, Lour., **23**, 126.  
     spp., **23**, 130.  
     *tinctorius*, Waldst., **23**, 129; men-  
     tioned, 130.  
     *utilis*, Decne., **23**, 128.  
     *virgatus*, Benth., **23**, 130.  
     *virgatus*, Maxim., **23**, 129.  
     *virgatus*, Roxb., **23**, 129; **36**,  
     507.  
     *Zizyphus*, Linn., **23**, 126.  
*Rhamphidia japonica*, Reichb. f., **36**,  
     44.  
*Rhaphidophora angustifolia*, Schott,  
     **36**, 185; mentioned, 186.  
     *formosana*, Engl., **36**, 507.  
     *hongkongensis*, Schott, **36**, 185;  
     mentioned, 186.  
     *lacera*, Hassk., **36**, 186.  
     *lanceifolia*, Schott, mentioned, **36**,  
     185.  
     *Peepla*, Benth., **36**, 185; men-  
     tioned, 186.  
     *pinnata*, Schott, **36**, 186.  
     *sylvestris*, Engl., **36**, 185.  
*Raphiolepis Loureiri*, Spreng., **23**,  
     264.  
     *Mertensii*, Sieb. & Zucc., **23**, 264.  
     *sinensis*, M. Roem., **23**, 264.  
     *spiralis*, G. Don, **23**, 264.  
*Rhaphis trivialis*, Lour., **36**, 368.  
*Rhapis flabelliformis*, Ait., **36**,  
     169.  
     *humilis*, Blume, **36**, 169.  
     *Kwanwortsik*, H. Wendl., **36**,  
     169.  
     *major*, Blume, **36**, 169.  
*Rhaponticum atriplicifolium*, DC., **23**,  
     468.  
     *pungens*, Franch. & Sav., **23**, 469.  
     *uniflorum*, DC., **23**, 470.  
*Rheum Alexandræ*, Batalin, **36**,  
     507.  
     *Barbarum*, Lour., **26**, 355.

- Rheum compactum*, *Linn.*, mentioned, 26, 355.  
*Delavayi*, *Franch.*, 36, 507.  
*Emodi*, *Wall.*, 26, 353.  
*kialense*, *Franch.*, 36, 507.  
*officinale*, *Baill.*, 26, 353; mentioned, 354.  
*palmatum*, *Linn.*, 26, 354.  
*Pichonii*, *Pierre*, mentioned, 26, 354.  
*pumilum*, *Maxim.*, 26, 354.  
*Rhabarbarum*, *Linn.*, mentioned, 26, 355.  
*Rhaponticum*, *Linn.*, mentioned, 26, 353, 354.  
*spiciforme*, *Royle*, 26, 355.  
*strictum*, *Franch.*, 36, 507.  
*undulatum*, *Lour.*, 26, 355.  
*uninerve*, *Maxim.*, 26, 355.  
*Rhexia heteranthera*, *Bonpl.*, 23, 300.  
 RHIZOPHOREÆ, 23, 293.  
*Rhodiola Kirilowi*, *Regel*, 23, 285.  
*Rhododendron adenopodum*, *Franch.*, 36, 507.  
*amœnum*, *Planch.*, 26, 26.  
*Annæ*, *Franch.*, 36, 507.  
*Anthopogon*, *D. Don*, 26, 24.  
*anthopogonoides*, *Maxim.*, 26, 19.  
*argyrophyllum*, *Franch.*, 36, 508.  
*atrovirens*, *Franch.*, 26, 19.  
*aucubæfolium*, *Hemsl. \**, 26, 19.  
*Augustinii*, *Hemsl. \**, 26, 19; 36, 508.  
*aureum*, *Franch.*, 36, 508.  
*auriculatum*, *Hemsl. \**, 26, 20.  
*barbatum*, *Wall.*, mentioned, 26, 20.  
*blepharocalyx*, *Franch.*, 36, 508.  
*Bodinieri*, *Franch.*, 36, 508.  
*Bonvaloti*, *Franch.*, 36, 508.  
*brachyanthum*, *Franch.*, 26, 20.  
*brevistylum*, *Franch.*, 36, 508.  
*Breyonii*, *Planch.*, 26, 25.  
*Buergeri*, *Miq.*, 26, 26.  
*bullatum*, *Franch.*, 26, 20; mentioned, 23.  
*Bureavi*, *Franch.*, 26, 21.  
*Burmanni*, *G. Don*, 26, 27.  
*Calleryi*, *Planch.*, 26, 26.  
*calophyton*, *Franch.*, 36, 508.  
*calycinum*, *Planch.*, 26, 28.  
*campylogynum*, *Franch.*, 26, 21.  
*capitatum*, *Maxim.*, 26, 21.  
 — *var.*, *Franch.*, 26, 23.  
*catawbiense*, *Michx.*, mentioned, 26, 22.  
*cephalanthum*, *Franch.*, 26, 21.  
*Championæ*, *Hook.*, 26, 21; mentioned, 32.  
*chartophyllum*, *Franch.*, 36, 508.  
*Rhododendron chionophyllum*, *Diels*, 36, 508.  
*ciliicalyx*, *Franch.*, 26, 21.  
*cœloneurum*, *Diels*, 36, 508.  
*concinnum*, *Hemsl. \**, 26, 21.  
*coriaceum*, *Franch.*, 36, 508.  
*costulatum*, *Franch.*, 36, 508.  
*crassum*, *Franch.*, 26, 22.  
*crinigerum*, *Franch.*, 36, 508.  
*crispiflorum*, *Planch.*, 26, 26.  
*cyanocarpum*, *Franch.*, 36, 508.  
*Danielsianum*, *Planch.*, 26, 27.  
*dauricum*, *Linn.*, 26, 22.  
 — *var. mucronulatum*, *Hance*, 26, 22.  
*Davidi*, *Franch.*, 36, 508.  
*decorum*, *Franch.*, 26, 22.  
*decumbens*, *D. Don*, 26, 26.  
*Delavayi*, *Franch.*, 26, 22.  
*dendrocharis*, *Franch.*, 36, 508.  
*detersile*, *Franch.*, 36, 508.  
*discolor*, *Franch.*, 36, 508.  
*Edgeworthii*, *Hook. f.*, mentioned, 26, 23.  
*ellipticum*, *Maxim.*, 26, 22.  
*Faberii*, *Hemsl. \**, 26, 22.  
*Fargesii*, *Franch.*, 36, 508.  
*Farreræ*, *Tate*, 26, 23.  
*fastigiatum*, *Franch.*, 26, 23.  
*flavidum*, *Franch.*, 36, 508.  
*floccigerum*, *Franch.*, 36, 508.  
*floribundum*, *Franch.*, 36, 508.  
*fokiense*, *Franch. & Finet*, 36, 508.  
*Fordii*, *Hemsl.*, 36, 508.  
*formosanum*, *Hemsl.*, 36, 508.  
*Fortunei*, *Lindl.*, 26, 23.  
*fragrans*, *Maxim.*, 26, 23.  
*glanduliferum*, *Franch.*, 36, 508.  
*gracilipes*, *Franch.*, 36, 508.  
*hæmatodes*, *Franch.*, 26, 24; mentioned, 23.  
*Hanceanum*, *Hemsl. \**, 26, 24.  
*Hancockii*, *Hemsl.*, 36, 508.  
*heliolepis*, *Franch.*, 26, 24.  
*Henryi*, *Hance*, 26, 24; mentioned, 32.  
*hypoglaucum*, *Hemsl. \**, 26, 24.  
*indicum*, *Sweet*, 26, 25; mentioned, 18, 31.  
 — *var. puniceum*, *Hort.*, 26, 27.  
*intricatum*, *Franch.*, 36, 508.  
*irroratum*, *Franch.*, 26, 26.  
*Kæmpferi*, *Planch.*, 26, 25.  
*Keiskei*, *Miq.*, mentioned, 26, 20.  
*kialense*, *Franch.*, 36, 508.  
*lacteum*, *Franch.*, 26, 26.  
*lateritium*, *Planch.*, 26, 25.  
*Latouchæ*, *Franch. & Finet*, 36, 508.  
*ledifolium*, *G. Don*, 26, 27.



- Rhododendron leucanthum*, Blume, 26, 27.  
*lucidum*, Franch., 36, 508.  
*lukiangense*, Franch., 36, 508.  
*lutescens*, Franch., 36, 508.  
*macranthum*, Sweet, 26, 25.  
*maculiferum*, Franch., 36, 508.  
*Mandarinorum*, Diels, 36, 508.  
*Mariæ*, Hance, 26, 27.  
*mekongense*, Franch., 36, 508.  
*micranthum*, Maxim., 26, 24.  
*micranthum*, Turcz., 26, 27.  
*microphyton*, Franch., 26, 28.  
*molle*, Sieb. & Zucc., 26, 30.  
*moupinense*, Franch., 36, 508.  
*mucronatum*, G. Don, 26, 27.  
*mucronulatum*, Turcz., 26, 32.  
*narcissiflorum*, Planch., mentioned, 26, 27, 28.  
*neriiflorum* Franch., 26, 28; mentioned, 25.  
*obtusum*, Planch., 26, 26.  
*Oldhami*, Maxim., 26, 28.  
*oleifolium*, Franch., 26, 28.  
*oreodoxa*, Franch., 36, 508.  
*ovatum*, Planch., 26, 28.  
*oxyphyllum*, Franch., 36, 508.  
*pachytrichum*, Franch., 36, 508.  
*parviflorum*, Adams, 26, 28.  
*parviflorum*, Ait., 26, 28.  
*phœniceum*, G. Don, 26, 28.  
*pittosporæfolium*, Hemsl. \*, 26, 29.  
*platypodium*, Diels, 36, 509.  
*polifolium*, Franch., 36, 509.  
*polycladum*, Franch., 26, 29.  
*polylepis*, Franch., 36, 509.  
*Potanini*, Batalin, 36, 509.  
*Prattii*, Franch., 36, 509.  
*Pritzelianum*, Diels, 36, 509.  
*Przewalskii*, Maxim., 26, 29.  
*puniceum*, Planch., 26, 27, 28.  
*racemosum*, Franch., 26, 29.  
*ramentaceum*, Planch., 26, 26.  
*ramosissimum*, Bur. & Franch., 36, 509.  
*rigidum*, Franch., 26, 29; mentioned, 20.  
*Rosthornii*, Diels, 36, 509.  
*rotundifolium*, David, 36, 509.  
*rubiginosum*, Franch., 26, 30; mentioned, 20.  
*rufescens*, Franch., 36, 509.  
*rufum*, Batalin, 36, 509.  
*saluenense*, Franch., 36, 509.  
*sanguineum*, Franch., 36, 509.  
*scabrifolium*, Franch., 26, 30.  
*Schlippenbachii*, Maxim., 26, 30.  
*selense*, Franch., 36, 509.  
*Sewiavini*, Maxim., 26, 30.
- Rhododendron siderophyllum*, Franch., 36, 509.  
*Sieboldii*, Miq., 26, 25.  
*Simiarum*, Hance, 26, 30.  
*Simsii*, Planch., 26, 25.  
*Souliei*, Franch., 36, 509.  
*sp. n. ?*, 26, 32.  
*spiciferum*, Franch., 36, 509.  
*spinuliferum*, Franch., 36, 509.  
*stamineum*, Franch., 26, 30; mentioned, 29.  
*strigillosum*, Franch., 36, 509.  
*sublanceolatum*, Miq., 26, 31.  
*sulfureum*, Franch., 26, 31.  
*sutchuense*, Franch., 36, 509.  
*taliense*, Franch., 26, 31.  
*Tashiroi*, Maxim., 26, 31.  
*tatsiense*, Franch., 36, 509.  
*Thompsoni*, Franch., 36, 509.  
*Thunbergii*, Planch., 26, 26.  
*thymifolium*, Maxim., 26, 31.  
*trichocladum*, Franch., 26, 31.  
*trichostomum*, Franch., 36, 509.  
*vernicosum*, Franch., 36, 509.  
*Viali*, Franch., 36, 509.  
*vittatum*, Planch., 26, 31.  
*Westlandi*, Hemsl. \*, 26, 31.  
*Weyrichii*, Maxim., 26, 32.  
*yanthinum*, Bur. & Franch., 36, 509.  
*yunnanense*, Franch., 26, 32.
- Rhodoleia Championi*, Hook., 23, 291.  
*formosa*, Champ., 23, 291.
- Rhodomyrtus tomentosa*, Hassk., 23, 295.
- Rhus ailanthoides*, Bunge, 23, 112.  
*chinensis*, Mill., 23, 147.  
*Cotinus*, Linn., 23, 146.  
*Delavayi*, Franch., 36, 509.  
*Henryi*, Diels, 36, 509.  
*hypoleuca*, Champ., 23, 146; mentioned, 147.  
*javanica*, Linn., 23, 147.  
*microlepis*, Turcz., 23, 146.  
*Potanini*, Maxim., 36, 509.  
*semialata*, Murr., 23, 146.  
*serrata*, Engl., 23, 146.  
*sinica*, Diels, 36, 509.  
*succedanea*, Linn., 23, 147; mentioned, 148.  
— var. *japonica*, Engl., 23, 147.  
*sylvestris*, Sieb. & Zucc., 23, 147.  
*Toxicodendron*, Linn., mentioned, 23, 148.  
— var., 36, 509.  
*trichocarpa*, Miq., 23, 148.  
*verniciifera*, DC., 23, 148.  
*Vernix*, Thunb., 23, 148.  
*Wallichii*, Hook. f., mentioned, 23, 148.

- Rhynchosia Dielsii*, *Harms*, **36**, 509.  
*Henryi*, *Hemsl.* \*, **23**, 196.  
*Henryi*, *Hemsl.*, **23**, 489.  
*himalensis*, *Benth.*, **36**, 509.  
*lutea*, *Dunn*, **36**, 509.  
*minima*, *DC.*, **23**, 196.  
*odorata*, *Hance*, **23**, 319.  
*sp.*, **23**, 197.  
*viscosa*, *DC.*, mentioned, **23**, 197.  
*volubilis*, *Lour.*, **23**, 196.  
*yunnanensis*, *Franch.*, **36**, 509.  
*Rhynchospermum jasminoides*, *Lindl.*,  
**26**, 99.  
*Rhynchospermum verticillatum*,  
*Reinw.*, **36**, 509.  
*Rhynchospora*, see *Rynchospora*.  
*Ribes*, *Linn.*, **23**, 278.  
*alpinum*, *Linn.*, **23**, 279.  
*atropurpureum*, *O. A. Mey.*, **26**, 279.  
*burejense*, *F. Schmidt*, **36**, 509.  
*chifuense*, *Hance*, **23**, 279.  
*Davidi*, *Franch.*, **36**, 509.  
*Diacantha*, *Pall.*, **36**, 509.  
*Fargesii*, *Franch.*, **36**, 509.  
*fasciculatum*, *Sieb. & Zucc.*, **23**, 279.  
*glaciale*, *Franch.*, **36**, 510.  
*glossularioides*, *Maxim.*, **23**, 279.  
*Henryi*, *Franch.*, **36**, 510.  
*longeracemosum*, *Franch.*, **36**, 510.  
*macrocalyx*, *Hance*, **23**, 279.  
*manshuricum*, *Maxim.*, **23**, 510.  
*Maximowiczii*, *Batalin*, **36**, 510.  
*Meyeri*, *Maxim.*, **23**, 279.  
*moupinense*, *Franch.*, **36**, 510.  
*multiflorum*, *Kit.*, **23**, 279.  
*nigrum*, *Linn.*, **23**, 279.  
*pachysandroides*, *Hook. f.*, **36**, 510.  
*petræum*, *Wulf.*, **23**, 280.  
*procumbens*, *Pall.*, **36**, 510.  
*pulchellum*, *Turez.*, **23**, 280.  
*Rosthornii*, *Diels*, **36**, 510.  
*stenocarpum*, *Maxim.*, **23**, 280.  
*tripartitum*, *Batalin*, **36**, 510.  
*Ricinus Apelta*, *Lour.*, **26**, 439.  
*communis*, *Linn.*, **26**, 443.  
*inermis*, var. *manchuriensis*, *Daniell*,  
**26**, 443.  
*Ricotia cantoniensis*, *Lour.*, **23**, 50.  
*Riedelia concatenata*, *DC.*, **23**, 91.  
*corchorifolia*, *DC.*, **23**, 91.  
*Ripidium japonicum*, *Trin.*, **36**, 348.  
*Robinia Amara*, *Lour.*, **23**, 202.  
*Chamlagu*, *L'Hérit.*, **23**, 163.  
*chinensis*, *Pers.*, **23**, 163.  
*flava*, *Lour.*, **23**, 202.  
*mitis*, *Lour.*, **23**, 200.  
*Rodgersia æsculifolia*, *Batalin*, **36**,  
510.  
*Henrici*, *Franch.*, **36**, 510.  
*pinnata*, *Franch.*, **36**, 510.  
*Rodgersia podophylla*, *A. Gray*, **23**,  
267.  
*tabularis*, *Hemsl.*, **36**, 510.  
*Roettlera aurea*, *Franch.*, **36**, 510.  
*Fargesii*, *Franch.*, **36**, 510.  
*mekongensis*, *Franch.*, **36**, 510.  
*tibetica*, *Franch.*, **36**, 510.  
*uniflora*, *Franch.*, **36**, 510.  
*yunnanensis*, *Franch.*, **36**, 510.  
*Rohdea japonica*, *Roth.*, **36**, 115.  
*Rosa acicularis*, *Lindl.*, **23**, 248.  
*alba*, *Lour.*, **23**, 248.  
*amoyensis*, *Hance*, **23**, 252.  
*amygdalifolia*, *Ser.*, **23**, 250.  
*anemonæflora*, *Fortune*, **23**, 247.  
*anemonæflora*, *Lindl.*, **23**, 248.  
*Banksiæ*, *R. Br.*, **23**, 248.  
*Beggeriana*, *Schrenk*, **36**, 510.  
*bengalensis*, *Pers.*, **23**, 250.  
*Biondii*, *Crép.*, **36**, 510.  
*bracteata*, *Wendl.*, **23**, 249.  
*Brunonii*, *Lindl.*, **23**, 252.  
*centifolia*, *Lour.*, **23**, 249.  
*cherokeensis*, *Donn*, **23**, 250.  
*cheusan glabra*, etc., *Petiver*, **23**,  
252.  
*chinensis*, *Jacq.*, **23**, 250.  
*cinnamomea*, var. *dahurica*, *Regel*,  
**23**, 249.  
*cucumerina*, *Tratt.*, **23**, 250.  
*Davidii*, *Crép.*, **36**, 510.  
*davurica*, *Pall.*, **23**, 249.  
*diversifolia*, *Vent.*, **23**, 250.  
*dubia*, *Carr.*, **23**, 252.  
*Fortunei*, *Crép.*, **23**, 247.  
*Fortuniana*, *Lindl. & Paxt.*, **23**,  
249.  
*fragaræflora*, *Ser.*, **23**, 248.  
*Giraldii*, *Crép.*, **36**, 510.  
*gymnocarpa*, *Nutt.*, **23**, 253.  
*Hystrix*, *Lindl.*, **23**, 250.  
*indica*, *Linn.*, **23**, 249; mentioned,  
250.  
*inermis*, *Roxb.*, **23**, 248.  
*intermedia*, *Carr.*, **23**, 252.  
*involutrata*, *Braam*, **23**, 249; men-  
tioned, 248 footnote.  
*koreana*, *Komarov*, **36**, 510.  
*lævigata*, *Michx.*, **23**, 250.  
*Linkii*, *Dehnhardt*, **23**, 253.  
*longifolia*, *Willd.*, **23**, 250.  
*Luciæ*, *Franch. & Rochebr.*, **23**,  
251; mentioned, 252, 253.  
*lucida*, *Lawr.*, **23**, 249.  
*Macartnea*, *Dum.-Cours.*, **23**, 249.  
*macrophylla*, *Lindl.*, **23**, 251.  
*microcarpa*, *Braam*, **23**, 252; men-  
tioned, 248 footnote.  
*microcarpa*, *Lindl.*, **23**, 251; men-  
tioned, 248.

- Rosa microphylla*, Roxb., 23, 252.  
*moschata*, Benth., 23, 251.  
*moschata*, Mill., 23, 252; mentioned, 248, 253; 36, 510.  
*multiflora*, Benth., 23, 252.  
*multiflora*, Thunb., 23, 252.  
*nankinensis*, Lour., 23, 250.  
*nepalensis*, Crép., 23, 252.  
*nivea*, DC., 23, 250.  
*odoratissima*, Sweet, 23, 250.  
*pimpinellifolia*, Linn., 23, 253; mentioned, 254.  
— var. *densiflora*, Crép., 23, 253.  
— var. *platyacantha*, Crép., 23, 253.  
*Prattii*, Hemsl., 36, 510.  
*Pseudo-indica*, Lindl., 23, 250.  
*rugosa*, Thunb., 23, 253.  
*semperflorens*, Curt., 23, 250.  
*sempervirens*, var. *anemoniflora*, Regel, 23, 247.  
*sericea*, Lindl., 23, 254.  
*Serræ*, Dehnhardt, 23, 249.  
*sinica*, Ait., 23, 250; mentioned, 251.  
*sinica*, Lindl., 23, 251.  
*sinica*, Linn., 23, 250.  
*Souliana*, Crép., 36, 510.  
*ternata*, Poir., 23, 250.  
*trifoliata*, Bosc, 23, 250.  
*triphylla*, Roxb., 23, 247.  
*Webbiana*, Wall., 36, 510.  
*xanthina*, Lindl., 23, 254; mentioned, 253.
- ROSACEÆ, 23, 217.**
- Roseoea alpina*, Royle, 36, 67.  
*capitata*, var. *purpurata*, Gagnep., 36, 67.  
— var. *scillifolia*, Gagnep., 36, 68.  
*cauleoides*, Gagnep., 36, 68.  
*Chamæleon*, Gagnep., 36, 68.  
*debilis*, Gagnep., 36, 68.  
*intermedia*, var. *anomala*, Gagnep., 36, 68.  
— var. *macrorhiza*, Gagnep., 36, 68.  
— var. *minuta*, Gagnep., 36, 68.  
*pentandra*, Roxb., 26, 265.  
*purpurea*, Royle, 36, 67.  
— var. *minor*, Wall., 36, 67.
- Rosmarinus officinalis*, Linn., mentioned, 26, 289.
- Rospidios vaccinioides*, A. DC., 26, 71.
- Rostellularia diffusa*, Nees, 26, 245.  
*procumbens*, Nees, 26, 246.
- Rostellularia trichochila*, Miq., 26, 249.
- Rotala densiflora*, Koehne, 23, 303.  
*indica*, Koehne, 23, 303.  
*rotundifolia*, Koehne, 23, 303.
- Rottboellia compressa*, Linn., 36, 361.  
— var. *fasciculata*, Hack., 36, 361.  
— var. *genuina*, Hack., 36, 361.  
— var. *japonica*, Hack., 36, 361.  
*exaltata*, Linn. f., 36, 362.  
*foliata*, Steud., 36, 362.  
*glandulosa*, Trin., mentioned, 36, 362.  
*latifolia*, Steud., 36, 362.  
*mollicoma*, Hance, 36, 362.  
*nigrescens*, Thw., mentioned, 36, 362.  
*setacea*, Roxb., 36, 402.
- Rottlera aurantiaca*, Hook. & Arn., 26, 440.  
*barbata*, Wall., 26, 439.  
*cantoniensis*, Spreng., 26, 439.  
*chinensis*, Juss., 26, 439.  
*cordifolia*, Benth., 26, 441.  
*Furetiana*, Baill., 26, 440.  
*paniculata*, A. Juss., 26, 439.  
*scabrifolia*, A. Juss., 26, 441.  
*tinctoria*, Roxb., 26, 440.  
*Zippelii*, Hassk., 26, 442.
- Roumea chinensis*, Wall. [Cat. n. 6681] ex Meisn., 26, 395.
- Rourea caudata*, Planch., 23, 150.  
*commutata*, Planch., mentioned, 23, 150.  
*humilis*, Blume, mentioned, 23, 150.  
*javanica*, Blume, mentioned, 23, 150.  
*microphylla*, Planch., 23, 149.  
*Millettii*, Planch., 23, 150.  
*santaloides*, Benth., 23, 150.  
*santaloides*, Wight & Arn., mentioned, 23, 150.
- Roxburghia gloriosoides*, Roxb., 36, 95.  
*japonica*, Blume, 36, 95.  
*javanica*, var. *australiana*, Benth., mentioned, 36, 95.  
*ruscifolia*, Zucc., 36, 95.  
*Stemona*, Steud., 36, 95.  
*viridiflora*, Smith, 36, 95.
- ROXBURGHACEÆ, 36, 94.**
- Rubia chinensis*, Regel, 23, 392.  
*cordifolia*, Linn., 23, 393.  
*gracilis*, Miq., 23, 395.  
*mitis*, Miq., 23, 393.  
*Schumanniana*, E. Pritz., 36, 510.

- Rubus abortivus*, O. Kuntze, 23, 229.  
*aceroides*, Miq., 23, 238.  
*adenanthus*, Finet & Franch., 36, 510.  
*allophyllus*, Hemsl., 36, 510.  
*althæoides*, Hance, 23, 229.  
*ampelinus*, Focke, 36, 510.  
*amphidasys*, Focke, 36, 510.  
*aralioides*, Hance, 23, 229.  
*Bambusarum*, Focke, 36, 510.  
*biflorus*, Buchen., 36, 510.  
*Bodinieri*, H. Léveillé & Vaniot, 36, 511.  
*Buergeri*, Miq., 36, 511.  
*Chaffangoni*, H. Léveillé & Vaniot, 36, 511.  
*chiliadenus*, Focke, 36, 511.  
*chinensis*, Franch., 36, 511.  
*chinensis*, Ser., 23, 237.  
*chinensis*, Thunb., 23, 235.  
*chroosepalus*, Focke, 36, 511.  
*cochinchinensis*, Tratt., 36, 511.  
*Cockburnianus*, Hemsl., 36, 511.  
*corchorifolius*, Linn., 23, 230; 36, 511.  
*coreanus*, Miq., 23, 230.  
*cratægifolius*, Bunge, 23, 230.  
*Davidianus*, O. Kuntze, 23, 230.  
*Delavayi*, Franch., 36, 511.  
*ellipticus*, Sm., mentioned, 23, 230; 36, 511.  
*eugenius*, Focke, 36, 511.  
*euleucus*, Focke, 36, 511.  
*Fargesii*, Franch., 36, 511.  
*flagelliflorus*, Focke, 36, 511.  
*flosculosus*, Focke, 36, 511.  
*Fockeanus*, Kurz, 36, 511.  
*foliolosus*, D. Don, mentioned, 23, 235; 36, 511.  
*Fordii*, Hance, 23, 231.  
*formosensis*, O. Kuntze, 23, 230.  
*fragarioides*, Bertol., 36, 511.  
*Gentilianus*, H. Léveillé & Vaniot, 36, 511.  
*Giraldianus*, Focke, 36, 511.  
*glaberrimus*, Champ., 23, 234.  
*glomeratus*, Blume, 23, 231.  
*Grayanus*, Maxim., 23, 231.  
*hakonensis*, Franch. & Sav., 23, 233.  
*Hanceanus*, O. Kuntze, 23, 231.  
*Hasskarlii*, Miq., 23, 231.  
*Henryi*, Hemsl. & O. Kuntze\*, 23, 231.  
*hibernus*, O. Kuntze, mentioned, 23, 231.  
*hupehensis*, Oliver, 36, 511.  
*hydrastifolius*, A. Gray, 23, 238.
- Rubus icbangensis*, Hemsl. & O. Kuntze\*, 23, 231.  
*idaeus*, Linn., 36, 511.  
*incisus*, Miq., 23, 238.  
*incisus*, Thunb., 23, 232.  
*innominatus*, S. Moore, 23, 231; mentioned, 233.  
*innoxius*, Focke, 36, 511.  
*irenæus*, Focke, 36, 511.  
*jambosoides*, Hance, 23, 232.  
*Jamini*, H. Léveillé & Vaniot, 36, 511.  
*kerriifolius*, H. Léveillé & Vaniot, 36, 511.  
*Kuntzeanus*, Hemsl.\* , 23, 232; mentioned, 233.  
*Lambertianus*, Ser., 23, 233.  
 — *var. glaber*, Hemsl.\* , 23, 233.  
*lasiocarpus*, Sm., 36, 511.  
*lasiostylus*, Focke, 36, 511.  
*leucanthus*, Hance, 23, 234.  
*Lobbianus*, Hook., mentioned, 23, 231.  
*loropetalus*, Franch., 36, 511.  
*lutescens*, Franch., 36, 511.  
*macilentus*, Camb., 36, 511.  
*malifolius*, Focke, 36, 511.  
*maximus*, O. Kuntze, mentioned, 23, 231.  
*mesogæus*, Focke, 36, 511.  
*microphyllus*, D. Don, 23, 235.  
*modicus*, Focke, 36, 511.  
*moluccanus*, Linn., 23, 234.  
*Monguilloni*, H. Léveillé & Vaniot, 36, 512.  
*multibracteatus*, H. Léveillé & Vaniot, 36, 512.  
*niveus*, Thunb., 36, 512.  
*ochlanthus*, Hance, 23, 233.  
*Oldhami*, Miq., 23, 236.  
*Oliveri*, Miq., 23, 230.  
*otophorus*, Franch., 36, 512.  
*pacatus*, Focke, 36, 512.  
*pacificus*, Hance, 23, 234.  
*palmatus*, Thunb., 23, 234.  
*paniculatus*, Sm., 23, 234.  
*paradoxus*, S. Moore, 23, 234.  
*Parkeri*, Hance, 23, 235.  
*parvifolius*, Linn., 23, 235.  
*peltatus*, Maxim., 36, 512.  
*phœnicolasius*, Maxim., 23, 235.  
*pileatus*, Focke, 36, 512.  
*pinnatisepalus*, Hemsl., 36, 512.  
*pirifolius*, Franch. = *pyrifolius*, 36, 512.  
*Playfairii*, Hemsl.\* , 23, 235.  
*poliophyllus*, O. Kuntze, 23, 236.  
*polytrichus*, Franch., 36, 512.

- Rubus pungens*, Camb., 23, 236.  
*purpureus*, Bunge, 23, 235.  
*pycnanthus*, Focke, 23, 233.  
*pyrifolius*, Sm., 36, 512.  
*rectangulifolius*, O. Kuntze, 23, 236.  
*reflexus*, Ker-Gawl., 23, 236; mentioned, 231.  
*reticulatus*, Wall., mentioned, 23, 238.  
*rosæfolius*, Sm., 23, 237.  
*rubifolius*, Sieb. & Zucc., 23, 238.  
*rugosus*, Maxim., 23, 230.  
*rugosus*, Sm., 23, 234.  
*saxatilis*, Linn., 23, 237.  
*sepalanthus*, Focke, 36, 512.  
*setchuenensis*, Franch., 36, 512.  
*Sieboldii*, Blume, 23, 237.  
*simplex*, Focke, 36, 512.  
*sorbifolius*, Maxim., 36, 512.  
*sozostylus*, Focke, 36, 512.  
*spinipes*, Hemsl., 36, 512.  
*sterilis*, O. Kuntze, mentioned, 23, 235.  
*Swinhooii*, Hance, 23, 237.  
*tagallus*, Cham. & Schlecht., 23, 237.  
*teledapos*, Focke, 36, 512.  
*tephrodes*, Hance, 23, 238.  
*thibetanus*, Franch., 36, 512.  
*Thunbergii*, Sieb. & Zucc., 23, 238.  
*trifidus*, Thunb., 23, 238.  
*triphyllus*, Thunb., 23, 235.  
*viburnifolius*, Franch., 36, 512.  
*villosus*, Thunb., 23, 230.  
*xanthocarpus*, Bur. & Franch., 36, 512.  
*xanthoneurus*, Focke, 36, 512.  
*yunanicus*, O. Kuntze, 23, 238.
- Ruellia chinensis*, Nees, 26, 238.  
*Cusia*, Buch.-Ham., 26, 240.  
*flagelliformis*, Roxb., 26, 237; mentioned, 238.  
*indigofera*, Griffith, 26, 240.  
*indigotica*, Fortune, 26, 240.  
*japonica*, Thunb., 26, 241.  
*repens*, Linn., 26, 238.  
*seclusa*, S. Moore, 26, 238.  
*tetrasperma*, Champ., 26, 242.
- Rumex Acetosa*, Linn., 26, 355.  
*Acetosella*, Linn., 26, 355.  
*egyptiacus*, Bieb., 26, 357.  
*aquaticus*, Linn., 26, 355.  
*chinensis*, Campd., 26, 357.  
*crispus*, Linn., 26, 356.  
*crispus*, Lour., 26, 357.  
*dentatus*, Linn., 26, 356; mentioned, 356, 358.  
*denticulatus*, Campd., 26, 357.  
*domesticus*, Hartm., mentioned, 26, 355; 36, 512.
- Rumex Fischeri*, Reichb., 26, 356.  
*hamatus*, Trev., 26, 357.  
*Hydrolapathum*, Huds., 26, 356.  
*japonicus*, Meisn., 26, 356.  
*longifolius*, Kunth, 26, 356.  
*Loureiri*, Campd., 26, 357.  
*Laederi*, Muentzer, 26, 356.  
*maritimus*, Linn., 26, 357.  
*Marschallianus*, Reichb., 26, 357.  
*nepalensis*, Spreng., 26, 357.  
*Patientia*, Linn., 26, 357.  
*persicarioides*, Maxim., 26, 357.  
*pulcher*, Linn., 26, 357; mentioned, 356, 358.  
*ramulosus*, Meisn., 26, 357.  
*Regelii*, F. Schmidt, 26, 356.  
*stenophyllus*, Ledeb., 36, 512.
- Rungia chinensis*, Benth., 26, 247.  
*parviflora*, Nees, 26, 247.
- Ruppia maritima*, Linn., 36, 197.  
*rostellata*, Koch, 36, 197.  
*sp.*, Griff., 26, 197.
- Ruta angustifolia*, Pers., 23, 102.  
*bracteosa*, DC., 23, 102.  
*chulepensis*, Linn., 23, 102.  
*dahurica*, DC., 23, 102.  
*graveolens*, Linn., mentioned, 23, 102.
- RUTACEÆ, 23, 102.
- Rynchospora alba*, Vahl, mentioned, 36, 260.  
*anomala*, Steud., 36, 236.  
*articulata*, Spreng., 36, 259.  
*aurea*, R. Br., 36, 259.  
*aurea*, Vahl, 36, 259.  
*capillacea*, Torr., mentioned, 36, 259.  
*chinensis*, Boeck., 36, 260.  
*Faberi*, C. B. Clarke\*, 36, 259.  
*glauca*, Boeck., 36, 260.  
*glauca*, Roem. & Schult., 36, 259.  
*glauca*, Vahl, 36, 259.  
 — *var. chinensis*, C. B. Clarke, 36, 259.  
*gracillima*, C. B. Clarke, 36, 260.  
*gracillima*, Thw., 36, 260.  
*Haenkei*, Presl, 36, 260.  
*Kamphoeveneri*, Boeck., 36, 260.  
*Knieskernii*, J. Carey, mentioned, 36, 259.  
*Lavarum*, Hook. & Arn., 36, 260.  
*laxa*, Benth., 36, 260.  
 — *var. chinensis*, C. B. Clarke, 36, 259.  
*laxa*, R. Br., 36, 259.  
*triflora*, Nees, 36, 260.
- Wallichiana*, C. B. Clarke, 36, 260.  
*Wallichiana*, Kunth, 36, 260.

- Sabia Bullockii*, Hance, **23**, 143.  
*japonica*, Maxim., **23**, 143.  
*leptandra*, Hook. f. & Thoms., **36**, 512.  
*limoniacea*, Wall., **23**, 144.  
*paniculata*, Seem., **23**, 144.  
*parviflora*, Wall., mentioned, **23**, 144.  
*purpurea*, Hook. f. & Thoms., **36**, 512.  
*Schumanniana*, Diels, **36**, 512.  
*Swinhoei*, Hemsl. \*, **23**, 144.  
*yunnanensis*, Franch., **36**, 512.  
**SABIACEÆ**, **23**, 143.  
*Saccharum arundinaceum*, Retz., **36**, 349.  
*japonicum*, Thunb., **36**, 347, 348.  
*Narenga*, Buch.-Ham., **36**, 349.  
*officinatum*, Linn., **36**, 349.  
*panicum*, Lam., **36**, 357.  
*polydactylon*, var. *a*, Thunb., **36**, 347.  
— var. *β*, Thunb., **36**, 348.  
*porphyrocomum*, Hack., **36**, 349.  
*procerum*, Roxb., **36**, 349.  
*rufipilum*, Steud., **36**, 350.  
*Sara*, Roxb., **36**, 349.  
*sinense*, Roxb., **36**, 349.  
*spicatum*, Linn., **36**, 343.  
*spicatum*, Thunb., **36**, 346.  
*spontaneum*, Linn., **36**, 349.  
— *subsp. a. indicum*, Hack., **36**, 349.  
*Saccolabium Fargesii*, Kränzl., **36**, 512.  
*hainanense*, Rolfe, **36**, 36.  
*Sageretia ferruginea*, Oliver, **36**, 512.  
*hamosa*, Brongn., **23**, 181.  
*oppositifolia*, Brongn., **36**, 512.  
*paucicostata*, Maxim., **36**, 512.  
*rugosa*, Hance, **23**, 133.  
*theezans*, Brongn., **23**, 133.  
*Sagina Linnæi*, Presl, **23**, 70.  
*maxima*, A. Gray, **23**, 70.  
*sinensis*, Hance, **23**, 70.  
*Sagittaria chinensis*, Sims, **36**, 190 ; mentioned, 191.  
*cordifolia*, Roxb., **36**, 190.  
*guyanensis*, Humb., **36**, 190.  
*macrophylla*, Bunge, **36**, 190 ; mentioned, 191.  
*obtusata*, Thunb., **36**, 190.  
*pygmæa*, Miq., **36**, 191.  
*sagittata*, Thunb., **36**, 190.  
*sagittifolia*, Hook. & Arn., **36**, 191.  
*sagittifolia*, Linn., **36**, 190.  
— var. *diversifolia*, Micheli, **36**, 190.  
— var. *longiloba*, Turcz., **36**, 190.  
*Sagittaria sagittifolia*, var. *oligocarpa*, Micheli, **36**, 191.  
*trifolia*, Linn. (pl. dub.), **36**, 191.  
*Saintlegeria gracilis*, Cordem., **26**, 367.  
*Salacia chinensis*, Linn., **23**, 125.  
*cochinchinensis*, Lour., **23**, 125.  
**SALICACEÆ**, **26**, 526.  
*Salicornia herbacea*, Linn., **36**, 512.  
*indica*, Willd., **26**, 328.  
*Salisburia adiantifolia*, Sm., **26**, 547.  
*macrophylla*, [Reyn.], C. Koch, **26**, 547.  
*Salix acutifolia*, Willd., **26**, 527.  
*alba*, Linn., **26**, 526.  
*amygdalina*, Linn., **26**, 533.  
*babylonica*, Linn., **26**, 526.  
*Bockii*, Seemen, **36**, 512.  
*brachystachys*, Franch. & Sav., **26**, 533.  
*cantoniensis*, Hance, **26**, 527.  
*Caprea*, Linn., **26**, 527.  
*Caprea* × *phyllicifolia*, **26**, 527.  
*cinerea*, Linn., **36**, 512.  
*Daltoniana*, Anderss., **26**, 528 ; mentioned, 529.  
— var. *Franchetiana*, Burkill \*, **26**, 528.  
— × *elegans*, **26**, 528.  
*daphnoides*, Villars, **26**, 527.  
— var. *acutifolia*, Anderss., **26**, 527.  
*densifoliata*, Seemen, **36**, 512.  
*dictyoneura*, Seemen, **36**, 512.  
*elegans*, Wall., **26**, 528 ; mentioned, 529.  
*eriophylla*, Anderss., **26**, 528.  
*Fargesii*, Burkill \*, **26**, 528.  
*floccosa*, Burkill \*, **26**, 529.  
*fragilis*, Linn., mentioned, **26**, 527.  
*gracilistyla*, Miq., **26**, 533.  
*Henryi*, Burkill \*, **26**, 530 ; mentioned, 529.  
*heterochroma*, Seemen, **36**, 512.  
*koreensis*, Anderss., **26**, 530.  
*Lindleyana*, Wall., **26**, 530.  
*longiflora*, Anderss., **26**, 530.  
— var. *albescens*, Burkill \*, **26**, 530.  
*Maximoviczii*, Komarov, **26**, 512.  
*Mesnyi*, Hance, **26**, 530 ; mentioned, 531, 534.  
*microphyta*, Franch., **26**, 531.  
*Miquelii*, Anderss., mentioned, **26**, 529.  
*moupinensis*, Franch., **26**, 531 ; mentioned, 529.  
*multinervis*, Komarov, **36**, 513.  
*myrtilloides*, Komarov, **36**, 513.  
*nigra*, Marsh., mentioned, **26**, 527.  
*obscura*, Anderss., **26**, 531.

- Salix Oldhamiana*, *A. Henry*, **26**, 533.  
*oreophila*, *Hook.*, **26**, 531.  
*pentandra*, *Linn.*, **26**, 531.  
*phylicifolia*, *Linn.*, **26**, 532.  
*populifolia*, *Anderss.*, **26**, 530.  
*purpurea*, *Linn.*, **26**, 532.  
— *var. stipularis*, *Franch.*, **26**, 532.  
*pycnostachya*, *Anderss.*, mentioned, **26**, 533.  
*repens*, *Linn.*, **26**, 534.  
*Rosthornii*, *Seemen*, **36**, 513.  
*Serpyllum*, *Anderss.*, mentioned, **26**, 531.  
*sikkimensis*, *Anderss.*, **26**, 532 ; mentioned, 529.  
*subpycnostachya*, *Burkill* \*, **26**, 532.  
*tetrasperma*, *Roxb.*, **26**, 533 ; mentioned, 531.  
*Thunbergiana*, *Blume*, **26**, 533.  
*triandra*, *Linn.*, **26**, 533 ; mentioned, 531.  
— *var. maxima*, *Debeaux*, **26**, 533.  
— *var. subdiscolor*, *Debeaux*, **26**, 534.  
*vagans*, *Komarov*, **36**, 513.  
*variegata*, *Franch.*, **26**, 534.  
*viminalis*, *Linn.*, **26**, 534 ; **36**, 513.  
*Wallichiana*, *var. grisea*, *Anderss.*, **26**, 534.  
*Warburgii*, *O. Seemen*, **26**, 534.  
*Salomonina aphylla*, *Griff.*, **23**, 59.  
*cantonensis*, *Lour.*, **23**, 59.  
*oblongifolia*, *DC.*, **23**, 59.  
*parasitica*, *Griff.*, **23**, 59.  
*Salsola collina*, *Pall.*, **26**, 330.  
*Kali*, *Linn.*, **26**, 330.  
— *var. spicata*, *Debeaux*, **26**, 330.  
— *var. tenuifolia*, *Moq.*, **26**, 330.  
— *var. vulgaris*, *auct.*, **26**, 330.  
*Soda*, *Linn.*, **26**, 330.  
*Salvia brevilabra*, *Franch.*, **36**, 513.  
*chinensis*, *Benth.*, **26**, 284.  
*Fortunei*, *Benth.*, **26**, 284.  
*hians*, *Royle*, **26**, 284.  
*japonica*, *Thunb.*, **26**, 284 ; mentioned, 287.  
— *var. bipinnata*, *Franch. & Sav.*, **26**, 284.  
— *var. erythrophylla*, *Hemsl. \**, **26**, 284.  
— *var. integrifolia*, *Franch. & Sav.*, **26**, 284.  
— *var. lanuginosa*, *Franch.*, **26**, 284.  
— *var. parvifoliola*, *Hemsl. \**, **26**, 285.  
— *var. ternata*, *Franch. & Sav.*, **26**, 284.  
*Salvia Mandarinorum*, *Diels*, **36**, 513.  
*Maximowicziana*, *Hemsl. \**, **26**, 285.  
*miltiorrhiza*, *Bunge*, **26**, 286.  
*minutiflora*, *Bunge*, **26**, 288.  
*officinalis*, *Linn.*, mentioned, **26**, 287.  
*Piasezkii*, *Maxim.*, **26**, 287.  
*pogonocalyx*, *Hance*, **26**, 286.  
*Prattii*, *Hemsl.*, **36**, 513.  
*Prionitis*, *Hance*, **26**, 287.  
*Przewalskii*, *Maxim.*, **26**, 287.  
*pygmæa*, *Matsum.*, **36**, 513.  
*Roborowskii*, *Maxim.*, **26**, 287.  
*scapiformis*, *Hance*, **26**, 287.  
*tatsienensis*, *Franch.*, **36**, 513.  
*tricuspis*, *Franch.*, **36**, 513.  
*umbratica*, *Hance*, **26**, 288.  
*yunnanensis*, *C. H. Wright*, **36**, 513.  
*Samara longifolia*, *Benth.*, **26**, 62.  
*obovata*, *Benth.*, **26**, 62.  
*Sambucus chinensis*, *Lindl.*, **23**, 348.  
*ebuloides*, *Desv.*, **23**, 348.  
*Ebulus*, *Linn.*, **23**, 348.  
*javanica*, *Blume*, **23**, 348 ; mentioned, 349.  
*Loureiriana*, *DC.*, **23**, 348.  
*nigra*, *Lour.*, **23**, 348.  
*nigra*, *Thunb.*, **23**, 349.  
*racemosa*, *Linn.*, **23**, 348.  
*Sieboldiana*, *Blume*, **23**, 349.  
*Thunbergiana*, *Blume*, **23**, 248.  
*Williamsii*, *Hance*, **23**, 349.  
*Samolus Valerandi*, *Linn.*, **26**, 59.  
SAMYDACEÆ, **23**, 311.  
*Sanguisorba canadensis*, *Linn.*, *var. latifolia*, *Ledeb.*, **23**, 246.  
*carnea*, *Franch.*, **23**, 247.  
*officinalis*, *Linn.*, **23**, 247.  
*rubra*, *Schrank*, **23**, 247.  
*tenuifolia*, *Fisch.*, **23**, 247.  
*Sanicula cærulescens*, *Franch.*, **36**, 513.  
*chinensis*, *Bunge*, **23**, 326.  
*clata*, *Buch.-Ham.*, **23**, 326.  
*europæa*, *Linn.*, **23**, 326.  
*haquetioides*, *Franch.*, **36**, 513.  
*lamelligera*, *Hance*, **23**, 326.  
*orthacantha*, *S. Moore*, **23**, 326.  
*rubriflora*, *F. Schmidt*, **23**, 327.  
*rugulosa*, *Diels*, **36**, 513.  
*tuberculata*, *Maxim.*, **23**, 327.  
*yunnanensis*, *Franch.*, **36**, 513.  
*Sansevieria carnea*, *Andr.*, **36**, 113.  
*sessiliflora*, *Ker-Gawl.*, **36**, 113.  
SANTALACEÆ, **26**, 408.  
SAPINDACEÆ, **23**, 138.  
*Sapindus abruptus*, *Lour.*, **23**, 139.  
*chinensis*, *Linn.*, **23**, 139.

- Sapindus Mukorossi*, Gaertn., 23, 139.  
*Sapium discolor*, Muell. Arg., 26, 445.  
     *rotundifolium*, Hemsl. \*, 26, 445.  
     *sebiferum*, Roxb., 26, 445; mentioned, 444.  
*Saponaria Vaccaria*, Linn., 23, 64.  
 SAPOTACEÆ, 26, 68.  
*Sarcandra chloranthoides*, Gardn., 26, 367.  
*Sarcanthus cerinus*, Rolfe \*, 36, 36.  
     *elongatus*, Rolfe \*, 36, 36.  
     *filiformis*, Lindl., mentioned, 36, 36, 37.  
     *Fordii*, Rolfe \*, 36, 37.  
     *formosanus*, Rolfe, 36, 37.  
     *honkongensis*, Rolfe, 36, 37.  
     *oxyphyllus*, Wall., mentioned, 36, 37.  
     *paniculatus*, Lindl., 36, 37.  
     *rostratus*, Lindl., 36, 37.  
     *rostratus*, Reichb. f., 36, 34.  
     *succisus*, Lindl., 36, 37.  
     *teretifolius*, Lindl., 36, 38; mentioned, 34.  
*Sarcochilus Arachnites*, Reichb. f., mentioned, 36, 35.  
     *aurifer*, Rolfe \*, 36, 34.  
     *hainanensis*, Rolfe, 36, 35.  
     *lunifer*, Benth., 36, 35.  
*Sarcococca Hookeriana*, Baill., 26, 418.  
     *pruniformis*, Lindl., 26, 418.  
     *salicifolia*, Baill., 26, 418.  
     *saligna*, Muell. Arg., 26, 418.  
     *trinervia*, Wight, 26, 418.  
*Sarcopyramis lanceolata*, Wall., 23, 302.  
     *nepalensis*, Wall., 23, 302.  
*Sarcosperma laurina*, Hook. f., 26, 68.  
     *pedunculata*, Hemsl. \*, 26, 68.  
*Sarcozygium xanthoxylum*, Bunge, 23, 97.  
*Saruma Henryi*, Oliver, 26, 360; 36, 513.  
*Sassafras officinalis*, Sieb., 26, 392.  
     *Parthenoxylon*, Nees, 26, 372.  
*Satyrium bracteatum*, Pers., 36, 54.  
     *nepalense*, D. Don, 36, 63.  
     *setchuenicum*, Kränzl., 36, 513.  
     *viride*, Linn., 36, 54.  
*Saurauja macrotricha*, Kurz, 23, 79.  
     *napaulensis*, DC., 36, 513.  
     *Oldhami*, Hemsl. \*, 23, 79.  
     *punduana*, Wall., 23, 79.  
     *tristyla*, DC., 23, 79.  
*Sauropsis chinensis*, Turcz., 26, 363.  
     *Cumingii*, C. DC., 26, 363.  
*Sauropus albicans*, Blume, 26, 420.  
*Saururus cernuus*, Thunb., 26, 363.  
*Saururus chinensis*, Baill., 26, 363.  
     *Loureiri*, Decne., 26, 363.  
     *natans*, Linn., 36, 193.  
*Saussurea affinis*, Spreng., 23, 463.  
     *alata*, DC., 23, 465.  
     *alaticipes*, Hemsl., 36, 513.  
     *alpina*, var. *leucophylla*, Hance, 23, 467.  
     *amara*, DC., 23, 463; mentioned, 466.  
     *amurensis*, Turcz., 23, 464.  
     *arenaria*, Maxim., 23, 463.  
     *auriculata*, Hemsl., 36, 513.  
     *auriculata*, Franch., 36, 513.  
     *Baroniana*, Diels, 36, 513.  
     *Bullockii*, Dunn, 36, 513.  
     *Bungei*, Benth. & Hook. f., 23, 463.  
     *carduiformis*, Franch., 36, 513.  
     *carthamoides*, Benth., 23, 463.  
     *caudata*, Franch., 36, 513.  
     *chetchozensis*, Franch., 36, 513.  
     *chinensis*, Sch.-Bip., 23, 465.  
     *ciliaris*, Franch., 36, 513.  
     *cirsioides*, Hemsl., 36, 513.  
     *compta*, Franch., 36, 513.  
     *conyzoides*, Hemsl., 36, 513.  
     *cordifolia*, Hemsl., 36, 513.  
     *crassifolia*, DC., 23, 463.  
     *crispa*, Vaniot, 36, 513.  
     *Davidii*, Franch., 23, 463.  
     *decurrens*, Hemsl., 36, 513.  
     *Delavayi*, Franch., 36, 513.  
     *dimorphæa*, Franch., 36, 513.  
     *discolor*, var. *elatior*, Maxim., 23, 464.  
     *dolichopoda*, Diels, 36, 513.  
     *Dutaillyana*, Franch., 36, 513.  
     *dzeurensis*, Franch., 36, 513.  
     *edulis*, Franch., 36, 514.  
     *epilobioides*, Maxim., 23, 464.  
     *eriocephala*, Franch., 36, 514.  
     *eriolepis*, Bunge, 23, 464.  
     *Fargesii*, Franch., 36, 514.  
     *flexuosa*, Franch., 36, 514.  
     *glomerata*, Poir., 23, 464.  
     *gossypiphora*, D. Don, 36, 514.  
     *graminea*, Dunn, 36, 514.  
     *grosseserrata*, Franch., 36, 514.  
     *Grumgrjmailo*, Winkl., 36, 514.  
     *Henryi*, Hemsl., 36, 514.  
     *hypoleuca*, Spreng., 23, 465, 466.  
     *intermedia*, Turcz., 23, 465.  
     *iodostegia*, Hance, 23, 464.  
     *japonica*, DC., 23, 464.  
     *japonica*, Franch., 23, 463.  
     *Katochæte*, Maxim., 23, 465.  
     *Kunthiana*, C. B. Clarke, 36, 514.  
     *lamprocarpa*, Hemsl. \*, 23, 465.  
     *lampsanifolia*, Franch., 36, 514.



- Saussurea lanuginosa*, *Vaniot*, **36**, 514.  
*Leontodon*, *Dunn*, **36**, 514.  
*linearis*, *Champ.*, **23**, 464.  
*lingulata*, *Franch.*, **36**, 514.  
*likiangensis*, *Franch.*, **36**, 514.  
*longifolia*, *Franch.*, **36**, 514.  
*macrota*, *Franch.*, **36**, 514.  
*manshurica*, *Komarov*, **36**, 514.  
*Medusa*, *Maxim.*, **23**, 466.  
*microcephala*, *Diels*, **36**, 514.  
*microcephala*, *Franch.* \*, **23**, 466.  
*minuta*, *Winkl.*, **36**, 514.  
*mollis*, *Franch.*, **36**, 514.  
*nigrescens*, *Maxim.*, **23**, 466.  
*nivea*, *Turcz.*, **23**, 464.  
*nobilis*, *Franch.*, **36**, 514.  
*obvallata*, *Wall.*, **36**, 514.  
*oligantha*, *Franch.*, **36**, 514.  
*otophylla*, *Hemsl.* \*, **36**, 514.  
*pachyneura*, *Franch.*, **36**, 514.  
*paleata*, *Maxim.*, **23**, 467.  
*pectinata*, *Bunge*, **23**, 467.  
— var. *amurensis*, *Maxim.*, **23**, 467.  
*peduncularis*, *Franch.*, **36**, 514.  
*phaantha*, *Maxim.*, **23**, 466.  
*populifolia*, *Hemsl.*, **36**, 514.  
*Przewalskii*, *Maxim.*, **23**, 467.  
*pulchella*, *Fisch.*, **23**, 464.  
*radiata*, *Franch.*, **36**, 514.  
*romuleifolia*, *Franch.*, **36**, 514.  
*Rosthornii*, *Diels*, **36**, 514.  
*salicifolia*, *DC.*, **23**, 467.  
— var. *chinensis*, *Maxim.*, **23**, 467.  
*saligna*, *Franch.*, **36**, 514.  
*scabrida*, *Franch.*, **36**, 514.  
*semilyrata*, *Bur. & Franch.*, **36**, 514.  
*serrata*, *DC.*, **23**, 467.  
*Souliei*, *Franch.*, **36**, 514.  
*sp.*, **23**, 468.  
*spatulifolia*, *Franch.*, **36**, 514.  
*Stella*, *Maxim.*, **23**, 468.  
*stricta*, *Franch.*, **36**, 514.  
*Sughoo*, *C. B. Clarke*, **36**, 514.  
*sutchuenensis*, *Franch.*, **36**, 514.  
*sylvatica*, *Maxim.*, **23**, 468.  
*Tanakæ*, *Maxim.*, **23**, 468.  
*tangutica*, *Maxim.*, **23**, 468.  
*taraxacifolia*, *Wall.*, **36**, 514.  
*tatsienensis*, *Franch.*, **36**, 514.  
*thibetica*, *Franch.*, **36**, 514.  
*triangulata*, *Trautv. & Mey.*, **23**, 468.  
*umbrosa*, *Komarov*, **36**, 514.  
*ussuriensis*, *Maxim.*, **23**, 468; mentioned, 467.  
— var. *odontolepis*, *Herder*, **23**, 467.  
*Saussurea vaginata*, *Dunn*, **36**, 514.  
*vestita*, *Franch.*, **36**, 515.  
*villosa*, *Franch.*, **36**, 515.  
*virgata*, *Franch.*, **36**, 515.  
*Woodiana*, *Hemsl.*, **36**, 515.  
*yunnanensis*, *Franch.*, **36**, 515.  
*Saxifraga aplophyloides*, *Franch.*, **36**, 515.  
*atrata*, *Engl.*, **23**, 266.  
*aurantiaca*, *Franch.*, **36**, 515.  
*brachyphylla*, *Franch.*, **36**, 515.  
*Candelabrum*, *Franch.*, **36**, 515.  
*cardiophylla*, *Franch.*, **36**, 515.  
*cernua*, *Linn.*, **23**, 266.  
*chinensis*, *Linn.*, **23**, 266.  
*chionophila*, *Franch.*, **36**, 515.  
*chrysantha*, *Franch.*, **36**, 515.  
*cortusæfolia*, *Sieb. & Zucc.*, **23**, 267.  
— var. *Fortunei*, *Maxim.*, **23**, 267.  
*crinalis*, *Franch.*, **36**, 515.  
*cuscutæformis*, *Lodd.*, **23**, 267.  
*Davidii*, *Franch.*, **36**, 515.  
*Delavayi*, *Franch.*, **23**, 267.  
*drabæformis*, *Franch.*, **36**, 515.  
*egregia*, *Engl.*, **23**, 267.  
*flabellifolia*, *Franch.*, **36**, 515.  
*Fortunei*, *Hance*, **23**, 267.  
*Fortunei*, *Hook.*, **23**, 267.  
*gemmigera*, *Engl.*, **36**, 515.  
*gemmipara*, *Franch.*, **36**, 515.  
*Gillii*, *Trimen*, **36**, 515.  
*Giraldiana*, *Engl.*, **36**, 515.  
*glaucophylla*, *Franch.*, **36**, 515.  
*hirculoides*, *Engl.*, **23**, 267.  
*Hirculus*, *Linn.*, **23**, 267.  
*hispidula*, *D. Don*, **36**, 515.  
*hypericoides*, *Franch.*, **36**, 515.  
*Josephi*, *Engl.*, **36**, 515.  
*likiangensis*, *Franch.*, **36**, 515.  
*longistyla*, *Franch.*, **36**, 515.  
*macrostigma*, *Franch.*, **36**, 515.  
*manshuriensis*, *Komarov*, **36**, 515.  
*melanocentra*, *Franch.*, **36**, 515.  
*micrantha*, *Franch.*, **36**, 515.  
*nana*, *Engl.*, **23**, 268.  
*nutans*, *Franch.*, **36**, 515.  
*oreophila*, *Franch.*, **36**, 515.  
*pekinensis*, *Maxim.*, **23**, 268.  
*peltata*, *Torr.*, mentioned, **23**, 269.  
*peplidifolia*, *Franch.*, **36**, 515.  
*petrophila*, *Franch.*, **36**, 515.  
*phænophylla*, *Franch.*, **36**, 515.  
*Przewalskii*, *Engl.*, **23**, 268.  
*punctata*, *Linn.*, **36**, 515.  
*purpurascens*, *Franch.*, **36**, 515.  
*Rossii*, *Oliver*, **23**, 268.  
*rotundifolia*, *Linn.*, **23**, 268.  
*rupicola*, *Franch.*, **36**, 515.

- Saxifraga sanguinea*, *Franch.*, **36**, 515.  
*sarmentosa*, *Linn.*, **23**, 268.  
 — var. *cuscutæformis*, *Ser.*, **23**, 267.  
*serpyllifolia*, var. *Pallasiana*, *Hance*, **23**, 269.  
*sibirica*, *Linn.*, **36**, 515.  
 sp. nova?, **23**, 269.  
*stellariæfolia*, *Franch.*, **36**, 516.  
*tabularis*, *Hemsl.* \*, **23**, 269.  
*tangutica*, *Engl.*, **26**, 269.  
 — var. *minutiflora*, *Engl.*, **23**, 269.  
*tellimoides*, *Maxim.*, mentioned, **23**, 269.  
*trinervia*, *Franch.*, **36**, 516.  
*tsangchanensis*, *Franch.*, **36**, 516.  
*unguiculata*, *Engl.*, **23**, 269.
- SAXIFRAGACEÆ, **23**, 265.
- Scabiosa Bretschneideri*, *Batalin*, **36**, 516.  
*cochinensis*, *Lour.*, **23**, 400.  
*Fischeri*, *DC.*, **23**, 400.
- Scævola hainanensis*, *Hance*, **26**, 2.  
*Koenigii*, *Vahl*, **26**, 2.  
*latevaga*, *Hance*, **26**, 2.  
*Lobelia*, *Benth.*, **26**, 2.
- Scepa chinensis*, *Champ.*, **26**, 429.
- Sceptrocnide macrostachya*, *Maxim.*, **36**, 516.
- Schanginia linifolia*, *C. A. Mey.*, **26**, 329.
- Schedonorus remotiflorus*, *Miq.*, **36**, 430.
- Schefflera Fargesii*, *Harms*, **36**, 516.  
*megalobotrya*, *Harms*, **36**, 516.
- Schima argentea*, *E. Pritz.*, **36**, 516.  
*Noronhæ*, *Reinw.*, **23**, 80.  
*superba*, *Gardn.*, **23**, 80.
- Schistocaryum*, *Franch.*, **36**, 516.  
*ciliare*, *Bur. & Franch.*, **36**, 516.  
*myosotideum*, *Franch.*, **36**, 516.  
*ovalifolium*, *Bur. & Franch.*, **36**, 516.
- Schistocodon Meyeni*, *Schau.*, **26**, 101.
- Schizandra axillaris*, *Hook.*, **36**, 516.  
*chinensis*, *Baill.*, **23**, 25.  
*elongata*, *Hook.*, **36**, 516.  
*glaucescens*, *Diels*, **36**, 516.  
*grandiflora*, *Hook.*, **36**, 516.  
*japonica*, *Hance*, **23**, 25.  
*propinqua*, *Hook. f. & Thoms.*, var., **36**, 516.
- Schizocapsa plantaginea*, *Hance*, **36**, 90.
- Schizocodon rotundifolius*, *Maxim.*, **26**, 34.
- Schizopepon dioicus*, *Cogn.*, **36**, 516.
- Schizophragma hydrangeoides*, *Sieb. & Zucc.*, **36**, 516.  
*integrifolia*, *Oliver*, **36**, 516.
- Schizostachyum chinense*, *Rendle* \*, **36**, 448.  
*Dumetorum*, *Munro*, **36**, 449.
- Schizotheca Hemprichii*, *Ehrenb.*, **26**, 563; **36**, 3.
- Schmidelia Cobbe*, *DC.*, **23**, 138.
- Schoberia glauca*, *Bunge*, **26**, 329.
- Schœnus apogon*, *Roem. & Schult.*, **36**, 261.  
*Brownii*, *Hook. f.*, **36**, 261.  
*coloratus*, *Linn.*, **36**, 223.  
 — var.  $\beta$ , *Linn.*, **36**, 209.  
*compressus*, *Linn.*, **36**, 248.  
*elatus*, *Boeck.*, **36**, 261.  
*falcatus*, *R. Br.*, **36**, 261.  
*imberbis*, *Poir.*, **36**, 261.  
*lithospermus*, *Linn.*, **36**, 265.  
*Mariscus*, *Linn.*, **36**, 262.  
*niveus*, *Linn.*, **36**, 224.  
*puberulus*, *C. A. Mey.*, **36**, 236.
- Schœpfia chinensis*, *Gardn. & Champ.*, **23**, 114.  
*jasminodora*, *Sieb. & Zucc.*, **23**, 114.
- Schœpflopsis chinensis*, *Miers*, **23**, 114.  
*jasminodora*, *Miers*, **23**, 114.
- Schollia crassifolia*, *Jacq.*, **26**, 116.
- Schubertia japonica*, *Spach*, **26**, 544.
- Scilla chinensis*, *Benth.*, **36**, 127.  
*japonica*, *Baker*, **36**, 127.  
*japonica*, *Thunb.*, **36**, 140.
- Scindapsus sinensis*, *Engl.*, **36**, 185.  
*sylvestris*, *Kunth*, **36**, 185.
- Scirpus acicularis*, *Linn.*, **36**, 225.  
*acuminatus*, *Poir.*, **36**, 230.  
*æstivalis*, *Retz.*, **36**, 231.  
*affinis*, *Roth*, **36**, 251.  
*afflatus*, *Benth.*, **36**, 226.  
*anceps*, *Willd.*, **36**, 231.  
*arvensis*, *Retz.*, **36**, 235.  
*atropurpureus*, *Retz.*, **36**, 226.  
*attenuatus*, *Franch.*, **36**, 226.  
*barbatus*, *Rottb.*, **36**, 247.  
*bispicatus*, *Roxb.*, **36**, 243.  
*capillaris*, *Linn.*, **36**, 247, 248.  
*capitatus*, *Linn.*, **36**, 227.  
*capsularis*, *Lour.*, **36**, 163.  
*Caricis*, *Retz.*, **36**, 248.  
*cephalotes*, *Jacq.*, **36**, 224.  
*chinensis*, *Munro*, **36**, 250, 253.  
*chinensis*, *Osbeck*, **36**, 253.  
*ciliaris*, *Linn.*, **36**, 256.  
*Cinnamometorum*, *Vahl*, **36**, 233.  
*cognatus*, *Hance*, **36**, 252.  
*complanatus*, *Retz.*, **36**, 231.  
*compressus*, *Pers.*, **36**, 248.  
*coniferus*, *Poir.*, **36**, 258.  
*corymbosus*, *Linn.*, **36**, 259.  
*cuspidatus*, *Roth*, **36**, 257.  
*cymosus*, *Lam.*, **36**, 232.

- Scirpus debilis*, Pursh, **36**, 248.  
*dichotomus*, Linn., **36**, 233, 234.  
*diphyllus*, Retz., **36**, 234.  
*dipsaceus*, Rottb., **36**, 235.  
*erectus*, Poir., **36**, 248.  
*Eriophorum*, Michx., **36**, 249.  
— var., **36**, 255.  
*ferrugineus*, Linn., **36**, 235.  
*filipes*, C. B. Clarke \*, **36**, 249.  
*fistulosus*, Poir., **36**, 227.  
*fuirenoides*, Maxim. \*, **36**, 250 ;  
mentioned, 253.  
*globulosus*, Retz., **36**, 237.  
*glomeratus*, Linn., **36**, 223.  
*glomeratus*, Retz., **36**, 244.  
*grossus*, Linn. f., **36**, 250.  
*hakonensis*, Franch. & Sav., **36**,  
228.  
*intricatus*, Linn., **36**, 209.  
*juncoides*, Roxb., **36**, 248.  
*kamtschaticus*, C. A. Mey., **36**, 228.  
*lacustris*, Linn., **36**, 250.  
*laxiflorus*, Thw., **36**, 230.  
*leptocarpus*, F. Muell., **36**, 257.  
*lithospermus*, Linn., **36**, 265.  
*luzonensis*, Presl, **36**, 248.  
*macrostachys*, Boeck., **36**, 251.  
*maritimus*, Linn., **36**, 251.  
— var. *affinis*, C. B. Clarke, **36**,  
251.  
— var., **36**, 251.  
*margaritiferus*, Boeck., **36**, 261.  
*Meyenii*, Nees, **36**, 251.  
*Michelianus*, Linn., **36**, 251 ; men-  
tioned, 208.  
*microcarpus*, Presl, **36**, 254.  
*miliaceus*, Linn., **36**, 242.  
*miliaceus*, Thunb., **36**, 237.  
*mucronatus*, Linn., **36**, 252.  
*nutans*, Retz., **36**, 241.  
*Onaei*, Franch. & Sav., **36**, 228.  
*palustris*, Linn., **36**, 227.  
*petasatus*, Maxim., **36**, 228.  
*pileatus*, Franch., **36**, 228.  
*plantagineus*, Franch., **36**, 229.  
*plantagineus*, Retz., **36**, 228.  
*plantaginoides*, Rottb., **36**, 228.  
*plicarrhachis*, Griseb., **36**, 230.  
*Pollichii*, Gren., **36**, 255.  
*polystachyus*, F. Muell., mentioned,  
**36**, 250.  
*polytrichoides*, Retz., **36**, 241.  
*quinquangularis*, Vahl, **36**, 242.  
*radicans*, Schkuhr, **36**, 252.  
*Rosthornii*, Diels, **36**, 253.  
*schœnoides*, Retz., **36**, 243.  
*sericeus*, Poir., **36**, 243.  
*setaceus*, Linn., **36**, 253.  
*squarrosa*, Poir., **36**, 244.  
*squarrosus*, Linn., **36**, 253.  
*Scirpus Stauntoni*, C. B. Clarke \*,  
**36**, 253.  
*strobilinus*, Roxb., **36**, 251.  
*subaphyllus*, Boeck., **36**, 254.  
*subcapitatus*, Thw., **36**, 254.  
*supinus*, Linn., **36**, 254.  
*sylvaticus*, Linn., **36**, 254 ; men-  
tioned, 249.  
— var. *radicans*, Willd., **36**, 252.  
*sylvatico-radicans* ×, Baenitz, **36**,  
253.  
*Tabernæmontani*, C. C. Gmel., **36**,  
251.  
*ternatensis*, Reinw., **36**, 254 ; men-  
tioned, 253.  
*tetragonus*, Poir., **36**, 246.  
*timorensis*, Kunth, **36**, 248.  
*tringulatus*, Roxb., **36**, 252.  
*trifidus*, Hance, **36**, 248.  
*triqueter*, Linn., **36**, 255.  
*tuberosus*, Roxb., **36**, 229.  
*uniglumis*, Link, **36**, 227.  
*Wichuraj*, Franch., **36**, 228.  
SCITAMINEÆ, **36**, 67.  
*Scleria annularis*, Kunth, **36**, 263.  
*aspera*, Boeck., **36**, 264, 266.  
*biflora*, Roxb., **36**, 263.  
*caricina*, Benth., **36**, 267.  
*chinensis*, Kunth, **36**, 263.  
*ciliaris*, Nees, **36**, 263.  
*Doederleiniana*, Boeck., **36**, 266.  
*elata*, Thw., **36**, 264.  
*filiformis*, Sw., **36**, 265 ; mentioned,  
265.  
*Græffeana*, Boeck., **36**, 265.  
*Harlandi*, Hance, **36**, 264.  
*hebecarpa*, Munro, **36**, 265.  
*hebecarpa*, Nees, **36**, 264.  
— var. *pubescens*, C. B. Clarke, **36**,  
265.  
*Hookeriana*, Boeck., **36**, 265.  
*japonica*, Steud., **36**, 264.  
*keyensis*, K. Schum., **36**, 266.  
*lævis*, Hance, **36**, 266.  
— var. *scaberrima*, Benth., **36**, 266.  
*lithosperma*, Sw., **36**, 265.  
*margaritifera*, Willd., **36**, 265.  
*Neesiana*, Hook. & Arn., **36**, 266.  
*pergracilis*, Kunth, **36**, 266.  
*polycarpa*, Boeck., **36**, 265.  
*pubescens*, Steud., **36**, 265.  
*purpurascens*, Benth., **36**, 264.  
*Radula*, Hance, **36**, 266 ; men-  
tioned, 265.  
*Ridleyi*, C. B. Clarke, **36**, 266.  
*Rinkiana*, Boeck., **36**, 264.  
*scrobiculata*, Moritzi, **36**, 263.  
*scrobiculata*, Nees, **36**, 266 ; men-  
tioned, 264, 267.  
*Stuedeliana*, Miq., **36**, 263.

- Scleria tessellata*, Benth., 36, 263.  
*tessellata*, Decne., 36, 266.  
*tessellata*, Willd., 36, 267; mentioned, 263.  
*vestita*, Boeck., 36, 265.  
*waigiouensis*, Steud., 36, 266.  
*Wichurii*, Boeck., 36, 264.  
*Scleromitrium angustifolium*, Benth., 23, 375.  
*sinense*, Miq., 23, 375.  
*Sclerostylis atalantoides*, Wight & Arn., 23, 110.  
*buxifolia*, Benth., 23, 110.  
*Hindsii*, Champ., 23, 110.  
*venosa*, Champ., 23, 110.  
*Scolopia acuminata*, Clos, 23, 57.  
*chinensis*, Briquet, 36, 516.  
*chinensis*, Clos, 23, 57.  
*crenata*, Clos, 23, 57.  
*Oldhami*, Hance, 23, 57.  
*Scoparia dulcis*, Linn., 26, 193.  
*gypsophiloides*, Walp., 26, 193.  
*Scopolia japonica*, Maxim., 26, 176.  
*lurida*, Dunal, mentioned, 26, 176.  
*physaloides*, Dunal, 26, 176.  
*sinensis*, Hemsl. \*, 26, 176.  
*tangutica*, Maxim., 26, 176.  
*Scorzonera*, Tourn., 23, 490.  
*albicaulis*, Bunge, 23, 488; mentioned, 490.  
*austriaca*, Willd., 23, 488.  
*divaricata*, Turcz., 23, 490.  
*macrosperma*, Turcz., 23, 488, 490.  
*mongolica*, Maxim., var., 36, 516.  
*parviflora*, Jacq., 23, 489.  
*radiata*, Fisch., 23, 489.  
*Scotanthus tubiflorus*, Naud., 23, 314.  
*Scrofella chinensis*, Maxim., 26, 179.  
*Scrophularia alata*, A. Gray, 36, 516.  
*Buergeriana*, Miq., 26, 179.  
*chinensis*, Linn., 26, 177.  
*Delavayi*, Franch., 36, 516.  
*diplodonta*, Franch., 36, 516.  
*Fargesii*, Franch., 36, 516.  
*Henryi*, Hemsl. \*, 26, 178.  
— var. *glabrescens*, Hemsl. \*, 26, 178.  
*kansuensis*, Batalin, 36, 516.  
*lateriflora*, Trautv., 26, 279.  
*Mandarinorum*, Franch., 36, 516.  
*microdonta*, Franch., 36, 516.  
*Moellendorffi*, Maxim., 26, 178.  
*Oldhami*, Oliver, 26, 179.  
*Souliei*, Franch., 36, 516.  
*spicata*, Franch., 36, 516.  
*yunnanensis*, Franch., 36, 516.  
SCROPHULARINEÆ, 26, 177.  
*Scutellaria adenophylla*, Miq., 26, 293.  
*amœna*, C. H. Wright, 36, 516.  
*angulosa*, Benth., 26, 293; mentioned, 297.  
*Scutellaria baikalensis*, Georgi, 26, 294.  
*barbata*, D. Don, 26, 296.  
*dependens*, Maxim., 26, 294; mentioned, 295.  
*formosana*, N. E. Br., 36, 516.  
*galericulata*, Linn., 26, 294.  
*grandiflora*, Adams, 26, 294.  
*hederacea*, Kunth & Bouché, 26, 295.  
*indica*, Linn., 26, 295.  
— var. *pekinensis*, Maxim., 26, 295.  
*japonica*, Burm. f., 26, 271.  
*japonica*, Morr. & Decne., 26, 295.  
*lanceolaria*, Miq., 26, 294.  
*leucodasys*, Miq., 26, 296.  
*luzonica*, Rolfe, 26, 296.  
*macrantha*, Fisch., 26, 294, 298.  
— var. *glabrescens*, Franch., 26, 294.  
— var. *pubescens*, Debeaux, 26, 294.  
*minor*, Linn., mentioned, 26, 294.  
*obtusifolia*, Hemsl. \*, 26, 296.  
*Oldhami*, Miq., 26, 294.  
*pekinensis*, Maxim., 26, 295.  
*rivularis*, Wall., 26, 296.  
*scandens*, D. Don, 26, 293.  
*sciaphila*, S. Moore, 26, 297.  
*scordifolia*, Fisch., 26, 294.  
*sessilifolia*, Hemsl. \*, 26, 297.  
*stenosiphon*, Hemsl. \*, 26, 297.  
*strigillosa*, Hemsl. \*, 26, 297.  
*violacea*, Heyne, 26, 298.  
*viscidula*, Bunge, 26, 298.  
*Scytalia chinensis*, Gaertn., 23, 139.  
*Sebastiana Chamælea*, Muell. Arg., 26, 445.  
*Sebifera glutinosa*, Lour., 26, 385.  
*Securidaca scandens*, Buch.-Ham., 23, 63.  
*tavoyana*, Wall., 23, 63.  
*Securinega fluggeoides*, Muell. Arg., 26, 426.  
*japonica*, Miq., 26, 426.  
*obovata*, Muell. Arg., 26, 427.  
*ramiflora*, Muell. Arg., 26, 426.  
*Sedum Aizoon*, Linn., 23, 282.  
— [var.] *latifolium*, Miq., 23, 285.  
*albo-roseum*, Baker, 23, 282.  
*Alfredi*, Hance, 23, 283.  
*algidum*, Ledeb., 23, 283.  
*alsinæfolium*, All., 23, 284.  
*Anacampseros*, Linn., 23, 283.  
*angustum*, Maxim., 23, 283.  
*bracteatum*, Diels, 36, 516.  
*chrysastrum*, Hance, 23, 283.  
*crassipes*, Wall., 36, 516.  
*discolor*, Franch., 36, 516.  
*drymarioides*, Hance, 23, 283.  
*drymarioides*, Maxim., 23, 284.

- Sedum dumulosum*, Franch., 23, 284.  
*elatinoides*, Franch., 23, 284.  
*erythrostictum*, Miq., 23, 283, 287.  
*Fabaria*, Koch, 23, 287.  
*Fabarium*, Lem., 23, 287.  
*filipes*, Hemsl. \*, 23, 284, 491.  
— var. *major*, Hemsl. \*, 26, 284, 285.  
*fimbriatum*, Franch., 23, 281.  
*formosanum*, N. E. Br., 23, 285.  
*glaciale*, Franch., 36, 516.  
*Henryi*, Diels., 36, 517.  
*hybridum*, var., Maxim., 23, 282.  
*japonicum*, Sieb., 23, 285; mentioned, 283.  
*kamtschaticum*, Fisch., 23, 285.  
*Kirilowi*, Regel, 23, 285.  
*leucocarpum*, Franch., 36, 517.  
*lineare*, Thunb., 23, 285.  
— var. *floribundum*, Miq., 23, 283.  
*macrolepis*, Franch., 36, 517.  
*malacophyllum*, Franch., 23, 282.  
*Maximowiczii*, Regel, 23, 282.  
*Middendorffianum*, Maxim., 36, 517.  
*mosoynense*, Franch., 36, 517.  
*multicaule*, Wall., mentioned, 23, 283.  
*nobile*, Franch., 36, 517.  
*obtusisepalum*, Franch., 36, 517.  
*platysepalum*, Franch., 36, 517.  
*polytrichoides*, Hemsl. \*, 23, 286, 492.  
*primuloides*, Franch., 36, 517.  
*Przewalskii*, Maxim., 23, 286.  
*Pseudo-Aizoon*, Debeaux, 23, 282.  
*purpurascens*, Koch, 23, 287.  
*purpureum*, Link, 23, 287.  
*ramosissimum*, Franch., 23, 281.  
*Roborowskii*, Maxim., 23, 286.  
*Rosthornianum*, Diels, 36, 517.  
*sarmentosum*, Bunge, 23, 286.  
*scabridum*, Franch., 36, 517.  
*Sheareri*, S. Moore, 23, 286.  
*sinicum*, Diels, 36, 517.  
*sp. aff. S. Sieboldii*, 23, 288.  
*spectabile*, Boreau, 23, 287.  
*stellariæfolium*, Franch., 23, 283.  
*stellatum*, Lour., 23, 283.  
*suboppositum*, Maxim., 23, 287.  
*subtile*, Debeaux, 23, 283.  
*subtile*, Miq., 23, 287.  
*tanguticum*, Maxim., 23, 283.  
*Tatarinowii*, Maxim., 23, 287.  
*Telephium*, Linn., 23, 287; mentioned, 283.  
*tenuifolium*, Franch., 36, 517.  
*uniflorum*, Hook. & Arn., 23, 288.  
*valerianoides*, Diels, 36, 517.  
*viviparum*, Maxim., 36, 517.
- Sedum vulgare*, Link, 23, 287.  
*yantaiense*, Debeaux, 23, 288.  
*yunnanense*, Franch., 36, 517.
- Seidlia*, Opiz, 36, 249.  
*radicans*, Opiz, 36, 252.  
*sylvatica*, Opiz, 36, 254.
- SELAGINEÆ, 26, 250.
- Selinum coreanum*, Boissieu, 36, 517.  
*cryptotænium*, Boissieu, 36, 517.  
*daucoides*, Franch., 36, 517.  
*Melanotilingia*, Boissieu, 36, 517.  
*Monnieri*, Linn., 23, 332.  
*Oliverianum*, Boissieu, 36, 517.
- Senebiera integrifolia*, DC., 23, 48.
- Senecillis Schmidtii*, Maxim., 23, 457.
- Senecio acerifolius*, Winkl., 36, 517.  
*aconitifolius*, Turcz., 23, 449; mentioned, 450, 458.  
*ægyptius*, Linn., 23, 452.  
*ainsliæflorus*, Franch., 36, 517.  
*altaicus*, Sch.-Bip., 23, 450.  
*ambraceus*, Turcz., 23, 450.  
*arachnanthus*, Franch., 36, 517.  
*argunensis*, Debeaux, 23, 450.  
*argunensis*, Turcz., 23, 450.  
*asperifolius*, Franch., 36, 517.  
*atroviolaceus*, Franch., 36, 517.  
*aurantiacus*, DC., 23, 451.  
*begoniæfolius*, Franch., 36, 517.  
*blattariæfolius*, Franch., 36, 517.  
*Bodinieri*, Vaniot, 36, 517.  
*bulbiferus*, Maxim., 23, 450.  
*cacaliæfolius*, Sch.-Bip., 23, 454.  
*cacaliformis*, Reichb., 23, 454.  
*campanulatus*, Franch., 36, 517.  
*campestris*, DC., 23, 450.  
*camptodontus*, Franch., 36, 517.  
*campylodes*, DC., 23, 457.  
*cannabifolius*, Less., 23, 456.  
*Caroli*, Winkl., 36, 517.  
*chinensis*, DC., 23, 457.  
*chrysanthemoides*, DC., 36, 517.  
*Clivorum*, Maxim., 23, 451.  
*concinuus*, Franch., 36, 517.  
*cyclaminifolius*, Franch., 36, 517.  
*cyclotus*, Bur. & Franch., 36, 517.  
*dahuricus*, Sch.-Bip., 23, 451.  
*Davidi*, Franch., 36, 517.  
*Delavayi*, Franch., 36, 517.  
*deltophyllus*, Maxim., 23, 451.  
*densiflorus*, Wall., 36, 517.  
*dianthus*, Franch., 36, 518.  
*dictyoneurus*, Franch., 36, 518.  
*dididymanthus*, Dunn, 36, 518.  
*discoideus*, Franch., 36, 518.  
*divaricatus*, Linn., 23, 447.  
*dodrans*, Winkl., 36, 518.  
*Dryas*, Dunn, 36, 518.  
*duciformis*, Winkl., 36, 518.

- Senecio Duclouxii*, *Dunn*, **36**, 518.  
*emodensis*, *Sch.-Bip.*, **23**, 454.  
*erythropappus*, *Bur. & Franch.*, **36**, 518.  
*euryophyllus*, *Winkl.*, **23**, 518.  
*Exul*, *Hance*, **23**, 451.  
*Faberi*, *Hemsl.* \*, **36**, 518.  
*farfaræfolius*, *Maxim.*, **23**, 452.  
*Farfugium*, *O. Koch*, **23**, 454.  
*Fargesii*, *Franch.*, **36**, 518.  
*fibrillosus*, *Dunn*, **36**, 518.  
*filiferus*, *Franch.*, **36**, 518.  
*ganpinensis*, *Vaniot*, **36**, 518.  
*Gentilianns*, *Vaniot*, **36**, 518.  
*glabellus*, *DC.*, **23**, 451.  
*glumaceus*, *Dunn*, **36**, 518.  
*Gynura*, *Winkl.*, **36**, 518.  
*Helianthus*, *Franch.*, **36**, 518.  
*Henrici*, *Vaniot*, **36**, 518.  
*Henryi*, *Hemsl.* \*, **23**, 452.  
*Hindsii*, *Benth.*, **23**, 457.  
*Hoi*, *Dunn*, **36**, 518.  
*ianthophyllus*, *Franch.*, **36**, 518.  
*Jacobæa*, *Linn.*, mentioned, **23**, 450.  
— var. *grandiflorus*, *Turcz.*, **23**, 450.  
*Jamesii*, *Hemsl.* \*, **23**, 453.  
*japonicus*, *Sch.-Bip.*, **23**, 453.  
*Kæmpferi*, *DC.*, **23**, 454.  
*kanaitzensis*, *Franch.*, **36**, 518.  
*kansuensis*, *Franch.*, **36**, 518.  
*Kaschkarowi*, *Winkl.*, **36**, 518.  
*kematongensis*, *Vaniot*, **36**, 518.  
*kialensis*, *Franch.*, **36**, 518.  
*Kirilowii*, *Turcz.*, **23**, 451.  
*koreanus*, *Komarov*, **36**, 518.  
*koualapensis*, *Franch.*, **36**, 518.  
*Labordei*, *Vaniot*, **36**, 518.  
*lankongensis*, *Franch.*, **36**, 518.  
*lapathifolius*, *Franch.*, **36**, 518.  
*latipes*, *Franch.*, **36**, 518.  
*Leucanthemum*, *Dunn*, **36**, 518.  
*leucocephalus*, *Franch.*, **36**, 518.  
*Leveillei*, *Vaniot*, **36**, 518.  
*liatroides*, *Winkl.*, **36**, 518.  
*Ligularia*, *Hook. f.*, **23**, 454; mentioned, 455.  
— var. *polycephalus*, *Hemsl.* \*, **23**, 455.  
*Lucorum*, *Franch.*, **36**, 518.  
*luticola*, *Dunn*, **36**, 518.  
*macranthus*, *C. B. Clarke*, **23**, 453.  
*Martini*, *Vaniot*, **36**, 518.  
*Maximowiczii*, *Franch.*, **36**, 518.  
*melanocephalus*, *Franch.*, **36**, 518.  
*microdontus*, *Bur. & Franch.*, **36**, 518.  
*monanthus*, *Diels*, **36**, 518.  
*mongolicus*, *Sch.-Bip.*, **23**, 455.  
*Senecio mosoynensis*, *Franch.*, **36**, 518.  
*nelumbifolius*, *Bur. & Franch.*, **36**, 518.  
*nemorensis*, *Linn.*, **23**, 455.  
*nigrocinctus*, *Franch.*, **36**, 519.  
*Nimborum*, *Franch.*, **36**, 519.  
*nobilis*, *Franch.*, **36**, 519.  
*Oldhamianus*, *Maxim.*, **23**, 455.  
*palmatus*, *Less.*, **23**, 458.  
*palmatus*, *Pall.*, **23**, 456.  
*paucinervis*, *Dunn*, **36**, 519.  
*phænicochætus*, *Franch.*, **36**, 519.  
*phalacrocarpus*, *Hance*, **23**, 456.  
*phyllolepis*, *Franch.*, **36**, 519.  
*Pierotii*, *Miq.*, **23**, 451.  
*plantaginifolius*, *Franch.*, **36**, 519.  
*platyglossus*, *Franch.*, **36**, 519.  
*pleurocaulis*, *Franch.*, **36**, 519.  
*Potanini*, *Winkl.*, **36**, 519.  
*pratensis*, *DC.*, **23**, 451.  
*Prattii*, *Hemsl.*, **36**, 519.  
*Principis*, *Franch.*, **36**, 519.  
*prionophyllus*, *Franch.*, **36**, 519.  
*Profundorum*, *Dunn*, **36**, 519.  
*Przewalskii*, *Maxim.*, **23**, 456; mentioned, 453.  
*Pseudochina*, *Andr.*, **23**, 457.  
*Pseudosonchus*, *Vaniot*, **36**, 519.  
*pteridophyllus*, *Franch.*, **36**, 519.  
*Putjatac*, *Winkl.*, **36**, 519.  
*quinquelobus*, *Hook. f. & Thoms.*, **36**, 519.  
*racemosus*, *Wall.*, **23**, 454.  
*renatus*, *Franch.*, **36**, 519.  
*Roborowskii*, *Maxim.*, **23**, 456.  
*rubescens*, *S. Moore*, **23**, 456.  
*ruficomus*, *Franch.*, **36**, 519.  
*rufipilis*, *Franch.*, **36**, 519.  
*Sagitta*, *Maxim.*, **23**, 456.  
*sagittatus*, *Sch.-Bip.*, **23**, 456.  
*Savatieri*, *Franch.*, **23**, 455.  
*scandens*, *Buch.-Ham.*, **23**, 457.  
*Schmidtii*, *Franch. & Sav.*, **23**, 457.  
*setchuenensis*, *Franch.*, **36**, 519.  
*sibiricus*, *C. B. Clarke*, **23**, 454.  
*solenoides*, *Dunn*, **36**, 519.  
*sonchifolius*, *Moench*, **23**, 449.  
*Souliei*, *Franch.*, **36**, 519.  
*sp. ?*, **23**, 458.  
*spathiphyllus*, *Franch.*, **36**, 519.  
*spathulæfolius*, *Turcz.*, **23**, 451.  
*speciosus*, *Willd.*, South African, **23**, 457.  
*Stauntoni*, *DC.*, **23**, 458.  
*stenocephalus*, *Maxim.*, **23**, 458; mentioned, 458.  
*stenoglossus*, *Franch.*, **36**, 519.  
*stipulatus*, *Wall.*, **23**, 463.  
*subensiformis*, *DC.*, **23**, 451.

- Senecio subspicatus*, *Bur. & Franch.*, **33**, 519.  
*Syneilesis*, *Franch. & Sav.*, **23**, 458; mentioned, 450.  
*taliensis*, *Franch.*, **36**, 519.  
*talongensis*, *Franch.*, **36**, 519.  
*tanguticus*, *Maxim.*, **23**, 458.  
*tatsienensis*, *Bur. & Franch.*, **36**, 519.  
*tenuipes*, *Franch.*, **36**, 519.  
*tongolensis*, *Franch.*, **36**, 519.  
*tricuspis*, *Franch.*, **36**, 519.  
*tsangchanensis*, *Franch.*, **36**, 519.  
*Turczaninowii*, *Walp.*, **23**, 449.  
*vellereus*, *Franch.*, **36**, 519.  
*Vespertilio*, *Franch.*, **36**, 519.  
*villiferus*, *Franch.*, **36**, 519.  
*Virgaurea*, *Maxim.*, **23**, 458.  
*Wightianus*, *DC.*, **23**, 457.  
*yesoënsis*, var., *Franch.*, **36**, 519.  
*yunnanensis*, *Franch.*, **36**, 519.  
*Serapias camtschatea*, *Steud.*, **36**, 40.  
*erecta*, *Thunb.*, **36**, 48.  
*falcata*, *Thunb.*, **36**, 48.  
*Seriola*, *D. Don*, **23**, 479.  
*Serissa Democritea*, *Baill.*, **23**, 391.  
*foetida*, *Comm.*, **23**, 391.  
*japonica*, *Thunb.*, **23**, 391.  
*Serpicula verticillata*, *Roxb.*, **36**, 1.  
*Serratula*, *Dill.*, **23**, 466.  
*atriplicifolia*, *Benth. & Hook. f.*, **23**, 468.  
*carthamoides*, *Buch.-Ham.*, **23**, 463.  
*centauroides*, *Linn.*, **23**, 469.  
*chinensis*, *S. Moore*, **23**, 469.  
*coronata*, *Linn.*, **23**, 469; **36**, 519.  
*japonica*, *Thunb.*, **23**, 464.  
*Martini*, *Vaniot*, **36**, 520.  
*multiflora*, *Lour.*, **23**, 469.  
*pungens*, *Franch. & Sav.*, **23**, 469.  
*radiata*, *Bieb.*, **23**, 469.  
*Scordium*, *Lour.*, **23**, 469.  
*tinctoria*, *Sieb.*, **23**, 463.  
*Sesamum indicum*, *Linn.*, **26**, 236.  
*Sesbania aculeata*, *Pers.*, **23**, 162.  
*ægyptiaca*, *Pers.*, **23**, 162.  
*Seseli bengalensis*, *Roxb.*, **23**, 331.  
*daucifolium*, *O. B. Clarke*, partim, **23**, 333.  
*Delavayi*, *Franch.*, **36**, 520.  
*Giraldii*, *Diels*, **36**, 520.  
*Libanotis*, *Koch*, **23**, 330.  
*Provostii*, *Boissieu*, **36**, 520.  
*sibiricum*, *Boiss.*, **23**, 330.  
*yunnanense*, *Franch.*, **36**, 520.  
*Sesuvium longifolium*, *Willd.*, **23**, 323.  
*Sesuvium Portulacastrum*, *Linn.*, **23**, 323; mentioned, 324.  
*repens*, *Willd.*, **23**, 323.  
*revolutifolium*, *Ort.*, **23**, 323.  
*Setaria Forbesiana*, *Hook. f.*, **36**, 334.  
*glauca*, *Beauv.*, **36**, 335.  
*italica*, *Beauv.*, **36**, 335.  
*macrostachya*, *H. B. K.*, mentioned, **36**, 334.  
— var. *Schultzii*, *Benth.*, **36**, 334.  
*mauritiana*, *Spreng.*, **36**, 336.  
*viridis*, *Beauv.*, **36**, 336.  
*Sethia indica*, *DC.* (errore *Lethia*), **23**, 96.  
*Severinia buxifolia*, *Ten.*, **23**, 110.  
*Sheareria nana*, *S. Moore*, **23**, 432.  
*Polii*, *Franch.*, **23**, 432.  
*Shortia rotundifolia*, *Makino*, **36**, 520.  
*sinensis*, *Hemsl.*, **36**, 520.  
*thibetica*, *Franch.*, **36**, 520.  
*Shuteria bicolor*, *Choisy*, **26**, 163.  
*longipes*, *Franch.*, **36**, 520.  
*sinensis*, *Hemsl.*, **36**, 520.  
*Sibbaldia erecta*, *Linn.*, **23**, 246.  
*procumbens*, *Linn.*, **36**, 520.  
*Sida Abutilon*, *Linn.*, **23**, 86.  
*acuta*, *Burm. f.*, **23**, 84.  
*asiatica*, *Linn.*, **23**, 86.  
*carpinifolia*, *Linn.*, **23**, 84.  
*chinensis*, *Retz.*, **23**, 85.  
*cordifolia*, *Linn.*, **23**, 85.  
*fallax*, *Walp.*, **23**, 85.  
*humilis*, *Willd.*, **23**, 85.  
*indica*, *Linn.*, **23**, 86.  
*philippica*, *DC.*, **23**, 85.  
*planicaulis*, *Linn.*, **23**, 84.  
*rhubifolia*, *Linn.*, **23**, 85.  
*rotundifolia*, *Hook. & Arn.*, **23**, 85.  
*spinosa*, *Linn.*, mentioned, **23**, 86.  
*Stantoniana*, *DC.*, **23**, 85.  
*stipulata*, *Cav.*, **23**, 85.  
*tiliaefolia*, *Fisch.*, **23**, 86.  
*Sideroxylon attenuatum*, *A. DC.*, **26**, 68.  
*ferrugineum*, *Hook. & Arn.*, **26**, 68.  
*Wightianum*, *Hook. & Arn.*, **26**, 68.  
*Siegesbeckia orientalis*, *Linn.*, **23**, 433.  
*Silene adenantha*, *Franch.*, **36**, 520.  
*aprica*, *Turcz.*, **23**, 64.  
*asclepiadea*, *Franch.*, **36**, 520.  
*cæspitosa*, *Bur. & Franch.*, **36**, 520.  
*capitata*, *Komarov*, **36**, 520.  
*cardiopetala*, *Franch.*, **36**, 520.  
*chinensis*, *Hort.*, **36**, 520.

- Silene conoidea*, Linn., 23, 65.  
*Delavayi*, Franch., 36, 520.  
*firma*, Sieb., 23, 65.  
*foliosa*, Maxim., 36, 520.  
*Fortunei*, Vis., 23, 65.  
*grandifolia*, Franch., 36, 520.  
*koreana*, Komarov, 36, 520.  
*langkongensis*, Franch., 36, 520.  
*lutea*, Franch., 36, 520.  
*macrostyla*, Maxim., 36, 520.  
*melanantha*, Franch., 36, 520.  
*melandryiformis*, Maxim., 23, 65.  
*napuligera*, Franch., 36, 520.  
*Oldhamiana*, Miq., 23, 64.  
*otodonta*, Franch., 36, 520.  
*pachyrrhiza*, Franch., 36, 520.  
*phænicodonta*, Franch., 36, 520.  
*platypetala*, Bur. & Franch., 36, 520.  
*platyphylla*, Franch., 36, 520.  
*Potanini*, Maxim., 36, 520.  
*pterosperma*, Maxim., 36, 520.  
*repens*, Patrin, 36, 520.  
*ruricunda*, Franch., 36, 520.  
*Scopulorum*, Franch., 36, 520.  
*sinensis*, Williams, 36, 520.  
*szechuensis*, Williams, 36, 520.  
*Tanakæ*, Maxim., 36, 520.  
*Tatarinowii*, Regel, 23, 65.  
*tenuis*, Willd., 36, 521.  
*trachyphylla*, Franch., 36, 521.  
*viscidula*, Franch., 36, 521.  
*yunnanensis*, Franch., 36, 521.  
*Siler divaricatum*, Benth. & Hook. f., 23, 332.  
SIMARUBEÆ, 23, 112.  
*Simmondsia californica*, Nutt., mentioned, 26, 419.  
*Sinapis brassicata*, Linn., 23, 47.  
*chinensis*, Linn., 23, 47.  
*juncea*, Linn., 23, 47.  
*pekinensis*, Lour., 23, 46.  
*Sindechites Henryi*, Oliver, 26, 100.  
*Siphonanthus indica*, Linn., 26, 262.  
*Siphonostegia chinensis*, Benth., 26, 202.  
*chinensis*, var., S. Moore, 26, 203.  
*læta*, S. Moore, 26, 203.  
*sp.*, S. Moore, 26, 203.  
*Sisymbrium atrovirens*, Hornem., 23, 40.  
*heteromallum*, C. A. Mey., 23, 45.  
*Irio*, Linn., 23, 45.  
*junceum*, Bieb., 23, 45.  
*Maximowiczii*, Palibin, 36, 521.  
*Piasezkii*, Maxim., 23, 45.  
*Sophia*, Linn., 23, 46.  
*Thalianum*, J. Gay, 36, 521.  
*Sium cicutæfolium*, C. C. Gmel., 23, 328.  
*græcum*, Linn., 23, 329.  
*latifolium*, Linn., mentioned, 23, 328.  
*lineare*, Michx., 23, 328.  
*Matsumuræ*, Boissieu, 36, 54.  
*Ninsi*, Linn., 36, 521.  
*nipponicum*, Maxim., 36, 521.  
*Sisarum*, Linn., 23, 329.  
*triternatum*, Miq., 23, 342.  
*Skimmia Fortunei*, Mast., 36, 521.  
*japonica*, Thunb., 36, 521.  
*Laureola*, Franch., 36, 521.  
*Sladenia celastrifolia*, Kurz, 23, 76.  
*Slateria Jaburan*, Sieb., 36, 77.  
*japonica*, Rich., 36, 78.  
*Sloanea hongkongensis*, Hemsl., 36, 521.  
*Smilacina*=*Tovaria*, 36, 521.  
*bifolia*, Schult., 36, 112.  
*Fargesii*, Diels, 36, 521.  
*hirta*, Maxim., 36, 111.  
*japonica*, A. Gray, 36, 110, 521.  
— var. *mandschurica*, Maxim., 36, 111.  
*Rossii*, Maxim., 36, 111.  
*stenoloba*, Diels, 36, 521.  
*tubifera*, Batalin, 36, 111, 521.  
*Smilax Bockii*, Warb., 36, 521.  
*brevipes*, Warb., 36, 521.  
*californica*, A. Gray, 36, 100.  
*calophylla*, Wall., 36, 96.  
— var. *concolor*, C. H. Wright\*, 36, 96.  
*China*, Linn., 36, 96.  
*cinerea*, Warb., 36, 521.  
*cocculoides*, Warb., 36, 521.  
*cyclophylla*, Warb., 36, 521.  
*Davidiana*, A. DC., 36, 97.  
*discotis*, Warb., 36, 521.  
*ferox*, Wall., 36, 96.  
*flaccida*, C. H. Wright, 36, 97.  
*Gaudichaudiana*, Kunth, 36, 95.  
*glabra*, Roxb., 36, 97.  
*glauco-china*, Warb., 36, 521.  
*herbacea*, var. *acuminata*, C. H. Wright\*, 36, 97.  
— var. *angusta*, C. H. Wright\*, 36, 97.  
— var. *intermedia*, C. H. Wright\*, 36, 97.  
— var. *nipponica*, Maxim., 36, 98.  
— var. *oblonga*, C. H. Wright\*, 36, 98.



- Smilax herbacea*, var. *Oldhami*, Maxim., 36, 100.  
 — var. *pubescens*, C. H. Wright\*, 36, 98.  
*hongkongensis*, Seem., 36, 95.  
*hypoglauca*, Benth., 36, 98.  
*japonica*, A. Gray, 36, 96.  
*lævis*, var. *ophirensis*, A. DC., 36, 99.  
 — var. *Parkii*, A. DC., 36, 99.  
*lanceifolia*, Roxb., var. *opaca*, A. DC., 36, 99.  
*lanceifolia*, Seem., 36, 99.  
*longipes*, Warb., 36, 521.  
*megalantha*, C. H. Wright, 36, 99.  
*menispermoidea*, A. DC., 36, 99.  
*microphylla*, C. H. Wright, 36, 99.  
*Nebelii*, Gilg, 36, 521.  
*nipponica*, Miq., 36, 98.  
*obtusa*, Benth., mentioned, 36, 99.  
*Oldhami*, Miq., 36, 100.  
 — var. *ussuriensis*, A. DC., 36, 100.  
*ovalifolia*, Roxb., 36, 100.  
*parvifolia*, Wall., mentioned, 36, 97.  
*pekingensis*, A. DC., 36, 100.  
*perfoliata*, Lour., 36, 100.  
*polycolea*, Warb., 36, 521.  
*riparia*, A. DC., 36, 100.  
*scobinicaulis*, C. H. Wright, 36, 100.  
*Sieboldi*, Miq., 36, 100.  
*stans*, Maxim., 36, 101.  
*stenopetala*, A. Gray, 36, 101; mentioned, 99.  
*trigona*, Warb., 36, 521.  
*ussuriensis*, A. DC., 36, 100.  
*vaginata*, Decne., 36, 101.  
*Smithia conferta*, Sm., 23, 170.  
*dichotoma*, Dalz., 23, 170.  
*geminiflora*, Roth, mentioned, 23, 170.  
*salsuginea* Hance, 23, 170.  
*sensitiva*, Ait., 23, 170.  
*yunnanensis*, Franch., 36, 521.  
*Soja hispida*, Moench, 23, 188.  
**SOLANACEÆ**, 26, 169.  
*Solanum athiopicum*, var. *violaceum*, Dunal, 26, 169 in note.  
*biflorum*, Lour., 26, 169; mentioned, 171.  
*Calleryanum*, Choisy, 26, 169.  
*chinense*, Dunal, 26, 171.  
*coagulans*, Forsk., 26, 170.  
*decemdentatum*, Roxb., 26, 169.  
*dichotomum*, Lour., 26, 170.  
*Dulcamara*, Linn., 26, 169.  
*Solanum Dulcamara*, var. *chinensis*, Dunal, 26, 169.  
*esculentum*, Dunal, 26, 169.  
*ferox*, Linn., 26, 170.  
*hainanense*, Hance, 26, 170.  
*immane*, Hance, 26, 170.  
*incanum*, Linn., 26, 170.  
*incanum*, Pluk., 26, 171.  
*indicum*, Linn., 26, 170.  
*lasiocarpum*, Dunal, 26, 170.  
*lyratum*, Thunb., 26, 169.  
*lysimachioides*, Wall., 26, 171.  
*macaonense*, Dunal, 26, 172.  
*macranthum*, Dunal, mentioned, 26, 172.  
*macranthum*, [Hook.] Bot. Mag., 26, 172.  
*macranthum*, Hort., 26, 172.  
*maroniense*, Poiteau, 26, 172.  
*Melongena*, Linn., mentioned, 26, 169.  
*nigrum*, Linn., 26, 171.  
*nodiflorum*, Jacq., 26, 171.  
*Osbeckii*, Choisy, 26, 169.  
*persicum*, Willd., 26, 170.  
*pittosporifolium*, Hemsl. \*, 26, 171.  
*sanctum*, Linn., 26, 170.  
*septemlobum*, Bunge, 26, 172.  
*torvum*, Sw., 26, 172.  
*verbascifolium*, Linn., 26, 172.  
*violaceum*, Ortega, 26, 171.  
*Wrightii*, Benth., South-American, 26, 172.  
*xanthocarpum*, Schrad. & Wendl., 26, 173.  
*Solena heterophylla*, Lour., 23, 318.  
*Solidago cantoniensis*, Lour., 23, 406.  
*decurrens*, Lour., 23, 406.  
*leiocarpa*, Miq., 23, 406.  
*Virgaurea*, Linn., 23, 406.  
*Sonchus arvensis*, Linn., 23, 48.  
 — var. *uliginosa*, Herd., 23, 48.  
*asper*, Vill., 23, 487.  
*brachyotus*, DC., 23, 487.  
*ciliatus*, Lam., 23, 487.  
*floridanus*, Lour., 23, 480.  
*lactuoides*, Bunge, 23, 487; mentioned, 414.  
*oleraceus*, Linn., 23, 487.  
*sibiricus*, Linn., 23, 484.  
*tataricus*, Linn., 23, 484.  
*uliginosus*, Bieb., 23, 487.  
*Sonerila elegans*, Wight, mentioned, 23, 301.  
*Fordii*, Oliver, 23, 301.  
*peperomiæfolia*, Oliver, 36, 521.  
*spp.?*, 23, 301, 302.  
*tenera*, R. Br., 23, 301.  
*Sophora alopecuroides*, Linn., 23, 201.

- Sophora angustifolia*, Sieb. & Zucc., **23**, 202.  
*fabacea*, Pall., **23**, 150.  
*flavescens*, Ait., **23**, 201, 202.  
*galegoides*, Pall., **23**, 202.  
*glabra*, Hassk., **23**, 203.  
*glauca*, Leschen., **36**, 521.  
*heptaphylla*, Linn., mentioned, **23**, 201.  
*japonica*, Linn., **23**, 202.  
*Kronei*, Hance, **23**, 202.  
*mollis*, R. Grak., **23**, 203.  
*Moorcroftiana*, Benth., **23**, 203; mentioned, 204.  
— var. *Davidii*, Franch., **23**, 203.  
*pachycarpa*, C. A. Mey., **23**, 203.  
*sororia*, Hance, **23**, 202.  
*tomentosa*, Linn., **23**, 203; mentioned, 201.  
*viciifolia*, Hance, **23**, 203.  
*Sophorocapnos pallida*, Turcz., **23**, 38.  
*Sopubia trifida*, Buch.-Ham., **26**, 202.  
*Sorbaria Kirilowii*, Maxim., **23**, 227.  
*sorbifolia*, Maxim., **23**, 227.  
*Sorbus Aria*, Crantz, **23**, 254.  
*Aucuparia*, Linn., **23**, 255.  
*pekinensis*, Koehne, **36**, 521.  
*Sorghum fulvum*, Beauv., **36**, 367.  
— var. *nitidum*, Hack., **36**, 367.  
*halepense*, Pers., **36**, 367.  
*nitidum*, Pers., **36**, 367.  
*vulgare*, Pers., **36**, 367.  
— var. *bicolor*, Hack., **36**, 368.  
— var. *nervosum*, Hack., **36**, 368.  
— var. *obovatum*, Hack., **36**, 368.  
*Souliea*, Franch., **36**, 521.  
*vaginata*, Franch., **36**, 521.  
*Southwellia nobilis*, Salisb., **23**, 89.  
*Spanioptilon lineare*, Less., **23**, 461.  
*Spathium chinense*, Lour., **26**, 363.  
*monostachyon*, Edgew., **36**, 192.  
*Spathodea Cauda-felina*, Hance, **26**, 235.  
*Spathoglottis Fortunei*, Lindl., **36**, 18.  
*pubescens*, Lindl., **36**, 18.  
*Spatholobus pulcher*, Dunn, **36**, 521.  
*suberectus*, Dunn, **36**, 521.  
*varians*, Dunn, **36**, 521.  
*Speirantha convallarioides*, Baker, **36**, 112.  
*Spenceria ramalana*, Trimen, **23**, 246.  
*Speranskia Henryi*, Oliver, **26**, 435.  
*tuberculata*, Baill., **26**, 436.  
*Spergularia media*, G. Don, **23**, 70.  
*Spermacoe articularis*, Linn., **23**, 392.  
*hedyotideia*, DC., mentioned, **23**, 373.  
*hispida*, Linn., **23**, 392.  
*Spermacoe lasiocarpa*, R. Br., **23**, 392.  
*philippensis*, Willd., **23**, 392.  
*scabra*, Willd., **23**, 392.  
*semierecta*, Roxb., **23**, 392.  
*sp.*, **23**, 392.  
*stricta*, Linn., **23**, 392.  
*Spermedietyon azureum*, Wall., **23**, 390.  
*suaveolens*, Roxb., **23**, 390.  
*Sphæranthus africanus*, Linn., **23**, 423.  
*chinensis*, Linn., Indian only, **23**, 423.  
*cochinensis*, Lour., **23**, 423.  
*microcephalus*, Willd., **23**, 423.  
*Sphærocaryum elegans*, Nees, **36**, 387.  
*Spheromorpha Centipeda*, DC., **23**, 440.  
*petiolaris*, DC., **23**, 423.  
*Sphaerophysa Salsula*, DC., **23**, 162.  
*Sphæroschænus Wallichii*, Arn., **36**, 261.  
*Sphaerostema japonica*, Sieb., **23**, 25.  
*Sphallerocarpus Cyminum*, Bess., **23**, 330.  
*Sphenoclea pongatium*, A. DC., **26**, 8.  
*zeylanica*, Gaertn., **26**, 8.  
*Sphenodesma acuminata*, Wight, **26**, 265.  
*pentandra*, Jack, **26**, 265.  
*unguiculata*, Schau., **26**, 265.  
*Spilanthus tinctorius*, Lour., **23**, 403.  
*Spinacia oleracea*, Linn., mentioned, **26**, 325.  
*Spinifex squarrosus*, Linn., **36**, 340.  
*Spinovitis Davidii*, Carr., **23**, 136.  
*Spiræa alpina*, Pall., **36**, 522.  
*angustiloba*, Turcz., **23**, 223.  
*anomala*, Batalin, **36**, 522.  
*Aruncus*, Linn., **23**, 223; **36**, 522;  
*Astilbe polyandra*, Hemsl., a state of, **23**, 489.  
*betulæfolia*, Pall., **23**, 223.  
*Blumei*, G. Don, **23**, 223; mentioned, 225.  
— var. *hirsuta*, Hemsl. \*, **23**, 224.  
— var. *latipetala*, Hemsl. \*, **23**, 224.  
— var. *rotundifolia*, Hemsl. \*, **23**, 224.  
*callosa*, Thunb., **23**, 225.  
*canescens*, D. Don, var., **36**, 522.  
*cantoniensis*, Lour., **23**, 224.  
*chamædryfolia*, Blume, **23**, 224.  
*chamædryfolia*, Cham. & Schlecht., **23**, 223.  
*chamædryfolia*, Thunb., **23**, 224.  
*chinensis*, Maxim., **23**, 225.  
*corymbosa*, Roxb., **23**, 224.

- Spiræa crenifolia*, var. *mongolica*, Maxim., **23**, 226.  
*dasyantha*, Bunge, **23**, 224; mentioned, 225, 227.  
*digitata*, Willd., **23**, 226.  
*Fortunei*, Planch., **23**, 226.  
*grandiflora*, Hook., **23**, 223.  
*Henryi*, Hemsl. \*, **23**, 225, 491.  
*incisa*, Carr., **23**, 228.  
*japonica*, Camb., **23**, 229.  
*japonica*, Linn., **23**, 225; **36**, 522.  
*japonica*, Sieb., **23**, 224.  
*Kirilowii*, Regel, **23**, 227.  
*lanceolata*, Poir., **23**, 224.  
*longigemmis*, Maxim., **23**, 226.  
*media*, var. *sericea*, Hance, **23**, 227.  
*mongolica*, Maxim., **23**, 226.  
*palmata*, Pall., **23**, 226.  
*prostrata*, Maxim., **23**, 226.  
*prunifolia*, Sieb. & Zucc., **23**, 224; mentioned, 228.  
*pubescens*, Lindl., **23**, 225.  
*pubescens*, Turcz., **23**, 227; mentioned, 225.  
*Reevesiana*, Lindl., **23**, 224.  
*Rosthornii*, E. Pritz., **36**, 522.  
*salicifolia*, Linn., **23**, 227.  
*sericea*, Turcz., **23**, 227.  
*sorbifolia*, Linn., **23**, 227.  
*sp.*, **23**, 228.  
*Thunbergii*, Sieb., **23**, 227.  
*trilobata*, Linn., **23**, 288.  
*Veitchi*, Hemsl., **36**, 522.  
*velutina*, Franch., **36**, 522.  
*vestita*, Wall., **36**, 522.  
*virgata*, Franch., **36**, 522.  
*yunnanensis*, Franch., **36**, 522.
- Spiranthes amœna*, Bunge, **36**, 41.  
*australis*, Lindl., **36**, 41.  
— var. *pubida*, Lindl., **36**, 41.  
*exigua*, Rolfe, **36**, 41.  
*pubida*, Lindl., **36**, 41.
- Spirodela polyrrhiza*, Schleid., **36**, 188.
- Spodiopogon albidus*, Benth., mentioned, **36**, 553.  
*aureus*, Hook. & Arn., **36**, 365.  
*cotulifer*, Hack, **36**, 351; mentioned, 353.  
*depauperatus*, Hack., mentioned, **36**, 353.  
*formosanus*, Rendle \*, **36**, 351.  
*obliquivalvis*, Nees, **36**, 365.  
*sagittifolia*, Rendle \*, **36**, 352.  
*sibiricus*, Trin., **36**, 353.
- Sponia amboinensis*, Decne., **26**, 451.  
*argentea*, Planch., **26**, 451.  
*argentea*, Seem., **26**, 451.  
*nudiflora*, Sieb. & Zucc., **26**, 452.  
*orientalis*, Planch., **26**, 451.
- Sponia Sampsoni*, Hance, **26**, 452.  
*velutina*, Planch., **26**, 451.  
*virgata*, Planch., **26**, 452.
- Spondias mangifera*, Willd., **23**, 149.
- Sporobolus ciliatus*, Presl, **36**, 388.  
*diander*, Beauv., **36**, 387.  
*elongatus*, Franch., **36**, 388.  
*Hancei*, Rendle \*, **36**, 307.  
*indicus*, R. Br., **36**, 388.  
*japonicus*, Maxim. \*, **36**, 388.  
*orientalis*, Kunth, mentioned, **36**, 388.  
*virginicus*, Kunth, **36**, 389.
- Sportella atalantoides*, Hance, **23**, 260.
- Stachyopogon pauciflorus*, Klotzsch, **36**, 76.  
*spicatus*, Klotzsch, **36**, 76.
- Stachys adulterina*, Hemsl. \*, **26**, 300.  
*affinis*, Bunge, **26**, 301, 302.  
*Artemisia*, Lour., **26**, 203.  
*arvensis*, Linn., **26**, 300.  
*aspera*, Michx., **26**, 300; mentioned, 302.  
*baicalensis*, Fisch., **26**, 301.  
*chinensis*, Bunge, **26**, 301.  
*japonica*, Miq., **26**, 301.  
*modica*, Hance, **26**, 301.  
*oblongifolia*, Benth., **26**, 301.  
*palustris*, Linn., **26**, 301, 302.  
— var. *hispida*, Miq., **26**, 301.  
— var. *hispida*, Regel, **26**, 301.  
*Sieboldi*, Miq., **26**, 301; mentioned, 300.  
*tuberifera*, Naud., **26**, 301, 302.
- Stachytarpha indica*, Schau., **26**, 251.  
*jamaicensis*, Schau., **26**, 251.
- Stachytarpheta indica*, Vahl, **26**, 251.  
*jamaicensis*, Vahl, **26**, 251.
- Stachyurus chinensis*, Franch., **36**, 522.  
*himalaicus*, Hook. f. & Thoms., **36**, 522.  
*præcox*, Sieb. & Zucc., **23**, 79.  
*salicifolius*, Franch., **36**, 522.  
*yunnanensis*, Franch., **36**, 522.
- Stalagmitis erosipetala*, Miq., **23**, 75.
- Stapelia chinensis*, Lour., **26**, 116.
- Staphylea Bumalda*, Sieb. & Zucc., **36**, 522.  
*holocarpa*, Hemsl., **36**, 522.  
*simplicifolia*, Gardn., **23**, 143.
- Statice aurea*, Linn., **26**, 34.  
*bicolor*, Bunge, **26**, 35.  
— var. *densiflora*, Bunge, **26**, 35.  
— var. *laxiflora*, Bunge, **26**, 35.  
*Bungeana*, Boiss., **26**, 35.  
*Fortuni*, Lindl., **26**, 35.  
*Franchetii*, Debeaux, **26**, 35.

- Statice japonica*, Sieb. & Zucc., 26, 35.  
*sinensis*, Girard, 26, 35.  
*Wrightii*, Hance, 26, 35.  
*Stauntonia chinensis*, DC., 23, 30.  
*hexaphylla*, Decne., 23, 30.  
*Stauranthera chiritæflora*, Oliver, 36, 522.  
*Stegosia cochinchinensis*, Lour., 36, 362.  
*Stellaria Alsine*, Hoffm., 23, 69.  
*bulbosa*, Wulf., mentioned, 23, 69.  
*Bungeana*, Fenzl, 23, 68.  
*chinensis*, Regel, 23, 67.  
*Davidi*, Hemsl. \*, 23, 67.  
*Delavayi*, Franch., 36, 522.  
*dichasioides*, Williams, 36, 522.  
*dichotoma*, Linn., 23, 67.  
*discolor*, Turcz., 23, 68.  
*diversifolia*, Maxim., 23, 68.  
— var. *gymnandra*, Franch., 23, 68.  
*ebracteata*, Komarov, 36, 522.  
*fecunda*, Hance, 23, 67.  
*florida*, Fisch., 36, 522.  
*glauca*, With., 23, 68.  
*graminea*, Linn., 23, 68.  
— var. *lanceolata*, Fenzl, 23, 68.  
*Henryi*, Williams, 36, 522.  
*heterophylla*, Hemsl. \*, 23, 68.  
*infracta*, Maxim., 36, 522.  
*leptophylla*, Hance, 23, 68.  
*longifolia*, Muhl., 36, 522.  
*media*, Cyr., 23, 68.  
*nemorum*, Linn., 23, 68.  
— var. *Bungeana*, Regel, 23, 68.  
*nutans*, Williams, 36, 522.  
*palustris*, Ehrh., 23, 68.  
*pilosa*, Franch., 36, 522.  
*radians*, Linn., 36, 522.  
*rhapnorrhiza*, Hemsl. \*, 23, 69.  
*rupestris*, Hemsl. \*, 23, 69.  
*saxatilis*, Buch.-Ham., 23, 69.  
*Souliei*, Williams, 36, 522.  
*uda*, Williams, 36, 522.  
*uliginosa*, Murr., 23, 69.  
*undulata*, Thunb., 23, 67.  
*vestita*, Kurz, 23, 69.  
*wushanensis*, Williams, 36, 522.  
*yunnanensis*, Franch., 36, 522.  
*Stellera*, 26, 399.  
*Chanæjasme*, Linn., 26, 401.  
*Stemodia capitata*, Benth., 26, 185.  
*gratioloides*, Benth., 26, 185.  
*Stemona acuta*, C. H. Wright, 36, 94.  
*erecta*, C. H. Wright, 36, 94.  
*japonica*, Franch. & Sav., 36, 95; mentioned, 94.  
*parviflora*, C. H. Wright, 36, 95.  
*Stemona sessiliflora*, Miq., mentioned, 36, 94.  
*tuberosa*, Lour., 36, 95.  
*Stenactis annua*, Less., 23, 418.  
*Stenocœlium divaricatum*, Turcz., 23, 332.  
*Stenophragma Thalianum*, Celak., 36, 522.  
*Stenosolenium saxatile*, Turcz., 26, 155.  
*Stephanandra chinensis*, Hance, 23, 228.  
*flexuosa*, Sieb. & Zucc., 23, 228.  
*incisa*, Zabel, 36, 522.  
*Stephania hernandifolia*, Walp., 23, 29.  
*longa*, Lour., 23, 29.  
*tetrandra*, S. Moore, 23, 30.  
*Stephanotis chinensis*, Champ., 26, 114.  
*Sterculia Balanghas*, Ait., 23, 89.  
*lanceolata*, Cav., 23, 89.  
*monosperma*, Vent., 23, 89.  
*nobilis*, Sm., 23, 89.  
*peltata*, G. Don, 23, 89.  
*platanifolia*, Linn., 23, 90.  
*pyriformis*, Bunge, 23, 90.  
STERCULIACEÆ, 23, 89.  
*Stereosantes*, Franch., 36, 522.  
*Delavayi*, Franch., 36, 522.  
*Souliei*, Franch., 36, 522.  
*yunnanensis*, Franch., 36, 522.  
*Stereospermum hypostictum*, Miq., var., 26, 236.  
*sinicum*, Hance, 26, 236.  
*sp.*, 26, 236.  
*Stillingfleetia sebifera*, Boj., 26, 445.  
*Stillingia discolor*, Champ., 26, 445.  
*japonica*, Sieb. & Zucc., 26, 446. 1  
*sebifera*, Michx., 26, 445.  
*sinensis*, Baill., 26, 445.  
*Stimpsonia chamædryoides*, Wright, 26, 46.  
*crispidens*, Hance, 26, 50.  
*Stipa Aliciae*, Kanitz, 36, 383.  
*Bungeana*, Trin., 36, 381.  
*capillata*, Linn., 36, 381.  
*consanguinea*, Trin. & Rupr., 36, 382.  
*Henryi*, Rendle \*, 36, 382.  
*mongholica*, Turcz., 36, 382.  
*pekinensis*, Hance, 36, 383.  
*sibirica*, Lam., 36, 383.  
*splendens*, Trin., mentioned, 36, 382.  
*Stipellaria breviovdes*, Benth., 26, 438.  
*Stranvæsia Calleryana*, Decne., 23, 264.  
*Davidiana*, Decne., 36, 522.  
*glaucescens*, Lindl., var., 36, 522.

- Stranvæsia undulata*, Decne., **23**, 264.  
*Stranvaisia digyna*, Sieb. & Zucc., **23**, 263.  
*Stratiotes acoroides*, Linn. f., **36**, 3.  
*alismoides*, Linn., **36**, 3.  
*Streblus asper*, Lour., **26**, 454.  
*cordatus*, Lour., **26**, 455.  
*Streptocarpus chinensis*, Franch., **36**, 523.  
*Streptocaulon calophyllum*, Wight, **26**, 101.  
*tomentosum*, Wight & Arn., **26**, 101.  
*Streptolirion longifolium*, Gagnep., **36**, 160.  
*volubile*, Edgew., **36**, 159.  
*Streptopus ajanensis*, Til., **36**, 523.  
*amplexicaulis*, Poir., **36**, 110.  
*amplexifolius*, DC., **36**, 110.  
*chinensis*, Sm., **36**, 143.  
*paniculatus*, Baker, **36**, 110.  
*parviflorus*, Franch., **36**, 523.  
*Striga coccinea*, Benth., **26**, 201.  
*hirsuta*, Benth., **26**, 201.  
*lutea*, Lour., **26**, 201.  
*Masuria*, Benth., **26**, 201.  
*parvula*, Miq., **26**, 201.  
*phænicea*, Benth., **26**, 201.  
*Strobilanthes apricus*, T. Anders., **26**, 239.  
*Championi*, T. Anders., **26**, 239.  
*consanguineus*, C. B. Clarke, mentioned, **26**, 241.  
*debilis*, Hemsl. \*, **26**, 239.  
*dimorphotrichus*, Hance, **26**, 240.  
*flaccidifolius*, Nees, **26**, 239; mentioned, 240.  
*flaccidus*, G. Mann, **26**, 239.  
*formosanus*, S. Moore, **26**, 240.  
*Henryi*, Hemsl. \*, **26**, 240.  
*aponicus*, Miq., **26**, 241.  
*latisepalus*, Hemsl. \*, **26**, 241.  
*oliganthus*, Miq., **26**, 241.  
*pentstemonoides*, T. Anders., **26**, 241.  
*radicans*, T. Anders., **26**, 242; mentioned, 239.  
*scaber*, Hance, **26**, 238.  
*Wallichii*, Nees, **26**, 242; mentioned, 241.  
*Strophanthus dichotomus*, var. *chinensis*, Ker-Gawl., **26**, 97.  
*divergens*, R. Grah., **26**, 97.  
*Strychnos angustiflora*, Benth., **26**, 121.  
*Colubrina*, var., Benth., **26**, 121.  
*paniculata*, Champ., **26**, 121.  
*Stuartia monadelphæa*, Sieb. & Zucc., **23**, 80.  
*Sturmia longipes*, Reichb. f., **36**, 7.  
*Sturmia nervosa*, Reichb. f., **36**, 7.  
**STYLIDIEÆ**, **26**, 1.  
*Stylidium chinense*, Lour., **23**, 344.  
*Kunthii*, Wall., **26**, 1.  
*sinicum*, Hance, **26**, 1.  
*uliginosum*, Sw., **26**, 1.  
*Stylis chinensis*, Poir., **23**, 344.  
*Stylocoryne attenuata*, Voigt, **23**, 380.  
*mollissima*, Walp., **23**, 381.  
*racemosa*, Cav., **23**, 382.  
*subsessilis*, A. Gray, **23**, 381.  
*Webera*, A. Rich., **23**, 380.  
*Stylodiscus trifoliatus*, Benn., **26**, 428.  
*Stylophorum japonicum*, Miq., **23**, 34.  
*Styphnolobium japonicum*, Schott, **23**, 202.  
**STYRACEÆ**, **26**, 72.  
*Styrax caloneurum*, Perk., **36**, 523.  
*dasyanthum*, Perk., **36**, 523.  
*formosanum*, Matsum., **36**, 523.  
*Fortunei*, Hance, **26**, 77.  
*grandiflorum*, Griff., **26**, 77.  
*Hemsleyanum*, Diels, **36**, 523.  
*hypoglaucum*, Perk., **36**, 523.  
*japonicum*, Sieb. & Zucc., **26**, 76.  
*macranthum*, Perk., **36**, 523.  
*Obassia*, Sieb. & Zucc., **26**, 76.  
*odoratissimum*, Champ., **26**, 76.  
*prunifolium*, Perk., **36**, 523.  
*serrulatum*, Roxb., **26**, 77.  
— var. *vestitum*, Hemsl. \*, **26**, 77.  
*sp. n.?*, **26**, 77.  
*suberifolium*, Hook. & Arn., **26**, 77.  
*Suæda australis*, Moq., **26**, 328.  
*glauca*, Bunge, **26**, 328; mentioned, 329.  
*indica*, Seem., **26**, 328.  
*linifolia*, Pall., **26**, 329.  
*maritima*, Dum., **26**, 329; mentioned, 328.  
*monoica*, Forsk., **26**, 329.  
*nudiflora*, Moq., **26**, 330.  
*Salsa*, Pall., **26**, 330.  
*Sugerokia japonica*, Miq., **36**, 140.  
*Swertia angustifolia*, Buch.-Ham., **26**, 138.  
*asarifolia*, Franch., **36**, 523.  
*bella*, Hemsl. \*, **26**, 138.  
*bifolia*, Batalin, **36**, 523.  
*binaculata*, C. B. Clarke, **26**, 139; mentioned, 140.  
*cærulea*, Royle, mentioned, **26**, 139.  
*calycina*, Franch., **36**, 523.  
*chinensis*, Franch., **26**, 139.  
*cuneata*, Wall., mentioned, **26**, 139.  
*Davidi*, Franch., **26**, 140; **36**, 523.  
*decora*, Franch., **36**, 523.

- Swertia Delavayi*, Franch., 36, 523.  
*diluta*, Benth., 26, 139.  
*dimorpha*, Franch., 36, 523.  
*erythrosticta*, Maxim., 26, 140.  
*gentianoides*, Franch., 36, 523.  
*gracilis*, Franch., 36, 523.  
*kouitchensis*, Franch., 36, 523.  
*Leduci*, Franch., 36, 523.  
*longipes*, Franch., 36, 523.  
*macrosperma*, C. B. Clarke, 36, 523.  
*membranifolia*, Franch., 36, 523.  
*Mussoti*, Franch., 36, 523.  
*oculata*, Hemsl. \*, 26, 140.  
*pubescens*, Franch., 36, 523.  
*pulchella*, Buch.-Ham., 26, 140.  
*punicea*, Hemsl. \*, 26, 140.  
*rotata*, Thunb., 26, 139.  
*stricta*, Franch., 36, 523.  
*Tashiroi*, Makino, 36, 523.  
*tetragona*, C. B. Clarke, 26, 141.  
*tetraptera*, Maxim., 26, 141.  
*vacillans*, Maxim., 26, 141.  
*Sycopsis sinensis*, Oliver, 36, 523.  
*Syllisium laxifolium*, Meyen & Schau., 26, 298.  
*Symphorema Jackianum*, Kurz, 26, 265.  
*unguiculatum*, Kurz, 26, 265.  
*Symphoglossum hastatum*, Turcz., 26, 105.  
*Symplocos acutangula*, Brand, 36, 523.  
*adenopus*, Hance, 26, 72.  
*alata*, Brand, 36, 523.  
*anomala*, Brand, 36, 523.  
*argentea*, Brand, 36, 523.  
*botryantha*, Franch., 26, 74; 36, 523.  
*caudata*, Wall., 26, 74.  
*congesta*, Benth., 26, 72.  
*crassifolia*, Benth., 26, 72.  
*cratægoides*, Buch.-Ham., 26, 72.  
*cuspidata*, Brand, 36, 523.  
*decora*, Hance, 26, 72.  
*ferruginea*, Roxb., 26, 72.  
*formosana*, Brand, 36, 523.  
*Fordii*, Hance, 26, 73.  
*Hamiltoniana*, Wall., 26, 74.  
*Henryi*, Brand, 36, 523.  
*Henscheli*, Benth., 26, 73.  
*japonica*, A. DC., 26, 73.  
— var. *crassifolia*, Benth., 26, 72.  
*lancifolia*, Sieb. & Zucc., 26, 73.  
*leptostachys*, Sieb. & Zucc., 26, 73.  
*leucophylla*, Brand, 36, 523.  
*lucida*, Sieb. & Zucc., 26, 73.  
*microcarpa*, Champ., 26, 73.  
*modesta*, Brand, 36, 523.  
*myrtacea*, Sieb. & Zucc., 26, 73.  
*Symplocos neriifolia*, Sieb. & Zucc., 26, 73.  
*paniculata*, Wall., 26, 72.  
*phyllocalyx*, C. B. Clarke, 26, 73.  
*propinqua*, Hance, 26, 74.  
*prunifolia*, Sieb. & Zucc., 26, 74.  
*racemosa*, Roxb., 26, 74.  
*ramosissima*, Wall., 26, 75.  
*setchuensis*, Brand, 36, 524.  
*sinica*, Ker-Gawl., 26, 74.  
— var. *vestita*, Hemsl. \*, 26, 75.  
*spectabilis*, Brand, 36, 524.  
*spicata*, Roxb., 26, 75.  
*sp. n. ?*, 26, 75.  
*stellaris*, Brand, 36, 524.  
*Swinhoeana*, Hance, 26, 74.  
*Tashiroi*, Matsum., 36, 524.  
*terminalis*, Brand, 36, 524.  
*urceolaris*, Hance, 26, 75.  
*yunnanensis*, Brand, 36, 524.  
*Synedrys ossea*, Lindl., 26, 510.  
*Syneilesis aconitifolia*, Maxim., 23, 449.  
*palmata*, Maxim., 23, 458.  
*Syringa amurensis*, Rupr., 26, 82.  
*chinensis*, Blume, 26, 82.  
*chinensis* ×, Willd., 26, 83.  
*correlata*, A. Br., 26, 83.  
*dubia*, Pers., 26, 83.  
*Emodi*, Wall., 26, 83.  
*microphylla*, Diels, 36, 524.  
*oblata*, Lindl., 26, 83.  
*pekinensis*, Rupr., 26, 82.  
*pubescens*, Turcz., 26, 83.  
— var., 36, 524.  
*rothomagensis*, A. Rich., 26, 83.  
*sempervirens*, Franch., 26, 84.  
*suspensa*, Thunb., 26, 82.  
*tomentella*, Bur. & Franch., 36, 524.  
*velutina*, Bur. & Franch., 36, 524.  
*velutina*, Komarov, 36, 524.  
*villosa*, Vahl, 26, 83.  
*vulgaris*, Linn., 26, 83.  
*yunnanensis*, Franch., 36, 524.  
*Syzygium buxifolium*, Hook. & Arn., 23, 298.  
*nervosum*, DC., 23, 297.  
*odoratum*, Hook. & Arn., 23, 297.  
*Szechenya lloydoides*, Kanitz, 36, 139.  
*Tabernaemontana coronaria*, R. Br., 26, 96.  
*mollis*, Hook. & Arn., 26, 96.  
*sp. ?*, 26, 96.  
*Tacca pinnatifida*, Forst., 36, 90.  
TACCACEÆ, 36, 90.  
*Tainia cordifolia*, Hook. f., 36, 20.  
*hongkongensis*, Rolfe, 36, 20.  
*macrantha*, Hook. f., 36, 21.

- Talauma obovata*, Hance, **23**, 23.  
*pumila*, Blume, **23**, 24.  
*Sieboldi*, Miq., **23**, 24.  
*Tamarindus indica*, Linn., **23**, 214.  
*officinalis*, Hook., **23**, 214.  
**TAMARISCINÆ**, **23**, 346.  
*Tamarix chinensis*, Lour., **23**, 346.  
*chinensis*, Sieb. & Zucc., **23**, 347.  
*gallica*, Linn., mentioned, **23**, 347.  
— var. *chinensis*, Ehrenb., **23**, 346.  
*indica*, Willd., **23**, 346.  
*juniperina*, Bunge, **23**, 347.  
*Pallasii*, Desv., **23**, 347; **36**, 524.  
*Tanacetum chinense*, A. Gray, **23**, 440.  
*myrianthum*, Franch., **36**, 524.  
*Pallasianum*, Trautv. & Mey., **36**, 524.  
*sibiricum*, Linn., **23**, 440.  
*Tapeinophallus Rivieri*, Baill., **36**, 182.  
*Tapiscia sinensis*, Oliver, **36**, 524.  
*Taraxacum ceratophorum*, DC., **23**, 478.  
*corniculatum*, DC., **23**, 478.  
*Dens-leonis*, Desf., **23**, 478.  
*officinale*, Web., **23**, 478.  
*sinense*, DC., **23**, 478.  
*Taxodium heterophyllum*, Brongn., **26**, 544.  
*japonicum*, Brongn., **26**, 544.  
— var. *heterophyllum*, Brongn., **26**, 544.  
*sinense*, Forbes, **26**, 544.  
*Taxotrophis*, Blume, sp.?, **26**, 454.  
*Taxus*, Tourn., **26**, 545.  
*baccata*, Linn., **26**, 546.  
*Harringtonia*, Knight, **26**, 545.  
*japonica*, Lodd., **26**, 546.  
*Tecoma grandiflora*, Loisel., **26**, 235.  
*Telanthera polygonoides*, Seem., **26**, 322.  
*Teloxys aristata*, Moq., **26**, 324.  
*Tenorea heterophylla*, Gasp., **26**, 466.  
*Tephrosia oraria*, Hance, **23**, 158, 489.  
*purpurea*, Pers., **23**, 158.  
*vestita*, Vogel, **23**, 158.  
*villosa*, Pers., **23**, 158.  
*Terminalia triptera*, Franch., **36**, 524.  
*Ternstroemia dubia*, Choisy, **23**, 75.  
*fragrans*, Choisy, **23**, 75.  
*japonica*, Thunb., **23**, 75.  
**TERNSTROEMIACEÆ**, **23**, 75.  
*Tetracentron sinense*, Oliver, **36**, 524.  
*Tetracera sarmentosa*, Vahl, **23**, 22.  
*Tetragonia expansa*, Murr., **23**, 323.  
*Tetramerium*, Nees, mentioned, **26**, 112.  
*Tetranthera citrata*, Nees, **26**, 379, 385.  
*citrifolia*, Benth., **26**, 385.  
*Cubeba*, Meisn., **26**, 380.  
*elongata*, Wall., **26**, 381.  
*floribunda*, Champ., **26**, 379.  
*japonica*, Spreng., **26**, 382.  
*laurifolia*, Jacq., **26**, 385.  
*monopetala*, Pers., **26**, 384.  
*polyantha*, Wall., **26**, 384.  
*Roxburghii*, Blume, **26**, 385.  
*sericea*, Wall., **26**, 386.  
*Tetrapanax papyrifera*, C. Koch, **23**, 341.  
*Tetrastigma Hemsleyanum*, Diels & Gilg, **36**, 524.  
*hypoglaucum*, Planch., **36**, 524.  
*obtectum*, Wall., **36**, 524.  
*Tetrathyrium subcordatum*, Benth., **23**, 291.  
*Teucrium albo-rubrum*, Hemsl. \*, **26**, 311.  
*bidentatum*, Hemsl. \*, **26**, 312, 313; **36**, 524.  
*Fortunei*, Benth., **26**, 314.  
*Franchetianum*, Diels, **36**, 524.  
*fulvum*, Hance, **26**, 314.  
*inflatum*, Sw., mentioned, **26**, 312.  
*japonicum*, Willd., **26**, 312.  
*nepetæfolium*, Benth., **26**, 264.  
*ningpoense*, Hemsl. \*, **26**, 313; mentioned, 314.  
*ornatum*, Hemsl. \*, **26**, 313.  
*palmatum*, Benth., **26**, 313.  
*Pernyi*, Franch., **26**, 314; mentioned, 313.  
*quadrifarium*, Buch.-Ham., **26**, 314.  
*stoloniferum*, Roxb., **26**, 314; mentioned, 312.  
*virginicum*, Thunb., **26**, 312.  
*Textoria japonica*, Miq., **23**, 343.  
*Teysmannia Hookeriana*, Miq., **26**, 97.  
*laxiflora*, Miq., **26**, 97.  
*Thalassia Hemprichii*, Aschers., **36**, 3.  
*Thalictrum actææfolium*, Sieb., mentioned, **23**, 10.  
*akanense*, Huth, **36**, 524.  
*alpinum*, Linn., **23**, 8.  
*amplissimum*, H. Lévillé & Vaniot, **36**, 524.  
*angustifolium*, Jacq., **36**, 524.  
*aquilegifolium*, Linn., **36**, 524.  
*baicalense*, Turcz., **23**, 8; mentioned, 10.  
*clematidifolium*, Franch., **36**, 524.  
*coraiense*, Matsum., **36**, 524.  
*coreanum*, H. Lévillé & Vaniot, **36**, 524.

- Thalictrum cultratum*, Wall., 36, 524.  
*Delavayi*, Franch., 36, 524.  
*diptero carpum*, Franch., 36, 524.  
*filamentosum*, Maxim., mentioned, 23, 10.  
*foeniculaceum*, Bunge, 23, 8.  
*fœtidum*, Linn., mentioned, 23, 10.  
*Fortunei*, S. Moore, 23, 8.  
*grandiflorum*, Maxim., 36, 524.  
*grandisepalum*, H. Léveillé & Vaniot, 36, 525.  
*hamatum*, Maxim., 36, 525.  
*ichangense*, Oliver, 36, 525.  
*Lecoyeri*, Franch., 36, 525.  
*leuconoton*, Franch., 36, 525.  
*macrorhynchum*, Franch., 36, 525.  
*majus*, Jacq., 23, 8.  
*microgynum*, Oliver, 36, 525.  
*minus*, Hook., 23, 8.  
*minus*, Linn., 23, 8.  
*mucronatum*, Ledeb., 23, 9.  
*oligandrum*, Maxim., 36, 525.  
*pallidum*, Franch., 36, 525.  
*petaloideum*, Linn., 23, 9.  
*Przewalskii*, Maxim., 23, 9.  
*punduanum*, Wall., 23, 10.  
*reticulatum*, Franch., 36, 525.  
*robustum*, Maxim., 36, 525.  
*scabrifolium*, Franch., 36, 525.  
*simplex*, Linn., 23, 9.  
*sinense*, Lour., 23, 9.  
*sp.*, 23, 10.  
*sparsiflorum*, Turcz., mentioned, 23, 10.  
*squarrosum*, Steph., 23, 9.  
*tenue*, Franch., 23, 9.  
*thibeticum*, Franch., 36, 525.  
*trichopus*, Franch., 36, 525.  
*trigynum*, Fisch., 23, 9.  
*tripeltatum*, Maxim., 36, 525.  
*tuberiferum*, H. Léveillé & Vaniot, 36, 525.  
*uncatum*, Maxim., 36, 525.  
*uncinulatum*, Franch., 36, 525.  
*virgatum*, Hook. f. & Thoms., 36, 525.
- Thamnosma*, Torr. & Frém., 23, 103.  
*Thea* = *Camellia*, 36, 525.  
*assamica*, Mast., 23, 83.  
*assimilis* Seem., 23, 80; 36, 525.  
*Bohea*, Linn., 23, 83.  
*cantoniensis*, Lour., 23, 83.  
*chinensis*, Sims, 23, 83.  
*cochinchinensis*, Lour., 23, 83.  
*cuspidata*, Kochs, 36, 525.  
*Edithæ*, Kuntze, 36, 525.  
*euryoides*, Booth, 36, 525.  
*Grijsii*, Kuntze, 36, 525.  
*hongkongensis*, Pierre, 36, 525.  
*japonica*, Kæmpf., 23, 83.  
*maliflora*, Seem., 23, 82.
- Thea punctata*, Kochs, 36, 525.  
*reticulata*, Pierre, 36, 525.  
*rosæflora*, Kuntze, 36, 525.  
*salicifolia*, Seem., 23, 82.  
*sinensis*, Linn., 23, 83.  
*speciosa*, Kochs, 36, 525.  
*spectabilis*, Kochs, 36, 525.  
*viridis*, Linn., 23, 83.
- Thela alba*, Lour., 26, 36.  
*coccinea*, Lour., 26, 36.
- Thelaia asarifolia*, Alef., 26, 33.  
*bracteosa*, Alef., 26, 33.  
*rotundifolia*, Alef., 26, 33.
- Thelasis elegans*, Blume, 36, 17.  
*elongata*, Blume, 36, 525.  
*hongkongensis*, Rolfe, 36, 39.  
*pygmæa*, Hance, 36, 39.  
*pygmæa*, Lindl., 36, 39.  
*triptera*, Reichb. f., 36, 39.
- Theligonum macranthum*, Franch., 26, 492.
- Themeda Forskalii*, Hack., 36, 377.  
*gigantea*, Hack., 36, 377.  
 — *subsp. caudata*, Hack., 36, 377.  
 — *subsp. villosa*, Hack., 36, 377.  
*triandra*, Forsk., 36, 377.  
 — *var. major*, *subvar. japonica*, Hack., 36, 378.
- Theobroma augusta*, Linn., 23, 91.
- Thermopsis alpina*, Ledeb., 36, 525.  
*chinensis*, Benth., 23, 150.  
*fabacea*, DC., 23, 150.  
*lanceolata*, R. Br., 23, 150.
- Thesium chinense*, Turcz., 26, 408.  
*longifolium*, Turcz., 36, 525.  
*psilotoides*, Hance, 26, 409.  
*rugulosum*, Bunge, 26, 408.
- Thladiantha calcarata*, C. B. Clarke, mentioned, 23, 316.  
*Davidi*, Franch., 36, 525.  
*dubia*, Bunge, 23, 316; mentioned, 317.  
*glabra*, Cogn., 36, 525.  
*Henryi*, Hemsl. \*, 23, 316.  
*longifolia*, Cogn., 36, 525.  
*maculata*, Cogn., 36, 526.  
*nudiflora*, Hemsl. \*, 23, 316, 492.  
*Oliveri*, Cogn., 36, 526.  
*sp. indescripta*?, 23, 317.  
*verrucosa*, Cogn., 36, 526.  
*villosula*, Cogn., 36, 526.
- Thlaspi arvense*, Linn., 23, 49.  
*Bursa-pastoris*, Linn., 23, 48.  
*yunnanensis*, Franch., 36, 526.
- Thoa edulis*, Willd., 26, 539.
- Thrixspermum luniferum*, Reichb., 36, 35.
- Thuarea sarmentosa*, Pers., 36, 340.
- Thunbergia fragrans*, Roxb., 26, 237.  
*grandiflora*, Roxb., 26, 237.



- Thunia Marshalliana*, Reichb. f., **36**, 18.  
*Thuya chinensis*, Hort., **26**, 540.  
*lineata*, Poir., **26**, 544.  
*orientalis*, Linn., **26**, 540.  
*pensilis*, Staunt., **26**, 544.  
*plicata*, D. Don, **26**, 540.  
*plicata*, Donn, mentioned, **26**, 540.  
*sutchuenensis*, Franch., mentioned, **26**, 540.  
**THYMELEACEÆ**, **26**, 395.  
*Thymus Serpyllum*, Linn., **26**, 282.  
*Thyrocarpus fulvescens*, Maxim., **26**, 149.  
*glochidiatus*, Maxim., **26**, 149.  
*Sampsoni*, Hance, **26**, 149.  
*Thysanolaena acarifera*, Arn., **36**, 340.  
*Agrostis*, Nees, **36**, 340.  
*Thysanospermum diffusum*, Champ., **23**, 371.  
*Thysanotus chinensis*, Benth., **36**, 119.  
*chrysanthereus*, F. Muell., **36**, 119.  
*chrysantherus*, Benth., **36**, 119.  
*Tilia argentea*, Boyer, **23**, 94.  
*Baroniana*, Diels, **36**, 526.  
*chinensis*, Maxim., **36**, 526.  
*cordata*, Mill., **36**, 526.  
*Henryana*, Szyszyl., **36**, 526.  
*mandshurica*, Rupr. & Maxim., **23**, 94.  
*Miqueliana*, Maxim., var., **36**, 526.  
*mongolica*, Maxim., **23**, 94.  
*Oliveri*, Szyszyl., **36**, 526.  
*paucicostata*, Maxim., **36**, 526.  
*pekinensis*, Rupr., **23**, 94.  
*Tuan*, Szyszyl., **36**, 526.  
**TILIACEÆ**, **23**, 92.  
*Tittmannia obovata*, Bunge, **26**, 183, 190.  
*ovata*, Reichb., **26**, 189.  
*stachydifolia*, Turcz., **26**, 183.  
*viscosa*, Reichb., **26**, 190.  
*Toddalea aculeata*, Pers., **23**, 108.  
*floribunda*, Wall., **23**, 108.  
*Tofieldia brevistyla*, Franch., **36**, 141.  
*divergens*, Bureau, **36**, 141.  
*iridacea*, Franch., **36**, 141.  
*macilenta*, Franch., **36**, 526.  
*nuda*, Maxim., **36**, 141.  
*setchuenensis*, Franch., **36**, 141.  
*yunnanensis*, Franch., **36**, 141.  
*Tomex japonica*, Thunb., **26**, 382.  
*Toona sinensis*, M. Roem., **23**, 114.  
*Torenia asiatica*, var. *concolor*, Hook. f., **26**, 187.  
*Bailloni*, Godefroy, **26**, 188.  
*Benthiana*, Hance, **26**, 187.  
*colorans*, Morr., **26**, 187.  
*concolor*, Lindl., **26**, 187.  
*Torenia cordifolia*, Hance, **26**, 189.  
*cordifolia*, Roxb., **26**, 187.  
*diffusa*, G. Don, **26**, 189.  
*edentula*, Benth., **26**, 189.  
*flava*, Buch.-Ham., **26**, 188.  
*Fordii*, Hook. f., **26**, 188.  
*lobelioides*, Hook. & Arn., **26**, 188.  
*longiflora*, Morr., **26**, 187.  
*oblonga*, Hance, **26**, 188.  
*oblonga*, Hook. & Arn., **26**, 188.  
*parviflora*, Buch.-Ham., **26**, 188.  
*peduncularia*, Benth., **26**, 188.  
*rubens*, Benth., **26**, 189.  
— var. *grandiflora*, Benth., **26**, 187.  
*vagans*, Roxb., **26**, 189.  
*Torilis Anthriscus*, Gmel., **23**, 337.  
*japonica*, DC., **23**, 337.  
*prætermissa*, Hance, **23**, 337.  
*scabra*, DC., **23**, 337.  
*Torreya Fargesii*, Franch., **26**, 546.  
*grandis*, Fortune, **26**, 546.  
*nucifera*, Sieb. & Zucc., **26**, 546.  
*Torricellia angulata*, Oliver, **36**, 526.  
*intermedia*, Harms, **36**, 526.  
*Torulinum confertum*, Buch.-Ham., **36**, 222.  
*ferox*, Kunth, **36**, 222.  
*Tournefortia argentea*, Linn., **26**, 146.  
*Arguzia*, Roem. & Schult., **26**, 147.  
— var. *angustifolia*, Franch., **26**, 147.  
— var. *latifolia*, Franch., **26**, 147.  
*ovata*, Wall., **26**, 146.  
*Sampsoni*, Hance, **26**, 146.  
*sarmentosa*, Lam., **26**, 146.  
*sibirica*, Linn., **26**, 147.  
*viridiflora*, Wall., **26**, 146.  
*Tovaria atropurpurea*, Franch., **36**, 110.  
*Delavayi*, Franch., **36**, 110.  
*Fargesii*, Franch., **36**, 110.  
*japonica*, Baker, **36**, 110.  
*Prattii*, Franch., **36**, 111.  
— var. *quadrifolia*, Franch., **36**, 111.  
— var. *robusta*, Franch., **36**, 111.  
*Rossii*, Baker, **36**, 111.  
*Souliei*, Franch., **36**, 111.  
*stenoloba*, Franch., **36**, 111.  
*tatsienensis*, Franch., **36**, 111.  
*tubifera*, C. H. Wright\*, **36**, 111.  
*yunnanensis*, Franch., **36**, 112.  
*Toxicarpus Wightianus*, Hook. & Arn., **26**, 101.  
*Trachelospermum axillare*, Hook. f., **36**, 526.

- Trachelospermum Bowringii*, *Hemsl.*, 26, 99.  
*divaricatum*, *Thunb.*, 36, 526.  
*jasminoides*, *Lem.*, 26, 99.  
*Trachycarpus excelsus*, *H. Wendl.*, 36, 168.  
*Fortunei*, *H. Wendl.*, 36, 168; mentioned, 167.  
*khasianus*, *H. Wendl.*, 36, 169.  
*Trachydium daucoides*, *Franch.*, 36, 526.  
*Delavayi*, *Franch.*, 36, 526.  
*hispidum*, *Franch.*, 36, 526.  
*novemjugum*, *C. B. Clarke*, 36, 526.  
*purpurascens*, *Franch.*, 36, 526.  
*Roylei*, *Lindl.*, 36, 526.  
*rubinerve*, *Franch.*, 36, 526.  
*viridiflorum*, *Franch.*, 36, 526.  
*Trachytella Actæa*, *DC.*, 23, 22.  
*aspera*, *DC.*, 23, 22.  
*Tradescantia axillaris*, *Roxb.*, 36, 157.  
*vaga*, *Blume*, mentioned, 36, 161.  
*vaga*, *Lour.* (sp. dub.), 36, 160.  
*Tragia cannabina*, *Linn. f.*, 26, 444.  
*Chamælea*, *Linn.*, 26, 446.  
*involuta*, *Linn.*, 26, 444.  
*Tragus racemosus*, *Scop.*, 36, 343.  
*tcheliensis*, *Debeaux*, 36, 343.  
*Trapa bicornis*, *Linn.*, 23, 311.  
*bispinosa*, *Roxb.*, 23, 311.  
*chinensis*, *Lour.*, 23, 311.  
*natans*, *Linn.*, 23, 311.  
*Trapella sinensis*, *Oliver*, 26, 236.  
*Trema amboinensis*, *Blume*, 26, 451.  
*orientalis*, *Blume*, 26, 451.  
*timorensis*, *Blume*, 26, 452.  
*virgata*, *Blume*, 26, 452.  
*Trentepohlia bifoliata*, *Boeck.*, 36, 209.  
*Tretocarya pratensis*, *Maxim.*, 26, 151.  
*sikkimensis*, *Oliver*, 36, 526.  
*Trewia nudiflora*, *Hance*, 26, 441.  
*Triadica sinensis*, *Lour.*, 26, 445.  
*Trianthema polyandra*, *Blume*, 23, 323, 324.  
*Tribulus cistoides*, *Linn.*, 23, 97.  
*terrestris*, *Linn.*, 23, 97.  
*Tricercandra Fortunei*, *A. Gray*, 26, 367.  
*quadrifolia*, *A. Gray*, 26, 368.  
*Trichelostylis complanata*, *Nees*, 36, 231, 246.  
*globulosa*, *Nees*, 36, 237.  
*miliacea*, *Nees*, 36, 239.  
*spp. nn.*, 36, 240, 246.  
*Trichosanthes Anguina*, *Linn.*, 23, 312.  
*chinensis*, *Ser.*, 23, 313.  
*Trichosanthes cucumerina*, *Linn.*, mentioned, 23, 313.  
*cucumeroides*, *Maxim.*, 23, 313.  
*dicælosperma*, *C. B. Clarke*, mentioned, 23, 314.  
*Kirilowii*, *Maxim.*, 23, 313; mentioned, 314.  
*multiloba*, *Miq.*, 23, 314.  
*palmata*, *C. B. Clarke*, 23, 313.  
*palmata*, *Hance*, 23, 313.  
*Rosthornii*, *Harms*, 36, 526.  
*sp.*, 23, 314.  
*Tricyrtis elegans*, *Wall.*, 36, 142.  
*formosana*, *Baker*, 36, 141.  
*lasiocarpa*, *Matsum.*, 36, 526.  
*latifolia*, *Maxim.*, 36, 141.  
*macropoda*, *Miq.*, 36, 142.  
*pilosa*, *Wall.*, 36, 142.  
*stolonifera*, *Matsum.*, 36, 526.  
*Tridesmis hispida*, *Lour.*, 26, 435.  
*tomentosa*, *Lour.*, 26, 434.  
*Trientalis*, *Rupp.*, 26, 55.  
*europæa*, *Linn.*, 26, 58.  
*Trifolium globosum*, *Lour.*, 23, 155.  
*indicum*, *Lour.*, 23, 155.  
*Lupinaster*, *Linn.*, 23, 155.  
*Triglochin maritimum*, *Linn.*, 36, 192.  
*palustre*, *Linn.*, 36, 192.  
*Trigonella cærulea*, *Ser.*, 23, 154.  
*ruthenica*, *Linn.*, 23, 153; correction, 489.  
*Trigonotis brevipes*, *Maxim.*, 26, 152.  
*mollis*, *Hemsl.* \*, 26, 153.  
*peduncularis*, *Benth.*, 26, 153.  
— *var. vestita*, *Hemsl.* \*, 26, 154.  
*petiolaris*, *Maxim.*, 26, 154; mentioned, 152.  
*Trilepis Royleana*, *Boeck.*, 36, 268.  
*Trillium erectum*, *Linn.*, mentioned, 36, 144.  
*Tschonoskii*, *Maxim.*, 36, 144.  
*Triosteum Fargesii*, *Franch.*, 36, 526.  
*himalayanum*, *Wall.*, 36, 526.  
*hirsutum*, *Wall.*, 23, 357.  
*intermedium*, *Diels & Graebn.*, 36, 526.  
*pinnatifidum*, *Maxim.*, 23, 357.  
*Rosthornii*, *Diels & Graebn.*, 36, 526.  
*sinuatum*, *Maxim.*, 23, 357.  
*Triphasia Aurantiola*, *Lour.*, 23, 109.  
*trifoliata*, *DC.*, 23, 109.  
*Triplostegia glandulifera*, *Wall.*, 23, 399.  
*glandulosa*, *DC.*, 23, 399.  
*grandiflora*, *Gagnep.*, 36, 526.  
*Tripogon chinensis*, *Hack.*, 36, 404.  
— *var. coreensis*, *Hack.*, 36, 405.

- Tripolium vulgare*, Nees, 23, 417.  
*Tripterospermum trinerve*, Blume, 26, 123.  
*Tripterygium Bullockii*, Hance, 23, 125.  
     *Wilfordii*, Hook., 23, 125; in Japan, 489.  
*Trisanthus cochinchinensis*, Lour., 23, 325.  
*Trisetum flavescens*, Beauv., 36, 399.  
     — *var. macranthum*, Hack., 36, 400.  
     *Henryi*, Rendle\*, 36, 400.  
     *subspicatum*, Beauv., 36, 400.  
*Triticum caninum*, Schreb., 36, 431.  
     *chinense*, Trin., 36, 432.  
     *ciliare*, Trin., 36, 432.  
     *Pseudo-agropyrum*, Griseb., 36, 432.  
     *repens*, Linn., 36, 432.  
     *strigosum*, Less., 36, 433.  
*Triumfetta angulata*, Lam., 23, 93.  
     *annua*, Linn., 36, 526.  
     *pilosa*, Roth, 23, 93.  
     *rhomboides*, Jacq., 23, 93.  
     *tomentosa*, Boj., 36, 527.  
*Trochostigma arguta*, Sieb. & Zucc., 23, 78.  
*Trollius asiaticus*, Linn., 23, 17.  
     — Linn. *var. affinis*, Regel, 23, 17.  
     *chinensis*, Bunge, 23, 17.  
     *Ledebourii*, Reichb., 36, 527.  
     *palmatum*, Hook. f. & Thoms., 36, 527.  
     *patulus*, Salisb., 36, 527.  
     *pumilis*, D. Don, 36, 527.  
     *ranunculoides*, Hemsl., 36, 527.  
*Trophis scandens*, Hook. f. & Arn., 26, 454.  
*Tropidia angulosa*, Lindl., 36, 39.  
     *curculigoides*, Benth., 36, 40.  
     *curculigoides*, Lindl., mentioned, 36, 40.  
     *formosana*, Rolfe, 36, 40.  
     *Govindooi*, Blume, 36, 39.  
     *grandis*, Hance, 36, 32; mentioned, 40.  
     *hongkongensis*, Rolfe, 36, 40.  
     *semilibera*, Blume, 36, 40.  
     *septemnervis*, Reichb. f., mentioned, 36, 40.  
*Tsuga Brunoniana*, Carr., 26, 556.  
     — *var. chinensis*, Franch., 26, 556.  
     *chinensis*, Mast. \*, 26, 556.  
     *Sieboldii*, Carr., 26, 556.  
     *yunnanensis*, Mast. \*, 26, 556.  
*Tulipa edulis*, Baker, 36, 138.  
     *erythronioides*, Baker, 36, 138.  
*Tulipa graminifolia*, Baker, 36, 138.  
     *ornithogaloides*, Fisch., 36, 139.  
*Tunga diandra*, Roxb., 36, 258.  
*Tunica Morrisii*, Walp., 23, 63.  
*Tupidanthus calyptratus*, Hook. f. & Thoms., 36, 527.  
*Tupistra chinensis*, Baker, 36, 114.  
     *chlorantha*, Baill., 36, 114.  
     *Delavayi*, Franch., 36, 114.  
     *Fargesii*, Baill., 36, 114.  
     *lorifolia*, Franch., 36, 114.  
     *viridiflora*, Franch., 36, 114.  
     *Wattii*, Hook. f., 36, 114.  
*Turczaninowia fastigiata*, DC., 23, 410.  
*Turpinia arguta*, Seem., 23, 143.  
     *nepalensis*, Wall., 23, 143.  
     *pomifera*, DC., 23, 143.  
*Turraea pubescens*, Hellen., 23, 113.  
*Turritis glabra*, Linn., 23, 42.  
     *hirsuta*, Linn., 23, 42.  
*Tussilago Anandria*, Linn., 23, 472.  
     *Bellidiastrum*, Linn., 23, 472.  
     *Farfara*, Linn., 23, 447.  
     *japonica*, Linn., 23, 454.  
     *lyrata*, Willd., 23, 473.  
     *Petasites*, Thunb., 23, 446.  
*Tylophora asthmatica*, Wight & Arn., *var.*, 26, 113.  
     *exilis*, Colebr., mentioned, 26, 111.  
     *hispida*, Decne., 26, 113.  
     *japonica*, Miq., mentioned, 26, 113.  
     *liukiensis*, Matsum., 36, 527.  
     *macrantha*, Hance, 26, 115.  
     *spp.*, 26, 113.  
*Typha angustata*, Bory, 36, 171.  
     *angustifolia*, Linn., 36, 172.  
     *Bungeana*, Presl, 36, 173.  
     *japonica*, Miq., 36, 173.  
     *latifolia*, Bretschn., 36, 173.  
     *latifolia*, Linn., 36, 172.  
     *Laxmanni*, Franch., 36, 172.  
     *Laxmanni*, Lepech., 36, 172.  
     *Laxmanni*, Ledeb., 36, 173.  
     *Martini*, H. Léveillé & Vaniot, 36, 527.  
     *Martini*, Jord., *var. Davidiana*, Kronfeld, 36, 172.  
     *minima*, Funck, 36, 172.  
     *minor*, Sm., 36, 173.  
     *orientalis*, Presl, 36, 173.  
     *Shuttleworthii*, Koch, *var. orientalis*, Rohrb., 36, 173.  
 TYPHACEÆ, 36, 171.  
*Typhonium divaricatum*, Decne., 36, 180; mentioned, 181.  
     *giganteum*, Engl., 36, 181.  
     — *var. Giraldii*, Baroni, 36, 181.  
     *javanicum*, Miq., 36, 181.  
     *yan m.*, Schott, mentioned, 36, 181.

- Typhonium Roxburghii*, Schott, mentioned, **36**, 181.
- Ulmaria palmata*, Focke, **36**, 527.
- Ulmus campestris*, Linn., **26**, 446.
- castaneifolia*, Hemsl. \*, **26**, 446.
- chinensis*, Loud., **26**, 448.
- chinensis*, Pers., **26**, 448.
- Davidiana*, Planch., **26**, 447; mentioned, 448.
- Keaki*, Sieb., **26**, 449.
- lancifolia*, Roxb., **26**, 447.
- macrocarpa*, Hance, **26**, 447.
- montana*, With., **26**, 448.
- parvifolia*, Jacq., **26**, 448.
- pumila*, Linn., **36**, 527.
- sp. nov. altera*, Maxim., **26**, 451.
- virgata*, Roxb., **26**, 448.
- UMBELLIFERE**, **23**, 324.
- Umbilicus fimbriatus*, Turcz., **23**, 281.
- leucanthus*, Ledeb., **23**, 281.
- malacophyllus*, DC., **23**, 282.
- ramosissimus*, Maxim., **23**, 281.
- spinusus*, DC., **23**, 282.
- thyrsiflorus*, DC., **23**, 281.
- Uncinia microglochis*, Spreng., **36**, 298.
- Unona chinensis*, DC., **23**, 27.
- discolor*, Vahl, **23**, 26.
- Lessertiana*, Dunal, **23**, 27.
- uncinata*, DC., **23**, 26.
- Uralepis fusca*, Steud., **36**, 411.
- Uraria comosa*, DC., **23**, 177.
- erinita*, Desv., **23**, 177.
- var. *macrostachya*, Wall., **23**, 177.
- hamosa*, Wall., **23**, 177.
- var. *sinensis*, Hemsl. \*, **23**, 177.
- lagopoides*, DC., **23**, 178.
- Lagopus*, DC., **36**, 527.
- picta*, Desv., **23**, 178; **36**, 527.
- sinensis*, Franch., **36**, 527.
- Urena chinensis*, Osbeck, **23**, 87.
- diversifolia*, Walp., **23**, 86.
- lobata*, Linn., **23**, 86.
- polyflora*, Lour., **23**, 86.
- procumbens*, Linn., **23**, 86.
- sinica xanthii facie*, Dill., **23**, 86.
- sinuata*, Linn., **23**, 87.
- Urostelma chinense*, Bunge, **26**, 110.
- Urostigma Benjaminia*, Miq., **26**, 458.
- insectorium*, Miq., **26**, 463.
- nervosum*, Miq., **26**, 464.
- nitidum*, Miq., **26**, 467.
- parvifolium*, Miq., **26**, 465.
- pisiferum*, Miq., **26**, 467.
- retusum*, Miq., **26**, 467.
- superbum*, Miq., **26**, 469.
- Urtica alienata*, Linn., **26**, 490.
- Urtica angulata*, Blume, **26**, 478.
- angustifolia*, Fisch., **26**, 472.
- angustifolia*, Buch.-Ham., **26**, 485.
- ardens*, Link, **26**, 472.
- cannabina*, Linn., **26**, 471.
- caudata*, Burm. f., **26**, 487.
- cordatifolia*, Steud., **26**, 471.
- dioica*, Thunb., **26**, 472.
- subvar. *hispida*, Wedd., **26**, 472.
- dioica*, var. *angustifolia*, Ledeb., **26**, 472.
- fissa*, E. Pritz., **36**, 527.
- “*foliis oppositis*,” Linn., **26**, 471.
- frutescens*, Thunb., **26**, 491.
- Hamiltoniana*, Wall., **26**, 478.
- heterophylla*, Vahl, **26**, 478.
- himalayensis*, Kunth & Bouché, **26**, 472.
- hirta*, Blume, **26**, 489.
- interrupta*, Linn., **26**, 473.
- laetevirens*, Maxim., **26**, 472.
- macrophylla*, Thunb., **26**, 485.
- macrostachya*, Wall., **26**, 485.
- microphylla*, Sw., **26**, 476.
- palmata*, Forsk., **26**, 475.
- parviflora*, Roxb., **26**, 472.
- penduliflora*, Wedd., **26**, 485.
- pilosiuscula*, Blume, **26**, 487.
- polystachya*, Wall., **26**, 488.
- pumila*, Linn., **26**, 478.
- scabra*, Blume, **26**, 495.
- scabrella*, Roxb., **26**, 489.
- scripta*, Buch.-Ham., **26**, 478.
- sp.*, **26**, 473.
- sp. nov.*, Maxim., **26**, 475.
- spicata*, Thunb., **26**, 488.
- squamigera*, Wall., **26**, 489.
- stipulosa*, Miq., **26**, 478.
- tenacissima*, Roxb., **26**, 486.
- Thunbergiana*, Sieb. & Zucc., **26**, 472.
- umbrosa*, Wall., **26**, 479.
- uragera*, Steud., **26**, 487.
- urens*, Linn., **26**, 472; mentioned, 473.
- utilis*, Hort., **26**, 486.
- viminea*, Wall., **26**, 491.
- virentula*, Wall., **26**, 472.
- URTICACEE**, **26**, 446.
- Utricularia bifida*, Linn., **26**, 222.
- cærulea*, Linn., **26**, 223.
- diantha*, A. DC., **26**, 222.
- diantha*, Roem. & Schult., **26**, 223.
- exoleta*, R. Br., **26**, 223.
- extensa*, Hance, **26**, 223.
- fasciculata*, Roxb., **26**, 223.
- flexuosa*, Vahl, **26**, 223.
- Harlandi*, Oliver, **26**, 223.

- Utricularia intermedia*, Hayne, 26, 223.  
*orbiculata*, Wall., 26, 224.  
*racemosa*, Wall., 26, 224.  
*uliginosa*, Benth., 26, 223.
- Uvaria badiiflora*, Hance, 23, 26.  
*calamistrata*, Hance, 23, 25.  
*microcarpa*, Champ., 23, 25.  
*odorata*, Lam., 23, 27.  
*platypetala*, Champ., 23, 26.  
*purpurea*, Blume, 23, 26.  
*rhodantha*, Hance, 23, 26.  
*suberosa*, Roxb., 23, 27.  
*synsepala*, Miq., 23, 26.  
*uncata*, Lour., 23, 26.
- Uvularia amplexifolia*, Linn., 36, 110.  
*chinensis*, Ker-Gawl., 36, 143; mentioned, 136.  
*cirrhusa*, Thunb., 36, 137.  
*sessilis*, Thunb., 36, 143.  
*viridescens*, Maxim., 36, 144.  
*Wallichii*, D. Don, 36, 142.
- VACCINIACEÆ**, 26, 14.
- Vaccinium bracteatum*, Thunb., 26, 14; mentioned, 15.  
*chinense*, Champ., 26, 14.  
*ciliatum*, Thunb., 26, 15.  
*Delavayi*, Franch., 36, 527.  
*Donianum*, Franch., 36, 527.  
*Donianum*, Miq., 26, 14.  
*Donianum*, Wight, mentioned, 26, 15.  
*Dunalianum*, Wight, mentioned, 26, 16.  
*fragile*, Franch., 36, 527.  
*Griffithianum*, Wight, mentioned, 26, 15.  
*Henryi*, Hemsl. \*, 26, 15.  
*hirtum*, Thunb., 36, 527.  
*iteophyllum*, Hance, 26, 15.  
*japonicum*, Miq., 26, 16.  
*lætum*, Diels, 36, 527.  
*Leschenaultii*, Wight, mentioned, 26, 14.  
*malaccense*, Wight, 26, 14.  
*Mandarinorum*, Diels, 36, 527.  
*moupinense*, Franch., 36, 527.  
*neilgherrense*, Wight, mentioned, 26, 14.  
*Oldhami*, Miq., 26, 14, 15.  
*pubicalyx*, Franch., 36, 527.  
*rotundifolium*, Wight, mentioned, 26, 14.  
*setosum*, C. H. Wright, 36, 527.  
*urceolatum*, Hemsl. \*, 26, 16.  
*Wrightii*, A. Gray, 26, 14.  
*yunnanense*, Franch., 36, 527.
- Valeriana chinensis*, Linn., 26, 317.  
*dubia*, Bunge, 23, 399.
- Valeriana Faberi*, Graebn., 36, 527.  
*flaccidissima*, Maxim., 36, 527.  
*flagellifera*, Batalin, 36, 527.  
*Hardwickii*, Wall., mentioned, 23, 399.  
*Harmsii*, Graebn., 36, 527.  
*helictes*, Graebn., 36, 527.  
*hiemalis*, Graebn., 36, 527.  
*officinalis*, Linn., 23, 399.  
*Rosthornii*, Graebn., 36, 527.  
*tangutica*, Batalin, 36, 527.  
*villosa*, Thunb., 23, 398.  
*Wallichii*, DC., 23, 399.
- VALERIANACEÆ**, 23, 396.
- Valerianella Olitoria*, Moench, mentioned, 23, 399.
- Vallisneria laxiflora*, Blume, 26, 97.  
*sinensis*, G. Don, 26, 101.
- Vallisneria alternifolia*, Roxb., 36, 2.  
*octandra*, Roxb., 36, 2.  
*spiralis*, Linn., 36, 2.
- Valoradia plumbaginoides*, Bunge, 26, 36.
- Vanda bicaudata*, Thw., 36, 34.  
*concolor*, Blume, 36, 35.  
*densiflora*, Lindl., mentioned, 36, 36.  
*furva*, Lindl., mentioned, 36, 36.  
*furva*, Lindl., partim, 36, 35.  
*hainanensis*, Rolfe, 36, 36.  
*parviflora*, Lindl., 36, 528.  
*recurva*, Hook., 36, 37.  
*rostrata*, Lodd., 36, 37.  
*Roxburghii*, R. Br., mentioned, 36, 36.  
 — var. *unicolor*, Hook., 36, 35.  
*teretifolia*, Lindl., 36, 38.  
*violacea*, Lindl., mentioned, 36, 36.
- Vandellia angustifolia*, Benth., 26, 189.  
*arridens*, Hance, 26, 191.  
*crustacea*, Benth., 26, 189; mentioned, 190.  
*erecta*, Benth., 26, 190.  
*hirsuta*, Buch.-Ham., 26, 190.  
*limosa*, Walp., 26, 190.  
*mollis*, Benth., 26, 190.  
*nummularifolia*, G. Don, 26, 190.  
*oblonga*, Benth., 26, 188.  
*obovata*, Bunge, 26, 183.  
*obovata*, Walp., 26, 190.  
*pedunculata*, Benth., 26, 191.  
*Pyxidaria*, Maxim., 26, 190.  
*scabra*, Benth., 26, 191.  
*stachydifolia*, Walp., 26, 184.  
*subcrenulata*, Miq., 26, 191.  
*urticifolia*, Hance, 26, 191.
- Vanieria chinensis*, Lour., 26, 470.
- Vanilla* sp., 36, 39.

- Vaniotia*, *H. Léveillé*, **36**, 528.  
*Martini*, *H. Léveillé*, **36**, 528.  
*Varronia sinensis*, *Lour.*, **26**, 143.  
*Vatica chinensis*, *Linn.*, **23**, 83.  
*Roxburghiana*, *Blume*, **23**, 83.  
*Ventilago elegans*, *Hemsl.*, **36**, 528.  
*leiocarpa*, *Benth.*, **23**, 125.  
*maderaspatana*, *Benth.*, **23**, 125.  
*Veratrina* *Bailloni*, *Franch.*, **36**,  
528.  
*Veratrum album*, *Linn.*, **36**, 147.  
— *var. viride*, *Baker*, **36**, 147.  
*bracteatum*, *Batalin*, **36**, 147.  
*Maackii*, *Regel*, **36**, 147.  
*Maximowiczii*, *Baker*, **36**, 147.  
*nigrum*, *Linn.*, **36**, 147.  
— *var. japonicum*, *Baker*, **36**,  
148.  
*parviflorum*, *Franch.*, **36**, 147.  
*viride*, *Ait.*, **36**, 147.  
*Verbascum Thapsus*, *Linn.*, **36**, 528.  
*Verbena officinalis*, *Linn.*, **26**, 252.  
*sp. nov.?*, *Hemsl.*, **26**, 252.  
VERBENACEÆ, **26**, 251.  
*Verbesina calendulacea*, *Linn.*, **23**,  
434.  
*chinensis*, *Linn.*, **23**, 432.  
*prostrata*, *Hook. & Arn.*, **23**, 434.  
*spicata*, *Lour.*, **23**, 434.  
*Vernicia montana*, *Lour.*, **26**, 433.  
*Vernonia ampla*, *Vaniot*, **36**, 528.  
*Andersoni*, *C. B. Clarke*, **23**, 400.  
*Bockiana*, *Diels*, **36**, 528.  
*bracteata*, *Wall.*, **23**, 400.  
*chinensis*, *Less.*, **23**, 401.  
*cinerea*, *Less.*, **23**, 401.  
*Clivorum*, *Hance*, **23**, 401.  
*congesta*, *Benth.*, **23**, 429.  
*conyzoides*, *Walp.*, **23**, 420.  
*Cumingiana*, *Benth.*, **23**, 400.  
*eriosematoides*, *Walp.*, **23**, 429.  
*esculenta*, *Hemsl. \**, **23**, 401.  
*exilis*, *Miq.*, **23**, 402.  
*Fargesii*, *Franch.*, **36**, 528.  
*flaccidissima*, *Maxim.*, **36**, 528.  
*Fortunei*, *Sch.-Bip.*, **23**, 420.  
*Gomphrena*, *Walp.*, **23**, 402.  
*gratiosa*, *Hance*, **23**, 402.  
*Hardwickii*, *Wall.*, **36**, 528.  
*Henryi*, *Dunn*, **36**, 528.  
*Kingii*, *C. B. Clarke*, **23**, 402.  
*Kroneana*, *Miq.*, **23**, 402.  
*Martini*, *Vaniot*, **36**, 528.  
*monosis*, *Franch.*, **36**, 528.  
*papillosa*, *Franch.*, **36**, 528.  
*saligna*, *DC.*, **23**, 402.  
*Seguini*, *Vaniot*, **36**, 528.  
*solanifolia*, *Benth.*, **23**, 402.  
*spelæicola*, *Vaniot*, **36**, 528.  
*subarborea*, *Vaniot*, **36**, 528.  
*Vernonia sylvatica*, *Dunn*, **36**, 528.  
*Veronica agrestis*, *Linn.*, **26**, 197.  
*alpina*, *Linn.*, **26**, 198.  
*Anagallis*, *Linn.*, **26**, 198.  
*angustifolia*, *Fisch.*, **26**, 200.  
*arvensis*, *Thunb.*, **26**, 197.  
*cana*, *Wall.*, **26**, 198.  
*Fargesii*, *Franch.*, **36**, 528.  
*galactites*, *Hance*, **26**, 200.  
*hederacea* [*Hemsl. lapsu = seq.*]  
*hederæfolia*, *Linn.*, mentioned, **26**,  
291.  
*hederæfolia*, *Miq.*, **26**, 197.  
*javanica*, *Blume*, **26**, 198.  
*laxa*, *Benth.*, **26**, 198.  
*longifolia*, *Linn.*, **26**, 199.  
— *var. subsessilis*, *Hook. f.*, **26**  
199.  
*Maddenii*, *Edgew.*, **26**, 198.  
*Murorum*, *Maxim.*, **26**, 199.  
*paniculata*, *Linn.*, **26**, 200.  
*peregrina*, *Linn.*, **26**, 199.  
*pirolæformis*, *Franch.*, **36**, 528.  
*serpyllifolia*, *Linn.*, **26**, 199.  
*sibirica*, *Linn.*, **26**, 200.  
*spicata*, *Linn.*, **26**, 199.  
*spuria*, *Linn.*, **26**, 200.  
*sutchuenensis*, *Franch.*, **26**, 528.  
*szechuanica*, *Batalin*, **36**, 528.  
*Thunbergii*, *A. Gray*, **26**, 199.  
*vandellioides*, *Maxim.*, **26**, 200.  
*virginica*, *Linn.*, **26**, 200.  
*Viburnum arborescens*, *Hemsl. \**, **23**,  
349.  
*arcuatum*, *Komarov*, **36**, 528.  
*Awabuki*, *C. Koch*, **23**, 354.  
*betulifolium*, *Batalin*, **36**, 528.  
*Bockii*, *Graebn.*, **36**, 528.  
*brachybotryum*, *Hemsl. \**, **23**,  
349.  
*buddleifolia*, *C. H. Wright*, **36**,  
528.  
*burejaticum*, *Regel & Herd.*, **23**,  
350.  
*burejanum*, *Herd.*, **23**, 350.  
*Carlesii*, *Hemsl. \**, **23**, 350.  
*ceanthoides*, *C. H. Wright*, **36**,  
528.  
*chinense*, *Hook. & Arn.*, **23**, 351;  
**26**, 256.  
*chinense*, *Zeyh.*, **23**, 354.  
*chinshanense*, *Graebn.*, **36**, 528.  
*cordifolium*, *Hook. f. & Thoms.*, **23**,  
352.  
*coriaceum*, *Blume*, **23**, 351.  
*cotinifolium*, *D. Don*, **23**, 356.  
*cotinifolium*, *Hance*, **23**, 356.  
*cuspidatum*, *Sieb. & Zucc.*, **23**  
356.  
*cylindricum*, *Buch.-Ham.*, **26** 351

- Viburnum Davidi*, Franch., 36, 528.  
*davuricum*, Maxim., 23, 350.  
*davuricum*, Pall., 23, 351.  
*dentatum*, Thunb., 23, 356.  
*Dielsii*, Graebn., 36, 528.  
*dilatatum*, Maxim., 23, 351.  
*dilatatum*, Thunb., 23, 351.  
*erosum*, A. Gray, 23, 351, 354.  
*erosum*, Thunb., 23, 351.  
— var. *ichangense*, Hemsl. \*, 23, 352.  
*erubescens*, Wall., 36, 528.  
*fallax*, Graebn., 36, 528.  
*Fordiæ*, Hance, 23, 352.  
*fragrans*, Bunge, 23, 352.  
*furcatum*, Blume, 23, 352.  
*glomeratum*, Maxim., 23, 352.  
*Hanceanum*, Maxim., 23, 353; mentioned, 349.  
*Henryi*, Hemsl. \*, 23, 353.  
*kansuense*, Batalin, 36, 528.  
*Keteleeri*, Carr., 23, 353.  
*lantanoïdes*, Michx., 23, 352.  
*lantanoïdes*, Miq., 23, 352.  
*lobophyllum*, Graebn., 36, 528.  
*lutescens*, Blume, 23, 353.  
*macrocephalum*, Fortune, 23, 353.  
*nervosum*, D. Don, mentioned, 23, 352.  
*nervosum*, Hook. & Arn., 23, 356.  
*odoratissimum*, Ker-Gawl., 23, 354; mentioned, 350.  
*eliganthum*, Batalin, 36, 528.  
*Opulus*, Linn., 23, 354.  
*pallidum*, Franch., 36, 528.  
*phlebotrichum*, Sieb. & Zucc., 23, 354.  
*plicatum*, A. Gray, 23, 352.  
*plicatum*, Thunb., 23, 356.  
*Prattii*, Graebn., 36, 528.  
*propinquum*, Hemsl. \*, 23, 355.  
*pubinerve*, Blume, 23, 354.  
*rectangulatum*, Graebn., 36, 528.  
*rhytidophyllum*, Hemsl. \*, 23, 355.  
*Rosthornii*, Graebn., 36, 528.  
*Sandankwa*, Hassk., 23, 355.  
*Sargenti*, Kochne, 36, 529.  
*schensianum*, Maxim., 23, 355.  
*sempervirens*, C. Koch, 23, 355.  
*setigerum*, Hance, 23, 356.  
*stuenense*, Miq., 23, 353.  
*stellulatum*, Franch., 36, 529.  
*sundaicum*, Miq., 23, 353.  
*sympodiale*, Graebn., 36, 529.  
*Timus*, Linn., mentioned, 23, 355.  
*tomentosum*, Hance, 23, 353.  
*tomentosum*, Thunb., 23, 356.  
*utile*, Hemsl. \*, 23, 356.  
*Veitchi*, C. H. Wright, 36, 529.
- Viburnum venulosum*, Benth., 23, 356.  
*Willeanum*, Graebn., 36, 529.  
*Wrightii*, Miq., mentioned, 23, 354.
- Vicia amœna*, Fisch., 23, 184.  
*angustifolia*, Roth, 23, 184.  
*chinensis*, Franch., 36, 529.  
*Cracca*, Linn., 23, 184.  
*gigantea*, Bunge, 23, 184.  
*hirsuta*, Koch, 23, 184.  
*megalotropis*, Ledeb., 36, 529.  
*Pseudo-Orobus*, Fisch. & Mey., 23, 185.  
*ramosissima*, Franch., 36, 529.  
*Rapunculus*, Debeaux, 23, 184.  
*sativa*, Linn., 23, 185.  
*tenera*, R. Grab., 36, 529.  
*tetrasperma*, Moench, 23, 185.  
*tridentata*, Bunge, 23, 185.  
*unijuga*, A. Br., 23, 186.  
*venosa*, Maxim., 23, 184.
- Vigna Catiang*, Miq., 23, 193.  
*lutea*, A. Gray, 23, 193.  
*sinensis*, Hassk., 23, 193.  
*vexillata*, Benth., 23, 193.
- Vilfa diandra*, Steud., 36, 387.  
*elongata*, Beaux., 36, 388.  
*indica*, Steud., 36, 388.
- Villarsia nymphoides*, Vent., 26, 142.
- Villebrunea frutescens*, Blume, 26, 491.  
*pedunculata*, Shirai, 36, 529.  
*scabra*, Wedd., 26, 491.
- Vinca rosea*, Linn., 26, 95.
- Vincetoxicum acuminatum*, Decne., 26, 104.  
*amplexicaule*, Sieb. & Zucc., 26, 104.  
*atratum*, Morr. & Decne., 26, 104.  
*chinense*, S. Moore, 26, 108.  
*japonicum*, Morr. & Decne., 26, 107.  
*macrophyllum*, Sieb. & Zucc., 26, 106.  
*mandshuricum*, Hance, 26, 107.  
*mongolicum*, Maxim., 26, 107.  
— var. *Hancockianum*, Hance, 26, 107.  
*sibiricum*, Decne., 26, 108.  
*sublanceolatum*, Maxim., 26, 106.  
*versicolor*, Decne., 26, 109.
- Viola acuminata*, Ledeb., 23, 52.  
*albida*, Palibin, 36, 529.  
*Averyi*, Kellogg, 23, 53.  
*biflora*, Baker & S. Moore, 23, 56.  
*biflora*, Linn., 23, 52.  
*Bisseti*, Maxim., mentioned, 23, 54.  
*bulbosa*, Maxim., 23, 52.  
*Camaleo*, Boissieu, 36, 529.

- Viola canina*, Linn., 23, 52.  
*chinensis*, G. Don, 23, 53.  
*confusa*, Champ., 23, 55.  
*dactyloides*, Roem. & Schult., 36, 529.  
*deltoidea*, Makino, 36, 529.  
*Davidi*, Franch., 36, 529.  
*Delavayi*, Franch., 36, 529.  
*dentariæfolia*, Boissieu, 36, 529.  
*diffusa*, Ging., 23, 52.  
*dissecta*, Ledeb., 23, 54.  
*distans*, Wall., 36, 529.  
*excisa*, Hance, 23, 56.  
*Fargesii*, Boissieu, 36, 529.  
*flavida*, Bur. & Franch., 36, 529.  
*glabrella*, Nutt., mentioned, 23, 56.  
*Gmeliniana*, Roem. & Schult., 36, 529.  
*Grayi*, Franch., 23, 55.  
*grypoceras*, A. Gray, 23, 55.  
*Henryi*, Boissieu, 36, 529.  
*hirta*, Linn., 23, 52.  
*hirtipes*, S. Moore, 23, 53.  
*Hookeri*, T. Thoms., 23, 53.  
*inconspicua*, Blume, mentioned, 23, 55.  
*japonica*, Langsd., 23, 53.  
— var. *pekinensis*, Maxim., 23, 53.  
*kamtschatica*, Ging., 23, 55.  
— var., *pekinensis*, Regel, 23, 53.  
*Leveillei*, Boissieu, 36, 529.  
*micrantha*, Turcz., 23, 52.  
*mongolica*, Franch., 23, 52.  
*moupinensis*, Franch., 36, 529.  
*odorata*, Linn., 23, 53.  
*palustris*, Franch., 36, 529.  
*Parkeri*, Champ., 23, 52.  
*Patrinii*, auct., 23, 53.  
*Patrinii*, DC., 23, 53.  
*phalacrocarpa*, Maxim., 36, 529.  
*pinnata*, Linn., 23, 54.  
— var. *chærophylloides*, Regel, 23, 54.  
*primulifolia*, Lour., 23, 53.  
*prionantha*, Bunge, 23, 53.  
*Raddeana*, Regel, 23, 54.  
*Rossii*, Hemsl. \*, 23, 54.  
*Rosthornii*, E. Pritz., 36, 529.  
*Selkirkii*, Pursh, 23, 55.  
*serpens*, Wall., 23, 55.  
*striatella*, Boissieu, 36, 529.  
*sylvatica*, var. *imberbis*, A. Gray, 23, 55.  
*sylvestris*, Kit., 23, 55.  
*tenuis*, Benth., 23, 52.  
*tricolor*, Linn., mentioned, 23, 55.  
*tuberifera*, Franch., 36, 529.  
*umbrosa*, Fries, 23, 55.  
*uniflora*, Linn., 23, 56.
- Viola urophylla*, Franch., 36, 529.  
*vaginata*, Maxim., 36, 529.  
*variegata*, Fisch., 23, 56.  
*verecunda*, A. Gray, 23, 56.  
*Websteri*, Hemsl. \*, 23, 56.
- VIOLARIÆ**, 23, 52.
- Viscum album*, Linn., 26, 407.  
*articulatum*, Benth., 26, 408.  
*articulatum*, Burm. f., 26, 407.  
*japonicum*, Thunb., 26, 407.  
*moniliforme*, Blume, 26, 407.  
*orientale*, Willd., 26, 408.  
*ovalifolium*, Wall., 26, 408.
- Visionia paniculata*, DC., 26, 92.  
*robusta*, DC., 26, 92.
- Vitex Agnus-castus*, Linn., mentioned, 26, 265.  
*cannabifolia*, Sieb. & Zucc., 26, 258.  
*heterophylla*, Roxb., 26, 257.  
*incisa*, Lam., 26, 257.  
— var. *heterophylla*, Franch., 26, 257.  
*Loureiri*, Hook. & Arn., 26, 257.  
*Negundo*, Linn., 26, 258.  
— var. *incisa*, C. B. Clarke, 26, 257, 258.  
*ovata*, Thunb., 26, 258.  
*Sampsoni*, Hance, 26, 258.  
*spicata*, Lour., 26, 258.  
*trifolia*, Linn., 26, 258.  
— var. *trifoliata*, Schau., 26, 258.  
— var. *unifoliata*, Schau., 26, 258.  
*urceolata*, C. B. Clarke, 26, 259.
- Vitis adstricta*, Hance, 23, 136.  
*amurensis*, Rupr., 23, 136.  
*angustifolia*, Benth., 23, 132.  
*armata*, Diels & Gilg, 36, 529.  
*betulifolia*, Diels & Gilg, 36, 529.  
*bryoniaefolia*, Bunge, 23, 136.  
*cantoniensis*, Hook. & Arn., 23, 132.  
*cantoniensis*, Seem., 23, 131.  
*Chiasii*, Carr., mentioned, 23, 137.  
*Coignetia*, Pulliat, 36, 529.  
*cordata*, Wall., 23, 135.  
*corniculata*, Benth., 23, 132; mentioned, 138.  
*ficifolia*, Bunge, 23, 134.  
*flexuosa*, Thunb., 23, 132.  
*formosana*, Hemsl., 36, 529.  
*Hancockii*, Hance, 23, 132.  
*Henryana*, Hemsl. \*, 23, 132.  
*heterophylla*, Thunb., 23, 133.  
— var., Regel, 23, 133.  
*himalayana*, Brandis, 23, 133; mentioned, 138.  
*inconstans*, Miq., 23, 133.



- Vitis indica*, Hook. & Arn., **23**, 134.  
*japonica*, Thunb., **23**, 134  
*Labrusca*, Linn., **23**, 134; mentioned, 135.  
*lanata*, Roxb., **23**, 134.  
*lecooides*, Maxim., **23**, 131, 132.  
*neurosa*, Kurz, **23**, 135.  
*orientalis*, Boiss., mentioned, **23**, 132.  
*pachyphylla*, Hemsl. \*, **23**, 135.  
*Pagnuccii*, Rom. du Caill., **36**, 529.  
*papillata*, Hance, **23**, 135.  
*parvifolia*, Roxb., **23**, 132.  
*pentagona*, Diels & Gilg, **36**, 229.  
*pentaphylla*, Miq., **23**, 136.  
*Piasezkii*, Maxim., **23**, 135.  
*planicaulis*, Hook. f., **23**, 135.  
*Potanini*, Maxim., **36**, 530.  
*repens*, Wight & Arn., **23**, 135.  
*Retordi*, Carr., mentioned, **23**, 137.  
*Romaneti*, Carr., **23**, 137; **36**, 530.  
*serianæfolia*, Maxim., **23**, 136; mentioned, 137.  
*sinica*, Miq., **23**, 133.  
*spp.*, **23**, 137, 138.  
*succisa*, Hance, **23**, 132.  
*tenuifolia*, Wight & Arn., **23**, 134.  
*Thunbergii*, Sieb. & Zucc., **23**, 134.  
*trichoclada*, Diels & Gilg, **36**, 530.  
*umbellata*, Hemsl. \*, **23**, 137.  
*vinifera*, Linn., **23**, 136.  
*Voinieriana*, Hort., **36**, 530.  
*Volkameria inermis*, Lour., **26**, 261.  
*japonica*, Thunb., **26**, 262.  
*Kämpferi*, Jacq., **26**, 262.  
*pumila*, Lour., **26**, 262.  
*Vrydagzynea nuda*, Blume, **36**, 42.
- Wahlenbergia agrestis*, A. DC., **26**, 4.  
*brevipes*, Hemsl., **36**, 530.  
*dehiscens*, A. DC., **26**, 4.  
*gracilis*, A. DC., **26**, 4.  
 — var. *miseræ*, Hemsl. \*, **26**, 4.  
*grandiflora*, Schrad., **26**, 5.  
*homallanthina*, A. DC., **26**, 5.  
*lavandulæfolia*, A. DC., **26**, 4.  
*marginata*, A. DC., **26**, 4.  
*quadrifolia*, A. DC., **26**, 4.  
*Sieberi*, A. DC., **26**, 4.  
*Waldsteinia sibirica*, Tratt., **23**, 239.  
*trifolia*, Rochel, **23**, 239.  
*Waltheria americana*, Linn., **23**, 91.  
*indica*, Linn., **23**, 91.  
*Wattakaka viridiflora*, Hassk., **26**, 115.
- Webera attenuata*, Hook. f., **23**, 380.  
*mollissima*, Benth., **23**, 381.  
*Wedelia biflora*, DC., **23**, 433.  
*calendulacea*, Less., **23**, 434.  
*prostrata*, Hemsl., **23**, 434.  
*urticæfolia*, DC., **23**, 434.  
*Weigela alba*, Carr., **23**, 369.  
*amabilis*, Hort., **23**, 369.  
 — *striata*, [Van Houtte] Flore des Serres, **23**, 369.  
*florida*, DC., **23**, 368.  
*Isolinæ*, [Van Houtte] Flore des Serres, **23**, 369.  
*pauciflora*, DC., **23**, 368.  
*rosea*, Lindl., **23**, 368.  
*Wendlandia glabrata*, DC., **23**, 371.  
*Henryi*, Oliver, **23**, 372; **36**, 530.  
*paniculata*, DC., **23**, 372.  
*uvariifolia*, Hance, **23**, 372.  
*Wickstrœmia* = seq.  
*Wikstrœma alpina*, var., Benth., **26**, 328.  
*alternifolia*, Batalin, **26**, 397.  
*angustifolia*, Hemsl. \*, **26**, 396.  
*canescens*, Meisn., **26**, 397.  
*Chamædaphne*, Meisn., **26**, 397; mentioned, 399.  
*chinensis*, Meisn., **26**, 397.  
*Forsteri*, Decne., **26**, 398.  
*gracilis*, Hemsl. \*, **26**, 397.  
*indica*, C. A. Mey., **26**, 398; mentioned, 400.  
*japonica*, Meisn., mentioned, **26**, 398.  
*linoides*, Hemsl. \*, **26**, 398.  
*micrantha*, Hemsl. \*, **26**, 399.  
*Monnula*, Hance, **26**, 399.  
*nutans*, Champ., **26**, 400.  
*obovata*, Hemsl. \*, **26**, 400.  
*pauciflora*, Franch. & Sav., **26**, 397.  
*retusa*, A. Gray, **36**, 530.  
*Spanoghii*, Decne., mentioned, **26**, 400.  
*stenantha*, Hemsl. \*, **26**, 400.  
*stenophylla*, E. Pritz., **36**, 530.  
*virgata*, Meisn., **26**, 397.  
 — var., Hemsl., **26**, 397.  
*viridiflora*, Meisn., **26**, 398.  
*Wistaria brachybotrys*, Sieb. & Zucc., **23**, 161.  
*chinensis*, DC., **23**, 161.  
*Consequana*, Loud., **23**, 161.  
*dubia*, Walp., **23**, 162.  
*floribunda*, DC., **23**, 161.  
*multijuga*, Van Houtte, **23**, 161.  
*sinensis*, Sieb. & Zucc., **23**, 161.  
*Wisteria*, Nutt., **23**, 160 = præc.  
*Wollastonia biflora*, DC., **23**, 434.

- Wollastonia prostrata*, Hook. & Arn., 23, 434.
- Woodfordia floribunda*, Salisb., 23, 304.  
*fruticosa*, Kurz, 23, 304.  
*uniflora*, Rich., 23, 304.
- Wrightia pubescens*, R. Br., 26, 97.
- Xanthium chinense*, Mill., 23, 433.  
*indicum*, Koen., 23, 433.  
*Strumarium*, Linn., 23, 433.
- Xanthoceras sorbifolia*, Bunge, 23, 140.
- Xanthopappus*, Winkl., 36, 530.  
*subacaulis*, Winkl., 36, 530.
- Xanthoxylum* = *Zanthoxylum*.
- Xeranthemum chinense*, Lour., 23, 459.
- Xylanche himalaica*, G. Beck, 36, 530.
- Xylophylla ramiflora*, Ait., 26, 426.
- Xylosma japonicum*, A. Gray, 23, 57.  
*racemosum*, Miq., 23, 57.  
*senticosum*, Hance, 23, 57.
- Xylosteum Maackii*, Rupr., 23, 364.
- XYRIDACEÆ**, 36, 150.
- Xyris bancana*, Miq., 36, 150; mentioned, 151.  
*pauciflora*, Willd., 36, 151.  
*schænoides*, Benth., 36, 150.  
*schænoides*, Mart., mentioned, 36, 151.
- Youngia ambigua*, DC., 23, 476.  
*chinensis*, DC., 23, 485.  
*chrysantha*, Maxim., 23, 481.  
*debilis*, DC., 23, 480.  
*dentata*, DC., 23, 480.  
*fastigiata*, DC., 23, 476.  
*gracilis*, Miq., 23, 476.  
*hastata*, Thunb., 23, 481.  
*humilis*, DC., 23, 476.  
*integra*, A. Gray, 23, 475.  
*japonica*, DC., 23, 475.  
*lanceolata*, DC., 23, 475.  
*napifolia*, DC., 23, 476.  
*Poosia*, DC., 23, 476.  
*runcinata*, DC., 23, 476.  
*serotina*, Maxim., 23, 481.  
*sonchifolia*, Maxim., 23, 401.  
*stricta*, DC., 23, 476.  
*Thunbergiana*, DC., 23, 475.
- Ypsilandra thibetica*, Franch., 36, 40.
- Yulania conspicua*, Spach, 23, 23.  
*japonica*, Spach, 23, 24.
- Zannichellia palustris*, Kunth, 36, 197.  
 — subsp. *pedicellata*, Hook. f., 36, 197.
- Zannichellia pedicellata*, Buch.-Ham., 36, 197.  
*pedunculata*, Reichb., 36, 197.
- Zanonia cissoides*, Wall., 23, 321.
- Zanthoxylum*, Linn., the host of  
*Loranthus odoratus*, 26, 400.  
*acanthopodium*, DC., 36, 530.  
*ailanthoides*, Sieb. & Zucc., 23, 105.  
*alatum*, Roxb., 23, 105.  
*Arnottianum*, Maxim., 36, 530.  
*Avicennæ*, DC., 23, 105.  
*Bretschneideri*, Maxim., 23, 105; mentioned, 104.  
*Bungei*, Hance, 23, 105.  
*Bungei*, Planch., 23, 105.  
*Clava-Herculis*, Lour., 23, 105.  
*cuspidatum*, Champ., 23, 106.  
*Daniellii*, Benn., 23, 104.  
*dimorphophyllum*, Hemsl., 36, 530.  
*dissitum*, Hemsl. \*, 23, 106.  
*echinocarpum*, Hemsl., 36, 530.  
*emarginellum*, Miq., mentioned, 23, 108; 36, 530.  
*fraxinoides*, Hemsl., 36, 530.  
*Lamarckianum*, Cham. & Schlecht., 23, 104.  
*lentiscifolium*, Champ., 23, 105.  
*mantschuricum*, Benn., 23, 107.  
*micranthum*, Hemsl., 36, 530.  
*multifoliolatum*, Hemsl., 36, 530.  
*multijugum*, Franch., 36, 530.  
*nitidum*, Bunge, 23, 106.  
*nitidum*, DC., 23, 106.  
*nitidum*, Wall., 23, 108.  
*oxyphyllum*, Edgew., 36, 530.  
*Piasezkii*, Maxim., 36, 530.  
*piperitum*, Benn., 23, 105.  
*piperitum*, DC., 23, 107.  
*planispinum*, Sieb. & Zucc., 23, 105.  
*podocarpum*, Hemsl. \*, 23, 107.  
*pteleaefolium*, Champ., 23, 104.  
*schinifolium*, Sieb. & Zucc., 23, 107.  
*setosum*, Hemsl. \*, 23, 107.  
*simulans*, Hance, 23, 106.  
*sp.*, 23, 108.  
*stenophyllum*, Hemsl., 36, 530.  
*trifoliatum*, Linn., 23, 339.  
*undulatifolium*, Hemsl., 36, 530.
- Zea Mays*, Linn., 36, 346.
- Zehneria Baueriana*, C. B. Clarke, 23, 318.  
*mucronata*, Miq., 23, 318.  
*mysorensis*, Arn., 23, 318.  
*umbellata*, Thw., 23, 318.
- Zelkova acuminata*, Planch., 26, 449.  
*Davidii*, Benth. & Hook. f., 26, 449.

- Zelkova Keaki*, Maxim., **26**, 449.  
*Zenobia cerasiflora*, H. Léveillé, **36**, 530.  
*Zerumbet speciosum*, Schrad., **36**, 72.  
*Zingiber*, Lestib., **36**, 70.  
*Zeuxine emarginata*, Lindl., **36**, 42.  
*formozana*, Rolfe, **36**, 42.  
*pallida*, Lindl., **36**, 42.  
*sulcata*, Lindl., **36**, 42.  
*Tripleura*, Lindl., **36**, 42.  
*Zingiber atrorubens*, Gagnep., **36**, 530.  
*confine*, Miq., **36**, 70.  
*corallinum*, Hance, **36**, 70.  
*integrilabrum*, Hance, **36**, 70.  
*officinale*, Rosc., **36**, 70.  
*roseum*, Rosc., **36**, 530.  
*sp. indscript.?*, **36**, 71.  
*striolatum*, Diels, **36**, 530.  
*Zerumbet*, Rosc., **36**, 70.  
*Zizania aquatica*, Linn., **36**, 345.  
*Zizyphus Jujuba*, Lam., **23**, 126.  
*Zizyphus Lotus*, Lam., **23**, 126.  
*nitida*, Roxb., **23**, 126.  
*sinensis*, Lam., **23**, 126.  
*soporifera*, Roem. & Schult., **23**, 126.  
*vulgaris*, Lam., **23**, 126; mentioned, 127.  
*Zornia angustifolia*, Sm., **23**, 171.  
*diphylla*, Pers., **23**, 171.  
*elegans*, Pers., **23**, 171.  
*lutescens*, Steud., **23**, 174.  
*Zostera marina*, Linn., **36**, 197.  
*Zoysia macrostachya*, Franch., **36**, 344.  
*pungens*, Munro, **36**, 344.  
*pungens*, Willd., **36**, 344.  
*sinica*, Hance, **36**, 344.  
*Zygadenus japonicus*, Miq., **36**, 147.  
*sibiricus*, A. Gray, **36**, 148.  
 ZYGOPHYLLÆ, **23**, 97.  
*Zygophyllum Loezii*, Kanitz, **36**, 530.  
*mucronatum*, Maxim., **23**, 97.

END OF THE THIRTY-SIXTH VOLUME.

## INTERNATIONAL CATALOGUE OF SCIENTIFIC LITERATURE.

---

THE ROYAL SOCIETY has been engaged for some years past in arranging for the publication of an International Catalogue of Scientific Literature, beginning from the 1st January, 1901. Each science will be represented in an annual volume containing lists arranged under authors and subjects, of all books and papers published during the year; these will be contributed through official channels of information—abroad, by direct control of the respective governments—at home, by means of the various Societies which devote themselves to particular sciences; those Societies whose domains overlap having arranged for mutual cooperation.

The collection of title-slips for the United Kingdom of Great Britain and Ireland as regards

### BOTANY

has been undertaken by the Council of the Linnean Society, and they appeal to all botanic workers for support in their endeavour to compile a complete record, by sending notices promptly of all botanic issues to the undersigned.

The fourth volume, for 1904, is in preparation.

B. DAYDON JACKSON,

*General Secretary, Linn. Soc.*

## NOTICE.

---

The enumeration of the Chinese Flora is completed by the issue of this double part, closing Volume XXXVI.

Attention to this announcement is specially requested, to prevent unnecessary application to the Librarian.

---

The CATALOGUE of the LIBRARY may be had on application. Price to Fellows, 5s. ; to the Public, 10s.

---

All communications relating to the general business of the Society should, as heretofore, be addressed to the SECRETARIES, but letters on library business only may be addressed to the "LIBRARIAN."